



Long-Term Water Quality Responses to Conservation Practices in Nested Coastal Plain Watersheds

George Vellidis

Partners

- **University of Georgia**
 - Susan Crow, Gary Hawkins
- **USDA-ARS Southeast Watershed Research Lab.**
 - David Bosch, Richard Lowrance, Dana Sullivan, Gary Feyereisen
- **USDA-NRCS**
 - Jimmy Bramblett
- **South Georgia Regional Development Center**
 - Emily Perry Davenport

Funding

- **USDA-CSREES Integrated Research, Education, and Extension CEAP Competitive Grants Program**
- **State and federal funds allocated to the University of Georgia Agricultural Experiment Stations**
- **CRIS project funds allocated to the USDA-ARS Southeast Watershed Research Laboratory**

Objectives



- To evaluate the effects of past and potential conservation practices on water quality in a coastal plain watershed;
- To evaluate social and economic factors influencing implementation and maintenance of these conservation practices; and
- Train and educate stakeholders about these issues and the effects that their actions have on watershed-scale water quality.

Objectives


- To evaluate the effects of past and potential conservation practices on water quality in a coastal plain watershed;
- To evaluate social and economic factors influencing implementation and maintenance of these conservation practices; and
- Train and educate stakeholders about these issues and the effects that their actions have on watershed-scale water quality.

Where?

- **Southeastern Coastal Plain**
 - **Suwannee River Basin**




Suwannee River Basin



- Representative of Coastal Plain ecoregions
- 60% in Georgia, 40% in Florida
- Priority watershed
- LREW

Little River Experimental Watershed



- 334 km² (82,500 ac)
- USDA-ARS regional experimental watershed
- Established in late 1960s
- 5th order stream

LREW Impairments

- **Main stem**
 - low DO
- **Tributaries**
 - low DO, fecal coliform, and sediment
- **Typical of impairments in Coastal Plain**
- **No point sources**

LREW Landcover


During late 1970s (windshield surveys)

- Agricultural land – 36% to 54%
- Row crops – 31% to 41%
- Most pasture land used for cow-calf operations
- Remainder of the watershed in upland pine forest
 - both natural regeneration and pine plantation



LREW Landcover

- **Forested riparian areas**
 - natural regeneration
- **Some landcover changes from 1975 to 2005**
- **Minor increases in suburban development**

Landsat Landcover in 1998



- Urban
- Pecan
- Forested
- Water/Wetland
- Crop
- Pasture



Landsat Landcover in 1998

Landcover	Percentage of Area
Urban	2.7
Pecan	0.5
Mature Deciduous	35.5
Mature Planted Pine	2.8
Mixed Deciduous and Pine	5.2
Young Planted Pine	0.3
Water	1.4
Wetland	9.1
Forested Wetland	2.8
Crop	24.7
Pasture	15.0

LREW Cropping History

- **Important changes – decrease in corn acreage and an increase in cotton**
- **Middle 1970's – major crops, roughly in order of acreage**
 - **corn, soybeans, peanuts, sorghum, tobacco, and vegetables**
 - **no cotton**

LREW Cropping History

Percent of Total Crops

	1995	1996	1997	1998	1999	2000
Cotton	54.1	55.8	59.7	57.1	61.5	64.5
Peanuts	43.6	39.7	36.9	34.8	38.5	34.7
Corn	2.3	4.5	3.4	8.1	0.0	0.7

- While small, vegetable acreage is increasing
- No soybeans grown in the watershed over the past few years


USDA Conservation Practices

-

USDA Conservation Practices

- **Late 1990s – present**
 - nutrient management
 - manure management
 - conservation tillage (cotton)
 - cover crops
 - filter strips
 - farm ponds

Inventory of Conservation Practices



- Available – historical database within LREW for 1985 - 2005
 - entered into GIS
- Extend record to 1975

Riparian Forests

Relevant Publications


- Lowrance, R., G. Vellidis, R.D. Wauchope, P. Gay, and D.D. Bosch. 1997. Herbicide transport in a riparian forest buffer system in the coastal plain of Georgia. *Transactions of the ASAE* 40(4):1047-1057.
- Perry, C.D., G. Vellidis, R. Lowrance, and D.L. Thomas. 1999. Watershed-scale water quality impacts of riparian forest management. *J. Water Resour. Plng. and Mgmt.*, ASCE, 125(3):117-125.
- Vellidis, G., R. Lowrance, P. Gay, and R.D. Wauchope. 2002. Herbicide transport in a restored riparian forest buffer system. *Transactions of the ASAE* 45(1):89-97.
- Vellidis G., R. Lowrance, P. Gay, R.W. Hill, and R.K. Hubbard. 2004. Nutrient transport in a restored riparian wetland. *Journal of Environmental Quality* 32(2):711-726.
- Gay, P., G. Vellidis, and J.J. Delfino. 2006. The Attenuation of Atrazine and It's Major Degradation Products in a Forested Riparian Buffer: I. Ground Water and Soils. *Transactions of the ASABE* (in review).
- Gay, P., G. Vellidis, and J.J. Delfino. 2006. The Attenuation of Atrazine and It's Major Degradation Products in a Forested Riparian Buffer: II. Surface Runoff Water. *Transactions of the ASABE* (in review).

Nutrient Retention by a Forested Riparian Buffer

Position	Water (m ³ /year)	Nutrients (kg/year)						
		NO ₃ -N	NH ₄ -N	TKN	TN	Cl	PO ₄ -P	TP
Original Total Entering	28,875	30.71	16.40	108.42	139.13	315.1	12.01	20.97
Original Total Leaving	22,125	5.15	5.99	38.73	43.88	240.2	3.23	5.41
Difference (In – Out)	4,430	25.55	10.41	69.69	95.25	75.16	8.77	15.56
Percent Retention	15.34	83.22	63.47	64.28	68.46	23.8	73.06	74.21
Total Leaving	29,047	6.77	7.86	50.84	57.61	315.4	4.25	7.10
Difference	-172	23.94	8.53	57.58	81.52	0.0	7.76	13.87
Percent Retention	-0.60	77.97	52.04	53.11	58.59	0.0	64.64	66.14


Vellidis G., R. Lowrance, P. Gay, R.W. Hill, and R.K. Hubbard. 2004. Nutrient transport in a restored riparian wetland. *Journal of Environmental Quality* 32(2):711-726.

Hydrology & Water Quality




- 8 nested watersheds with v-notch weirs
- 334 km² (82,500 ac) at Station B
- continuous hydrologic record since 1974
- mostly continuous water quality record since 1974





30-Year NO₃-N Concentrations at Station B



Hypothesis – Nitrogen

- Establishment of trees (under CRP) on highly erodible crop land led to a change in N transport from LREW with larger decreases for watersheds with higher levels of tree establishment.

Hypotheses – Nitrogen

- Establishment of trees (under CRP) on highly erodible crop land led to a change in N transport from LREW with larger decreases for watersheds with higher levels of tree establishment.
- Decrease in corn and increase in cotton production led to decreases in N transport from LREW.

Hypothesis – Buffers

- Riparian forest buffers left voluntarily by farmers have a larger positive impact on water quality than some practices installed under USDA cost-share programs.

Hypotheses – Buffers

- Riparian forest buffers left voluntarily by farmers have a larger positive impact on water quality than some practices installed under USDA cost-share programs.
- Farmers have left RFBs for a wide variety of reasons.

Implementation

- **Assess water quality effects of conservation practices at multiple temporal and spatial scales in LREW**
- **Provide a geographic prioritization for future implementation in the watershed and similar watersheds**
- **Cumulative Effects Analysis**

Cumulative Effects Analysis (CEA)

- Recommended by the President's Council on Environmental Quality (CEQ)


“CEA allows analysts to determine the cumulative environmental consequences of actions by delineating the cause-and-effect relationships between multiple actions and the resources, ecosystems, and human communities of concern.”

(CEQ, 1997)

CEA → Conceptual Model

- **Build a Conceptual Model to:**
 - determine the cumulative water quality consequences of conservation practices

Wetland Restoration for Sediment Yield Reduction: A Conceptual Model



Develop Conceptual Model

- **Accumulate relevant information**
- **Panel of Experts**
 - **biophysical interactions in coastal plain watersheds**
 - **socioeconomic factors influencing the adoption, implementation, and maintenance of conservation practices**

Implement Conceptual Model

- **STELLA – process model**
 - www.iseesystems.com
- **GIS to aggregate and visualize**
- **SWAT and REMM to test findings**
- **Recruiting Post-Doc**

Thank you for your attention !!



For more information:

**Dr. George Vellidis
Biological & Agricultural Engineering Department
University of Georgia
Tifton GA 31793-0748**

voice: 229.386.3912 fax: 229.386.3958

e-mail: yiorgos@uga.edu