


Introduction

The Des Lacs 8-Digit Hydrologic Unit Code (HUC) (09010002) sub-basin includes land in Canada and the United States. There are approximately 614,600 acres in the United States (North Dakota) portion of the sub-basin. This sub-basin is located in Souris-Red-Rainy Region, Souris Sub-Region.

This report addresses only the portion located within North Dakota. The Des Lacs is approximately 614,600 acres covering parts of four counties (Burke, Mountrail, Renville, and Ward) in North Dakota. Of the 614,600 acres, Ward contains 48%, Burke 44%, Renville 4%, and Mountrail 4%. There are approximately 450 farms in the sub-basin. The following two maps show the entire sub-basin and also the portion of the sub-basin located within North Dakota.

This sub-basin encompasses commodities ranging from canola, wheat, barley, sunflowers, flax, alfalfa, and pulse crops (peas & lentils) to beef cattle, swine, and bees.

Conservation assistance is provided by four Natural Resources Conservation Service (NRCS) Service Centers and two Resource Conservation & Development (RC&D) Offices.



*Produced by
the Natural
Resources
Planning Staff
Bismarck, ND*

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.


Physical Description

The following table and map show land use / land cover within the sub-basin.

Land Use/ Land Cover (<i>National Resources Inventory [NRI]</i>) ¹	Acres	Percent of HUC
Forestland	2,000	*
Cropland	443,600	72 %
Conservation Reserve Program (CRP) Land ² ^a	2,900	1 %
Tame Grass/Hayland	11,900	2 %
Pastureland	7,700	1 %
Rangeland	80,100	13 %
Urban/Farmstead/ Transportation Land	30,800	5 %
Water/Wetlands	9,600	2 %
Federal Lands	26,000	4 %
North Dakota HUC Totals ^b	614,600	100% *
<p><i>* Less than one percent of total acres. See below for special considerations. a: Estimate from Farm Service Agency records and include CRP/CREP. b: Totals may not add due to rounding and small unknown acreages.</i></p>		
Irrigated Land (<i>Farm Services Agency</i>) ³	0	<1%

Physical Description – Continued


Land Use/Land Cover Map



The above map was developed from U.S. Geologic Survey's (USGS) ND Gap Analysis Program data.⁴


Physical Description – Continued

The sub-basin is part of the Souris-Red-Rainy River Region - Souris Sub-Region. The drainage pattern flows to the southeast ending at the Souris River, which then flows north into Canada. The following map shows the relief for the sub-basin.⁵



Physical Description – Continued

The following map is a plot of 1961-1990 annual average precipitation contours from National Oceanic and Atmospheric Administration (NOAA) Cooperative Stations and (where appropriate) USDA-NRCS Snowpack Telemetry (SNOTEL) Stations. Christopher Daly used the PRISM (Parameter-elevation Regressions on Independent Slopes Model) model to generate the gridded estimates from which this map was derived; the modeled grid was approximately 4x4 km latitude/longitude, and was resampled to 2x2 km using a Gaussian filter. Mapping was performed by Jenny Weisberg and Nathaniel DeYoung. Funding was provided by USDA-NRCS National Water and Climate Center. (4/20/98)



Physical Description – Continued

The North Dakota Department of Health collects water quality data on major water bodies. The following table shows the total miles of streams and acres of lakes/reservoirs within the sub-basin and also the miles and acres with a water quality limitation. A map showing the Total Maximum Daily Load (TMDL) waters within the watershed follows the table. TMDL is the amount of a particular pollutant that a particular stream, lake, estuary, or other waterbody can "handle" without violating State water quality standards.

		Units	Des Lacs Sub-basin ⁶	Des Lacs Impaired Water Quality (303d) ⁷	Percent Impaired* Des Lacs
Water Quality Data <i>*Percent of Total Miles and acres in HUC</i>	Total – Major Water bodies				
	Rivers/Streams	Miles	715.9	71.1	9.9
	Lakes/Reservoirs	Acres	155.0	150.8	97.3



Physical Description – Continued

The following two tables show feeding operations, permitted operations, and livestock numbers. The first table lists the number of animal feeding operations and animals as tracked by the North Dakota Department of Health. The second table shows livestock numbers for all cattle, beef cows, dairy cows, hogs and pigs, and sheep and lambs. These livestock numbers were extrapolated from 2002 Agricultural Census county data to 8-digit HUC's.

Animal Feeding Facilities – North Dakota Department of Health Permit⁸					
Animal Type	Dairy	Beef	Swine	Other	Total
Number of Animal Feeding Operations	3	0	4	0	7
Number of Animals	270	0	2,680	0	2,950
Number of State Permitted Operations					4

Livestock Numbers (rounded to nearest 100)⁹					
	Cattle and Calves	Beef Cows	Dairy Cows	Hogs and Pigs	Sheep and Lambs
North Dakota	1,873,200	982,300	34,500	138,800	114,000
Des Lacs	16,900	10,200	200	400	500
Des Lacs as a percent of North Dakota	0.9%	1.0%	0.6%	0.3%	0.4%

Physical Description – Continued

Common Resource Areas (CRAs) are geographical areas where resource concerns, problems, or treatments are similar. Landscape conditions, soil, climate, human considerations, and other natural resource information were used to determine the geographic boundaries. CRAs are subsets of Major Land Resource Areas. The following map¹⁰ shows the CRAs for Des Lacs sub-basin with the descriptions below.


53A.1 – Northern Dark Brown

Glaciated Plains: The Northern Dark Brown Glaciated Plains are gently undulating low relief till plains with many knobs and temporary and seasonal wetlands. The soils are mainly medium textured, well drained, and have a frigid temperature regime. Most of this area is in farms and ranches. Average annual precipitation is 300 to 350 mm. Average annual temperature is 3 to 5 degrees C. Average freeze free period is 90 to 120 days.

53B.1 Central Dark Brown

Glaciated Plains: The Central Dark Brown Glaciated Plains are nearly level to rolling with steeper areas along rivers. Land use is a mosaic of cropland and rangeland. Soil textures are dominantly loamy in glacial till, sandy in outwash areas, and clayey in lacustrine areas. Most soils are moderately deep or deep, well drained or moderately well drained, and have a frigid temperature regime


55A.1 – Northern Black Glaciated Drift Plains: The Northern Black Glaciated Drift Plain is a nearly level to undulating landscape composed of glacial till and lacustrine sediments. Temporary and seasonal wetlands are numerous throughout the area. Agriculture is limited by a very short growing season and the coldest January temperatures in the Northern Plains.



Soil Productivity¹¹

The northern and southwestern areas of the Des Lacs sub-basin are dominated by soils with moderate to poor productivity indexes (PI). Many of these soils are high in sodium which is evident by the wide expanse of clay pan soils. The south central areas above the Souris River breaks have soils with a high PI.

The term "Productivity Index" used in this document reflects soil properties and the inherent production capacity of the soil to produce spring wheat



Common Land Unit

The entire sub-basin has the common land unit digitized by Farm Services Agency (FSA).

Resource Concerns


One of the goals of NRCS is to help quantify the types and amounts of resources that may be of concern in an area. This helps identify priority areas for the types and amounts of assistance to be given to a particular watershed.

The following table shows the different projects, plans, studies, and assessments conducted within the sub-basin.

Watershed Projects, Plans, Studies and Assessments			
NRCS Watershed Projects		NRCS Watershed Plans, Studies & Assessments	
Name	Status	Name	Status
None	NA	Des Lacs- Souris River Basin Study	Completed 1992
		Northgate Dam Watershed – Stony Run Riparian Wetland Assessment	Completed 2003
NDDH TMDLs		Soil Conservation District Assessments and Studies	
Number Listed		Name	Status
Lakes/Reservoirs - 1	Streams – 1	None	NA
EPA 319 Watershed Projects			
Name		Status	
Northgate Dam Watershed		Ongoing	

Soil

- NRI estimates indicate cultivated cropland acreage experiencing erosion rates above sustainable levels decreased to 77,200 acres in 1997, as compared to 217,100 acres in 1987.
- Through NRCS programs many farmers and ranchers have applied conservation practices to reduce the effects of both wind and water erosion. From 1982 to 1997, the average wind erosion rate reduced from 5.4 tons/acre/year to 2.4 tons/acre/year on all cultivated cropland. The average water erosion rate reduced from 1.4 tons/acre/year to 0.9 tons/acre/year on cultivated cropland.
- Fine textured clayey and sandy soils still require conservation practices to control excessive wind erosion.
- Soil erosion and low organic matter remain resource concerns.




Resource Concerns - Continued

Soil - Continued

- Soil health on rangeland sites as it relates to the stability of redistributing and losing nutrients and organic matter.
- More utilization of soil fertility testing is needed to reduce nutrient loading in the soil profile.
- Salt build-up in soil profile remains a production issue.

Water

- **Aquifers**¹² - There are four glacial drift aquifers (Burlington, Columbus, Kenmare, and Lignite) underlying the Des Lacs sub-basin.



- **Wellhead Protection Areas**¹³ – There are six protection areas located in the sub-basin. They are designated to protect the municipal water supply for the cities of Flaxton, Kenmare, Lignite, Portal, and the service area of the Upper Souris WUA – System I.

Resource Concerns - Continued

Water - Continued

- A stream section of the Des Lacs River, from Lower Des Lacs Lake downstream to the confluence of the Souris River, was placed on the 303(d) list for total fecal coliform.
- Northgate Dam reservoir is on the 303(d) list for loss of the designated uses of "Fish and Other Aquatic Biota" and "Recreation". These impairments are due to excessive nutrients/eutrophication, low dissolved oxygen, and excessive sediment/siltation.
- Conservation practices used to address these water quality issues include grazing management, erosion control, nutrient and ag waste management, and riparian buffers.
- Lack of adequate riparian buffer width and health are impacting water quality and stream health.
- Conventional tillage systems continue to hinder efficient water use on dry land cropland.
- Salinity in surface water is a concern in the upper end of the sub-basin.
- Season long grazing on or near water courses is a water quality concern.

Air

- Objectionable odors are minimal with some feedlot or wintering areas having noticeable odor during certain times of the year.
- Visibility is reduced during winter months from blowing snow.
- A better awareness of green house gases and their impact is needed; especially as it relates to the Canadian coal fired power plant located near Estevan, Saskatchewan.

Plants

- Maintaining rangeland productivity, health, and vigor are of major concern.
- Past utilization of non adapted or suited species has resulted in invasive plant species among native plant communities and declining forage yields.
- There are concerns about controlling invasive weeds and maintaining good pasture condition.
- Noxious weeds and poor range condition reduce productivity for livestock and wildlife.
- Maintaining forage quality and palatability are needed to sustain the livestock base and provide plant community sustainability.


Resource Concerns - Continued

Animals

- Inadequate food, cover/shelter and water is a concern for maintaining wildlife populations.
- Habitat fragmentation, including loss of old field and farmstead windbreaks, has resulted in lost cover and shelter for certain wildlife species as well as domestic livestock.
- Animals that are threatened and endangered can be seen in the following table of threatened and endangered species.

Federally Listed Threatened And Endangered Species			
Species Category	Threatened	Endangered	Candidate
Mammals	Piping Plover	Gray Wolf	None
Birds	None	Whooping Crane	None
Fish	None	None	None
Invertebrates	None	None	Dakota Skipper
Plants	None	None	None
Critical Habitat – Piping Plover			

- The adjacent map shows land owned by the US Fish and Wildlife Service (FWS). The FWS land is separated into the following designations: National Wildlife Refuge and Waterfowl Production Areas. The FWS controls over 30,300 acres within the watershed.





Census and Social Data¹⁴

Number of Farms: 450

Number of Operators:

- Average Age: 54
- Full-Time Operators: 73%
- Part-Time Operators: 27%



Limited Resource and Beginning Farmer

Approximately 5.5 percent of the operators are minority producers. Limited Resource Farmers are estimated at 12 percent. Although rather low percentages, these facts point to the potential need for special technical assistance targeted to reach people who (1) may lack experience with government farm programs, (2) have good stewardship intentions but lack management skills, and (3) lack the time to visit an NRCS field office and seek assistance.

All data is provided "as is." There are no warranties, express or implied, including warranty of fitness for a particular purpose, accompanying this document. Use for general planning purposes only.



Des Lacs 09010002

8-Digit Hydrologic Unit Profile

September 2007

References

- ¹ USDA-NRCS, NRI data.
- ² USDA-Farm Services Agency, Common Land Unit GIS data layer, 2005.
- ³ USDA-Farm Services Agency, Common Land Unit GIS data layer, 2005.
- ⁴ USDI-US Geologic Services, ND GAP analysis data, 2005.
- ⁵ USDA-NRCS, Natural Resources Planning Staff, 30 meter Relief Data GIS data layer, 2002.
- ⁶ ND Department of Health, Environmental Health Section, Water Quality Division, National Hydrography GIS layers, June 2006.
- ⁷ ND Department of Health, Environmental Health Section, Water Quality Division, List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota, 2006.
- ⁸ ND Department of Health, Environmental Health Section, Water Quality Division, Animal Feeding Operations Program data, 2006.
- ⁹ 2002 Census of Agriculture, North Dakota, State and County Data Volume 1, Geographic Area Series Part 34, U.S. Department of Agriculture, National Agricultural Statistics Service, June 2004. (County data was prorated to HUC by the percent of a HUC in a county.)
- ¹⁰ USDA-NRCS, Natural Resources Planning Staff, Common Resource Area GIS data layer, 2004.
- ¹¹ USDA-NRCS, Natural Resources Planning Staff, Soils Productivity GIS data layer, 2006.
- ¹² ND Department of Health, Environmental Health Section, Water Quality Division, Ambient Ground Water Monitoring Program data, 1997.
- ¹³ ND Department of Health, Environmental Health Section, Water Quality Division, Source Water Protection Program data, 2003.
- ¹⁴ 2002 Census of Agriculture, North Dakota, State and County Data Volume 1, Geographic Area Series Part 34, U.S. Department of Agriculture, National Agricultural Statistics Service, June 2004. (County data was prorated to HUC by the percent of a HUC in a county.)