

UNITED STATES DEPARTMENT OF AGRICULTURE
NATURAL RESOURCES CONSERVATION SERVICE

STATE: Utah

SITE TYPE: Forestland

ECOLOGICAL SITE NAME: Mountain Loam (Ponderosa pine)

SITE NUMBER: 048AY417UT

MLRA: 048A

Original Site Description: Author: GSC, DJS

Date: 01/19/1994

Revised Site Description: Author:

Date:

Approved by: Title:

Signed:

Date:

Ecological Site Definition - A distinctive kind of land, with specific physical characteristics, which differs from other kinds of land in its ability to produce a distinctive kind and amount of vegetation, and in its response to management.

A. PHYSICAL CHARACTERISTICS

(description narrative of this particular site)

1. SOILS

Depth: 20-60 inches

Surface Textures: Sandy Loam to Loam

Surface Fragments(<=3" % cover, >3" % cover):

Subsurface Textures: Loam to Clay

Subsurface Fragments(<=3" % vol, >3" % vol): <25%

Geologic Parent Materials: Mixed Alluvium and Colluvium from Sedimentary Rock

Moisture Regime:

Temperature Regime:

Runoff: Slight

Permeability(min-max):

Drainage Class(min-max):

Water Erosion Hazard:

Wind Erosion Hazard:

Electrical Conductivity (EC in mmhos/cm):

Sodium Adsorption Ration (SAR):

Soil Reaction (1:1 water):

Soil Reaction (0.1 M CaCl₂):

pH Range: Slightly Acid-Moderately Alkaline

Available Water Capacity (inches):

Major Soils Associated With This Site:

Soil Survey Area:

Brycan VFSL – fine-loamy, mixed Cumulic Haploborolls

Tukuhnik L – fine, montmorillonitic Typic Argiborolls

Hogg FSL – fine, mixed Mollic Eutroboralfs

Tomaski L – fine, montmorillonitic Typic Argiborolls

Additional information may be found in Section II of the Field Office Technical Guide.

Site Type: Forestland
 Ecological Site Name: Mountain Loam (Ponderosa pine)
 Site Number: 048AY417UT

2. PHYSIOGRAPHIC FEATURES

Landform and Position: Gently to Moderately Steep Mountain Slopes and Mesa
 Tops
 Aspect:

	<u>Minimum</u>	<u>Maximum</u>
Slope:	2	50
Elevation:	4000	8000
Flooding:		
Frequency:		
Duration:		
Ponding:		
Depth (inches):		
Frequency:		
Duration:		
Water Table Depth:		

3. FOREST COMMUNITY TYPE

Overstory: ponderosa pine (*Pinus ponderosa*)
 Understory: Gambel oak, mountain big sagebrush
 Site Index: 37 to 40

B. CLIMATIC FEATURES

Mean Annual Precipitation (inches): 25-35
 Mean Annual Air Temperature: 34-38
 Mean Annual Soil Temperature: 42-47
 Frost Free Period (days): 50-75
 Freeze Free Period (days): 0-0

Temperature and Moisture Distribution:

Temp	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
High												
Mean												
Low												

PPT	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
High												
Mean												
Low												

Climate Stations: St. ID.:

Location:

Period:

Site Type: Forestland

Ecological Site Name: Mountain Loam (Ponderosa pine)

Site Number: 048AY417UT

From: To:

(Includes factors such as storm intensity, precipitation dependability, origin and pattern of storms, driest and wettest months, orographic effects, etc.)

Influencing Water Features (if any):

Wetland Description(Cowardin System) System Subsystem Class

Stream Types(Rosgen System) System

C. PLANT COMMUNITY CHARACTERISTICS

1. Potential Plant Community Description and Ecological Factors

(Includes dominant vegetative aspect, cool-season and warm-season components, typical plant spacing, etc.)

a. Nature of Forest Community

The overstory tree canopy cover is about 35-40 percent. Common understory plants are Gambel oak, muttongrass, Geyer sedge, and bottlebrush squirreltail. Understory composition by air-dry weight is approximately 45 percent perennial grasses and grasslike plants, 20 percent forbs, and 35 percent shrubs. Understory production ranges from 900 pounds per acre in favorable years to about 300 pounds per acre in unfavorable years. Understory production includes the total annual production of all species within 4 ½ feet of the ground surface.

b. Productivity Rating of Major Understory Species:

c. **Productivity Rating Index:** This rating provides an index to the relative importance of species in the understory community as affected by overstory canopy cover.

Productivity Index

1	Always present:	More than 50% of total understory production
2	Always present:	25 to 50% of total understory production
3	Generally present:	10-24% of total understory production
4	Frequently present:	5-9% of total understory production
5	Occasionally present:	1-5% of total understory production
6	Rarely present:	Less than 1% of total understory production

2. Plant Community Composition by Overstory Canopy Class

Site Type: Forestland

Ecological Site Name: Mountain Loam (Ponderosa pine)

Site Number: 048AY417UT

Common Name	National Symbol	0-10%	11-20%	21-35%	36-60%
Gambel oak	QUGA	4	3	2	2
Indian ricegrass	ACHY	2	2	3	4
Muttongrass	POFE	2	3	3	3
Geyer sedge	CAGE2	3	3	3	3
Bottlebrush squirreltail	ELEL5	3	3	4	4
Needleandthread	HECO26	4	4	5	5
Prairie junegrass	KOMA	5	4	4	4
Mountain snowberry	SYOR2	4	4	3	3
Creeping Oregon grape	MARE11	4	3	3	3
Mountain big sagebrush	ARTRV	2	3	4	4

3. Plant Community Annual Production

At the highest potential similarity index, this site will produce approximately the following amount of air-dry herbage, expressed as pounds/acre:

Total Average Understory Production by Overstory Canopy Class (lbs./acre air-dry weight)

	Open 0-10%	Sparse 11-20%	Medium 21-35%	Dense 36-60%
Favorable Year	1200	1000	800	600
Average Year	900	700	500	400
Unfavorable Year	600	450	300	150

4. Ground Cover and Structure

a. Vegetative

Vegetation Type	Percent Canopy Cover	Height Range	Percent Basal Area Cover
Grasses & Grass-like (perennial)			
Forbs (perennial)			
Shrubs			
Trees			
Cryptogams			

b. Other

Litter	
Coarse Fragments	
Bare Ground	

5. Ecological Dynamics of the Site

a. Herbaceous:

Vegetation is dominated by grasses and forbs under full sunlight. This stage is experienced after a major disturbance such as crown fire or tree harvest. Skeleton forest (dead trees) remaining after fire or residual trees left following harvest have little or no effect on the composition and production of the herbaceous vegetation.

b. Shrub-Herbaceous:

Herbaceous vegetation and woody shrubs dominate the site. Various amounts of tree seedlings (less than 20 inches in height) may be present up to the point where they are obviously a major component of the vegetal structure.

c. Sapling:

In the absence of disturbance, the tree seedlings develop into saplings (20 inches to 4.5 feet in height) with a range in canopy cover of about 5 to 10 percent. Vegetation consists of grasses, forbs, and shrubs in association with tree saplings.

d. Immature Forest:

The visual aspect and vegetal structure are dominated by ponderosa pine greater than 4.5 feet in height. Seedlings and saplings are present in the understory. Understory vegetation is moderately influenced by a tree overstory canopy about 10 to 20 percent. Lodgepole pine and quaking aspen are principle seral associates.

e. Mature Forest:

The visual aspect and vegetal structure are dominated by ponderosa pine that have reached or are near maximal heights for the site. Tree diameters of 30-50 inches and heights of 90 to 130 feet are common. Tree canopy cover ranges from 20 to 40 percent. Understory vegetation is strongly influenced by tree competition, overstory shading, duff accumulation, etc. Few seedlings and/or saplings of the major overstory tree species occur in the understory.

f. Climax Forest:

In the absence of wildfire or other naturally occurring disturbances, the tree canopy on this site can become very dense. This stage is dominated by trees that have reached maximal heights for the site. Understory vegetation is sparse to absent due to tree competition, overstory shading, duff accumulation, etc. Tree canopy cover is at a maximum for the site and is commonly greater than 50 percent.

6. Productivity Capacity

Productivity Class: 1.0

CMAI:	10. to 25.	cu ft/ac/yr
	0.7 to 1.7	cu m/ha/yr

Fuelwood Production:

33 to 45 cords per acre per year for stands averaging 27 to 35 feet in height and 50 years of age. Firewood is commonly measured in cords, or a stacked unit equivalent to 128 cubic feet. Assuming an average of 90 cubic

Site Type: Forestland

Ecological Site Name: Mountain Loam (Ponderosa pine)

Site Number: 048AY417UT

feet of solid volume wood per cord, there are about 211,500 British thermal units (BTU's) per cubic foot or about 19 million BTU's of heat value in a cord of ponderosa pine.

Tree volume per acre: 3000 to 4000 cubic feet/acre/year for stands averaging 27 to 35 feet in height and 50 years of age.

Plant Communities & Transitional Pathways

(Show a steady state diagram with influences to move from one steady state to another)

7. Plant Growth Curves

	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
Percent Growth												
Name												
ID Number												
Description												

8. Aspect Differences Near MLRA Boundaries

(Give related range sites in MLRA's above and below)

9. Associated Sites Within MLRA

(Give site name and number)

10. Correlated Sites in Other States

(Give site name and number)

D. MAJOR USES OF THIS SITE

1. Forage Products

a. Livestock Grazing

This site is suited to cattle and sheep grazing during the summer and fall. Livestock will often concentrate on this site taking advantage of the shade and shelter offered by the tree overstory. Many areas are not used because of steep slopes or lack of adequate water. Attentive grazing management is required due to steep slopes and erosion hazards. Harvesting trees under a sound management program can open up the tree canopy to allow increased production of understory species desirable for grazing.

b. Initial Stocking Rates

Stocking rates vary in accordance with such factors as kind and class of grazing animal, season of use, and fluctuation in climate. Actual use records for individual sites, together with a determination of the

Site Type: Forestland

Ecological Site Name: Mountain Loam (Ponderosa pine)

Site Number: 048AY417UT

degree to which the sites have been grazed and an evaluation of trend in site condition, offer the most reliable basis for developing initial stocking rates.

Selection of initial stocking rates for given grazed units is a planning decision. This decision should be made only after careful consideration of the total resources available, evaluation of alternatives for use and treatment, and establishment of objectives by the decisionmaker.

c. Forage Value Rating (P) Preferred, (D) Desirable, (U) Undesirable

Common Name	National Symbol	Relative Forage Value for:			
		Cattle	Horses	Sheep	Deer
Gambel oak	QUGA	D	U	D	P
Indian ricegrass	ACHY	P	P	P	D
Muttongrass	POFE	P	P	P	D
Geyer sedge	CAGE2	P	P	U	D
Bottlebrush squirreltail	ELEL5	D	D	D	D
Needleandthread	HECO26	P	P	D	D
Prairie junegrass	KOMA	D	D	D	D
Mountain snowberry	SYOR2	U	U	D	D
Creeping Oregon grape	MARE11	U	U	U	D
Mountain big sagebrush	ARTRV	U	U	D	D
Ponderosa pine	PIPO	U	U	U	D

d. Guide to Forage Quality(Plant preference by season)

Species	Oct-Nov	Dec-Feb	Mar-May	Jun-Sep

VG = Very Good G = Good F = Fair P = Poor

2. Wildlife

a. Site Factors Influencing Management

b. List of Potential Species Present

Wildlife species seeking food and cover in this forest site include elk, mule deer, bear, porcupine, snowshoe hare, owl, and woodpecker.

This is a short list of the more common species found. Many other species are present as well and migratory birds are present at times.

c. Guide to Forage Preference of Managed Wildlife Species

Wildlife Species →				
Plant Species ↓	Use	Season	Use	Season

Site Type: Forestland

Ecological Site Name: Mountain Loam (Ponderosa pine)

Site Number: 048AY417UT

Use - A = preferred or desirable

B = some use, but less important

C = little use or used occasionally

Season - F = Fall (Oct-Nov)

W = Winter (Dec-Feb)

Sp. = Spring (Mar-May)

Su. = Summer (Jun-Sep)

3. Recreational Uses

4. Limitations and Considerations

- Potential for sheet and rill erosion is moderate to severe depending on slope.
- Moderate to severe equipment limitations on steeper slopes and on sites having extreme surface stoniness.
- Proper spacing is the key to a well managed multiple use and multi-product forest.

5. Essential Requirements

- Adequately protect from uncontrolled burning.
- Protect soils from accelerated erosion.
- Apply proper grazing management practices (see management guides)

6. Silvicultural Practices

- Harvest cut selectively or in small patches (size dependent upon site conditions) to enhance forage production.
 - Thinning and improvement cutting – removal of poorly formed, diseased, and low vigor trees.
 - Harvest cutting – selectively harvest surplus trees to achieve desired spacing. Save large, healthy, full-crowned trees. Do not select only “high grade” trees during harvest.
- Prescription burning program – to maintain desired canopy cover and manage site reproduction.
- Selective tree removal on suitable sites – to enhance forage production and manage site reproduction.
- Pest Control – use necessary and approved control for specific pests or diseases.
- Fire hazard – fire is usually not a problem in mature grazed stands.

8. Other uses

E. THREATENED AND ENDANGERED SPECIES

1. Plants

