


SOIL LEGEND*

- 1** LOAMY AND SANDY SOILS ON BOTTOM LANDS AND STREAM TERRACES
Alda-Boel-Barney association: Nearly level, somewhat poorly drained and poorly drained loamy soils that formed in alluvium; on bottom lands
- 2** Brocksburg-Simeon-Thurman association: Nearly level to gently sloping, well drained to excessively drained loamy and sandy soils that formed in loamy and sandy material; on stream terraces
- 3** SILTY SOILS ON BOTTOM LANDS, FOOT SLOPES, AND STREAM TERRACES
Gibbon-Muir-Zook association: Nearly level, well drained, somewhat poorly drained, and poorly drained silty soils that formed in alluvium; on bottom lands and stream terraces
- 4** Muir-Grigston-Cozad association: Nearly level and very gently sloping, well drained silty soils that formed in loess, alluvium, or colluvium; on stream terraces, foot slopes, and bottom lands
- 5** Hall-Muir-Hobbs association: Nearly level and very gently sloping, well drained silty soils that formed in alluvium and loess; on stream terraces, foot slopes, and bottom lands
- 6** Hobbs-Kezan-Muir association: Nearly level and very gently sloping, well drained and poorly drained silty soils that formed in alluvium and loess; on bottom lands, foot slopes, and stream terraces

- 7** SILTY SOILS ON UPLAND BREAKS
Uly-Coly association: Moderately steep to very steep, well drained to excessively drained silty soils that formed in loess; on upland breaks
- 8** Crofton-Monona association: Gently sloping to very steep, well drained to excessively drained silty soils that formed in loess; on upland breaks
- 9** SILTY, NEARLY LEVEL TO STEEP SOILS ON UPLANDS
Ponca-Sharpsburg association: Nearly level to steep, well drained and moderately well drained silty soils that formed in loess; on uplands
- 10** Sharpsburg association: Nearly level to strongly sloping, moderately well drained silty soils that formed in loess; on uplands
- 11** SILTY, NEARLY LEVEL TO STRONGLY SLOPING SOILS ON UPLANDS
Hastings-Butler association: Nearly level to strongly sloping, well drained and somewhat poorly drained silty soils that formed in loess; on uplands
- 12** Butler-Hastings association: Nearly level to gently sloping, somewhat poorly drained and well drained silty soils that formed in loess; on uplands
- 13** SILTY AND LOAMY SOILS ON UPLANDS
Sharpsburg-Steinauer-Pawnee association: Gently sloping to very steep, moderately well drained to excessively drained silty and loamy soils that formed in loess and glacial till; on uplands

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE
UNIVERSITY OF NEBRASKA
CONSERVATION AND SURVEY DIVISION
GENERAL SOIL MAP
BUTLER COUNTY, NEBRASKA



SET MISSING
28, 29, 30, 31, 32

SECTIONALIZED TOWNSHIP

6	5	4	3	2	1
7	8	9	10	11	12
18	17	16	15	14	13
19	20	21	22	23	24
30	29	28	27	26	25
31	32	33	34	35	36

Each area outlined on this map consists of more than one kind of soil. The map is thus meant for general planning rather than a basis for decisions on the use of specific tracts.

*Texture terms refer to the surface layer of the major soils.