

**Estimated Total Tons Lost per Year due to Wind Induced Erosion on Cultivated Cropland  
Ten Counties\* in Oregon by NRCS Basin\***

National Resources Inventory Results  
*Estimated for Non-Federal Lands Only*

NRCS Basin Name	1982		1987		1992		1997	
	Estimates	Error	Estimates	Error	Estimates	Error	Estimates	Error
<b>--- Tons/Year ---</b>								
John Day/Umatilla	5,895,500	463,100	5,636,500	482,000	4,035,200	400,400	4,060,700	438,600
Deschutes	1,027,800	62,500	762,200	57,500	794,400	77,100	861,800	69,900
Snake River	350,800	86,300	340,700	62,400	288,100	65,000	246,200	75,200
High Desert	305,900	164,900	295,200	108,500	243,600	113,300	229,900	114,200
NRCS Basins where wind erosion data not collected due to low levels of wind erosion.								
Central Coast/Upper Willamette	-	-	-	-	-	-	-	-
Lower Willamette	-	-	-	-	-	-	-	-
North Coast	-	-	-	-	-	-	-	-
Southwest Oregon	-	-	-	-	-	-	-	-
<b>Oregon Totals</b>	<b>7,580,000</b>	<b>501,000</b>	<b>7,034,600</b>	<b>500,800</b>	<b>5,361,300</b>	<b>428,100</b>	<b>5,398,600</b>	<b>464,600</b>

\* See Details Page

NOTE: Data may not add to totals because of rounding. The error referred to in the table is the standard error of the estimate. You can obtain the margin of error at the 95% confidence limit by multiplying the error by 1.96.

Standard errors greater than 50% of the estimate indicate a low level of statistical reliability.