

Introduction

The Lower Malheur 8-Digit Hydrologic Unit Code (HUC) subbasin is comprised of 605,000 acres in Malheur County. Seventy-five percent of the subbasin is rangeland, twenty percent is hayland and pastureland, three percent used for grain crops, and the remainder includes areas used for row crops, wetlands, and small farms. There are six permitted Confined Animal Feeding Operations (CAFOs) and over 35,000 permitted animals in the subbasin. Major resource concerns include concentrated flow, streambank, and irrigation-induced erosion; invasive and noxious weeds; insufficient water to meet livestock, wildlife, and irrigation needs; impaired water quality; and loss of wildlife habitat. High costs, unreliable markets, and inadequate incentives limit conservation adoption among the farmers and ranchers in the Lower Malheur subbasin.

There are 221 operations and 362 farmers and ranchers in the subbasin. Most operators are well educated, aware of local resource concerns, and good stewards of the natural resources. Unfortunately, the perceived expense and risk of implementing conservation limit its adoption. There is a need for additional risk-reducing incentives and greater community support for conservation to increase the diffusion of conservation in the Lower Malheur subbasin.



The Ontario NRCS Service Center, Malheur County Soil and Water Conservation District, and Malheur Watershed Council provide much of the conservation assistance in the subbasin.

Profile Contents

[Introduction](#)
[Physical Description](#)
[Land Use Map & Precipitation Map](#)
[Common Resource Area](#)

[Resource Concerns](#)
[Census and Social Data](#)
[Progress/Status](#)
[Footnotes/Bibliography](#)

Relief Map



Physical Description

[Back to Contents](#)

ALL NUMBERS IN THIS PROFILE ARE FOR OREGON ONLY

Land Cover/Land Use (NLCD ^{1/2})	Ownership - (2003 Draft BLM Surface Map Set ¹)							
	Public		Private		Tribal		Totals	%
	Acres	%	Acres	%	Acres	%		
Forest	*	---	*	---	0	0%	*	---
Grain Crops	*	---	15,700	3%	0	0%	16,000	3%
Conservation Reserve Program Land ^a	0	0%	0	0%	0	0%	0	0%
Grass/Pasture/Hay	71,900	12%	50,800	8%	0	0%	122,700	20%
Orchards/Vineyards	0	0%	0	0%	0	0%	0	0%
Row Crops	*	---	*	---	0	0%	*	---
Shrub/Rangelands	384,200	64%	69,500	11%	0	0%	453,700	75%
Water/Wetlands/Developed/Barren	*	---	*	---	0	0%	*	---
Oregon HUC Totals ^b	459,400	76%	145,500	24%	0	0%	604,900	100%

*: Less than 1 percent of total acres. See below for special considerations.

a: Estimate from Farm Service Agency records and includes CRP/CREP.

b: Totals are approximate due to rounding and small unknown acreages.



Special Considerations for This 8-Digit HUC:

- None

	Type of Land	ACRES	% of Irrigated Lands	% of HUC
Irrigated Lands (1997 NRI ³ Estimates for Non-Federal Lands Only)	Cultivated Cropland	37,100	46%	6%
	Uncultivated Cropland	31,100	38%	5%
	Pastureland	13,000	16%	2%
	Total Irrigated Lands	81,200	100%	13%

(Continued on the following pages)


[Back to Contents](#)



Common Resource Area Map

[Back to Contents](#)

Only the major units are described below - for descriptions of all units within the HUC, go to: <http://ice.or.nrcs.usda.gov/website/cra/viewer.htm>



10.1 – Central Rocky and Blue Mountains Foothills - Warm Dry Blue and Seven Devils Mountains Foothills:

This unit is between the Blue and Wallowa Mountains in Oregon and the northwestern part of the Snake River Plain. It is characterized by rangeland soils on hills and mountains associated with basalt and exposed tuffaceous sediment. The Cascade Range and the Blue and Wallowa Mountains block any maritime influence, creating a continental climate. As a result, plants are subject to wide temperature ranges, a high rate of evapotranspiration, and high early-season moisture stress. The dominant soils are those of the Brogan, Simas, Ruckles, and Ruclick series. The temperature regime is mesic, and the moisture regime is aridic. The mean annual precipitation is 9 to 12 inches. The vegetation is Wyoming big sagebrush and bluebunch wheatgrass (warm, dry climate).

10.16 – Central Rocky and Blue Mountains Foothills – Cool, Moist Blue Mountains Foothills:

This unit is characterized by rangeland soils on hills and mountains associated with basalt. It is similar to the Lava Fields unit except that this unit has higher precipitation and a xeric soil moisture regime. The temperature regime is frigid. The mean annual precipitation is 12 to 20 inches. The dominant soils are those of the Ateron, Durkee, Menbo, Merlin, and Observation series. The vegetation is dominantly mountain big sagebrush and Idaho fescue (cool, moist climate).

11.1 – Snake River Plains - Treasure Valley:

This unit is characterized by irrigated cropland, pastureland, and rapidly growing cities, suburbs, and industries. Many canals, reservoirs, and diversions are present. Aridic soils are dominant; irrigation is required to grow commercial crops. Surface water quality has been significantly affected by channel alteration, dams, irrigation return flow, and urban, industrial, and agricultural pollution. Crops include wheat, barley, alfalfa, sugar beets, potatoes, and beans. Crop diversity is greater, temperatures are warmer, and the mean frost free season is longer in this unit than in other units. Population density is much greater in this unit than in the nearby rangeland-dominated units.

11.7 – Snake River Plains - Dry Unwooded Alkaline Foothills:

This shrub- and grass-covered unit is higher and more rugged than adjacent valley units. Unlike other units, this unit consists of terrace deposits of alkaline lacustrine and supports a unique flora. Shallow and moderately deep soils over a cemented pan are common. The potential natural vegetation is saltbush-greasewood and sagebrush steppe. Presently, cheatgrass and crested wheatgrass are also common, and the unit is used for livestock grazing.

Physical Description – Continued


[Back to Contents](#)

		ACRES	ACRE-FEET			
Irrigated Adjudicated Water Rights (OWRD ⁴)	Surface	10,930	33,170			
	Well	6,873	20,623			
	Total Irrigated Adjudicated Water Rights	17,803	53,794			
Stream Flow Data	USGS 13233300 MALHEUR RIVER BELOW NEVADA DAM, NEAR VALE, OR	Total Avg. Yield	149,936			
		May – Sept. Yield	37,077			
		MILES	PERCENT			
Stream Data ⁵	Total Miles – Major (100K Hydro GIS Layer)	357	---			
	303d/TMDL Listed Streams (DEQ)	46	13%			
	Anadromous Fish Presence (StreamNet)	0	0%			
	Bull Trout Presence (StreamNet)	0	0%			
*Percent of Total Miles of Streams in HUC						
		ACRES	PERCENT			
Land Cover/Use ²	Forest	178	0%			
	Grain Crops	1,169	3%			
	Grass/Pasture/Hay	8,001	20%			
	Orchards/Vineyards	0	0%			
	Row Crops	323	0%			
	Shrub/Rangelands – Includes CRP Lands	30,182	75%			
	Water/Wetlands/Developed/Barren	657	2%			
	Total Acres of 100-foot Stream Buffers	40,511	---			
Land Capability Class	1 – slight limitations	18,900	19%			
	2 – moderate limitations	27,000	27%			
	3 – severe limitations	36,200	37%			
	4 – very severe limitations	14,500	15%			
	5 – no erosion hazard, but other limitations	0	0%			
	6 – severe limitations; unsuitable for cultivation; limited to pasture, range, forest	2,100	2%			
	7 – very severe limitations; unsuitable for cultivation; limited to grazing, forest, wildlife habitat	0	0%			
	8 – miscellaneous areas; limited to recreation, wildlife habitat, water supply	0	0%			
	Total Croplands & Pasturelands	98,700	---			
Confined Animal Feeding Operations – Oregon CAFO Permit – 12/2004						
Animal Type	Dairy	Feedlot	Poultry	Swine	Mink	Other
No. of Permitted Farms	4	2	0	0	0	0
No. of Permitted Animals	3,865	33,100	0	0	0	0

Resource Concerns

[Back to Contents](#)


Tons of Soil Loss by Water Erosion: Due to the limited amount of non-Federal cropland and pastureland within this HUC, no reliable NRI soil loss estimates are available.



- ❖ Most of the listed stream miles exceed State water quality standards for temperature. Elevated stream temperatures may be due to inadequate riparian shade, stream channel widening, and other anthropogenic or natural causes.
- ❖ Conservation practices that can be used to address these water quality issues include irrigation water management, grazing management, and use of riparian buffers.

Watershed Projects, Plans, Studies, and Assessments			
NRCS Watershed Projects ⁶		NRCS Watershed Plans, Studies, and Assessments ⁷	
Name	Status	Name	Status
None	None	None	None
ODEQ TMDL's ⁸		ODA Agricultural Water Quality Management Plans ⁹	
Name	Status	Name	Status
None	None	Malheur	Completed
OWEB Watershed Council ¹⁰		Watershed Council Assessments ¹¹	NWPCC Subbasin Plans and Assessments ¹⁸
Owyhee Watershed Council Malheur Watershed Council	Malheur Basin Watershed Action Plan and Assessment		Malheur

(Continued on page 8)



Map Footnote [417](#)

FEDERALLY LISTED THREATENED AND ENDANGERED SPECIES ¹²	
THREATENED SPECIES	CANDIDATE SPECIES
Mammals - Canada lynx	Birds – Yellow-billed cuckoo
Birds – Bald eagle	Amphibians and Reptiles – Columbia spotted frog
Fish – Bull trout, Lahontan cutthroat trout	PROPOSED SPECIES - None
Plants – Howell's spectacular thelypody, Malheur wire-lettuce	
ESSENTIAL FISH HABITAT ¹³ - None	

Resource Concerns - Continued

[Back to Contents](#)

Resource Concerns/Issues by Land Use								
SWAPA +H Concerns	Specific Resource Concern/Issue	Grass\Pasture\ Hay	Grain Crops	Row Crops	Perennial Crops (Orch/Vine/ Berries)	Shrub/Range	Forest	Confined Feeding Operations
Soil Erosion	Concentrated Flow or Gully					X		
	Streambank					X		
	Irrigation Induced	X	X	X				
Soil Condition	Tilth, Crusting, Infiltration, Organic Matter		X	X				
Soil Contamination	Excess Fertilizers and Pesticides			X				X
Water Quantity	Water Management for Irrigated Land	X	X	X				
	Water Management for Nonirrigated Land					X		
Water Quality, Groundwater	Pesticides			X				
	Nutrients and Organics		X	X				
Water Quality, Surface	Pesticides			X				
	Nutrients and Organics		X	X				X
	Suspended Sediments and Turbidity	X	X	X		X		
	Pathogens		X	X				X
Air Quality	Undesirable Odors from Agricultural Sources							X
Plant Suitability	Site and Intended Use Suitability	X						
	Invasive and Noxious Weeds	X	X	X		X		
Plant Condition	Productivity, Health, and Vigor	X				X		
Plant Management	Establishment, Growth, and Harvest					X		
Animal Habitat, Domestic	Water - Quantity and Quality					X		
	Management							
Animal Habitat, Wildlife	Food, Cover ,and/or Shelter	X	X	X		X		
Human, Economics	Land Use Constraints/Restrictions							X
	High Risk and Uncertainty			X				X
	High Capital/Financial Costs	X	X	X		X		X
	High Management Level Required			X				X
	Low or Unreliable Profitability	X	X	X				
Human, Political	Inadequate Availability of Cost-Share Programs					X		
	High Degree of Controversy							X

Pasture/Hay

- Better irrigation water management is practiced in areas used for alfalfa than in areas of pasture.
- In some areas of pasture, a lack of proper grazing management has lead to its poor condition.
- Areas of pasture commonly are adjacent to streams, which can contribute to streambank erosion and sedimentation as a result of loss of riparian vegetation.

Grain and Row Crops

- Most grain is produced in rotation with other crops (potatoes, onions, corn, alfalfa, etc.)
- Irrigation-induced erosion may occur on fields used for crops such as potatoes or corn.
- Surface-irrigated areas of grain are also prone to irrigation-induced erosion.
- Surface irrigation of crops generates tailwater returns to area streams and drains that are high in content of nutrients and sediment.
- Water conservation is always a concern with irrigated crops, but irrigation water management is better in areas used for row crops than it is in areas used as pasture.

Confined Animal Feeding Operations (CAFOs)

- Livestock manure, pathogens, and odors are continuing issues for CAFOs.
- Winter feeding of cow-calf herds can generate erosion and contribute to poor water quality.

Rangeland

- Rangeland can become infested with noxious weeds, annual grasses, and shrubs due to inadequate forage and grazing management.
- Loss of riparian vegetation contributes to the warming and nutrient-loading of streams.


Census and Social Data^{/14}

[Back to Contents](#)

Number of Farms: 221

Number of Operators: 362

- Full-Time Operators: **130**
- Part-Time Operators: **232**



Estimated Level of Willingness and Ability to Participate in Conservation^{/15}: **Moderate to High**



Most operators in the Lower Malheur subbasin are well educated, aware of local resource concerns, and likely to have conservation plans; have adopted some conservation practices; and understand the economic and environmental benefits of conservation. Most recommended conservation practices can be implemented incrementally and are compatible with local management systems and equipment. The perceived high capital costs of conservation and risks associated with intense irrigated agriculture discourage many operators from adopting conservation systems.

Additional financial incentives or other risk-reducing incentives may increase the adoption of conservation in the subbasin.

Evaluation of Social Capital^{/16}: **Moderate**

Social capital and the ability of the community to solve problems and support conservation are estimated to be moderate. Recent trends indicate that the population of the subbasin is increasing slightly. The primary occupation of new landowners commonly is non-agricultural and not resource based. People moving to the area commonly do so for the rural, high-quality lifestyle and relatively inexpensive housing and property. Newcomers to the area tend to look at the natural resources as recreational opportunities, not as a means for making a living. In part, this has resulted in community interest shifting from agricultural and natural resource concerns to issues related to improving schools, transportation, health services, and so on.


Until resource management and agriculture regain the attention of the community, it is unlikely that the community will be a significant partner in the diffusion of conservation in the agricultural community.



Progress/Status

[Back to Contents](#)

PRMS Data	FY99	FY00	FY01	FY02	FY03	Avg/Year	Total
Total Conservation Systems Planned (Acres)	312	1,078	343	0	64	359	1,797
Total Conservation Systems Applied (Acres)	0	1,078	216	207	1,245	549	2,746
Conservation Treatment (Acres)							
Waste Management	2	1	0	1	0	1	4
Buffers	0	0	0	0	0	0	0
Erosion Control	353	925	69	184	187	344	1,718
Irrigation Water Management	468	187	20	85	131	178	891
Nutrient Management	173	792	180	136	466	349	1,747
Pest Management	0	792	0	0	0	158	792
Prescribed Grazing	0	0	0	0	300	60	300
Trees & Shrubs	0	0	0	0	0	0	0
Conservation Tillage	0	769	0	0	0	154	769
Wildlife Habitat	0	168	0	0	13	36	181
Wetlands	0	2	0	0	0	0	2



Estimates are based on information received from local conservationists in the watershed.

Progress over the last 5 years has been focused on:

- ~ Erosion control and irrigation water management in areas of grain and row crops.
 - ~ Nutrient and pest management.
 - ~ Conservation tillage.
 - ~ Wildlife habitat management, including use of buffers, trees, and shrubs in riparian areas.
- ❖ Most grain and row crop producers practice conservation cropping and residue management.
 - ❖ Most hay producers use good irrigation water management, but adequate grazing and water management commonly is lacking on pastures.
 - ❖ Most livestock operations are at the progressive level. Focus has been placed on meeting State CAFO regulations. High capital cost has hindered conservation adoption to attain the RMS level.
 - ❖ Most private, non-industrial woodlots are scattered throughout areas of rangeland and are not primarily managed for timber production.

Lands Removed from Production through Farm Bill Programs

- ❖ Conservation Reserve Program (CRP): **None**
- ❖ Wetland Restoration Program (WRP): **None**
- ❖ Conservation Reserve Enhancement Program (CREP): **None**

Footnotes/Bibliography

[Back to Contents](#)

All data is provided "as is." There are no warranties, express or implied, including the warranty of fitness for a particular purpose, accompanying this document. Use for general planning purposes only.

1. Ownership Layer – Source: The 1:24,000 scale public ownership layer is the land ownership/management for public entities, including Federal, Tribal, State, and local entities. This is a seamless, statewide Oregon Public Ownership vector layer composed of fee ownership of lands by Federal, State, Tribal, county, and city agencies. The layer is comprised of the best available data compiled at 1:24,000 scale or larger, and the line work matches GCDB boundary locations and ORMAP standards where possible. The layer is available from the State of Oregon GIS Service Center: <http://www.gis.state.or.us/data/alphalist.html>. For current ownership status, consult official records at appropriate Federal, State, and county offices. Ownership classes grouped to calculate Federal ownership vs. non-Federal ownership by the Water Resources Planning Team.
2. National Land Cover Dataset (NLCD) - Originator: U.S. Geological Survey (USGS); Publication date: 19990631; Title: Oregon Land Cover Data Set, Edition: 1; Geospatial data presentation form: Raster digital data; Publisher: U.S. Geological Survey, Sioux Falls, SD, USA; Online linkage: <http://edcwww.cr.usgs.gov/programs/lccp/nationallandcover.html>; Abstract: These data can be used in a geographic information system (GIS) for any number of purposes, such as assessing wildlife habitat, water quality, pesticide runoff, land use change, etc. The State data sets are provided with a 300-meter buffer beyond the State border to facilitate combining the State files into larger regions.
3. ESTIMATES FROM THE 1997 NRI DATABASE (REVISED DECEMBER 2000) REPLACE ALL PREVIOUS REPORTS AND ESTIMATES. Comparisons made using data published for the 1982, 1987, or 1992 NRI may produce erroneous results. This is because of changes in statistical estimation protocols and because all data collected prior to 1997 were simultaneously reviewed (edited) as 1997 NRI data were collected. All definitions are available in the glossary. In addition, this December 2000 revision of the 1997 NRI data updates information released in December 1999 and corrects a computer error discovered in March 2000. For more information: <http://www.nrcs.usda.gov/technical/NRI/>
4. Irrigated Adjudicated Water Rights – Water Rights Information System (WRIS), Oregon Water Resources Department, <http://www.wrd.state.or.us/maps/wrlexport.shtml>
5. StreamNet is a cooperative venture of the Pacific Northwest's fish and wildlife agencies and tribes and is administered by the [Pacific States Marine Fisheries Commission](#). StreamNet provided data and data services in support of the region's fish and wildlife program and other efforts to manage and restore the region's aquatic resources. Official StreamNet website: <http://www.streamnet.org/>
6. Natural Resources Conservation Service, Watershed Projects Planned and Authorized, <http://www.nrcs.usda.gov/programs/watershed/Purpose>.
7. Natural Resources Conservation Service, Watershed Plans, Studies, and Assessments completed, http://www.nrcs.usda.gov/programs/watershed/Surveys_Plng.html#Watershed%20Surveys%20and%20Plan
8. Oregon Department of Environmental Quality Total Maximum Daily Loads, <http://www.deq.state.or.us/wq/TMDLs/TMDLs.htm>
9. Oregon Department of Agriculture, Agricultural Water Quality Management Plans, http://www.oregon.gov/ODA/NRD/water_agplans.shtml

Footnotes/Bibliography Continued

[Back to Contents](#)

All data is provided "as is." There are no warranties, express or implied, including the warranty of fitness for a particular purpose, accompanying this document. Use for general planning purposes only.

10. Oregon Watershed Enhancement Board, <http://oregon.gov/OWEB/WSHEDS/index.shtml>
11. Watershed Assessments completed by local watershed councils following the Oregon Watershed Assessment Manual, http://oregon.gov/OWEB/docs/pubs/ws_assess_manual.shtml.
12. NRCS Field Office Technical Guide, Section II, Threatened and Endangered List.
13. Magnuson-Stevens Fishery Conservation and Management Act, Public Law 94-265. As amended through October 11, 1996.
14. Data were taken from the 2002 Agricultural Census and adjusted by percent of HUC in the county or by percent of zip code area in the HUC, depending on the level of data available. Data were also taken from the U.S. Population Census, 2000.
15. Conservation participation was estimated using NRCS Social Sciences Technical Note 1801, [Guide for Estimating Participation in Conservation](#), 2004. Four categories of indicators were evaluated: Personal characteristics, farm structural characteristics, perceptions of conservation, and community context. Estimates are based on information received from local conservationists in the watershed.
16. Social capital is an indicator of the community's ability and willingness to work together to solve problems. A high amount of social capital helps a community to be physically healthy, socially progressive, and economically vigorous. A low amount of social capital typically results in community conflict, lack of trust and respect, and unsuccessful attempts to solve problems. The evaluation is based on NRCS Technical Report Release 4.1, March, 2002: [Adding Up Social Capital: An Investment in Communities](#). Local conservationists provided information to measure social capital. Scores range from 0 to 76.
17. [Surface and Groundwater Resource Protection Map](#)
 - a. 2002 303d Listed Streams designated by Oregon Department of Environmental Quality and approved by the Environmental Protection Agency, Section 303d Clean Water Act, <http://www.deq.state.or.us/wq/303dlist/303dpage.htm>
 - b. Groundwater Management Areas designated by the Oregon Department of Environmental Quality, Oregon Revised Statutes – Ground Water ORS 468B.150 to ORS 468B.190, <http://www.deq.state.or.us/wq/groundwa/wqgw.htm>
 - c. Groundwater Restricted Areas designated by Oregon Water Resources Commission, Oregon Department of Water Resources, http://egov.oregon.gov/OWRD/PUBS/aquabook_protections.shtml
 - d. The Sole Source Aquifer (SSA) Protection Program is authorized by Section 1424(e) of the Safe Drinking Water Act of 1974 (Public Law 93-523, 42 U.S.C. 300 et. seq), <http://www.epa.gov/safewater/ssanp.html>
18. Subbasin assessments and plans are developed by local groups (SWCDs, watershed councils, tribes, and others) as part of the Northwest Power and Conservation Council's fish and wildlife program in the Columbia River Basin. This program is funded and implemented by the Bonneville Power Administration. <http://www.nwcouncil.org/fw/subbasinplanning/Default.htm>.