

Legend

soils_csp_chester

MUNAME

	Carmichael loam, 0 to 2 percent slopes		Reybold loam, 2 to 5 percent slopes
	Christiana silt loam, 0 to 5 percent slopes		Reybold loam, 5 to 10 percent slopes
	Collington fine sandy loam, 2 to 5 percent slopes		Reybold-Queponco complex, 0 to 2 percent slopes
	Collington fine sandy loam, 5 to 15 percent slopes		Reybold-Urban land complex, 0 to 5 percent slopes
	Corsica mucky loam, 0 to 2 percent slopes		Rosedale loamy sand, 2 to 5 percent slopes
	Crosiadore silt loam, 0 to 2 percent slopes		Sassafras loam, 0 to 2 percent slopes
	Downer sandy loam, 0 to 2 percent slopes		Sassafras loam, 2 to 5 percent slopes
	Downer sandy loam, 2 to 5 percent slopes		Sassafras sandy loam, 0 to 2 percent slopes
	Downer sandy loam, 5 to 10 percent slopes		Sassafras sandy loam, 2 to 5 percent slopes
	Elkton silt loam, 0 to 2 percent slopes		Sassafras sandy loam, 5 to 10 percent slopes
	Evesboro loamy sand, 0 to 5 percent slopes		Sassafras soils, 10 to 15 percent slopes, moderate
	Fallsington sandy loam, 0 to 2 percent slopes		Udortheents, 0 to 10 percent slopes
	Fallsington-Blackiston complex, 0 to 2 percent slopes		Udortheents, 10 to 30 percent slopes
	Fallsington-Udortheents-Urban Land Complex, 0 to 5 percent slopes		Udortheents, waste substratum, 0 to 2 percent slopes
	Greenwich loam, 0 to 2 percent slopes		Udortheents-Mattapex-Urban land complex, 0 to 8 percent slopes
	Greenwich loam, 2 to 5 percent slopes		Unicorn loam, 0 to 2 percent slopes
	Hambrook sandy loam, 0 to 2 percent slopes		Unicorn loam, 2 to 5 percent slopes
	Hambrook sandy loam, 2 to 5 percent slopes		Water
	Hambrook-Fallsington-Woodstown complex, 0 to 5 percent slopes		Westphalia soils, 10 to 30 percent slopes
	Hambrook-Udortheents-Urban Land Complex, 0 to 5 percent slopes		Whitemarsh silt loam, 0 to 2 percent slopes
	Hammonton loamy sand, 0 to 2 percent slopes		Woodstown loam, 0 to 2 percent slopes
	Hammonton sandy loam, 0 to 2 percent slopes		Woodstown loam, 2 to 5 percent slopes
	Hammonton sandy loam, 2 to 5 percent slopes		Woodstown sandy loam, 0 to 2 percent slopes
	Hammonton-Fallsington-Longmarsh complex, 0 to 5 percent slopes		Woodstown sandy loam, 2 to 5 percent slopes
	Hurlock sandy loam, 0 to 2 percent slopes		Zekiah sandy loam, frequently flooded
	Ingleside loamy sand, 0 to 2 percent slopes		Zekiah-Codorus complex, 0 to 2 percent slopes, frequently flooded
	Ingleside loamy sand, 2 to 5 percent slopes		
	Ingleside sandy loam, 0 to 2 percent slopes		
	Ingleside sandy loam, 2 to 5 percent slopes		
	Ingleside sandy loam, 5 to 10 percent slopes		
	Ingleside-Hammonton-Fallsington complex, 0 to 10 percent slopes		
	Ingleside-Udortheents-Urban Land Complex, 0 to 5 percent slopes		
	Kentuck mucky silt loam, 0 to 2 percent slopes		
	Keyport silt loam, 0 to 2 percent slopes		
	Keyport silt loam, 2 to 5 percent slopes		
	Lenni sandy loam, 0 to 2 percent slopes		
	Lenni silt loam, 0 to 2 percent slopes		
	Longmarsh and Indiantown soils, frequently flooded		
	Matapeake silt loam, 0 to 2 percent slopes		
	Matapeake silt loam, 2 to 5 percent slopes		
	Mattapex silt loam, 0 to 2 percent slopes		
	Mattapex silt loam, 2 to 5 percent slopes		
	Mattapex silt loam, 5 to 10 percent slopes, severe		
	Mullica sandy loam, 0 to 2 percent slopes		
	Nassawango silt loam, 0 to 2 percent slopes		
	Nassawango silt loam, 2 to 5 percent slopes		
	Othello silt loam, 0 to 2 percent slopes		
	Pineyneck loam, 0 to 2 percent slopes		
	Pineyneck loam, 2 to 5 percent slopes		
	Pits, sand and gravel, 0 to 5 percent slopes		
	Reybold loam, 0 to 2 percent slopes		