



Spokane  
ICCS

## Northwest Team Estimated Average Annual Wind Erosion on Cultivated Cropland

		< 2 tons/acre /year	2.0 - 4.9 tons/acre /year	5.0 to 9.9 tons/acre /year	10.0 + tons/acre /year	Total
1982	Estimated	105,700	0	0	0	105,700
1982	Error *	19,000	0	0	0	19,000
1987	Estimated	88,000	0	0	0	88,000
1987	Error *	16,800	0	0	0	16,800
1992	Estimated	98,600	0	0	0	98,600
1992	Error *	15,700	0	0	0	15,700
1997	Estimated	86,200	0	0	0	86,200
1997	Error *	14,900	0	0	0	14,900


Estimates may not total because of rounding.

\* The error referred to in the table is the standard error of the estimate.

(To obtain the margin of error at the 95% confidence limit multiply the error by 1.96.)

Geographic Area of the Northwest Team of Washington State:

<b>Clallam</b>	<b>King</b>	<b>San Juan</b>	<b>Whatcom</b>
<b>Island</b>	<b>Kitsap</b>	<b>Skagit</b>	
<b>Jefferson</b>	<b>Pierce</b>	<b>Snohomish</b>	



Wind erosion on cultivated cropland is not a problem in northwestern Washington.