

UNITED STATES DEPARTMENT OF AGRICULTURE  
NATURAL RESOURCES CONSERVATION SERVICE  
NEW MEXICO PLANT MATERIALS CENTER  
LOS LUNAS, NEW MEXICO

and

NEW MEMCO STATE UNIVERSNY  
AGRICULTURAL SCIENCE CENTER AT LOS LUNAS  
LOS LUNAS, NEW MEXICO

NOTICE OF RELEASE OF A SELECTION OF CANE BLUESTEM  
SELECTED CLASS OF GERMPLASM

The Natural Resources Conservation Service (NRCS), United States Department of Agriculture and New Mexico State University (NMSP Agricultural Science Center at Los Lunas announce the release of a selected class of cane bluestem (*Bothriochloa barbinodis*) (Heter) for the southwestern United States.

As a selected release, this germplasm win be referred to as the Grant germplasm cane bluestem. It has been assigned the NRCS accession number 9066390. Grant germplasm is released as a selected class of certified seed (natural track).

The alternative release procedure is justified because existing commercial sources of cane bluestem are inadequate. The commercial sources that do exist have not been selected for performance under agronomic conditions. Propagation material of this species is needed for ecosystem restoration and enhancement and for beautification in urban and rural landscaping situations. The potential for immediate use is high. No commercial cultivars of cane bluestem have been previously released.

**Collection Site Information:** Grant germplasm was originally collected in 1982. Accessions were collected by USDA-NRCS field office personnel from native stands in New Mexico and Arizona (Table 1).

**Table 1.** Collection sites of accessions making up the composite Grant germplasm of cane bluestem.

Accession Number	County & State	Elevation (ft)	MLRA	Collector
1. 399204	Pima AZ	3496		M Lamoreaux
2. 9032338	Grant NM	6197	36	H. Bray
3. 9032398	Socorro NM	4659	42	D. Reasner
4. 9032342	Grant NM	4898	36	H. Bray
5. 9032343	Grant NM	4678	36	H. Bray
6. 9032344	Grant NM	5279	36	H. Bray

**Description:** Grant germplasm cane bluestem is a coarse warm-season bunch grass. It is a large robust grass with foliage well distributed along the stem. Blades are 2-7 cm broad and 25-30 cm or more long. Plants may be 1.5 -2 m tall, depending upon environmental conditions. Flowering is from June to July.

Cane bluestem has been observed to be remarkably drought tolerant and is well adapted to southwestern rangeland. It has been classified as fair to good forage for cattle and wildlife. Cane bluestem has produced yields of approximately 4 metric tons/ha annually under dryland conditions. When green the total protein of cane bluestem forage is about 10%, however when dormant forage protein may drop to 4.5%. When grown under similar conditions cane bluestem and switchgrass (*Panicum virgatum* L.) compared favorably. Under drought conditions, cane bluestem produced more forage than switchgrass.

**Methods of Breeding and/or Selection:** Grant germplasm cane bluestem was selected from 21 accessions collected from New Mexico and Arizona. Accessions were individual populations that were collected by USDA-NRCS field office staff. From the initial accessions, six were selected for superior forage and seed yield. Equal amounts of seed from each accession were combined. The composite mix was seeded to become Grant germplasm.

**Environmental Impact Assessment:** Grant germplasm cane bluestem is a selection of naturally occurring germplasm. Grant germplasm did not meet the assessment of a plant, which could become invasive based on guidelines adopted by the NRCS Plant Materials Program.

**Anticipated Conservation Use:** The potential uses of the Grant germplasm include erosion control, wildlife food/cover, restoration of disturbed sites, increasing plant diversity of rangelands and for low water use beautification of urban and rural landscapes.

**Anticipated Area of Adaptation:** Cane bluestem is naturally occurring from Oklahoma to Arizona and Texas, and south into Mexico. It commonly grows on open rangelands, mesas, dry rocky or sandy slopes and open plains.

**Availability of Plant Materials:** Breeder and/or foundation seed will be maintained at the NRCS New Mexico Plant Materials Center. Seed will be distributed to interested growers through the New Mexico Crop Improvement Association.

**References:**

1. Bernardon, A.E., D.L. Huss and W.G. McCully. 1967. Effects of herbage removal on seedling development in cane bluestem.
2. Allred, K.W. 1997. A field guide to the grasses of New Mexico. 2<sup>nd</sup> ed. New Mexico State University. Las Cruces, NM.
3. Gould, F.W. 1975. The grasses of Texas. Texas A&M Press. College Station, TX.
4. Gould, F.W. 1977. Grasses of the Southwestern United States. University of Arizona Press. Tuscon, AZ.
5. Humphrey, R. R. 1960. Arizona range grasses their description, forage value and management. Arizona Agricultural Experiment Station Bulletin 298.
6. Koshi, P.T., H.V. Eck, J..Stubbendieck and W.G. McCully. 1977. Cane bluestems: Forage yield, forage quality and water-use efficiency. Journal of Range Management. 30(3):190-193.
7. Martin, W.C. and C.R. Hutchins. 1980 A flora of New Mexico. J. Cramer. Valduz, West Germany

**Prepared by:**

E. Rarnona Garner, USDA-NRCS New Mexico Plant Materials Center, 1036 Miner Street, SW, Los Lunas, New Mexico 87031

Michelle Jespersen, NMSU Agricultural Science Center at Los Lunas, 1036 Miller Street, SW, Los Lunas, New Mexico 87031

Danny Goodson, USDA-NRCS New Mexico Plant Materials Center, 1036 Miller Street, SW, Los Lunas, New Mexico 87031

Signatures for release of:  
Grant Germplasm Cane bluestem (*Bothriochloa barbinodis*)

---



*for* Rosendo Trevino III  
State Conservationist  
United States Department of Agriculture  
Natural Resources Conservation Service  
Albuquerque, New Mexico

3/23/01

Date



I. Miley Gonzalez  
Associate Dean/Director Agricultural Experiment Station  
New Mexico State University  
Las Cruces, New Mexico

3/20/01

Date



*for* Diane Gelbund  
Director, Ecological Science Division  
United State Department of Agriculture  
Natural Resources Conservation Service  
Washington, D.C.

5/10/01

Date