

Animal Enhancement Activity – ANM03 - Incorporate native grasses and/or legumes to 15% or more of herbage dry matter productivity



Enhancement Description

Improve pasture by increasing native grasses and/or legumes to 15% of herbage dry matter (productivity by weight) using adapted species and varieties, appropriate seeding rates, and timing of seeding. Pastures containing about 15% native grasses and/or legumes by weight dry matter are approximately equal to 30% foliar cover.

Land Use Applicability

Pastureland

Benefits

Enhancing existing pasture by incorporating native grasses and legumes can provide:

1. Improved forage quality and quantity
2. Improved soil fertility (legumes fix nitrogen in the soil), increase organic matter
3. Increased plant diversity and promote wildlife habitat
4. Additional forage during seasonal slump periods
5. Extended grazing season
6. Food source for pollinating insects

Conditions Where Enhancement Applies

This enhancement only applies to acres of pasture land use that DO NOT currently have a mixed stand of native grasses and/or legumes.

Criteria

A written grazing management plan that outlines specific goals and objectives, including:

1. Utilize adapted species, seeding rates and seeding dates according to local NRCS practice standards.
2. Determine species composition before and after seeding. Species composition must be 15% or more of native grasses and/or legumes.
3. If legumes are incorporated, a current soil test is required. Apply lime and fertilizer to facilitate establishment and persistence of legumes as required by the current soil test report.
4. Livestock stocking rates that will allow for proper forage utilization.

Note: Bloat can be a risk to grazing livestock where legumes make up greater than 50% of the total forage. Legumes with the highest likelihood to cause bloat include white clover, alfalfa, annual medics and Persian clover. Red clover, crimson clover and subterranean clover would be classified as moderately likely to cause bloat, while berseem clover and arrowleaf clover are low risks for causing bloat. Legumes that don't cause bloat are birdsfoot trefoil, sainfoin and



crownvetch. Livestock producers grazing alfalfa aftermath in the fall months should be cautioned of bloat, especially following a killing frost. The recommendation for grazing frost killed alfalfa is to wait 5 to 7 days after the killing frost (less than 28 degrees Fahrenheit) before grazing. This will allow the live tissue to fully break down, minimizing the soluble leaf proteins, and making a much safer feed base for ruminant livestock. If bloat is a concern, there are several precautions that can be taken. (A technical reference sheet will be available to address these issues).

Adoption Requirements

This enhancement is considered adopted when the subject pasture acre(s) contain 30% or more foliar coverage of native grasses and/or legumes.

Documentation Requirements

1. A written planting specifications plan identifying:
 - a. Plant species' to be seeded,
 - b. Seeding rates and dates,
 - c. Site preparations and planting method, and
 - d. Amounts of fertilizer and lime to be applied.
2. Map showing locations where seeding activity is applied.
3. Copy of the grazing management plan.

References

Ball, D.M., C.S. Hoveland and G.D. Lacefield. 2007. Southern Forages, 4th Edition. International Plant Nutrition Institute, Norcross, GA.

Bartholomew, P.W. 2005. Comparison of Conventional and Minimal Tillage for Low-input Pasture Improvement. Online-Forage and Grazinglands – Plant Management Network.

Ruffin, B.G. 1994. Controlling Bloat in Cattle. Alabama Cooperative Extension System, Pub. ANR-148.

USDA-NRCS. 2010. Conservation Practice Standard: Forage and Biomass Planting-Code 512.



United States Department of Agriculture
Natural Resources Conservation Service

IDAHO ADDENDUM 2013

Animal Enhancement Activity – ANM03 – *Incorporate Native Grasses and/or Legumes to 15% or More of Herbage Dry Matter Productivity*

Additional guidance for incorporating native grasses and legumes into the forage base:

Appropriate native grasses and/or legumes will depend on site conditions and intended use. Potential native grasses for pasture include western wheatgrass, thickspike wheatgrass, basin wildrye, and big bluegrass. Native legumes available include basalt milkvetch and Utah or northern sweetvetch. A desirable native forb to include is western yarrow.

Refer to the following guidance for more information on selecting and establishing native grasses and legumes that fit your conditions and needs.

Idaho NRCS Plant Materials Technical Note 10, *Pasture and Range Seedings*. ftp://ftp-fc.sc.egov.usda.gov/ID/programs/technotes/tn10_pasture_1011.pdf

Idaho NRCS Plant Materials Technical Note 11, *Pasture – Species Selections and Grazing Management Guidelines*. ftp://ftp-fc.sc.egov.usda.gov/ID/programs/technotes/tn11_species_selection.pdf

Idaho NRCS Plant Materials Technical Note 24, *Conservation Plant Species for the Intermountain West*. ftp://ftp-fc.sc.egov.usda.gov/ID/programs/technotes/tn24_seed_species_1011.pdf

Idaho NRCS Plant Material Technical Note 24 Supplement: *Intermountain Planting Guide*, USDA-ARS Forage and Range Research Lab/Utah State Extension, AG 510. <ftp://ftp-fc.sc.egov.usda.gov/ID/programs/technotes/tn24supplement.pdf>

University of Idaho Bulletin 547, *Idaho Forage Handbook*.
<http://info.ag.uidaho.edu/forage/index.html>.

**This activity may NOT be used with the following enhancements:
ANM21**

**Potential duplicate practices:
512 – Forage and Biomass Planting, 528 – Prescribed Grazing**