



United States
Department of
Agriculture



United States
Department of
the Interior



Natural
Resources
Conservation
Service



National Park
Service

Soil Survey of Manzanar National Historic Site, California



How To Use This Soil Survey

This publication consists of text, tables, and a map. The text includes descriptions of detailed soil map units and provides an explanation of the information presented in the tables. It also includes a glossary of terms used in the text and tables and a list of references.

The detailed soil map can be useful in planning the use and management of small areas. To find information about your area of interest, locate that area on the map sheet. Note the map unit symbols that are in that area. Go to the Contents, which lists the map units by symbol and name and shows where each map unit is described.

The Contents shows which table has data on a specific land use for each detailed soil map unit. Also see the Contents for sections of this publication that may address your specific needs.

National Cooperative Soil Survey

This soil survey is a publication of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service has leadership for the Federal part of the National Cooperative Soil Survey. This survey was made cooperatively by the United States Department of Agriculture, Natural Resources Conservation Service, and the United States Department of the Interior, National Park Service.

The soil map in this survey may be copied without permission. Enlargement of this map, however, could cause misunderstanding of the detail of mapping. If enlarged, the map does not show the small areas of contrasting soils that could have been shown at a larger scale.

Literature Citation

The correct citation for this survey is as follows:

United States Department of Agriculture, Natural Resources Conservation Service, and the United States Department of the Interior, National Park Service. 2013. Soil survey of Manzanar National Historic Site, California. (Accessible online at: http://soils.usda.gov/survey/printed_surveys/)

Cover Caption

The reconstructed guard tower in Manzanar National Historic Site, located in an area of Winnedumah fine sandy loam, 0 to 2 percent slopes. View is to the west.

Additional information about the Nation's natural resources is available online from the Natural Resources Conservation Service at <http://www.nrcs.usda.gov/>.

Nondiscrimination Statement

Nondiscrimination Policy

The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers, employees, and applicants for employment on the basis of race, color, national origin, age, disability, sex, gender identity, religion, reprisal, and where applicable, political beliefs, marital status, familial or parental status, sexual orientation, whether all or part of an individual's income is derived from any public assistance program, or protected genetic information. The Department prohibits discrimination in employment or in any program or activity conducted or funded by the Department. (Not all prohibited bases apply to all programs and/or employment activities.)

To File an Employment Complaint

If you wish to file an employment complaint, you must contact your agency's EEO Counselor (<http://directives.sc.egov.usda.gov/33081.wba>) within 45 days of the date of the alleged discriminatory act, event, or personnel action. Additional information can be found online at http://www.ascr.usda.gov/complaint_filing_file.html.

To File a Program Complaint

If you wish to file a Civil Rights program complaint of discrimination, complete the USDA Program Discrimination Complaint Form, found online at http://www.ascr.usda.gov/complaint_filing_cust.html or at any USDA office, or call (866) 632-9992 to request the form. You may also write a letter containing all of the information requested in the form. Send your completed complaint form or letter by mail to U.S. Department of Agriculture; Director, Office of Adjudication; 1400 Independence Avenue, S.W.; Washington, D.C. 20250-9419; by fax to (202) 690-7442; or by email to program.intake@usda.gov.

Persons with Disabilities

If you are deaf, are hard of hearing, or have speech disabilities and you wish to file either an EEO or program complaint, please contact USDA through the Federal Relay Service at (800) 877-8339 or (800) 845-6136 (in Spanish).

If you have other disabilities and wish to file a program complaint, please see the contact information above. If you require alternative means of communication for program information (e.g., Braille, large print, audiotape, etc.), please contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

Supplemental Nutrition Assistance Program

For additional information dealing with Supplemental Nutrition Assistance Program (SNAP) issues, call either the USDA SNAP Hotline Number at (800) 221-5689, which is also in Spanish, or the State Information/Hotline Numbers (<http://directives.sc.egov.usda.gov/33085.wba>).

All Other Inquiries

For information not pertaining to civil rights, please refer to the listing of the USDA Agencies and Offices (<http://directives.sc.egov.usda.gov/33086.wba>).

Contents

How To Use This Soil Survey	iii
Preface	ix
Introduction	1
How This Survey Was Made	1
Detailed Soil Map Units	3
488047—Cartago gravelly loamy sand, 0 to 2 percent slopes	4
488048—Cartago gravelly loamy sand, 2 to 5 percent slopes	5
488091—Goodale-Cartago complex, 0 to 2 percent slopes	6
488173—Lubkin gravelly loamy sand, 5 to 15 percent slopes	8
488211—Morey family-Winnedumah-Rindge family complex, 0 to 2 percent slopes	10
488326—Winnedumah fine sandy loam, 0 to 2 percent slopes	13
Use and Management of the Soils	15
Interpretive Ratings	15
Rating Class Terms	15
Numerical Ratings	15
Land Capability Classification	16
Prime and Other Important Farmland	17
Hydric Soils	18
Range Sites and Ecological Sites	19
Landscape, Parent Material, and Range/Ecological Site	21
Rangeland	21
Land Management	22
Recreation	24
Engineering	25
Dwellings and Small Commercial Buildings	26
Roads and Streets, Shallow Excavations, and Landscaping	27
Sewage Disposal	28
Source of Gravel and Sand	29
Source of Reclamation Material, Roadfill, and Topsoil	29
Ponds and Embankments	30
Soil Properties	33
Engineering Properties	33
Physical Soil Properties	34
Erosion Properties	35
Total Soil Carbon	36
Chemical Soil Properties	37
Water Features	37
Soil Features	39
Formation and Classification of the Soils	41
Factors of Soil Formation	41
Classification of the Soils	49
References	53
Glossary	55

Tables	65
Table 1.—Soil Legend	66
Table 2.—Land Capability Classification	67
Table 3.—Prime and Other Important Farmland	67
Table 4.—Hydric Soils	68
Table 5.—Range/Ecological Sites	69
Table 6.—Landform, Parent Material, and Range/Ecological Site	70
Table 7.—Rangeland Productivity	71
Table 8.—Land Management, Part I (Planting)	72
Table 8.—Land Management, Part II (Hazard of Erosion and Suitability for Roads)	73
Table 8.—Land Management, Part III (Site Preparation)	74
Table 8.—Land Management, Part IV (Site Restoration)	75
Table 9.—Recreation, Part I (Camp and Picnic Areas)	76
Table 9.—Recreation, Part II (Trail Management)	77
Table 10.—Dwellings and Small Commercial Buildings	78
Table 11.—Roads and Streets, Shallow Excavations, and Landscaping	79
Table 12.—Sewage Disposal	80
Table 13.—Source of Gravel and Sand	81
Table 14.—Source of Reclamation Material, Roadfill, and Topsoil	82
Table 15.—Ponds and Embankments	83
Table 16.—Engineering Properties	84
Table 17.—Physical Soil Properties	86
Table 18.—Erosion Properties	87
Table 19.—Total Soil Carbon	88
Table 20.—Chemical Soil Properties	89
Table 21.—Water Features	90
Table 22.—Soil Features	93
Table 23.—Taxonomic Classification of the Soils	94
Table 24.—Soil Classification Key	95
Appendix	97

Issued 2013

Preface

This soil survey was developed in conjunction with the National Park Service's Soil Inventory and Monitoring Program and is intended to serve as the official source document for soils occurring within Manzanar National Historic Site.

This soil survey contains information that affects current and future land use planning in the park. It contains predictions of soil behavior for selected land uses. The survey highlights soil limitations, actions needed to overcome the limitations, and the impact of selected land uses on the environment. It is designed to meet the needs of the National Park Service and its partners to better understand the properties of the soils in the park and the effects of these soil properties on various natural ecological characteristics. This knowledge can help the National Park Service and its partners to understand, protect, and enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. The information in this report is intended to identify soil properties that are used in making various land use or land treatment decisions. Statements made in this report are intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are shallow to bedrock. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

These and many other soil properties that affect land use are described in this soil survey. The location of each map unit is shown on the detailed soil map. Each soil in the survey area is described, and information on specific uses is given. Help in using this publication and additional information are available at the local office of the Natural Resources Conservation Service or the park office for Manzanar National Historic Site.

Soil Survey of Manzanar National Historic Site, California

United States Department of Agriculture, Natural Resources Conservation Service, and United States Department of the Interior, National Park Service

How This Survey Was Made

This soil survey document was prepared by the National Park Service's Soil Inventory and Monitoring Program to provide information about the soils and miscellaneous areas within Manzanar National Historic Site. The document was written and reviewed by the Natural Resources Conservation Service staff.

The data for Manzanar National Historic Site was taken from SSURGO data certified on March 2, 2010. Six map units comprise the soil legend for the park. The original, larger soil survey of Benton-Owens Valley Area, Parts of Inyo and Mono Counties, California (CA802) was initiated in 1984. Fieldwork for the project commenced shortly thereafter and continued until 1987. Final correlation for CA802 was completed in May 1996.

During the soil survey, range sites and soil component relationships were observed. Soil-range site correlation concepts were established to help in designing the map units. Soil and plant specialists tested the concepts during mapping and collected field documentation at numerous points across the landscape. Descriptions of the range sites in the park are given in the Appendix. They will be rewritten as ecological sites at some time in the future.

The information in this report includes a description of the soils and miscellaneous areas and their location and a discussion of their suitability, limitations, and management for specified uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of native plants; and the kinds of bedrock. They dug many holes to study the soil profile, which is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

The soils and miscellaneous areas in the survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by

Soil Survey of Manzanar National Historic Site, California

an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units).

Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date. Soil scientists observed in this park that hydrology had been altered in some areas, resulting in soils that were much drier than the conditions under which they formed.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they delineated the boundaries of these bodies on digital imagery and identified each as a specific map unit.

Detailed Soil Map Units

The map units delineated on the detailed soil map in this survey represent the soils. The map unit descriptions in this section, along with the map, can be used to determine the suitability and potential of a unit for specific uses. They also can be used to plan the management needed for those uses.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. The contrasting components are mentioned in the map unit descriptions. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives the principal hazards and limitations to be considered in planning for specific uses.

Soils that have profiles that are almost alike make up a *soil series*. All the soils of a series have major horizons that are similar in composition, thickness, and arrangement. The soils of a given series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The

name of a soil phase commonly indicates a feature that affects use or management. For example, Cartago gravelly loamy sand, 0 to 2 percent slopes, is a phase of the Cartago series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes. A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Goodale-Cartago complex, 0 to 2 percent slopes, is an example.

Table 1 lists each map unit in the park, its major and minor components, and the percentage of each component in the unit. Map unit composition may not total 100 percent. Other tables give properties of the soils and the limitations, capabilities, and potentials for many uses. The Glossary defines many of the terms used in describing the soils or miscellaneous areas.

488047—Cartago gravelly loamy sand, 0 to 2 percent slopes

Map Unit Setting

Major land resource area (MLRA): 29—Southern Nevada Basin and Range

Elevation: 3,795 to 4,395 feet

Mean annual precipitation: 6 to 8 inches

Mean annual air temperature: 57 to 61 degrees F

Frost-free period: 150 to 200 days

Map Unit Composition

Cartago and similar soils: 85 percent

Dissimilar minor components: 1 percent

Description of Cartago Soil

Classification

Soil taxonomic classification: Sandy, mixed, thermic Xeric Torriorthents

Range site name/ecological site name and identification: Sandy 5-8" P.Z.

(R029XG011CA)

Setting

Landform: Alluvial fans

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Tread

Slope range: 0 to 2 percent

Down-slope shape: Linear

Across-slope shape: Linear

Representative aspect: East

Aspect range: North to southeast (clockwise)

Soil temperature class: Thermic

Soil temperature regime: Thermic

Soil moisture class: Aridic (torric)

Properties and Qualities

Runoff: Negligible

Parent material: Alluvium derived from granite

Restrictive feature(s): None within a depth of 60 inches

Frequency of flooding: Rare

Frequency of ponding: None

Depth to water table: More than 72 inches
Drainage class: Somewhat excessively drained
Shrink-swell potential: Low (about 1.5 LEP)
Salinity maximum: Not saline
Sodicity maximum: Not sodic
Calcium carbonate equivalent (maximum weight percentage): 0
Available water capacity: Low (about 4.3 inches)

Interpretive Groups

Land capability subclass: 6e-4 (nonirrigated); 3s-1 (irrigated)
Meets hydric soil criteria: No
Hydrologic soil group: A

Vegetation

Existing plants: Fourwing saltbush, Nevada jointfir, winterfat, spiny hopsage, Indian ricegrass, Fremont's dalea, and bottlebrush squirreltail

Typical Profile

0 to 14 inches; gravelly loamy sand
14 to 33 inches; gravelly loamy sand and very gravelly loamy sand
33 to 53 inches; gravelly sandy loam
53 to 60 inches; loamy fine sand

Minor Components

Unnamed soils

Percent of map unit: 1 percent
Landform: Drainageways
Representative aspect: North
Meets hydric soil criteria: Yes

488048—Cartago gravelly loamy sand, 2 to 5 percent slopes

Map Unit Setting

Major land resource area (MLRA): 29—Southern Nevada Basin and Range
Elevation: 3,795 to 4,395 feet
Mean annual precipitation: 6 to 8 inches
Mean annual air temperature: 57 to 61 degrees F
Frost-free period: 150 to 200 days

Map Unit Composition

Cartago and similar soils: 85 percent
Dissimilar minor components: 1 percent

Description of Cartago Soil

Classification

Soil taxonomic classification: Sandy, mixed, thermic Xeric Torriorthents
Range site name/ecological site name and identification: Gravelly Loamy Sand 5-8"
P.Z. (R029XG018CA)

Setting

Landform: Alluvial fans
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Tread
Slope range: 2 to 5 percent

Soil Survey of Manzanar National Historic Site, California

Down-slope shape: Linear
Across-slope shape: Linear
Representative aspect: Northeast
Aspect range: North to east (clockwise)
Soil temperature class: Thermic
Soil temperature regime: Thermic
Soil moisture class: Aridic (torric)

Properties and Qualities

Runoff: Negligible
Parent material: Alluvium derived from granite
Restrictive feature(s): None within a depth of 60 inches
Frequency of flooding: Rare
Frequency of ponding: None
Depth to water table: More than 72 inches
Drainage class: Somewhat excessively drained
Shrink-swell potential: Low (about 1.5 LEP)
Salinity maximum: Not saline
Sodicity maximum: Not sodic
Calcium carbonate equivalent (maximum weight percentage): 0
Available water capacity: Low (about 4.3 inches)

Interpretive Groups

Land capability subclass: 6e-4 (nonirrigated); 3s-1 (irrigated)
Meets hydric soil criteria: No
Hydrologic soil group: A

Vegetation

Existing plants: White bursage, bud sagebrush, shadscale, cattle saltbush, winterfat, and desert needlegrass

Typical Profile

0 to 14 inches; gravelly loamy sand
14 to 33 inches; gravelly loamy sand and very gravelly loamy sand
33 to 53 inches; gravelly sandy loam
53 to 60 inches; loamy fine sand

Minor Components

Unnamed soils

Percent of map unit: 1 percent
Landform: Drainageways
Representative aspect: North
Meets hydric soil criteria: Yes

488091—Goodale-Cartago complex, 0 to 2 percent slopes

Map Unit Setting

Major land resource area (MLRA): 29—Southern Nevada Basin and Range
Elevation: 3,700 to 4,500 feet
Mean annual precipitation: 4 to 8 inches
Mean annual air temperature: 57 to 61 degrees F
Frost-free period: 180 to 200 days

Map Unit Composition

Goodale and similar soils: 60 percent

Soil Survey of Manzanar National Historic Site, California

Cartago and similar soils: 25 percent
Dissimilar minor components: 1 percent

Description of Goodale Soil

Classification

Soil taxonomic classification: Sandy-skeletal, mixed, thermic Xeric Torriorthents
Range site name/ecological site name and identification: Gravelly Loamy Sand 5-8"
P.Z. (R029XG018CA)

Setting

Landform: Fan terraces and alluvial fans
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Tread
Slope range: 0 to 2 percent
Down-slope shape: Linear
Across-slope shape: Linear
Representative aspect: Northeast
Aspect range: North to east (clockwise)
Soil temperature class: Thermic
Soil temperature regime: Thermic
Soil moisture class: Aridic (torric)

Properties and Qualities

Runoff: Negligible
Parent material: Alluvium derived from granite
Restrictive feature(s): None within a depth of 60 inches
Frequency of flooding: Rare
Frequency of ponding: None
Depth to water table: More than 72 inches
Drainage class: Somewhat excessively drained
Shrink-swell potential: Low (about 1.5 LEP)
Salinity maximum: Not saline
Sodicity maximum: Not sodic
Calcium carbonate equivalent (maximum weight percentage): 0
Available water capacity: Very low (about 1.6 inches)

Interpretive Groups

Land capability subclass: 7s-1 (nonirrigated)
Meets hydric soil criteria: No
Hydrologic soil group: A

Vegetation

Existing plants: White bursage, bud sagebrush, shadscale, cattle saltbush, winterfat, and desert needlegrass

Typical Profile

0 to 2 inches; very gravelly loamy sand
2 to 16 inches; gravelly loamy sand and very gravelly loamy sand
16 to 22 inches; extremely gravelly loamy sand
22 to 60 inches; extremely stony loamy sand

Description of Cartago Soil

Classification

Soil taxonomic classification: Sandy, mixed, thermic Xeric Torriorthents
Range site name/ecological site name and identification: Gravelly Loamy Sand 5-8"
P.Z. (R029XG018CA)

Setting

Landform: Fan terraces and alluvial fans
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Tread
Slope range: 0 to 2 percent
Down-slope shape: Linear
Across-slope shape: Linear
Representative aspect: Northeast
Aspect range: North to east (clockwise)
Soil temperature class: Thermic
Soil temperature regime: Thermic
Soil moisture class: Aridic (torric)

Properties and Qualities

Runoff: Negligible
Parent material: Alluvium derived from granite
Restrictive feature(s): None within a depth of 60 inches
Frequency of flooding: Rare
Frequency of ponding: None
Depth to water table: More than 72 inches
Drainage class: Somewhat excessively drained
Shrink-swell potential: Low (about 1.5 LEP)
Salinity maximum: Not saline
Sodicity maximum: Not sodic
Calcium carbonate equivalent (maximum weight percentage): 0
Available water capacity: Very low (about 3.0 inches)

Interpretive Groups

Land capability subclass: 7e-4 (nonirrigated)
Meets hydric soil criteria: No
Hydrologic soil group: A

Vegetation

Existing plants: White bursage, bud sagebrush, shadscale, cattle saltbush, winterfat, and desert needlegrass

Typical Profile

0 to 10 inches; gravelly loamy sand
10 to 60 inches; cobbly loamy sand and gravelly loamy sand

Minor Components

Unnamed soils

Percent of map unit: 1 percent
Landform: Drainageways
Representative aspect: North
Meets hydric soil criteria: Yes

488173—Lubkin gravelly loamy sand, 5 to 15 percent slopes

Map Unit Setting

Major land resource area (MLRA): 29—Southern Nevada Basin and Range
Elevation: 3,795 to 5,400 feet
Mean annual precipitation: 4 to 10 inches

Soil Survey of Manzanar National Historic Site, California

Mean annual air temperature: 55 to 61 degrees F

Frost-free period: 150 to 200 days

Map Unit Composition

Lubkin and similar soils: 85 percent

Dissimilar minor components: 2 percent

Description of Lubkin Soil

Classification

Soil taxonomic classification: Loamy-skeletal, mixed, thermic Xeric Haplargids

Range site name/ecological site name and identification: Gravelly Loamy Sand 5-8"
P.Z. (R029XG018CA)

Setting

Landform: Fan terraces

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Tread

Slope range: 5 to 15 percent

Down-slope shape: Linear

Across-slope shape: Linear

Representative aspect: Northeast

Aspect range: North to east (clockwise)

Soil temperature class: Thermic

Soil temperature regime: Thermic

Soil moisture class: Aridic (torric)

Properties and Qualities

Runoff: Low

Parent material: Alluvium derived from granite

Restrictive feature(s): None within a depth of 60 inches

Frequency of flooding: Rare

Frequency of ponding: None

Depth to water table: More than 72 inches

Drainage class: Well drained

Shrink-swell potential: Low (about 1.5 LEP)

Salinity maximum: Not saline

Sodicity maximum: Not sodic

Calcium carbonate equivalent (maximum weight percentage): 0

Available water capacity: Low (about 3.1 inches)

Interpretive Groups

Land capability subclass: 6e-7 (nonirrigated)

Meets hydric soil criteria: No

Hydrologic soil group: A

Vegetation

Existing plants: White bursage, bud sagebrush, shadscale, cattle saltbush, winterfat, and desert needlegrass

Typical Profile

0 to 5 inches; gravelly loamy sand

5 to 26 inches; very stony sandy loam

26 to 46 inches; very cobbly loamy sand

46 to 60 inches; very gravelly loamy sand

Minor Components

Unnamed soils

Percent of map unit: 2 percent
Landform: Drainageways
Representative aspect: North
Meets hydric soil criteria: Yes

488211—Morey family-Winnedumah-Rindge family complex, 0 to 2 percent slopes

Map Unit Setting

Major land resource area (MLRA): 29—Southern Nevada Basin and Range
Elevation: 3,795 to 3,995 feet
Mean annual precipitation: 5 to 6 inches
Mean annual air temperature: 57 to 61 degrees F
Frost-free period: 180 to 225 days

Map Unit Composition

Morey family and similar soils: 40 percent
Winnedumah and similar soils: 25 percent
Rindge family and similar soils: 20 percent

Description of Morey Family Soil

Classification

Soil taxonomic classification: Fine-silty, mixed, thermic Typic Argiaquolls
Range site name/ecological site name and identification: Saline Meadow (R029XG002CA)

Setting

Landform: Stream terraces and alluvial fans
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Tread
Slope range: 0 to 2 percent
Down-slope shape: Linear
Across-slope shape: Linear
Representative aspect: Northeast
Aspect range: North to east (clockwise)
Soil temperature class: Thermic
Soil temperature regime: Thermic
Soil moisture class: Aquic

Properties and Qualities

Runoff: Medium
Parent material: Organic material and/or alluvium derived from mixed sources
Restrictive feature(s): None within a depth of 60 inches
Frequency of flooding: None
Frequency of ponding: None
Depth to water table: More than 72 inches
Drainage class: Well drained
Shrink-swell potential: Moderate (about 4.5 LEP)
Salinity maximum: Very slightly saline (about 3.0 mmhos/cm)
Sodicity maximum: Sodium adsorption ratio of 3.0

Soil Survey of Manzanar National Historic Site, California

Calcium carbonate equivalent (maximum weight percentage): 2

Available water capacity: High (about 10.8 inches)

Interpretive Groups

Land capability subclass: 7e-5 (nonirrigated); 2s-5 (irrigated)

Meets hydric soil criteria: Yes

Hydrologic soil group: C

Vegetation

Existing plants: Cattle saltbush, Torrey's saltbush, inland saltgrass, Baltic rush, and alkali sacaton

Typical Profile

0 to 5 inches; silt loam

5 to 18 inches; clay loam

18 to 48 inches; clay loam

48 to 60 inches; sandy clay loam

Description of Winnedumah Soil

Classification

Soil taxonomic classification: Fine-loamy, mixed, thermic Xeric Haplargids

*Range site name/ecological site name and identification: Sodid Fan 5-8" P.Z.
(R029XG005CA)*

Setting

Landform: Stream terraces and fan terraces

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Tread

Slope range: 0 to 2 percent

Down-slope shape: Linear

Across-slope shape: Linear

Representative aspect: Northeast

Aspect range: North to east (clockwise)

Soil temperature class: Thermic

Soil temperature regime: Thermic

Soil moisture class: Aridic (torric)

Properties and Qualities

Runoff: Medium

Parent material: Alluvium derived from mixed sources

Restrictive feature(s): None within a depth of 60 inches

Frequency of flooding: None

Frequency of ponding: None

Depth to water table: More than 72 inches

Drainage class: Well drained

Shrink-swell potential: Moderate (about 4.5 LEP)

Salinity maximum: Very slightly saline (about 3.0 mmhos/cm)

Sodicity maximum: Sodium adsorption ratio of 7.0

Calcium carbonate equivalent (maximum weight percentage): 3

Available water capacity: High (about 9.5 inches)

Interpretive Groups

Land capability subclass: 7e-6 (nonirrigated); 3e-3 (irrigated)

Meets hydric soil criteria: No

Hydrologic soil group: B

Vegetation

Existing plants: Shadscale, Torrey's saltbush, inland saltgrass, black greasewood, and alkali sacaton

Typical Profile

0 to 3 inches; loam
3 to 44 inches; loam
44 to 60 inches; sandy clay loam

Description of Rindge Family Soil

Classification

Soil taxonomic classification: Euic, thermic Typic Medisaprists
Range site name/ecological site name and identification: Wetland (R029XG003CA)

Setting

Landform: Stream terraces and fan terraces
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Tread
Slope range: 0 to 2 percent
Down-slope shape: Linear
Across-slope shape: Linear
Representative aspect: Northeast
Aspect range: North to east (clockwise)
Soil temperature class: Thermic
Soil temperature regime: Thermic
Soil moisture class: Aquic

Properties and Qualities

Runoff: Low
Parent material: Organic material over alluvium derived from mixed sources
Restrictive feature(s): None within a depth of 60 inches
Frequency of flooding: None
Frequency of ponding: None
Depth to water table: About 18 to 36 inches (see table 21)
Drainage class: Poorly drained
Shrink-swell potential: Moderate (about 4.5 LEP)
Salinity maximum: Very slightly saline (about 3.0 mmhos/cm)
Sodicity maximum: Sodium adsorption ratio of 7.0
Calcium carbonate equivalent (maximum weight percentage): 3
Available water capacity: High (about 10.0 inches)

Interpretive Groups

Land capability subclass: 7w-2 (nonirrigated); 3w-2 (irrigated)
Meets hydric soil criteria: Yes
Hydrologic soil group: C

Vegetation

Existing plants: Spikerush, common reed, bulrush, and cattail

Typical Profile

0 to 34 inches; mucky loam
34 to 55 inches; loam
55 to 60 inches; sandy clay loam

488326—Winnedumah fine sandy loam, 0 to 2 percent slopes

Map Unit Setting

Major land resource area (MLRA): 29—Southern Nevada Basin and Range

Elevation: 3,670 to 3,900 feet

Mean annual precipitation: 4 to 6 inches

Mean annual air temperature: 57 to 61 degrees F

Frost-free period: 180 to 220 days

Map Unit Composition

Winnedumah and similar soils: 85 percent

Dissimilar minor components: 6 percent

Description of Winnedumah Soil

Classification

Soil taxonomic classification: Fine-loamy, mixed, thermic Xeric Haplargids

Range site name/ecological site name and identification: Sodic Fan 5-8" P.Z.
(R029XG005CA)

Setting

Landform: Stream terraces and lakebeds (relict)

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Tread and talf

Slope range: 0 to 2 percent

Down-slope shape: Linear

Across-slope shape: Linear

Representative aspect: Northeast

Aspect range: North to east (clockwise)

Soil temperature class: Thermic

Soil temperature regime: Thermic

Soil moisture class: Aridic (torric)

Properties and Qualities

Runoff: Low

Parent material: Alluvium derived from mixed and/or lacustrine deposits

Restrictive feature(s): None within a depth of 60 inches

Frequency of flooding: None

Frequency of ponding: None

Depth to water table: About 48 to 60 inches (see table 21)

Drainage class: Somewhat poorly drained

Shrink-swell potential: Moderate (about 4.5 LEP)

Salinity maximum: Moderately saline (about 12.0 mmhos/cm)

Sodicity maximum: Sodium adsorption ratio of 57.0

Calcium carbonate equivalent (maximum weight percentage): 3

Available water capacity: Moderate (about 6.8 inches)

Interpretive Groups

Land capability subclass: 7s-6 (nonirrigated)

Meets hydric soil criteria: No

Hydrologic soil group: B

Vegetation

Existing plants: Shadscale, Torrey's saltbush, inland saltgrass, black greasewood, and alkali sacaton

Typical Profile

0 to 14 inches; fine sandy loam

14 to 29 inches; fine sandy loam and loam

29 to 38 inches; sandy clay loam

38 to 60 inches; fine sandy loam

Minor Components

Morey family soils

Percent of map unit: 3 percent

Landform: Alluvial fans

Representative aspect: North

Meets hydric soil criteria: Yes

Rindge family soils

Percent of map unit: 3 percent

Landform: Alluvial fans

Representative aspect: North

Meets hydric soil criteria: Yes

Use and Management of the Soils

This soil survey is an inventory and evaluation of the soils in Manzanar National Historic Site. It can be used to adjust land uses to the limitations and potentials of natural resources and the environment. Also, it can help to prevent soil-related failures in land uses.

In preparing a soil survey, soil scientists, conservationists, engineers, and others collect extensive field data about the nature and behavioral characteristics of the soils. They collect data on erosion, droughtiness, flooding, and other factors that affect various soil uses and management. Field experience and collected data on soil properties and performance are used as a basis in predicting soil behavior.

Information in this section can be used to plan the use and management of soils as rangeland and as sites for buildings, sanitary facilities, highways and other transportation systems, and recreational facilities. It can be used to identify the potentials and limitations of each soil for specific land uses and to help prevent construction failures caused by unfavorable soil properties.

Planners and others using soil survey information can evaluate the effect of specific land uses on productivity and on the environment in all or part of the park. The survey can help planners to maintain or create a land use pattern in harmony with the natural soil.

Contractors can use this survey to locate sources of sand and gravel, roadfill, and topsoil. They can use it to identify areas where bedrock, wetness, or very firm soil layers can cause difficulty in excavation.

Health officials, highway officials, engineers, and others may also find this survey useful. The survey can help them plan the safe disposal of wastes and locate sites for pavements, sidewalks, campgrounds, playgrounds, and trees and shrubs.

Interpretive Ratings

The interpretive tables in this survey rate the soils in the park for various uses. Many of the tables identify the limitations that affect specified uses and indicate the severity of those limitations. The ratings in these tables are both verbal and numerical.

Rating Class Terms

Rating classes are expressed in the tables in terms that indicate the extent to which the soils are limited by all of the soil features that affect a specified use or in terms that indicate the suitability of the soils for the use. Thus, the tables may show limitation classes or suitability classes. Terms for the limitation classes are *not limited*, *slightly limited*, *somewhat limited*, and *very limited*. The suitability ratings are expressed as *well suited*, *moderately well suited*, *poorly suited*, and *unsuited* or as *good*, *fair*, and *poor*.

Numerical Ratings

Numerical ratings in the tables indicate the relative severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate

gradations between the point at which a soil feature has the greatest negative impact on the use and the point at which the soil feature is not a limitation. The limitations appear in order from the most limiting to the least limiting. Thus, if more than one limitation is identified, the most severe limitation is listed first and the least severe one is listed last.

Land Capability Classification

Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels—capability class, subclass, and unit (USDA-SCS, 1961).

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

Class 1 soils have slight limitations that restrict their use.

Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.

Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.

Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.

Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral, for example, 2*e*. The letter *e* shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and *c*, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by *w*, *s*, or *c* because the soils in class 5 are subject to little or no erosion. They have other limitations that restrict their use to pasture, rangeland, forestland, wildlife habitat, or recreation.

The capability classification of map units in this park is given in the section “Detailed Soil Map Units” and in table 2.

Prime and Other Important Farmland

Table 3 lists the map units in the park that are considered important farmlands. Important farmlands consist of prime farmland, unique farmland, and farmland of statewide or local importance. This list does not constitute a recommendation for a particular land use.

In an effort to identify the extent and location of important farmlands, the Natural Resources Conservation Service, in cooperation with other interested Federal, State, and local government organizations, has inventoried land that can be used for the production of the Nation's food supply.

Prime farmland is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil quality, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. The water supply is dependable and of adequate quality. Prime farmland is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

For some of the soils identified in the table as prime farmland, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures.

A recent trend in land use in some areas has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

Unique farmland is land other than prime farmland that is used for the production of specific high-value food and fiber crops, such as citrus, tree nuts, olives, cranberries, and other fruits and vegetables. It has the special combination of soil quality, growing season, moisture supply, temperature, humidity, air drainage, elevation, and aspect needed for the soil to economically produce sustainable high yields of these crops when properly managed. The water supply is dependable and of adequate quality. Nearness to markets is an additional consideration. Unique farmland is not based on national criteria. It commonly is in areas where there is a special microclimate, such as the wine country in California.

In some areas, land that does not meet the criteria for prime or unique farmland is considered to be *farmland of statewide importance* for the production of food, feed, fiber, forage, and oilseed crops. The criteria for defining and delineating farmland of statewide importance are determined by the appropriate State agencies. Generally, this land includes areas of soils that nearly meet the requirements for prime farmland and that economically produce high yields of crops when treated and managed according to acceptable farming methods. Some areas may produce as high a yield

as prime farmland if conditions are favorable. Farmland of statewide importance may include tracts of land that have been designated for agriculture by State law.

In some areas that are not identified as having national or statewide importance, land is considered to be *farmland of local importance* for the production of food, feed, fiber, forage, and oilseed crops. This farmland is identified by the appropriate local agencies. Farmland of local importance may include tracts of land that have been designated for agriculture by local ordinance.

Hydric Soils

Table 4 lists the map unit components that are rated as hydric soils in the park. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; USDA-NRCS, 2010).

The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for all of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils, under natural conditions, are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 2002). These criteria are used to identify map unit components that normally are associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (Soil Survey Staff, 1999) and "Keys to Soil Taxonomy" (Soil Survey Staff, 2010) and in the "Soil Survey Manual" (Soil Survey Division Staff, 1993).

If soils are wet enough for a long enough period of time to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils are specified in "Field Indicators of Hydric Soils in the United States" (USDA-NRCS, 2010).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units that are dominantly made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units dominantly made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

The criteria for hydric soils are represented by codes in the table (for example, 2B3). Definitions for the codes are as follows:

1. All Histels except for Folistels and Histosols except for Folist.
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Historthels great group, Histoturbels great group, Pachic subgroups, or Cumulic subgroups that:
 - A. are somewhat poorly drained and have a water table at the surface (0.0 feet) during the growing season, or
 - B. are poorly drained or very poorly drained and have either:
 - 1) a water table at the surface (0.0 feet) during the growing season if textures are coarse sand, sand, or fine sand in all layers within a depth of 20 inches, or
 - 2) a water table at a depth of 0.5 foot or less during the growing season if saturated hydraulic conductivity (K_{sat}) is equal to or greater than 6.0 in/hr in all layers within a depth of 20 inches, or
 - 3) a water table at a depth of 1.0 foot or less during the growing season if saturated hydraulic conductivity (K_{sat}) is less than 6.0 in/hr in any layer within a depth of 20 inches.
3. Soils that are frequently ponded for periods of long or very long duration during the growing season.
4. Soils that are frequently flooded for periods of long or very long duration during the growing season.

Range Sites and Ecological Sites

Plant communities are largely dependent on the soil, climate, topography, aspect, and slope of the landscape, as well as other abiotic features. To better understand these soil-plant interactions and the effects of selected management practices, the Natural Resources Conservation Service classifies forestlands and rangelands into ecological sites. While the vegetation information for Manzanar National Historic Site is included in this report, true ecological sites have not been developed for this park. Included are traditional range site descriptions. An explanation of ecological site is provided below for future reference as the site names in this document will remain the same for the future ecological sites.

Landscapes of native vegetation are divided into ecological sites for the purposes of inventory, evaluation, and management. An ecological site, as defined for rangeland, is a distinctive kind of land with specific physical characteristics that differs from other kinds of land in its ability to produce a distinctive kind and amount of vegetation.

An ecological site is the product of all the environmental factors responsible for its development, including parent material, landscape, climate, soils, living organisms, hydrology, fire, and time in place. Ecological site descriptions contain information on each of these environmental factors. Included are brief descriptions of: a) physiographic and climatic features; b) major identifiable plant community types that may occupy the site, including the reference plant community; c) total annual production; d) ecological dynamics of the plant communities; e) soils and their main properties; and f) site interpretations and general management considerations for wildlife, hydrology, recreation, fire, esthetics, and restoration or revegetation.

The reference plant community for a site has evolved under natural ecological processes and disturbances and is considered to be at the highest natural site potential under the current climate. It has developed on the site as a result of all site-forming factors and is best adapted to the unique combination of environmental factors associated with the site. Natural disturbances, such as fire, drought, herbivory, and flooding, were inherent in the development and maintenance of the reference plant community. Plant communities that have been subject to anthropogenic disturbances or physical site deterioration or have been protected from the natural disturbances do not typify the reference state and may exist in a stable or steady state that is different from the reference plant community.

Soil Survey of Manzanar National Historic Site, California

The reference plant community of an ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, the productivity and occurrence of individual species vary. Special boundaries of the communities can be recognized by characteristic patterns of species composition, association, and community structure. Generally, one species or group of species dominates the site and the stability within the natural dynamics or disturbances of the site allows the species to be used as the factor that distinguishes one site from another.

At times, the extent of the less frequently occurring plants may increase on a site or plants not formerly occurring in the reference community may invade the site. The presence or abundance of these plants may fluctuate greatly because of the ability of the plants to adapt to the differences in the microenvironment, weather conditions, soil alterations, or human actions. Using these species for site identification can be misleading; thus they should not be used to differentiate sites.

The following ecological site inventory methods are used in determining the characteristic plant communities of an ecological site:

1. Identification and evaluation of reference and/or relict sites with similar plant communities and associated soils.
2. Interpolation and extrapolation of plant, soil, and climatic data from existing historic reference areas along a continuum to other points on that continuum for which no suitable reference community is available.
3. Evaluation and comparison of the same ecological site that occurs in different areas but that has experienced different levels of disturbance and management. Further comparison is made with areas that are not disturbed.
4. Evaluation and interpretation of research data dealing with the ecology, management, and soils in areas of the plant communities.
5. Review of historical accounts, survey and military records, and botanical literature of the areas.

The initial description of the reference state should be considered an approximation subject to modification as additional knowledge is gained or discovered.

Plant communities change along environmental gradients. When changes in soils, aspect, topography, or moisture conditions are abrupt, the plant community boundaries will be reasonably distinct. Boundaries are less distinct where the plant communities change gradually over wide environmental gradients of relatively uniform soils and topography. Thus, the need for site differentiation may not be readily apparent until the cumulative impact of soil, topography, hydrology, or climate is examined over a broad area. Frequently, such differences are reflected first in production and second in the kinds and proportions of a plant species making up the core of the plant community. In some cases, the boundaries that are drawn between ecological sites along a continuum of closely related soils and a gradually changing climate are somewhat arbitrary.

The following criteria are used to differentiate one ecological site from another:

1. Significant differences in the species or species groups that are in the characteristic plant community.
2. Significant differences in the relative proportion of species or species groups in the characteristic plant community.
3. Significant differences in the total annual production or site index of the characteristic plant community.
4. Soil factors that determine plant production and composition, the hydrology of the site, and the functioning of the ecological process of the water cycle, mineral cycles, and energy flow.

Differences in kind, proportion, and production of plants are the result of differences in soil, topography, climate, and other environmental factors. Slight variations in these factors are not criteria for site differentiation. Individual environmental factors

are frequently associated with significant differences in reference plant communities. For differentiation into a distinct site to occur, the differences in the environmental factors must be great enough to affect the kinds, amounts, and proportions of the plant community.

Forestland is a spatially defined site where the reference community has at least 25 percent canopy cover. The reference community is the present-day climax community that most resembles the forest conditions prior to European contact. It developed with natural disturbances such as drought, fire, and insects. Several other plant communities may be present during the seral stages of development. Vegetation on forestland provides many habitat components, assists in controlling soil erosion, is suitable for grazing or browsing by wildlife, and offers scenic and recreational opportunities. Forestland is environmentally and economically important. For more information about NRCS national forestry policies, see the NRCS “National Forestry Manual,” which is available online at <http://soils.usda.gov/technical/nfmanual/>.

The reference community for a rangeland ecological site does not have the potential to produce at least 25 percent canopy cover. Several other plant communities may be present during phases of development or altered conditions. Vegetation on rangeland provides many habitat components, assists in controlling soil erosion, is suitable for grazing or browsing by wildlife and domestic animals, and offers scenic and recreational opportunities. Rangeland is environmentally and economically important.

Table 5 lists the map unit symbol and each map unit component’s name and percent of map unit alongside the range/ecological site name, site type (forestland or rangeland), and range/ecological site number. Approved ecological site descriptions are available online at <http://esis.sc.egov.usda.gov/>. These descriptions are dynamic documents that are constantly updated as new research and data is gained; thus, the online version will be the most recent version of the descriptions.

Landscape, Parent Material, and Range/Ecological Site

Table 6 displays information related to the range/ecological sites that are correlated to each soil in a map unit.

Percent of the map unit is the extent of the named soil in the map unit.

Slope is the inclination of the land surface from the horizontal. Percentage of slope is the vertical distance divided by horizontal distance, then multiplied by 100. Thus, a slope of 20 percent is a drop of 20 feet in 100 feet of horizontal distance. The table shows the low and high range of slope for the named component or soil.

Elevation is the height of an object or area on the earth’s surface in reference to a fixed reference point, such as mean sea level. The typical low and high range of elevation is displayed for each soil.

MAP is the mean annual precipitation for areas of the soil in the map unit.

Landform is a specific shape of the earth in the area where a soil typically occurs. Examples are a mountain summit and a valley bottom.

Parent material is the material in which soils formed. Examples are the underlying geological material (including bedrock), a surficial deposit (such as volcanic ash), and organic material. Soils inherit their chemical and physical properties from the parent material.

Range/ecological site name and number is the range/ecological site name and unique reference number that are correlated to the named soil in the map unit.

Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management is based on the relationship between the soils and vegetation and water.

Table 7 shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, and unfavorable years. An explanation of the column headings in table 7 follows.

A *range/ecological site* is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on a site is typified by an association of species that differs from that of other sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the "Field Office Technical Guide," which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, normal, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the "National Range and Pasture Handbook," which is available in local offices of the Natural Resources Conservation Service.

The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

Land Management

In table 8, parts I through IV, interpretive ratings are given for various aspects of land management. The ratings are both verbal and numerical.

Some rating class terms indicate the degree to which the soils are suited to a specified land management practice. *Well suited* indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. *Moderately suited* indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable, and fair performance can be expected.

Some maintenance is needed. *Poorly suited* indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. *Unsuited* indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the specified land management practice (1.00) and the point at which the soil feature is not a limitation (0.00).

Rating class terms for *fire damage* and *seedling mortality* are expressed as low, moderate, and high. Where these terms are used, the numerical ratings indicate gradations between the point at which the potential for fire damage or seedling mortality is highest (1.00) and the point at which the potential is lowest (0.00).

Rating class terms for *hazard of erosion* are expressed as slight, moderate, severe, and very severe. Where these terms are used, the numerical ratings indicate gradations between the point at which the potential for erosion is highest (1.00) and the point at which the potential is lowest (0.00).

The paragraphs that follow indicate the soil properties considered in rating the soils for land management practices.

Ratings in the columns *suitability for hand planting* and *suitability for mechanical planting* are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately suited, poorly suited, or unsuited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column *soil rutting hazard* are based on depth to a water table, rock fragments on or below the surface, the Unified classification, depth to a restrictive layer, and slope. Ruts form as a result of the operation of planting equipment. The hazard is described as slight, moderate, or severe. A rating of *slight* indicates that the soil is subject to little or no rutting, *moderate* indicates that rutting is likely, and *severe* indicates that ruts form readily.

Ratings in the column *hazard of erosion* are based on slope and on soil erodibility factor K. The soil loss is caused by sheet or rill erosion in areas where 50 to 75 percent of the surface has been exposed by different kinds of disturbance. The hazard is described as slight, moderate, severe, or very severe. A rating of *slight* indicates that erosion is unlikely under ordinary climatic conditions; *moderate* indicates that some erosion is likely and that erosion-control measures may be needed; *severe* indicates that erosion is very likely and that erosion-control measures, including revegetation of bare areas, are advised; and *very severe* indicates that significant erosion is expected, loss of soil productivity and off-site damage are likely, and erosion-control measures are costly and generally impractical.

Ratings in the column *hazard of erosion on roads and trails* are based on the soil erodibility factor K, slope, and content of rock fragments. The ratings apply to unsurfaced roads and trails. The hazard is described as slight, moderate, or severe. A rating of *slight* indicates that little or no erosion is likely; *moderate* indicates that some erosion is likely, that the roads or trails may require occasional maintenance, and that simple erosion-control measures are needed; and *severe* indicates that significant erosion is expected, that the roads or trails require frequent maintenance, and that costly erosion-control measures are needed.

Ratings in the column *suitability for roads (natural surface)* are based on slope, rock fragments on the surface, plasticity index, content of sand, the Unified classification, depth to a water table, ponding, flooding, and the hazard of soil slippage. The ratings

indicate the suitability for using the natural surface of the soil for roads. The soils are described as well suited, moderately suited, or poorly suited to this use.

Ratings in the column *suitability for mechanical site preparation (deep)* are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column *suitability for mechanical site preparation (surface)* are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 1 foot is considered in the ratings.

Ratings in the column *potential for damage to soil by fire* are based on texture of the surface layer, content of rock fragments and organic matter in the surface layer, thickness of the surface layer, and slope. The soils are described as having a low, moderate, or high potential for this kind of damage. The ratings indicate an evaluation of the potential impact of prescribed fires or wildfires that are intense enough to remove the duff layer and consume organic matter in the surface layer.

Ratings in the column *potential for seedling mortality* are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality.

Recreation

The soils of the park are rated in table 9, parts I and II, according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the table are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in table 9 can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas. The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Foot traffic and equestrian trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Mountain bike and off-road vehicle trails require little or no site preparation. They are not covered with surfacing material or vegetation. Considerable compaction of the soil material is likely. The ratings are based on the soil properties that influence erodibility, trafficability, dustiness, and the ease of revegetation. These properties are stoniness, depth to a water table, ponding, slope, flooding, and texture of the surface layer.

Engineering

This section provides information for planning land uses related to urban development and to water management. Soils are rated for various uses, and the most limiting features are identified. Ratings are given for building site development, landscaping, sanitary facilities, construction materials, and water management. The ratings are based on observed performance of the soils and on the estimated data and test data in the "Soil Properties" section.

Information in this section is intended for land use planning, for evaluating land use alternatives, and for planning site investigations prior to design and construction. The information, however, has limitations. For example, estimates and other data generally apply only to that part of the soil between the surface and a depth of 5 to 7 feet. Because of the map scale, small areas of different soils may be included within the mapped areas of a specific soil.

The information is not site specific and does not eliminate the need for onsite investigation of the soils or for testing and analysis by personnel experienced in the design and construction of engineering works.

Government ordinances and regulations that restrict certain land uses or impose specific design criteria were not considered in preparing the information in this section.

Local ordinances and regulations should be considered in planning, in site selection, and in design.

Soil properties, site features, and observed performance were considered in determining the ratings in this section. During the fieldwork for this soil survey, determinations were made about particle-size distribution, liquid limit, plasticity index, soil reaction, depth to bedrock, hardness of bedrock within 5 to 7 feet of the surface, soil wetness, depth to a water table, ponding, slope, likelihood of flooding, natural soil structure aggregation, and soil density. Data were collected about kinds of clay minerals, mineralogy of the sand and silt fractions, and the kinds of adsorbed cations. Estimates were made for erodibility, permeability, corrosivity, shrink-swell potential, available water capacity, and other behavioral characteristics affecting engineering uses.

This information can be used to evaluate the potential of areas for residential, commercial, and recreational uses; make preliminary estimates of construction conditions; evaluate alternative routes for roads, streets, highways, pipelines, and underground cables; evaluate alternative sites for septic tank absorption fields and sewage lagoons; plan detailed onsite investigations of soils and geology; locate potential sources of gravel, sand, earthfill, and topsoil; plan drainage systems, ponds, and other structures for soil and water conservation; and predict performance of proposed small structures and pavements by comparing the performance of existing similar structures on the same or similar soils.

The information in the tables, along with the soil map, the soil descriptions, and other data provided in this survey, can be used to make additional interpretations.

Some of the terms used in this soil survey have a special meaning in soil science and are defined in the Glossary.

Dwellings and Small Commercial Buildings

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. Table 10 shows the degree and kind of soil limitations that affect dwellings and small commercial buildings.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the

load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Roads and Streets, Shallow Excavations, and Landscaping

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. Table 11 shows the degree and kind of soil limitations that affect local roads and streets, shallow excavations, and landscaping.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the

amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Landscaping requires soils on which turf, trees, and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

Sewage Disposal

Table 12 shows the degree and kind of soil limitations that affect septic tank absorption fields and sewage lagoons. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 72 inches or between a depth of 24 inches and a restrictive layer is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Saturated hydraulic conductivity (K_{sat}), depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, saturated hydraulic conductivity (K_{sat}), depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Saturated hydraulic conductivity (K_{sat}) is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used

as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a K_{sat} rate of more than 14 micrometers per second are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

Source of Gravel and Sand

Table 13 gives information about the soils as potential sources of gravel and sand. Normal compaction, minor processing, and other standard construction practices are assumed.

Gravel and sand are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. Only the likelihood of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the bottom layer of the soil contains sand or gravel, the soil is considered a likely source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness. The ratings are for the whole soil, from the surface to a depth of about 6 feet.

The soils are rated *good*, *fair*, or *poor* as potential sources of sand and gravel. A rating of *good* or *fair* means that the source material is likely to be in or below the soil. The bottom layer and the thickest layer of the soils are assigned numerical ratings. These ratings indicate the likelihood that the layer is a source of sand or gravel. The number 0.00 indicates that the layer is a poor source. The number 1.00 indicates that the layer is a good source. A number between 0.00 and 1.00 indicates the degree to which the layer is a likely source.

Source of Reclamation Material, Roadfill, and Topsoil

Table 14 gives information about the soils as potential sources of reclamation material, roadfill, and topsoil. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated *good*, *fair*, or *poor* as potential sources of reclamation material, roadfill, and topsoil. The features that limit the soils as sources of these materials are specified in the table. Numerical ratings between 0.00 and 0.99 are given after the specified features. These numbers indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability

of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments. The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Ponds and Embankments

Table 15 gives information on the soil properties and site features that affect water management. The degree and kind of soil limitations are given for pond reservoir areas; embankments, dikes, and levees; and aquifer-fed excavated ponds. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the saturated hydraulic conductivity (K_{sat}) of the soil and the depth to

fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. Embankments that have zoned construction (core and shell) are not considered. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of 5 or 6 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects trafficability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, K_{sat} of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Soil Properties

Data relating to soil properties are collected during the course of the soil survey.

Soil properties are ascertained by field examination of the soils and by laboratory index testing of some benchmark soils. Established standard procedures are followed. During the survey, many shallow borings are made and examined to identify and classify the soils and to delineate them on the soil maps. Samples are taken from some typical profiles and tested in the laboratory to determine particle-size distribution, plasticity, and compaction characteristics.

Estimates of soil properties are based on field examinations, on laboratory tests of samples from the survey area, and on laboratory tests of samples of similar soils in nearby areas. Tests verify field observations, verify properties that cannot be estimated accurately by field observation, and help to characterize key soils.

The estimates of soil properties are shown in tables. They include engineering properties, physical and chemical properties, and pertinent soil and water features.

Engineering Properties

Table 16 gives the engineering classifications and the range of engineering properties for the layers of each soil in the park.

Depth to the upper and lower boundaries of each layer is indicated.

Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. "Loam," for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, "gravelly."

Classification of the soils is determined according to the Unified soil classification system (ASTM, 2005) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 2004).

The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection.

If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement,

the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage.

Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and *plasticity index* (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination.

Physical Soil Properties

Table 17 shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the park. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, saturated hydraulic conductivity (K_{sat}), plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earthmoving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at $1/3$ - or $1/10$ -bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute linear extensibility, shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water

and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Permeability (K_{sat}) refers to the ability of a soil to transmit water or air. The term “permeability,” as used in soil surveys, indicates saturated hydraulic conductivity (K_{sat}). The estimates in the table indicate the rate of water movement, in inches per hour, when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Shrink-swell potential is the potential for volume change in a soil with a loss or gain in moisture. Volume change occurs mainly because of the interaction of clay minerals with water and varies with the amount and type of clay minerals in the soil. The size of the load on the soil and the magnitude of the change in soil moisture content influence the amount of swelling of soils in place. Laboratory measurements of swelling of undisturbed clods were made for many soils. For others, swelling was estimated on the basis of the kind and amount of clay minerals in the soil and on the basis of measurements of similar soils.

If the shrink-swell potential is rated moderate to very high, shrinking and swelling can cause damage to buildings, roads, and other structures. Special design is often needed.

Shrink-swell potential classes are based on the change in length of an unconfined clod as moisture content is increased from air-dry to field capacity. The classes are *low*, a change of less than 3 percent; *moderate*, 3 to 6 percent; *high*, 6 to 9 percent; and *very high*, greater than 9 percent.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In this table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion Properties

Table 18 shows estimates of some erosion factors that affect a soil's potential for different uses. These estimates are given for each layer of every soil for K factors and are given as one rating for the entire soil for the T factor, the wind erodibility group, and the wind erodibility index. Values are reported for each soil in the park. Estimates are based on field observations and on test data for these and similar soils.

Erosion factors are shown in the table as the K factor (K_w and K_f) and the T factor. Soil erosion factors K_w and K_f quantify soil detachment by runoff and raindrop impact. These erosion factors are indexes used to predict the long-term average soil loss from sheet and rill erosion under crop systems and conservation techniques. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and K_{sat} . Values

of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

The procedure for determining the Kf factor is outlined in Agriculture Handbook 703, "Predicting Soil Erosion by Water: A Guide to Conservation Planning with the Revised Universal Soil Loss Equation (RUSLE)," USDA, Agricultural Research Service, 1997.

Depth to the upper and lower boundaries of each layer is indicated.

Erosion factor Kw indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments. In horizons where total rock fragments are 15 percent or more, by volume, the Kw factor is always less than the Kf factor.

Erosion factor Kf indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size. Soil horizons that do not have rock fragments are assigned equal Kw and Kf factors.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind and/or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are described in the "National Soil Survey Handbook."

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

Total Soil Carbon

Table 19 gives estimates of total soil carbon. Soil carbon occurs as organic and inorganic carbon.

Soil organic carbon (SOC) is carbon (C) in soil that originated from a biological source, such as plants, animals, or micro-organisms. SOC is found in both organic and mineral soil layers. The term "soil organic carbon" refers only to the carbon occurring in soil organic matter (SOM). Soil organic carbon makes up about one-half the weight of soil organic matter. The rest of SOM is mostly oxygen, nitrogen, and hydrogen.

Soil inorganic carbon (SIC) is carbon found in soil carbonates, typically as calcium carbonate layers in the soil or as clay-sized fractions throughout the soil. Carbonates in soils are most common in areas where evaporation rates exceed precipitation, as is the case in most desert environments. Typically, the carbonates accumulated from carbonatic dust or from solution during periods of wetter climates. Soil inorganic carbon also occurs in soils that formed in marl in all regions of the country.

The SOC and SIC contents are reported in kilograms per square meter to a depth of 2 meters or to a representative depth of either hard bedrock or a cemented horizon. The SOC and SIC values are on a whole soil basis, corrected for rock fragments.

SOC can be an indicator of overall soil fertility and soil quality that affects ecosystem function. SOM is the main reservoir for most plant nutrients, such as phosphorus and nitrogen. Managing for SOC by managing for SOM increases the content of these elements and improves soil resiliency.

Soil organic matter binds soil particles together and thus increases soil porosity and water infiltration and allows better root penetration and waterflow into the soil. Greater inflow of water reduces the hazard of erosion and the rate of surface water runoff.

Greater SOC levels improve not only soil quality but also the quality of air and water. Soil acts as a filter and improves water quality. Fertile soils that support plant life remove CO₂ from the atmosphere and increase oxygen levels through photosynthesis. Maintaining the level of soil organic carbon reduces C release into the atmosphere and thus can lessen the effects of global warming.

SIC influences the types of plants that will grow. High SIC levels are commonly associated with a higher soil pH, which limits the types of plants that will thrive.

Like SOM, soil carbonates, the source of SIC, also bind soil particles together. They fill voids in the soil and thus can reduce soil porosity. Compacted soil carbonates may restrict root penetration and waterflow into the soil.

Chemical Soil Properties

Table 20 shows estimates of some chemical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the park. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable cations that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. The ability to retain cations reduces the hazard of ground-water pollution.

Soil reaction is a measure of acidity or alkalinity. The pH of each soil horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced saturated hydraulic conductivity (K_{sat}) and aeration, and a general degradation of soil structure.

Water Features

Table 21 gives estimates of various soil water features. The estimates are used in land use planning that involves engineering considerations.

Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained

Soil Survey of Manzanar National Historic Site, California

soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

If a soil is assigned to a dual hydrologic group (A/D, B/D, or C/D), the first letter is for drained areas and the second is for undrained areas.

The *months* in the table indicate the portion of the year in which a water table, ponding, and/or flooding is most likely to be a concern.

Water table refers to a saturated zone in the soil. The water features table indicates, by month, depth to the top (*upper limit*) and base (*lower limit*) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table.

Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The table indicates *surface water depth* and the *duration* and *frequency* of ponding. Duration is expressed as *very brief* if less than 2 days, *brief* if 2 to 7 days, *long* if 7 to 30 days, and *very long* if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. *None* means that ponding is not probable; *rare* that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); *occasional* that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and *frequent* that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding is the temporary inundation of an area caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and *frequency* are estimated. Duration is expressed as *extremely brief* if 0.1 hour to 4 hours, *very brief* if 4 hours to 2 days, *brief* if 2 to 7 days, *long* if 7 to 30 days, and *very long* if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. *None* means that flooding is not probable; *very rare* that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); *rare* that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); *occasional* that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); *frequent* that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and *very frequent* that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of

flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

Soil Features

Table 22 gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A *restrictive layer* is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness of the restrictive layer, which significantly affects the ease of excavation. *Depth to top* is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, saturated hydraulic conductivity (K_{sat}), content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as *low*, *moderate*, or *high*, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as *low*, *moderate*, or *high*. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

Formation and Classification of the Soils

This section relates the soils in Manzanar National Historic Site to the major factors of soil formation and describes the system of soil classification.

Factors of Soil Formation

By Susan Burlew Southard, Natural Resources Conservation Service.

Soil covers the surface of the earth as a three-dimensional body of varying thickness and is made up of varying proportions of organic and mineral material, pore space with gases, and water. Soils differ in their appearance, productivity, and management requirements due to their chemical and physical properties. The characteristics and properties of soils are determined by physical and chemical processes that result from the interaction of five soil-forming factors. These factors of soil formation are interdependent, and few generalizations can be made regarding any one factor unless the effects of the other factors are known. The term “pedogenesis” is often used to connote the processes of soil formation.

The interacting soil-forming factors are parent material, climate, organisms, relief or topography, and time (Jenny, 1941). Parent material is the source material in which soils formed. Soils are influenced by the texture and structure of the parent material and its mineralogical and chemical composition. Climate is predominantly temperature and kind and amount of precipitation. Climate is also the seasonal distribution and range of temperature and precipitation. Organisms are the plants and other organisms living in and on the soil, including humans. Time refers to how long the soil-forming factors have been operating on a particular landscape. Relief or topography is the shape and elevation of the landscape. It affects internal and external soil properties, such as soil drainage, aeration, susceptibility to erosion, and the soil’s exposure to the sun and wind.

The processes of soil formation are a sequence of often overlapping events, involving biogeochemical reactions that are energized by climate and spatially related to relief or topography (Buol and others, 2011). The physical and chemical properties of a soil are altered by these reactions over time. The influence of any one of these factors varies among all parks and within localities of a particular park. Soils may differ significantly from place to place in a park and within very short distances as a result of complex interactions among the five soil-forming factors. In some instances parks may have vast stretches of the same type of soil because of uniform soil-forming factors.

Parent Material

The unconsolidated mass in which soils form is called parent material. Mineral soil parent material is a product of weathering of underlying bedrock in place or weathering of material that has been transported. Organic soils form in place from the accumulation and decomposition of plant material, such as wood, leaves, or aquatic plants. Weathering refers to the chemical and physical disintegration and decomposition of parent material. Few soils weather directly from the underlying rocks. More commonly, soils form in materials that have moved in from elsewhere. Soils

generally have a dominant kind of parent material but are influenced by other types of parent material. Material may have been moved only a few feet by gravity (colluvial parent material) or transported long distances by wind (eolian parent material) or water (alluvial parent material). Soils are said to have residual parent material if they formed directly from underlying rocks (residuum) or, in the case of most organic soils, from an *in situ* plant source. Soils formed in residuum may have the same general chemistry as the original rocks or plant material depending on the degree of weathering that has occurred.

Eolian parent material includes windblown sand. A considerable amount of windblown sand is around the construction areas and roads in Manzanar National Historic Site (Manzanar NHS) where the soils were disturbed. The soils in the park have a high content of sand and are subject to wind erosion and transport.

Loess, which has been blown for long distances, consists mainly of silt-sized particles. A human-induced source of loess in the vicinity of Manzanar NHS is the exposed alkaline lake bed of Owens Lake, which was drained starting in 1913 by diversions of water from the Owens River and its tributaries to southern California. The salt-rich loess travels both north and south on turbulent winds that are funneled through Owens Valley by the adjacent Sierra Nevada on the west and the White-Inyo Range and Coso Range on the east (Reheis, 2011). The alkaline loess is composed of a variety of minerals, including those derived from sodium chloride, sodium sulfate, sodium bicarbonate, and sodium carbonate. Internees at Manzanar war camp described a “fine white dust” that got in their food and water (Levine, 1995). Table 17 shows the low contents of silt and clay of the major soils in the vicinity of the park.

Alluvium has different textures, depending on whether the water moves quickly or slowly, the type of rock, and the size of the stream or river. Fast-moving water deposits gravel, rocks, and sand. Slow-moving water deposits fine textured material (clay and silt) as sediments in the water settle out. According to the county soil survey encompassing Manzanar NHS, coarse granitic alluvium is the dominant parent material on the alluvial fans of the Sierra Nevada and the northern part of the White Mountains (USDA-NRCS, 2002). The soils on the west side of the valley formed in granitic alluvium deposited in connection with glacial activity in the mountains. As the glaciers melted and retreated, stream-transported glacial outwash and debris flows created alluvial fans. The alluvial fans are composed of varying amounts of angular boulders, stones, cobbles, gravel, and soil material. Rock fragments in the soils become more rounded the further they travel. Figure 1 shows the Block 34 mess hall meditation garden. The garden appears to have rounded cobbles (foreground) from the local area. The standing, more angular and less rounded rocks were probably brought from the nearby mountains to the park by internees.

An example of soils mapped in Manzanar NHS that formed in alluvium is Cartago. Cartago soils formed from granitic alluvium where alluvial fans join the valley floor. These soils have sandy surface horizons and a gravelly subsoil. The gravel were deposited by fast-moving water carrying a heavy bed load of gravel. The deposits were laid down from drainages flowing eastward out of the Sierra Nevada to the Owens River. Cartago soils are mapped near the reconstructed barracks in map unit 488047 (fig. 2).

Mixed alluvium in the context of the local soil survey indicates parent material sources from the Inyo and White Mountains as well as the Sierra Nevada (USDA-NRCS, 2002). Carbonates observed in map units 488211 and 488326 indicate calcareous alluvium contributions from the White and Inyo Mountains (see table 19). The soils in map units 488211 and 488326 formed in mixed alluvium. The soil inorganic carbon (SIC) in these soils (Morey family, Rindge family, and Winnedumah soils) is derived from calcium carbonate. Rainwater containing small amounts of carbon dioxide percolates from the surface into the soil profile and dissolves calcium carbonate, thus producing calcium bicarbonate. Calcium



Figure 1.—The mess hall's Japanese meditation garden in Manzanar National Historic Site. The garden was created from local, well rounded rocks (in left foreground) and angular rocks (in background) that were hauled from nearby mountains.

bicarbonate in solution moves downward through the soil each time it becomes wet; calcium carbonate is deposited when the soil is dry. Calcium carbonate is disseminated throughout the profile or occurs as coatings on rock fragments.

Soils in the park that formed dominantly from granitic alluvium, such as Cartago, Goodale, and Lubkin, have low contents of clay and silt (see table 17). As a result, these soils have a low available water-holding capacity (fig. 3). Crops grown in soils with a low available water-holding capacity need frequent irrigation. Efforts to rejuvenate surviving fruit trees from the war-time Manzanar orchards will require extra water due to a low available water-holding capacity and local, human-altered hydrologic conditions.

Climate

Differences in climate can result in differences in soils. Temperature and moisture, the two most commonly measured features of climate, influence soil formation. Weathering is most active when soils are moist and warm because these soil conditions are conducive to rapid chemical reactions and increased biological activity. Cooler temperatures result in slower chemical reactions. While average temperatures and precipitation are important in determining soil properties, the extremes of climate in any given locale also play a major role in soil formation.

During periods of rainfall or snowmelt, water carrying dissolved or suspended solids moves through the soil in a process called leaching. The leaching process becomes active with the onset of rainfall or snowmelt. Different temperatures and moisture amounts cause different patterns of weathering and leaching in the soil. Seasonal and daily changes in temperature affect moisture effectiveness, biological activity, rates of chemical reactions, and kinds of vegetation. The soil climate of Manzanar NHS is fairly warm but dry, so leaching is limited.



Figure 2.—The reconstructed barracks in Manzanar National Historic Site, located on Cartago soils derived from granitic alluvium from the Sierra Nevada mountains (background).

Past climates in the survey area have fluctuated between periods that were drier and colder than the present climate to periods that were warmer and wetter. Glacial activity in the Sierras Nevada led to the deposition of debris on the alluvial fans and valley floors. While the terrain of the park is relatively flat, the surrounding terrain results in cold air drainage to the lower elevations of the valley. Manzanar NHS also has a very dry climate since it is in the rain shadow of the Sierra Nevada (fig. 4).

Wind redistributes sand, salts, and other particles in arid and semiarid regions. Except for Morey and Rindge family soils, all the soils in the park have high wind erodibility due to loamy sand surface layers (see table 18). Wind erodibility group 2 is the second highest group. Internees at the Manzanar Relocation Center experienced local windblown sands infiltrating their living quarters (Houston, 1973). They also experienced clay- and silt-sized salts blowing in from the dry lake bed of Owens Lake.

Organisms

Plants, animals, micro-organisms, and humans affect the formation of soils. Plants capture solar energy, a fundamental driver of many soil processes, via photosynthesis and transfer that energy to the soil. Fungi and bacteria are the primary organisms that decompose organic matter and add nutrients to the soil. Animals and micro-organisms

mix soils and form burrows and pores. Plant roots open channels in the soils. Abandoned tunnels commonly are filled with loose material from the overlying horizons and transmit water more readily than the surrounding undisturbed soil material.

Different types of roots also have different effects on soils. Grass roots are fibrous and decompose easily, adding organic matter to the soil. Fine grass roots can extend below the surface for many feet. Plant roots also help to develop soil structure and aggregate stability. Vegetation increases soil stability by protecting the surface against wind and water erosion. Micro-organisms affect chemical exchanges between roots and soil.

Humans also can mix the soil extensively. The surface horizons of soils in Manzanar NHS have been mixed by human activities. Humans have also changed soils by diverting water from them or onto them (fig. 5).

Water diverted onto soils derived from granite accelerates erosion as these soils have low cohesiveness (fig. 6). Low cohesiveness results in soils caving in. Table 11 shows that the soils in Manzanar NHS have severe limitations for shallow excavations due to unstable walls. This is why the mess hall root cellar in the park is lined with rocks (fig. 7).

Rindge Family soils formed predominantly from organic material mixed with mineral material, specifically from decomposed plant material that accumulated under water near springs and around seeps. Water-tolerant plants, such as sedges, rushes, and mosses, grow around the water sources. Plant residue slowly accumulates around the springs and seeps because the permanently wet soil conditions retard oxidation of the plant material and slow the process of decomposition. The result is the formation of mounds. In Manzanar NHS, the springs and seeps are associated with a geologic fault. Rindge family soils are the only soils identified in the park that have a water table within 4 feet of the soil surface during some months (see table 21). Table 19 shows the



Figure 3.—Orchards planted during World War II in Manzanar National Historic Site. The restoration of these orchards will require irrigation due to the soil's low available water-holding capacity.

Soil Survey of Manzanar National Historic Site, California




Figure 4.—A stylized model of the landscape surrounding Manzanar National Historic Site. The park lies in the rain shadow of the Sierra Nevada. View is to the north.




Figure 5.—An image showing the system of water diversions (in blue) that has altered the hydrology of all the soils in Manzanar National Historic Site. The park boundary is in green, and the park auto tour route is in red. View is to the west. (Provided by the National Park Service using a Google Earth image.)

high levels of soil organic carbon (SOC) of Rindge family soils. The 59 kg/m² of SOC (calculated to a depth of 152 centimeters) in Rindge family soils equates to 266 tons of SOC per acre. Rindge family soils used to be much wetter but their water regime has been altered due to water diversion. Present-day Rindge family soils are not as wet



Figure 6.—An area of highly erodible soils that are easily gullied by overland water flow.

as in their natural state. In some areas, they have dried out completely and are now subject to wind erosion because of the low bulk-density organic material.

The influence of organic residual parent material on soil is a major factor in the development of ecological niches in a park. Native vegetation depends on climate, drainage and topography, and biological factors. Other factors affecting vegetation are soil density, depth, chemistry, temperature, and moisture. In Manzanar NHS, Morey family, Rindge family, and Winnedumah soils vary in soil chemistry, which helps determine ecological niches in their natural state. These conditions are reflected in the range sites assigned to each soil (see table 6). Morey family soils are in the Saline Meadow site, Rindge family soils are in the Wetland site, and Winnedumah soils are in the Sodic Fan site. Vegetation on the Saline Meadow site is typically alkali sacaton, inland saltgrass, and some Baltic rush. Vegetation on the Wetland site typically is bulrush, spikerush, and broadleaf cattail. Vegetation on the Sodic Fan site is predominantly Torrey's saltbush. Descriptions of the range sites in Manzanar NHS, including data on native vegetation, are provided in the Appendix. These range sites will be rewritten as ecological sites by NRCS in the future.

The different drainage conditions in the three major soils of map unit 488211 result in a range of electrical conductivity from 2 to 16 mmhos/cm and a range of sodium adsorption ratios (a measure of the predominance of sodium in the soil) from 1 to 13 in the surface layer. Sediment and salts accumulate as a result of the water runoff from weathered rock and eroding soils in the surrounding mountains. Evapotranspiration and restricted subsurface drainage resulting from a high water table promote the accumulation of salt and sodium in the soils.

Both Morey family and Rindge family soils are hydric based on prior wet conditions and not on their present altered hydrologic condition (see table 4). These soils are associated with the treed areas in Manzanar NHS. Figure 8 shows these soils as a black area. The southwest- to northeast-trending swath of darker soils across the park



Figure 7.—The mess hall’s root cellar in Manzanar National Historic Site. Because it was built in an area of sandy soils that easily cave in, the cellar was lined with rocks.

is associated with a geologic fault (USDI-NPS, 2011). Organic matter, evidenced by the dark color, accumulated in the soils under previous, wetter conditions.

Topography and Relief

Topography refers to the shape of the landscape, and relief refers to differences in elevation. The overall landscape in a park, whether it consists of flat depressions or sloping alluvial fans, is the result of erosion and depositional processes. These processes may have occurred in response to changes in climate, glaciation, and/or tectonic activities. Cyclic periods of landscape stability and instability influence the types of soils that form on the landscape.

The youngest geomorphic surfaces generally are alluvial fans, flood plains, and basin floors associated with the major rivers and streams where alluvium has been deposited. They make up the geographic setting of Manzanar NHS. The survey area is slightly sloping to the east, and relief is limited. The proximity to the Owens River drainage defines the landscape, and a fault line that traverses the park is responsible for local wetness (fig. 9).

Map unit 488211 is comprised of three soils (Ridge family, Morey family, and Winnedumah) that are all very deep and have variable drainage that is determined by landscape position and topography. This unit is mapped along the treed swath across the western portion of Manzanar NHS (see figure 9).

Rindge and Morey family soils formed in climates that were much wetter than today’s. Their hydrology has been altered by human activities. Surface water has been diverted for decades, and the ground water has little recharge.

Time

Time for parent material, climate, organisms, and topography to interact with the soil also is a soil-forming factor. Over time, soils exhibit features that reflect the interaction

of other soil-forming factors. Recently deposited materials, such as those deposited by a flood, exhibit no features from soil development activities and have properties that are mostly inherited from the new material. The previous soil surface and underlying horizons become buried. The time clock resets for these soils. The different horizons in a soil profile and the degree of development can be directly related to time. Terraces above the active flood plain, while similar in origin to the flood plain, are old, abandoned flood plains. Soils on these terraces exhibit more horizon development. Cartago and Goodale soils on active alluvial fans have few distinctive characteristics and no diagnostic subsurface horizons.

Lubkin and Winnedumah soils, which have argillic diagnostic subsurface horizons, are on stable terraces and have had time to develop distinctive profile characteristics. An argillic horizon is a zone of clay accumulation. Winnedumah soils are mapped around the entrance to Manzanar NHS (fig. 10).

Classification of the Soils

Soils are named and classified on the basis of physical and chemical properties in their horizons (layers). Color, texture, structure, and other properties of the soil to a depth of 2 meters are used to key the soil into a classification system. This system helps people to use soil information and also provides a common language for scientists.

Soils and their horizons differ from one another, depending on how and when they formed. Soil scientists use the five soil-forming factors to help predict where different soils may occur. The degree and expression of the soil horizons reflect the extent of interaction of the soil-forming factors with one or more of the soil-forming processes (Simonson, 1959).

When mapping soils, a soil scientist looks for areas with similar soil-forming factors to find similar soils. The properties of the soils are described. Soils with the same kind of properties are given taxonomic names. Soils are classified, mapped, and interpreted on the basis of various kinds of soil horizons and their arrangement. The distribution of




Figure 8.—An image of Manzanar National Historic Site showing the distribution of the organic matter in soils. The darker area on the west side of the park consists of Rindge family, Morey family, and Winnedumah soils that have high organic matter contents due to historic wetness. (Google Earth image)

soil orders corresponds with the general patterns of the soil-forming factors within the park.

The system of soil classification used by the National Cooperative Soil Survey has six categories (Soil Survey Staff, 1999 and 2010). Beginning with the broadest, these categories are the order, suborder, great group, subgroup, family, and series. Classification is based on soil properties observed in the field or inferred from those observations or from laboratory measurements. The categories are defined in the following paragraphs.

ORDER. Soil taxonomy at the highest hierarchical level identifies 12 soil orders. The names for the orders and taxonomic soil properties relate to Greek, Latin, or other root words that reveal something about the soil. The differences among orders reflect the dominant soil-forming processes and the degree of soil formation. Each order is identified by a word ending in *sol*. An example is Aridisol, meaning arid soil.

SUBORDER. Each order is divided into suborders primarily on the basis of properties that influence soil genesis and are important to plant growth or properties that reflect the most important variables within the orders. Sixty-four suborders are recognized at the next level of classification. The last syllable in the name of a suborder indicates the order. An example is Argid (*Arg*, meaning clay, plus *id*, from Aridisol).

GREAT GROUP. Each suborder is divided into great groups on the basis of close similarities in kind, arrangement, and degree of development of pedogenic horizons; soil moisture and temperature regimes; type of saturation; and base status. There are about 300 great groups. Each great group is identified by the name of a suborder and by a prefix that indicates a property of the soil. An example is Haplargids (*Hapl*, meaning minimal horizonation, plus *argid*, the suborder of the Aridisols that has a zone of clay accumulation).

SUBGROUP. There are more than 2,400 subgroups. Each great group has a typic subgroup. The typic subgroup is the central concept of the great group; it is not



Figure 9.—An area of map unit 488211 along a fault line in the northwestern area of Manzanar National Historic Site. Trees grow along the fault line where a water table existed. Surface soil colors are noticeably darker.



Figure 10.—The entrance to Manzanar National Historic Site, located on Winnedumah soils. Winnedumah soils formed under conditions that were wetter than those of today due to altered drainage. They have a zone of clay accumulation below a depth of 30 inches.

necessarily the most extensive. Other subgroups are intergrades or extragrades. Intergrades are transitions to other orders, suborders, or great groups. Extragrades have some properties that are not representative of the great group but do not indicate transitions to any other taxonomic class. Each subgroup is identified by one or more adjectives preceding the name of the great group. The adjective *Xeric* identifies the subgroup that has a summer precipitation pattern. An example is Xeric Haplargids.

FAMILY. Families are established within a subgroup on the basis of physical and chemical properties and other characteristics that affect management. Generally, the properties for family placement are those of horizons below a traditional agronomic plow depth. Among the properties and characteristics considered are particle-size class, mineralogy class, cation-exchange activity class, soil temperature regime, soil depth, and reaction class. A family name consists of the name of a subgroup preceded by terms that indicate soil properties. An example is loamy-skeletal, mixed, thermic Xeric Haplargids.

SERIES. The soil series is the lowest category in the soil classification system. The series consists of soils within a family that have horizons similar in color, texture, structure, reaction, consistence, mineral and chemical composition, and arrangement in the profile.

Most parks are mapped to the series level. The names of soil series are selected by the soil scientists during the course of mapping. The series names are commonly geographic place names or are coined. The Lubkin series is classified as loamy-skeletal, mixed, thermic Xeric Haplargids.

Table 23 indicates the order, suborder, great group, subgroup, and family of the soil series in the park. Table 24 displays the classification as a key sorted by soil order.

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Buol, S.W., R.J. Southard, R.C. Graham, and P.A. McDaniel. 2011. Soil genesis and classification. 6th edition.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Houston, W.J. 1973. Farewell to Manzanar.

Jenny, Hans. 1941. Factors of soil formation.

Levine, Ellen. 1995. A fence away from freedom: Japanese Americans and World War II. 1st edition.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Reheis, Marith C. 2011. Owens (Dry) Lake, California: A human-induced dust problem. U.S. Geological Survey.

<http://geochange.er.usgs.gov/sw/impacts/geology/owens/>

Simonson, Roy W. 1959. Outline of a generalized theory of soil genesis. Soil Science Society of America Proceedings 23:152-156.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18.

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service. U.S. Department of Agriculture Handbook 436.

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service.

Soil Survey of Manzanar National Historic Site, California

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. ESIS user guide.
<http://esis.sc.egov.usda.gov/Welcome/pgESDWelcome.aspx>

United States Department of Agriculture, Natural Resources Conservation Service. National forestry handbook. <http://soils.usda.gov/technical/nfhandbook/>

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. <http://soils.usda.gov/technical/nfmanual/>

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook.
<http://www.glti.nrcs.usda.gov/technical/publications/nrph.html>

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI.
<http://soils.usda.gov/technical/>

United States Department of Agriculture, Natural Resources Conservation Service. PLANTS database. National Plant Data Center. <http://plants.usda.gov>

United States Department of Agriculture, Natural Resources Conservation Service. 2002. Soil survey of Benton-Owens Valley Area, Parts of Inyo and Mono Counties, California.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

United States Department of Agriculture, Natural Resources Conservation Service. 2010. Field indicators of hydric soils in the United States, Version 7.0. L.M. Vasilas, G.W. Hurt, and C.V. Noble, editors. USDA-NRCS in cooperation with the National Technical Committee for Hydric Soils.

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210.

United States Department of the Interior, National Park Service. 2011. Personal communication with the staff at Manzanar National Historic Site.

Glossary

Aeration, soil. The exchange of air in soil with air from the atmosphere. The air in a well aerated soil is similar to that in the atmosphere; the air in a poorly aerated soil is considerably higher in carbon dioxide and lower in oxygen.

Aggregate, soil. Many fine particles held in a single mass or cluster. Natural soil aggregates, such as granules, blocks, or prisms, are called peds. Clods are aggregates produced by tillage or logging.

Alkali (sodic) soil. A soil having so high a degree of alkalinity (pH 8.5 or higher) or so high a percentage of exchangeable sodium (15 percent or more of the total exchangeable bases), or both, that plant growth is restricted.

Alluvial fan. The fanlike deposit of a stream where it issues from a gorge upon a plain or of a tributary stream near or at its junction with its main stream.

Alluvium. Material, such as sand, silt, or clay, deposited on land by streams.

Alpha,alpha-dipyridyl. A dye that when dissolved in 1N ammonium acetate is used to detect the presence of reduced iron (Fe II) in the soil. A positive reaction indicates a type of redoximorphic feature.

Aquic conditions. Current soil wetness characterized by saturation, reduction, and redoximorphic features.

Argillic horizon. A subsoil horizon characterized by an accumulation of illuvial clay.

Aspect. The direction in which a slope faces.

Available water capacity (available moisture capacity). The capacity of soils to hold water available for use by most plants. It is commonly defined as the difference between the amount of soil water at field moisture capacity and the amount at wilting point. It is commonly expressed as inches of water per inch of soil. The capacity, in inches, in a 60-inch profile or to a limiting layer is expressed as:

Very low	0 to 3
Low	3 to 6
Moderate.....	6 to 9
High	9 to 12
Very high.....	more than 12

Base saturation. The degree to which material having cation-exchange properties is saturated with exchangeable bases (sum of Ca, Mg, Na, and K), expressed as a percentage of the total cation-exchange capacity.

Bedrock. The solid rock that underlies the soil and other unconsolidated material or that is exposed at the surface.

Canopy. The leafy crown of trees or shrubs. (See Crown.)

Cation. An ion carrying a positive charge of electricity. The common soil cations are calcium, potassium, magnesium, sodium, and hydrogen.

Cation-exchange capacity. The total amount of exchangeable cations that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. The term, as applied to soils, is synonymous with base-exchange capacity but is more precise in meaning.

Clay. As a soil separate, the mineral soil particles less than 0.002 millimeter in diameter. As a soil textural class, soil material that is 40 percent or more clay, less than 45 percent sand, and less than 40 percent silt.

- Claypan.** A slowly permeable soil horizon that contains much more clay than the horizons above it. A claypan is commonly hard when dry and plastic or stiff when wet.
- Climax plant community.** The stabilized plant community on a particular site. The plant cover reproduces itself and does not change so long as the environment remains the same.
- Coarse textured soil.** Sand or loamy sand.
- Colluvium.** Soil material or rock fragments, or both, moved by creep, slide, or local wash and deposited at the base of steep slopes.
- Complex, soil.** A map unit of two or more kinds of soil or miscellaneous areas in such an intricate pattern or so small in area that it is not practical to map them separately at the selected scale of mapping. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas.
- Control section.** The part of the soil on which classification is based. The thickness varies among different kinds of soil, but for many it is that part of the soil profile between depths of 10 inches and 40 or 80 inches.
- Corrosion.** Soil-induced electrochemical or chemical action that dissolves or weakens concrete or uncoated steel.
- Crown.** The upper part of a tree or shrub, including the living branches and their foliage.
- Culmination of the mean annual increment (CMAI).** The average annual increase per acre in the volume of a stand. Computed by dividing the total volume of the stand by its age. As the stand increases in age, the mean annual increment continues to increase until mortality begins to reduce the rate of increase. The point where the stand reaches its maximum annual rate of growth is called the culmination of the mean annual increment.
- Depth, soil.** Generally, the thickness of the soil over bedrock. Very deep soils are more than 60 inches deep over bedrock; deep soils, 40 to 60 inches; moderately deep, 20 to 40 inches; shallow, 10 to 20 inches; and very shallow, less than 10 inches.
- Drainage class (natural).** Refers to the frequency and duration of wet periods under conditions similar to those under which the soil formed. Alterations of the water regime by human activities, either through drainage or irrigation, are not a consideration unless they have significantly changed the morphology of the soil. Seven classes of natural soil drainage are recognized—*excessively drained, somewhat excessively drained, well drained, moderately well drained, somewhat poorly drained, poorly drained, and very poorly drained*. These classes are defined in the “Soil Survey Manual.”
- Drainage, surface.** Runoff, or surface flow of water, from an area.
- Ecological site.** An area where climate, soil, and relief are sufficiently uniform to produce a distinct natural plant community. An ecological site is the product of all the environmental factors responsible for its development. It is typified by an association of species that differ from those on other ecological sites in kind and/or proportion of species or in total production.
- Eluviation.** The movement of material in true solution or colloidal suspension from one place to another within the soil. Soil horizons that have lost material through eluviation are eluvial; those that have received material are illuvial.
- Eolian soil material.** Earthy parent material accumulated through wind action; commonly refers to sandy material in dunes or to loess in blankets on the surface.
- Erosion.** The wearing away of the land surface by water, wind, ice, or other geologic agents and by such processes as gravitational creep.
Erosion (geologic). Erosion caused by geologic processes acting over long geologic periods and resulting in the wearing away of mountains and the building

up of such landscape features as flood plains and coastal plains. Synonym: natural erosion.

Erosion (accelerated). Erosion much more rapid than geologic erosion, mainly as a result of human or animal activities or of a catastrophe in nature, such as a fire, that exposes the surface.

Escarpment. A relatively continuous and steep slope or cliff breaking the general continuity of more gently sloping land surfaces and resulting from erosion or faulting. Synonym: scarp.

Fertility, soil. The quality that enables a soil to provide plant nutrients, in adequate amounts and in proper balance, for the growth of specified plants when light, moisture, temperature, tilth, and other growth factors are favorable.

Fill slope. A sloping surface consisting of excavated soil material from a road cut. It commonly is on the downhill side of the road.

Fine textured soil. Sandy clay, silty clay, or clay.

Flood plain. A nearly level alluvial plain that borders a stream and is subject to flooding unless protected artificially.

Fluvial. Of or pertaining to rivers; produced by river action, as a fluvial plain.

Forb. Any herbaceous plant not a grass or a sedge.

Forest cover. All trees and other woody plants (underbrush) covering the ground in a forest.

Forest type. A stand of trees similar in composition and development because of given physical and biological factors by which it may be differentiated from other stands.

Gravel. Rounded or angular fragments of rock as much as 3 inches (2 millimeters to 7.6 centimeters) in diameter. An individual piece is a pebble.

Gravelly soil material. Material that has 15 to 35 percent, by volume, rounded or angular rock fragments, not prominently flattened, as much as 3 inches (7.6 centimeters) in diameter.

Ground water. Water filling all the unblocked pores of the material below the water table.

Hard bedrock. Bedrock that cannot be excavated except by blasting or by the use of special equipment that is not commonly used in construction.

Hill. A natural elevation of the land surface, rising as much as 1,000 feet above surrounding lowlands, commonly of limited summit area and having a well defined outline; hillsides generally have slopes of more than 15 percent. The distinction between a hill and a mountain is arbitrary and is dependent on local usage.

Horizon, soil. A layer of soil, approximately parallel to the surface, having distinct characteristics produced by soil-forming processes. In the identification of soil horizons, an uppercase letter represents the major horizons. Numbers or lowercase letters that follow represent subdivisions of the major horizons. An explanation of the subdivisions is given in the "Soil Survey Manual." The major horizons of mineral soil are as follows:

O horizon.—An organic layer of fresh and decaying plant residue.

A horizon.—The mineral horizon at or near the surface in which an accumulation of humified organic matter is mixed with the mineral material. Also, a plowed surface horizon, most of which was originally part of a B horizon.

E horizon.—The mineral horizon in which the main feature is loss of silicate clay, iron, aluminum, or some combination of these.

B horizon.—The mineral horizon below an A horizon. The B horizon is in part a layer of transition from the overlying A to the underlying C horizon. The B horizon also has distinctive characteristics, such as (1) accumulation of clay, sesquioxides, humus, or a combination of these; (2) prismatic or blocky structure; (3) redder or browner colors than those in the A horizon; or (4) a combination of these.

C horizon.—The mineral horizon or layer, excluding indurated bedrock, that is little affected by soil-forming processes and does not have the properties typical of the overlying soil material. The material of a C horizon may be either like or unlike that in which the solum formed. If the material is known to differ from that in the solum, an Arabic numeral, commonly a 2, precedes the letter C.

Cr horizon.—Soft, consolidated bedrock beneath the soil.

R layer.—Consolidated bedrock beneath the soil. The bedrock commonly underlies a C horizon, but it can be directly below an A or a B horizon.

Hydrologic soil groups. Refers to soils grouped according to their runoff potential.

The soil properties that influence this potential are those that affect the minimum rate of water infiltration on a bare soil during periods after prolonged wetting when the soil is not frozen. These properties are depth to a seasonal high water table, the infiltration rate and permeability after prolonged wetting, and depth to a very slowly permeable layer. The slope and the kind of plant cover are not considered but are separate factors in predicting runoff.

Illuviation. The movement of soil material from one horizon to another in the soil profile. Generally, material is removed from an upper horizon and deposited in a lower horizon.

Infiltration. The downward entry of water into the immediate surface of soil or other material, as contrasted with percolation, which is movement of water through soil layers or material.

Infiltration capacity. The maximum rate at which water can infiltrate into a soil under a given set of conditions.

Infiltration rate. The rate at which water penetrates the surface of the soil at any given instant, usually expressed in inches per hour. The rate can be limited by the infiltration capacity of the soil or the rate at which water is applied at the surface.

Intake rate. The average rate of water entering the soil under irrigation. Most soils have a fast initial rate; the rate decreases with application time. Therefore, intake rate for design purposes is not a constant but is a variable depending on the net irrigation application. The rate of water intake, in inches per hour, is expressed as follows:

Less than 0.2	very low
0.2 to 0.4	low
0.4 to 0.75	moderately low
0.75 to 1.25	moderate
1.25 to 1.75	moderately high
1.75 to 2.5	high
More than 2.5	very high

K_{sat} . Saturated hydraulic conductivity. (See Permeability.)

Leaching. The removal of soluble material from soil or other material by percolating water.

LEP. See Linear extensibility percent.

Linear extensibility (LE). Refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. Linear extensibility is used to determine the shrink-swell potential of soils. It is an expression of the volume change between the water content of the clod at $1/3$ - or $1/10$ -bar tension (33kPa or 10kPa tension) and oven dryness. Volume change is influenced by the amount and type of clay minerals in the soil. The volume change is the percent change for the whole soil. If it is expressed as a fraction, the resulting value is COLE, coefficient of linear extensibility.

Linear extensibility percent. Refers to the percent change in linear extensibility.

Liquid limit. The moisture content at which the soil passes from a plastic to a liquid state.

Loam. Soil material that is 7 to 27 percent clay particles, 28 to 50 percent silt particles, and less than 52 percent sand particles.

Loess. Fine grained material, dominantly of silt-sized particles, deposited by wind.

Low strength. The soil is not strong enough to support loads.

Medium textured soil. Very fine sandy loam, loam, silt loam, or silt.

Mineral soil. Soil that is mainly mineral material and low in organic material. Its bulk density is more than that of organic soil.

Miscellaneous area. An area that has little or no natural soil and supports little or no vegetation.

Moderately coarse textured soil. Coarse sandy loam, sandy loam, or fine sandy loam.

Moderately fine textured soil. Clay loam, sandy clay loam, or silty clay loam.

Neutral soil. A soil having a pH value of 6.6 to 7.3. (See Reaction, soil.)

Nutrient, plant. Any element taken in by a plant essential to its growth. Plant nutrients are mainly nitrogen, phosphorus, potassium, calcium, magnesium, sulfur, iron, manganese, copper, boron, and zinc obtained from the soil and carbon, hydrogen, and oxygen obtained from the air and water.

Organic matter. Plant and animal residue in the soil in various stages of decomposition. The content of organic matter in the surface layer is described as follows:

Very low	less than 0.5 percent
Low	0.5 to 1.0 percent
Moderately low.....	1.0 to 2.0 percent
Moderate.....	2.0 to 4.0 percent
High	4.0 to 8.0 percent
Very high.....	more than 8.0 percent

Pan. A compact, dense layer in a soil that impedes the movement of water and the growth of roots. For example, *hardpan*, *fragipan*, *claypan*, *plowpan*, and *traffic pan*.

Parent material. The unconsolidated organic and mineral material in which soil forms.

Ped. An individual natural soil aggregate, such as a granule, a prism, or a block.

Pedon. The smallest volume that can be called "a soil." A pedon is three dimensional and large enough to permit study of all horizons. Its area ranges from about 10 to 100 square feet (1 square meter to 10 square meters), depending on the variability of the soil.

Percolation. The movement of water through the soil.

Permeability. The quality of the soil that enables water or air to move downward through the profile. The rate at which a saturated soil transmits water is accepted as a measure of this quality. In soil physics, the rate is referred to as "saturated hydraulic conductivity," which is defined in the "Soil Survey Manual." In line with conventional usage in the engineering profession and with traditional usage in published soil surveys, this rate of flow continues to be expressed as "permeability." Terms describing permeability, measured in inches per hour, are as follows:

Extremely slow.....	0.0 to 0.01 inch
Very slow	0.01 to 0.06 inch
Slow	0.06 to 0.2 inch
Moderately slow.....	0.2 to 0.6 inch
Moderate.....	0.6 inch to 2.0 inches
Moderately rapid.....	2.0 to 6.0 inches
Rapid	6.0 to 20 inches
Very rapid.....	more than 20 inches

Phase, soil. A subdivision of a soil series based on features that affect its use and management, such as slope, stoniness, and flooding.

pH value. A numerical designation of acidity and alkalinity in soil. (See Reaction, soil.)

Plasticity index. The numerical difference between the liquid limit and the plastic limit; the range of moisture content within which the soil remains plastic.

Plastic limit. The moisture content at which a soil changes from semisolid to plastic.

- Plowpan.** A compacted layer formed in the soil directly below the plowed layer.
- Ponding.** Standing water on soils in closed depressions. Unless the soils are artificially drained, the water can be removed only by percolation or evapotranspiration.
- Poorly graded.** Refers to a coarse grained soil or soil material consisting mainly of particles of nearly the same size. Because there is little difference in size of the particles, density can be increased only slightly by compaction.
- Potential native plant community.** See Climax plant community.
- Potential rooting depth (effective rooting depth).** Depth to which roots could penetrate if the content of moisture in the soil were adequate. The soil has no properties restricting the penetration of roots to this depth.
- Productivity, soil.** The capability of a soil for producing a specified plant or sequence of plants under specific management.
- Profile, soil.** A vertical section of the soil extending through all its horizons and into the parent material.
- Rangeland.** Land on which the potential natural vegetation is predominantly grasses, grasslike plants, forbs, or shrubs suitable for grazing or browsing. It includes natural grasslands, savannas, many wetlands, some deserts, tundras, and areas that support certain forb and shrub communities.
- Reaction, soil.** A measure of acidity or alkalinity of a soil, expressed in pH values. A soil that tests to pH 7.0 is described as precisely neutral in reaction because it is neither acid nor alkaline. The degrees of acidity or alkalinity, expressed as pH values, are:

Ultra acid.....	less than 3.5
Extremely acid	3.5 to 4.4
Very strongly acid	4.5 to 5.0
Strongly acid	5.1 to 5.5
Moderately acid	5.6 to 6.0
Slightly acid.....	6.1 to 6.5
Neutral	6.6 to 7.3
Slightly alkaline.....	7.4 to 7.8
Moderately alkaline.....	7.9 to 8.4
Strongly alkaline.....	8.5 to 9.0
Very strongly alkaline.....	9.1 and higher

- Redoximorphic concentrations.** Nodules, concretions, soft masses, pore linings, and other features resulting from the accumulation of iron or manganese oxide. An indication of chemical reduction and oxidation resulting from saturation.
- Redoximorphic depletions.** Low-chroma zones from which iron and manganese oxide or a combination of iron and manganese oxide and clay has been removed. These zones are indications of the chemical reduction of iron resulting from saturation.
- Redoximorphic features.** Redoximorphic concentrations, redoximorphic depletions, reduced matrices, a positive reaction to alpha,alpha-dipyridyl, and other features indicating the chemical reduction and oxidation of iron and manganese compounds resulting from saturation.
- Relief.** The elevations or inequalities of a land surface, considered collectively.
- Residuum (residual soil material).** Unconsolidated, weathered or partly weathered mineral material that accumulated as consolidated rock disintegrated in place.
- Rock fragments.** Rock or mineral fragments having a diameter of 2 millimeters or more; for example, pebbles, cobbles, stones, and boulders.
- Root zone.** The part of the soil that can be penetrated by plant roots.
- Runoff.** The precipitation discharged into stream channels from an area. The water that flows off the surface of the land without sinking into the soil is called surface runoff. Water that enters the soil before reaching surface streams is called ground-water runoff or seepage flow from ground water.
- Saline soil.** A soil containing soluble salts in an amount that impairs growth of plants. A saline soil does not contain excess exchangeable sodium.

- Sand.** As a soil separate, individual rock or mineral fragments from 0.05 millimeter to 2.0 millimeters in diameter. Most sand grains consist of quartz. As a soil textural class, a soil that is 85 percent or more sand and not more than 10 percent clay.
- Sandstone.** Sedimentary rock containing dominantly sand-sized particles.
- Saprolite.** Unconsolidated residual material underlying the soil and grading to hard bedrock below.
- Saturation.** Wetness characterized by zero or positive pressure of the soil water. Under conditions of saturation, the water will flow from the soil matrix into an unlined auger hole.
- Sedimentary rock.** Rock made up of particles deposited from suspension in water. The chief kinds of sedimentary rock are conglomerate, formed from gravel; sandstone, formed from sand; shale, formed from clay; and limestone, formed from soft masses of calcium carbonate. There are many intermediate types. Some wind-deposited sand is consolidated into sandstone.
- Series, soil.** A group of soils that have profiles that are almost alike. All the soils of a series have horizons that are similar in composition, thickness, and arrangement.
- Shale.** Sedimentary rock formed by the hardening of a clay deposit.
- Silt.** As a soil separate, individual mineral particles that range in diameter from the upper limit of clay (0.002 millimeter) to the lower limit of very fine sand (0.05 millimeter). As a soil textural class, soil that is 80 percent or more silt and less than 12 percent clay.
- Siltstone.** Sedimentary rock made up of dominantly silt-sized particles.
- Similar soils.** Soils that share limits of diagnostic criteria, behave and perform in a similar manner, and have similar conservation needs or management requirements for the major land uses in the survey area.
- Site index.** A designation of the quality of a forest site based on the height of the dominant stand at an arbitrarily chosen age. For example, if the average height attained by dominant and codominant trees in a fully stocked stand at the age of 50 years is 75 feet, the site index is 75.
- Slope.** The inclination of the land surface from the horizontal. Percentage of slope is the vertical distance divided by horizontal distance, then multiplied by 100. Thus, a slope of 20 percent is a drop of 20 feet in 100 feet of horizontal distance.
- Sodic (alkali) soil.** A soil having so high a degree of alkalinity (pH 8.5 or higher) or so high a percentage of exchangeable sodium (15 percent or more of the total exchangeable bases), or both, that plant growth is restricted.
- Sodicity.** The degree to which a soil is affected by exchangeable sodium. Sodicity is expressed as a sodium adsorption ratio (SAR) of a saturation extract, or the ratio of Na^+ to $\text{Ca}^{++} + \text{Mg}^{++}$. The degrees of sodicity and their respective ratios are:
- | | |
|---------------|----------------|
| Slight..... | less than 13:1 |
| Moderate..... | 13-30:1 |
| Strong | more than 30:1 |
- Sodium adsorption ratio (SAR).** A measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration.
- Soft bedrock.** Bedrock that can be excavated with trenching machines, backhoes, small rippers, and other equipment commonly used in construction.
- Soil.** A natural, three-dimensional body at the earth's surface. It is capable of supporting plants and has properties resulting from the integrated effect of climate and living matter acting on earthy parent material, as conditioned by relief over periods of time.
- Soil separates.** Mineral particles less than 2 millimeters in equivalent diameter and ranging between specified size limits. The names and sizes, in millimeters, of separates recognized in the United States are as follows:

Soil Survey of Manzanar National Historic Site, California

Very coarse sand	2.0 to 1.0
Coarse sand	1.0 to 0.5
Medium sand	0.5 to 0.25
Fine sand	0.25 to 0.10
Very fine sand	0.10 to 0.05
Silt	0.05 to 0.002
Clay.....	less than 0.002

Solum. The upper part of a soil profile, above the C horizon, in which the processes of soil formation are active. The solum in soil consists of the A, E, and B horizons. Generally, the characteristics of the material in these horizons are unlike those of the material below the solum. The living roots and plant and animal activities are largely confined to the solum.

Stone line. A concentration of coarse fragments in a soil. Generally, it is indicative of an old weathered surface. In a cross section, the line may be one fragment or more thick. It generally overlies material that weathered in place and is overlain by recent sediment of variable thickness.

Stones. Rock fragments 10 to 24 inches (25 to 60 centimeters) in diameter if rounded or 15 to 24 inches (38 to 60 centimeters) in length if flat.

Stony. Refers to a soil containing stones in numbers that interfere with or prevent tillage.

Structure, soil. The arrangement of primary soil particles into compound particles or aggregates. The principal forms of soil structure are—*platy* (laminated), *prismatic* (vertical axis of aggregates longer than horizontal), *columnar* (prisms with rounded tops), *blocky* (angular or subangular), and *granular*. *Structureless* soils are either *single grained* (each grain by itself, as in dune sand) or *massive* (the particles adhering without any regular cleavage, as in many hardpans).

Subsoil. Technically, the B horizon; roughly, the part of the solum below plow depth.

Substratum. The part of the soil below the solum.

Subsurface layer. Any surface soil horizon (A, E, AB, or EB) below the surface layer.

Surface layer. The soil ordinarily moved in tillage, or its equivalent in uncultivated soil, ranging in depth from 4 to 10 inches (10 to 25 centimeters). Frequently designated as the “plow layer,” or the “Ap horizon.”

Surface soil. The A, E, AB, and EB horizons, considered collectively. It includes all subdivisions of these horizons.

Terrace. An embankment, or ridge, constructed across sloping soils on the contour or at a slight angle to the contour. The terrace intercepts surface runoff so that water soaks into the soil or flows slowly to a prepared outlet. A terrace in a field generally is built so that the field can be farmed. A terrace intended mainly for drainage has a deep channel that is maintained in permanent sod.

Terrace (geologic). An old alluvial plain, ordinarily flat or undulating, bordering a river, a lake, or the sea.

Texture, soil. The relative proportions of sand, silt, and clay particles in a mass of soil. The basic textural classes, in order of increasing proportion of fine particles, are *sand*, *loamy sand*, *sandy loam*, *loam*, *silt loam*, *silt*, *sandy clay loam*, *clay loam*, *silty clay loam*, *sandy clay*, *silty clay*, and *clay*. The sand, loamy sand, and sandy loam classes may be further divided by specifying “coarse,” “fine,” or “very fine.”

Tilth, soil. The physical condition of the soil as related to tillage, seedbed preparation, seedling emergence, and root penetration.

Topsoil. The upper part of the soil, which is the most favorable material for plant growth. It is ordinarily rich in organic matter and is used to topdress roadbanks, lawns, and land affected by mining.

Upland. Land at a higher elevation, in general, than the alluvial plain or stream terrace; land above the lowlands along streams.

Weathering. All physical and chemical changes produced in rocks or other deposits at or near the earth's surface by atmospheric agents. These changes result in disintegration and decomposition of the material.

Well graded. Refers to soil material consisting of coarse grained particles that are well distributed over a wide range in size or diameter. Such soil normally can be easily increased in density and bearing properties by compaction. Contrasts with poorly graded soil.

Tables

Soil Survey of Manzanar National Historic Site, California

Table 1.—Soil Legend

(Map unit composition may not total 100 percent)

Map unit symbol and map unit name	Components in map unit	Percent of map unit
488047: Cartago gravelly loamy sand, 0 to 2 percent slopes-----	Cartago	85
	Unnamed	1
488048: Cartago gravelly loamy sand, 2 to 5 percent slopes-----	Cartago	85
	Unnamed	1
488091: Goodale-Cartago complex, 0 to 2 percent slopes-----	Goodale	60
	Cartago	25
	Unnamed	1
488173: Lubkin gravelly loamy sand, 5 to 15 percent slopes-----	Lubkin	85
	Unnamed	2
488211: Morey family-Winnedumah-Rindge family complex, 0 to 2 percent slopes-----	Morey family	40
	Winnedumah	25
	Rindge family	20
488326: Winnedumah fine sandy loam, 0 to 2 percent slopes-----	Winnedumah	85
	Morey family	3
	Rindge family	3

Soil Survey of Manzanar National Historic Site, California

Table 2.—Land Capability Classification

(Land capability classification is a system of grouping soils primarily on the basis of their capability to produce common cultivated crops and pasture plants without deteriorating over a long period of time)

Map unit symbol and component name	Land capability	
	N	I
488047: Cartago-----	6e-4	3s-1
488048: Cartago-----	6e-4	3s-1
488091: Goodale-----	7s-1	---
Cartago-----	7e-4	---
488173: Lubkin-----	6e-7	---
488211: Morey family-----	7e-5	2s-5
Winnedumah-----	7e-6	3e-3
Rindge family-----	7w-2	3w-2
488326: Winnedumah-----	7s-6	---

Table 3.—Prime and Other Important Farmland

(Only the soils considered prime or important farmland are listed. Urban or built-up areas of the soils listed are not considered prime or important farmland. If a soil is prime or important farmland only under certain conditions, the conditions are indicated in the column "Farmland classification")

Map unit symbol	Map unit name	Farmland classification
488047	Cartago gravelly loamy sand, 0 to 2 percent slopes	Prime farmland if irrigated
488048	Cartago gravelly loamy sand, 2 to 5 percent slopes	Prime farmland if irrigated
488211	Morey family-Winnedumah-Rindge family complex, 0 to 2 percent slopes	Prime farmland if irrigated

Soil Survey of Manzanar National Historic Site, California

Table 4.-Hydric Soils

(This report lists only those map unit components that are rated as hydric. Definitions of hydric criteria codes are included at the bottom of the report)

Map unit symbol and map unit name	Component	Percent of map unit	Landform	Hydric soils criteria			
				Hydric criteria code	Meets saturation criteria	Meets flooding criteria	Meets ponding criteria
488047: Cartago gravelly loamy sand, 0 to 2 percent slopes	Unnamed	1	drainageways	4	No	Yes	No
488048: Cartago gravelly loamy sand, 2 to 5 percent slopes	Unnamed	1	drainageways	4	No	Yes	No
488091: Goodale-Cartago complex, 0 to 2 percent slopes	Unnamed	1	drainageways	4	No	Yes	No
488173: Lubkin gravelly loamy sand, 5 to 15 percent slopes	Unnamed	2	drainageways	4	No	Yes	No
488211: Morey family-Winnedumah-Rindge family complex, 0 to 2 percent slopes	Morey family	40	stream terraces, alluvial fans	2B3	Yes	No	No
	Rindge family	20	stream terraces, fan terraces	1, 2B3	Yes	No	No
488326: Winnedumah fine sandy loam, 0 to 2 percent slopes	Morey family	3	alluvial fans	2B3	Yes	No	No
	Rindge family	3	alluvial fans	2B3	Yes	No	No

Explanation of hydric criteria codes:

1. All Histels (except for Folistels), and Histosols (except for Folists), which are, by definition, saturated.
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Historthels great group, Histoturbels great group, Pachic subgroups, or Cumulic subgroups that:
 - A. are somewhat poorly drained and have a water table at the surface (0.0 feet) during the growing season, or
 - B. are poorly drained or very poorly drained and have either:
 - 1.) a water table at the surface (0.0 feet) during the growing season if textures are coarse sand, sand, or fine sand in all layers within a depth of 20 inches, or
 - 2.) a water table at a depth of 0.5 foot or less during the growing season if permeability is equal to or greater than 6.0 in/hr in all layers within a depth of 20 inches, or
 - 3.) a water table at a depth of 1.0 foot or less during the growing season if permeability is less than 6.0 in/hr in any layer within a depth of 20 inches.
3. Soils that are frequently ponded for periods of long or very long duration during the growing season.
4. Soils that are frequently flooded for periods of long or very long duration during the growing season.

Soil Survey of Manzanar National Historic Site, California

Table 5.-Range/Ecological Sites

Map unit symbol, soil name, and percent of map unit	Range/ecological site name	Site type	Range/ecological site ID
488047: Cartago (85%)-----	Sandy 5-8" P.Z.	Rangeland	R029XG011CA
488048: Cartago (85%)-----	Gravelly Loamy Sand 5-8" P.Z.	Rangeland	R029XG018CA
488091: Goodale (60%)-----	Gravelly Loamy Sand 5-8" P.Z.	Rangeland	R029XG018CA
Cartago (25%)-----	Gravelly Loamy Sand 5-8" P.Z.	Rangeland	R029XG018CA
488173: Lubkin (85%)-----	Gravelly Loamy Sand 5-8" P.Z.	Rangeland	R029XG018CA
488211: Morey family (40%)-----	Saline Meadow	Rangeland	R029XG002CA
Winnedumah (25%)-----	Sodic Fan 5-8" P.Z.	Rangeland	R029XG005CA
Rindge family (20%)-----	Wetland	Rangeland	R029XG003CA
488326: Winnedumah (85%)-----	Sodic Fan 5-8" P.Z.	Rangeland	R029XG005CA

Table 6.-Landform, Parent Material, and Range/Ecological Site

(Component percents may not add up to 100. MAP is the mean annual precipitation)

Map unit symbol and soil name	Percent of map unit	Slope	Elevation	MAP	Landform	Parent material	Range/ecological site name and number
	Pct	Pct	Ft	In			
488047: Cartago-----	85	0-2	3799-4400	6-8	Alluvial fan	Alluvium derived from granite	Sandy 5-8" P.Z., R029XG011CA
488048: Cartago-----	85	2-5	3799-4400	6-8	Alluvial fan	Alluvium derived from granite	Gravelly Loamy Sand 5-8" P.Z., R029XG018CA
488091: Goodale-----	60	0-2	3701-4501	4-8	Alluvial fan and fan terrace	Alluvium derived from granite	Gravelly Loamy Sand 5-8" P.Z., R029XG018CA
Cartago-----	25	0-2	3701-4501	4-8	Alluvial fan and fan terrace	Alluvium derived from granite	Gravelly Loamy Sand 5-8" P.Z., R029XG018CA
488173: Lubkin-----	85	5-15	3799-5400	4-10	Fan terrace	Alluvium derived from granite	Gravelly Loamy Sand 5-8" P.Z., R029XG018CA
488211: Morey family-----	40	0-2	3799-3999	5-6	Alluvial fan and stream terrace	Organic material and/or alluvium derived from mixed sources	Saline Meadow, R029XG002CA
Winnedumah-----	25	0-2	3799-3999	5-6	Fan terrace and stream terrace	Alluvium derived from mixed sources	Sodic Fan 5-8" P.Z., R029XG005CA
Rindge family-----	20	0-2	3799-3999	5-6	Fan terrace and stream terrace	Organic material over alluvium derived from mixed sources	Wetland, R029XG003CA
488326: Winnedumah-----	85	0-2	3671-3901	4-6	Lakebed (relict) and stream terrace	Alluvium derived from mixed and/or lacustrine deposits	Sodic Fan 5-8" P.Z., R029XG005CA

Soil Survey of Manzanar National Historic Site, California

Table 7.-Rangeland Productivity

Map unit symbol and soil name	Range/ecological site name and symbol	Total dry-weight production		
		Favorable year	Normal year	Unfavorable year
		<u>Lb/acre</u>	<u>Lb/acre</u>	<u>Lb/acre</u>
488047: Cartago-----	Sandy 5-8" P.Z. (R029XG011CA)	700	550	300
488048: Cartago-----	Gravelly Loamy Sand 5-8" P.Z. (R029XG018CA)	500	300	150
488091: Goodale-----	Gravelly Loamy Sand 5-8" P.Z. (R029XG018CA)	500	300	150
Cartago-----	Gravelly Loamy Sand 5-8" P.Z. (R029XG018CA)	500	300	150
488173: Lubkin-----	Gravelly Loamy Sand 5-8" P.Z. (R029XG018CA)	500	300	150
488211: Morey family-----	Saline Meadow (R029XG002CA)	3,300	2,200	1,000
Winnedumah-----	Sodic Fan 5-8" P.Z. (R029XG005CA)	1,500	1,200	700
Rindge family-----	Wetland (R029XG003CA)	5,000	3,500	2,500
488326: Winnedumah-----	Sodic Fan 5-8" P.Z. (R029XG005CA)	1,500	1,200	700

Soil Survey of Manzanar National Historic Site, California

Table 8.—Land Management, Part I (Planting)

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Suitability for hand planting		Suitability for mechanical planting		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Well suited		Moderately suited Rock fragments	0.50	Slight Strength	0.10
488048: Cartago-----	85	Well suited		Moderately suited Rock fragments	0.50	Slight Strength	0.10
488091: Goodale-----	60	Moderately suited Sandiness	0.50	Moderately suited Rock fragments Sandiness	0.50 0.50	Slight Strength	0.10
Cartago-----	25	Moderately suited Sandiness	0.50	Moderately suited Rock fragments Sandiness	0.50 0.50	Moderate Low strength	0.50
488173: Lubkin-----	85	Moderately suited Rock fragments	0.50	Moderately suited Rock fragments Slope	0.50 0.50	Moderate Low strength	0.50
488211: Morey family-----	40	Well suited		Well suited		Severe Low strength	1.00
Winnedumah-----	25	Well suited		Well suited		Severe Low strength	1.00
Rindge family-----	20	Well suited		Well suited		Severe Low strength	1.00
488326: Winnedumah-----	85	Well suited		Well suited		Severe Low strength	1.00

Soil Survey of Manzanar National Historic Site, California

Table 8.-Land Management, Part II (Hazard of Erosion and Suitability for Roads)

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Hazard of erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Slight		Slight		Well suited	
488048: Cartago-----	85	Slight		Slight		Well suited	
488091: Goodale-----	60	Slight		Slight		Moderately suited Sandiness	0.50
Cartago-----	25	Slight		Slight		Moderately suited Sandiness	0.50
488173: Lubkin-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
488211: Morey family-----	40	Slight		Slight		Moderately suited Low strength	0.50
Winnedumah-----	25	Slight		Slight		Moderately suited Low strength	0.50
Rindge family-----	20	Slight		Slight		Moderately suited Low strength	0.50
488326: Winnedumah-----	85	Slight		Slight		Moderately suited Low strength	0.50

Soil Survey of Manzanar National Historic Site, California

Table 8.—Land Management, Part III (Site Preparation)

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Suitability for mechanical site preparation (deep)		Suitability for mechanical site preparation (surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Well suited		Well suited	
488048: Cartago-----	85	Well suited		Well suited	
488091: Goodale-----	60	Well suited		Well suited	
Cartago-----	25	Well suited		Well suited	
488173: Lubkin-----	85	Poorly suited Rock fragments	0.50	Poorly suited Rock fragments	0.50
488211: Morey family-----	40	Well suited		Well suited	
Winnedumah-----	25	Well suited		Well suited	
Rindge family-----	20	Well suited		Well suited	
488326: Winnedumah-----	85	Well suited		Well suited	

Soil Survey of Manzanar National Historic Site, California

Table 8.-Land Management, Part IV (Site Restoration)

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	High Texture/rock fragments	1.00	Low	
488048: Cartago-----	85	High Texture/rock fragments	1.00	Low	
488091: Goodale-----	60	High Texture/surface depth/rock fragments	1.00	Low	
Cartago-----	25	High Texture/rock fragments	1.00	Low	
488173: Lubkin-----	85	High Texture/rock fragments	1.00	Low	
488211: Morey family-----	40	Low Texture/rock fragments	0.10	Low	
Winnedumah-----	25	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50
Rindge family-----	20	Low Texture/rock fragments	0.10	High Soil reaction	1.00
488326: Winnedumah-----	85	Low Texture/rock fragments	0.10	High Soil reaction Salinity	1.00 1.00

Soil Survey of Manzanar National Historic Site, California

Table 9.—Recreation, Part I (Camp and Picnic Areas)

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Camp areas		Picnic areas	
		Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Very limited Flooding Too sandy Gravel	1.00 0.42 0.41	Somewhat limited Too sandy Gravel	0.42 0.41
488048: Cartago-----	85	Very limited Flooding Too sandy Gravel	1.00 0.42 0.41	Somewhat limited Too sandy Gravel	0.42 0.41
488091: Goodale-----	60	Very limited Flooding Gravel Too sandy	1.00 1.00 0.79	Very limited Gravel Too sandy	1.00 0.79
Cartago-----	25	Very limited Flooding Too sandy Gravel	1.00 0.84 0.33	Somewhat limited Too sandy Gravel	0.84 0.33
488173: Lubkin-----	85	Very limited Flooding Large stones content Too sandy Slope Gravel	1.00 1.00 0.79 0.16 0.02	Very limited Large stones content Too sandy Slope Gravel	1.00 0.79 0.16 0.02
488211: Morey family-----	40	Somewhat limited Slow water movement	0.21	Somewhat limited Slow water movement	0.21
Winnedumah-----	25	Not rated		Not rated	
Rindge family-----	20	Somewhat limited Slow water movement Depth to saturated zone	0.21 0.07	Somewhat limited Slow water movement Depth to saturated zone	0.21 0.03
488326: Winnedumah-----	85	Very limited Sodium content Salinity	1.00 1.00	Very limited Sodium content Salinity	1.00 1.00

Soil Survey of Manzanar National Historic Site, California

Table 9.-Recreation, Part II (Trail Management)

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Foot traffic and equestrian trails		Mountain bike and off-road vehicle trails	
		Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Somewhat limited Too sandy	0.42	Somewhat limited Too sandy	0.42
488048: Cartago-----	85	Somewhat limited Too sandy	0.42	Somewhat limited Too sandy	0.42
488091: Goodale-----	60	Somewhat limited Too sandy	0.79	Somewhat limited Too sandy	0.79
Cartago-----	25	Somewhat limited Too sandy	0.84	Somewhat limited Too sandy	0.84
488173: Lubkin-----	85	Very limited Large stones content Too sandy	1.00 0.79	Very limited Large stones content Too sandy	1.00 0.79
488211: Morey family-----	40	Not limited		Not limited	
Winnedumah-----	25	Not rated		Not rated	
Rindge family-----	20	Not limited		Not limited	
488326: Winnedumah-----	85	Not limited		Not limited	

Soil Survey of Manzanar National Historic Site, California

Table 10.—Dwellings and Small Commercial Buildings

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
488048: Cartago-----	85	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
488091: Goodale-----	60	Very limited Flooding Large stones	1.00 0.12	Very limited Flooding Large stones	1.00 0.12	Very limited Flooding Large stones	1.00 0.12
Cartago-----	25	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
488173: Lubkin-----	85	Very limited Flooding Large stones Slope	1.00 0.63 0.16	Very limited Flooding Large stones Slope	1.00 0.63 0.16	Very limited Flooding Slope Large stones	1.00 1.00 0.63
488211: Morey family-----	40	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
Winnedumah-----	25	Not rated		Not rated		Not rated	
Rindge family-----	20	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
488326: Winnedumah-----	85	Not limited		Somewhat limited Depth to saturated zone	0.35	Not limited	

Soil Survey of Manzanar National Historic Site, California

Table 11.—Roads and Streets, Shallow Excavations, and Landscaping

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Somewhat limited Flooding	0.40	Very limited Unstable excavation walls	1.00	Somewhat limited Droughty Gravel	0.90 0.41
488048: Cartago-----	85	Somewhat limited Flooding	0.40	Very limited Unstable excavation walls	1.00	Somewhat limited Droughty Gravel	0.90 0.41
488091: Goodale-----	60	Somewhat limited Flooding Large stones	0.40 0.12	Very limited Unstable excavation walls Large stones	1.00 0.12	Very limited Droughty Gravel	1.00 1.00
Cartago-----	25	Somewhat limited Flooding	0.40	Very limited Unstable excavation walls	1.00	Very limited Droughty Gravel	1.00 0.33
488173: Lubkin-----	85	Somewhat limited Large stones Flooding Slope	0.63 0.40 0.16	Very limited Unstable excavation walls Large stones Slope	1.00 0.63 0.16	Somewhat limited Droughty Large stones Slope Gravel	0.97 0.38 0.16 0.02
488211: Morey family-----	40	Very limited Low strength Shrink-swell	1.00 0.50	Somewhat limited Unstable excavation walls	0.10	Not limited	
Winnedumah-----	25	Not rated		Not rated		Not rated	
Rindge family-----	20	Somewhat limited Depth to saturated zone	0.03	Very limited Depth to saturated zone Unstable excavation walls	1.00 0.10	Somewhat limited Depth to saturated zone	0.03
488326: Winnedumah-----	85	Somewhat limited Frost action	0.50	Somewhat limited Depth to saturated zone Unstable excavation walls	0.35 0.10	Very limited Salinity Sodium content	1.00 1.00

Soil Survey of Manzanar National Historic Site, California

Table 12.—Sewage Disposal

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Somewhat limited Flooding	0.40	Very limited Seepage Flooding	1.00 0.40
488048: Cartago-----	85	Somewhat limited Flooding	0.40	Very limited Seepage Flooding Slope	1.00 0.40 0.32
488091: Goodale-----	60	Very limited Filtering capacity Flooding Large stones	1.00 0.40 0.12	Very limited Seepage Flooding	1.00 0.40
Cartago-----	25	Very limited Filtering capacity Flooding	1.00 0.40	Very limited Seepage Flooding	1.00 0.40
488173: Lubkin-----	85	Somewhat limited Large stones Flooding Slope	0.63 0.40 0.16	Very limited Seepage Slope Large stones Flooding	1.00 1.00 0.93 0.40
488211: Morey family-----	40	Very limited Slow water movement	1.00	Not limited	
Winnedumah-----	25	Not rated		Not rated	
Rindge family-----	20	Very limited Depth to saturated zone Slow water movement	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 0.53
488326: Winnedumah-----	85	Very limited Slow water movement Depth to saturated zone	1.00 0.84	Very limited Seepage Depth to saturated zone	1.00 0.17

Soil Survey of Manzanar National Historic Site, California

Table 13.—Source of Gravel and Sand

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Gravel source		Sand source	
		Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Poor		Fair	
		Bottom layer	0.00	Bottom layer	0.02
		Thickest layer	0.00	Thickest layer	0.03
488048: Cartago-----	85	Poor		Fair	
		Bottom layer	0.00	Bottom layer	0.02
		Thickest layer	0.00	Thickest layer	0.03
488091: Goodale-----	60	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
Cartago-----	25	Poor		Fair	
		Bottom layer	0.00	Bottom layer	0.10
		Thickest layer	0.00	Thickest layer	0.10
488173: Lubkin-----	85	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.03
		Bottom layer	0.06	Bottom layer	0.10
488211: Morey family-----	40	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
Winnedumah-----	25	Not rated		Not rated	
Rindge family-----	20	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
488326: Winnedumah-----	85	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00

Soil Survey of Manzanar National Historic Site, California

Table 14.—Source of Reclamation Material, Roadfill, and Topsoil

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Source of reclamation material		Roadfill source		Topsoil source	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.12 0.30	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments)	0.00 0.30 0.50
488048: Cartago-----	85	Poor Wind erosion Low content of organic matter Too sandy	0.00 0.12 0.30	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments)	0.00 0.30 0.50
488091: Goodale-----	60	Poor Stone content Droughty Too sandy	0.00 0.00 0.02	Poor Stones	0.00	Poor Hard to reclaim (rock fragments) Rock fragments Too sandy	0.00 0.00 0.00 0.02
Cartago-----	25	Poor Too sandy Wind erosion Droughty	0.00 0.00 0.00	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.12
488173: Lubkin-----	85	Poor Wind erosion Droughty Low content of organic matter	0.00 0.01 0.12	Poor Stones Cobble content	0.00 0.70	Poor Hard to reclaim (rock fragments) Rock fragments Slope	0.00 0.00 0.00 0.84
488211: Morey family-----	40	Fair Low content of organic matter Water erosion	0.12 0.90	Poor Low strength Shrink-swell	0.00 0.74	Good	
Winnedumah-----	25	Not rated		Not rated		Fair Rock fragments	0.97
Rindge family-----	20	Poor Too alkaline Sodium content	0.00 0.78	Fair Wetness	0.76	Fair Organic matter content low Wetness Sodium content	0.02 0.76 0.78
488326: Winnedumah-----	85	Poor Wind erosion Sodium content Low content of organic matter	0.00 0.00 0.12	Good		Poor Sodium content Salinity	0.00 0.00

Soil Survey of Manzanar National Historic Site, California

Table 15.—Ponds and Embankments

(Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table)

Map unit symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
488047: Cartago-----	85	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Depth to water	1.00
488048: Cartago-----	85	Very limited Seepage Slope	1.00 0.08	Very limited Seepage	1.00	Very limited Depth to water	1.00
488091: Goodale-----	60	Very limited Seepage	1.00	Very limited Seepage Large stones	1.00 0.12	Very limited Depth to water	1.00
Cartago-----	25	Very limited Seepage	1.00	Very limited Seepage	1.00	Very limited Depth to water	1.00
488173: Lubkin-----	85	Very limited Seepage Slope	1.00 1.00	Very limited Seepage Large stones	1.00 0.63	Very limited Depth to water	1.00
488211: Morey family-----	40	Somewhat limited Seepage	0.04	Somewhat limited Piping	0.35	Very limited Depth to water	1.00
Winnedumah-----	25	Somewhat limited Seepage	0.04	Not rated		Not rated	
Rindge family-----	20	Somewhat limited Seepage	0.72	Very limited Seepage Hard to pack Depth to saturated zone	1.00 1.00 0.95	Somewhat limited Slow refill Unstable excavation walls Depth to saturated zone	0.28 0.10 0.02
488326: Winnedumah-----	85	Very limited Seepage	1.00	Very limited Piping Salinity	1.00 0.50	Somewhat limited Depth to saturated zone Salinity and saturated zone Unstable excavation walls	0.96 0.78 0.10

Table 16.—Engineering Properties

(Absence of an entry indicates that data were not estimated)

Map unit symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					in	in						
	In				Pct	Pct					Pct	
488047: Cartago-----	0-14	Gravelly loamy sand	SM, SP-SM, GM, GP-GM	A-2, A-1	0	0-5	55-80	50-75	25-55	10-25	0-0	NP
	14-33	Very gravelly loamy sand, gravelly loamy sand	SM, SP-SM, GM, GP-GM	A-3, A-2, A-1	0	0-5	35-80	30-75	15-55	5-25	0-0	NP
	33-53	Gravelly sandy loam	SC-SM, SM, GC-GM, GM	A-2, A-1	0	0-5	55-80	50-75	30-50	15-30	15-20	NP-5
	53-60	Loamy fine sand	SP-SM, SM	A-3, A-2, A-1	0	0-5	90-100	85-100	45-75	5-15	0-0	NP
488048: Cartago-----	0-14	Gravelly loamy sand	SM, SP-SM, GM, GP-GM	A-2, A-1	0	0-5	55-80	50-75	25-55	10-25	0-0	NP
	14-33	Very gravelly loamy sand, gravelly loamy sand	SM, SP-SM, GM, GP-GM	A-3, A-2, A-1	0	0-5	35-80	30-75	15-55	5-25	0-0	NP
	33-53	Gravelly sandy loam	SC-SM, SM, GC-GM, GM	A-2, A-1	0	0-5	55-80	50-75	30-50	15-30	15-20	NP-5
	53-60	Loamy fine sand	SP-SM, SM	A-3, A-2, A-1	0	0-5	90-100	85-100	45-75	5-15	0-0	NP
488091: Goodale-----	0-2	Very gravelly loamy sand	SM, SP-SM, GM, GP-GM	A-1	0-5	0-5	40-60	30-50	20-30	5-20	0-0	NP
	2-16	Very gravelly loamy sand, gravelly loamy sand	SM, SP-SM, GM, GP-GM	A-3, A-2, A-1	0-5	0-5	35-80	30-75	15-55	5-20	0-0	NP
	16-22	Extremely gravelly loamy sand	GP-GM, GP	A-1	0-5	0-5	20-30	15-25	5-15	0-5	0-0	NP
	22-60	Extremely stony loamy sand	GP-GM, GP	A-1	40-55	10-20	20-25	15-20	5-15	0-5	0-0	NP
Cartago-----	0-10	Gravelly loamy sand	SP-SM, SM	A-1	0-5	0-5	65-80	50-75	30-50	10-20	0-0	NP
	10-60	Gravelly loamy sand, cobble loamy sand	SP-SM, SM	A-1	0-5	10-30	65-80	50-75	30-50	10-20	0-0	NP

Table 16.—Engineering Properties—Continued

Map unit symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 in	3-10 in	4	10	40	200		
	In				Pct	Pct					Pct	
488173:												
Lubkin-----	0-5	Gravelly loamy sand	SM	A-2, A-1	0-15	0-15	65-90	55-80	30-60	15-25	0-0	NP
	5-26	Very bouldery sandy loam, very stony sandy loam, very stony loamy sand	GM	A-2	10-15	10-40	55-65	45-55	35-45	25-35	20-30	NP-5
	26-46	Very bouldery loamy sand, very stony loamy sand, very cobbly loamy sand	SM, SP-SM, GM, GP-GM	A-1	10-40	10-40	55-65	45-55	25-45	10-15	0-0	NP
	46-60	Very gravelly loamy sand	SP-SM, GP-GM	A-1	0-15	0-15	35-60	25-50	15-30	5-10	0-0	NP
488211:												
Morey family----	0-5	Silt loam	CL-ML, CL	A-4	0	0	100	100	90-100	70-90	20-25	5-10
	5-18	Silt loam, clay loam, loam	CL	A-6	0	0	100	95-100	85-100	60-90	25-40	10-20
	18-48	Clay loam, loam, silt loam	CL	A-6	0	0	100	95-100	85-100	60-90	25-40	10-20
	48-60	Clay loam, sandy clay loam	CL	A-7	0	0	80-100	75-100	70-95	55-80	40-45	20-25
Winnedumah-----	0-3	Loam	CL-ML, CL	A-4	---	0	100	100	85-95	60-75	20-30	5-10
	3-44	Loam, sandy clay loam	SC, SM, CL, ML	A-6, A-4	---	0	80-100	75-100	60-90	40-65	30-40	5-15
	44-60	Sandy clay loam	SM, SC-SM, SC	A-4	---	0	80-100	75-100	60-90	35-50	25-35	5-10
Rindge family---	0-34	Mucky loam	CL-ML, CL	A-4	0	0	100	100	85-95	60-75	25-30	5-10
	34-55	Loam, clay loam	CL-ML, CL	A-6, A-4	0	0	80-100	75-100	65-95	50-75	25-35	5-15
	55-60	Sandy clay loam	SC, SC-SM, CL, CL-ML	A-6, A-4, A-2	0	0	80-100	75-100	60-90	30-55	25-35	5-15
488326:												
Winnedumah-----	0-14	Fine sandy loam	SC-SM, SM, CL-ML, ML	A-4	0	0	100	95-100	70-85	40-55	20-25	NP-5
	14-29	Loam, fine sandy loam	SM, ML	A-4	0	0	100	95-100	75-95	45-70	20-35	NP-10
	29-38	Sandy clay loam	SC, CL	A-6	0	0	100	95-100	79-90	35-55	25-40	10-20
	38-60	Fine sandy loam	SC-SM, SM, CL-ML, ML	A-4	0	0	100	95-100	70-85	40-55	20-25	NP-5

Table 17.—Physical Soil Properties

(Sand, silt, and clay values are shown either as a range or as a representative value)

Map unit symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permeability	Available water capacity	Shrink- swell potential	Organic matter
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct
488047:									
Cartago-----	0-14	79	16	2-8	1.55-1.65	6.0-20.0	0.05-0.07	0.0-2.9	0.0-0.5
	14-33	79	16	2-8	1.55-1.65	6.0-20.0	0.03-0.07	0.0-2.9	0.0-0.5
	33-53	67	23	8-12	1.50-1.60	2.0-6.0	0.08-0.11	0.0-2.9	0.0-0.5
	53-60	79	16	2-8	1.55-1.65	6.0-20.0	0.06-0.09	0.0-2.9	0.0-0.5
488048:									
Cartago-----	0-14	79	16	2-8	1.55-1.65	6.0-20.0	0.05-0.07	0.0-2.9	0.0-0.5
	14-33	79	16	2-8	1.55-1.65	6.0-20.0	0.03-0.07	0.0-2.9	0.0-0.5
	33-53	67	23	8-12	1.50-1.60	2.0-6.0	0.08-0.11	0.0-2.9	0.0-0.5
	53-60	79	16	2-8	1.55-1.65	6.0-20.0	0.06-0.09	0.0-2.9	0.0-0.5
488091:									
Goodale-----	0-2	84	9	5-10	1.60-1.70	6.0-20.0	0.02-0.04	0.0-2.9	0.5-1.0
	2-16	84	9	5-10	1.60-1.70	6.0-20.0	0.02-0.07	0.0-2.9	0.0-0.5
	16-22	84	9	5-10	1.60-1.70	6.0-20.0	0.01-0.03	0.0-2.9	0.0-0.5
	22-60	84	9	5-10	1.60-1.70	6.0-20.0	0.01-0.03	0.0-2.9	0.0-0.5
Cartago-----	0-10	84	9	3-10	1.60-1.70	6.0-20.0	0.04-0.06	0.0-2.9	0.0-0.5
	10-60	84	9	3-10	1.60-1.70	6.0-20.0	0.04-0.06	0.0-2.9	0.0-0.5
488173:									
Lubkin-----	0-5	84	9	5-10	1.60-1.70	6.0-20.0	0.05-0.07	0.0-2.9	0.5-1.0
	5-26	67	19	10-18	1.55-1.70	2.0-6.0	0.04-0.07	0.0-2.9	0.0-0.5
	26-46	84	9	5-10	1.60-1.75	6.0-20.0	0.03-0.06	0.0-2.9	0.0-0.5
	46-60	84	9	5-10	1.60-1.75	6.0-20.0	0.03-0.05	0.0-2.9	0.0-0.5
488211:									
Morey family----	0-5	14	71	12-18	1.45-1.55	0.6-2.0	0.15-0.20	0.0-2.9	4.0-10
	5-18	27	45	20-35	1.40-1.55	0.2-0.6	0.14-0.21	3.0-5.9	2.0-4.0
	18-48	27	45	20-35	1.40-1.55	0.2-0.6	0.14-0.21	3.0-5.9	0.0-0.5
	48-60	47	15	35-40	1.40-1.50	0.2-0.6	0.15-0.21	6.0-8.9	0.0-0.5
Winnedumah-----	0-3	43	40	15-20	1.45-1.55	0.6-2.0	0.13-0.17	0.0-2.9	1.0-5.0
	3-44	38	36	22-30	1.45-1.55	0.2-0.6	0.14-0.18	3.0-5.9	1.0-2.0
	44-60	55	17	20-35	1.45-1.55	0.2-0.6	0.14-0.18	0.0-2.9	0.0-0.5
Rindge family----	0-34	43	38	15-22	0.20-0.30	0.6-2.0	0.14-0.18	0.0-2.9	22-35
	34-55	39	37	18-30	1.50-1.60	0.2-0.6	0.16-0.19	3.0-5.9	2.0-8.0
	55-60	57	18	20-30	1.45-1.55	0.2-0.6	0.14-0.18	3.0-5.9	1.0-2.0
488326:									
Winnedumah-----	0-14	65	20	12-18	1.35-1.55	2.0-6.0	0.08-0.12	0.0-2.9	2.0-5.0
	14-29	43	38	12-25	1.30-1.65	0.6-2.0	0.08-0.14	0.0-2.9	0.5-1.0
	29-38	55	17	20-35	1.45-1.55	0.2-0.6	0.08-0.11	3.0-5.9	0.0-0.5
	38-60	65	20	12-18	1.15-1.50	2.0-6.0	0.11-0.14	0.0-2.9	0.0-0.5

Soil Survey of Manzanar National Historic Site, California

Table 18.—Erosion Properties

(Entries under "Erosion factors" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

Map unit symbol and soil name	Depth (inches)	Erosion factors			Wind erodi- bility group	Wind erodi- bility index	
		Kw	Kf	T			
488047: Cartago-----	0-14	.10	.24	5	2	134	
	14-33	.10	.28				
	33-53	.15	.28				
	53-60	.32	.32				
488048: Cartago-----	0-14	.10	.24	5	2	134	
	14-33	.10	.28				
	33-53	.15	.28				
	53-60	.32	.32				
488091: Goodale-----	0-2	.05	.15	5	3	86	
	2-16	.05	.15				
	16-22	.05	.15				
	22-60	.05	.15				
	Cartago-----	0-10	.10	.15	5	2	134
		10-60	.05	.15			
488173: Lubkin-----	0-5	.10	.17	5	2	134	
	5-26	.10	.24				
	26-46	.05	.15				
	46-60	.05	.15				
488211: Morey family-----	0-5	.43	.43	5	5	56	
	5-18	.37	.37				
	18-48	.43	.43				
	48-60	.37	.37				
	Winnedumah-----	0-3	.28	.28	5	4L	86
		3-44	.37	.37			
		44-60	.37	.37			
	Rindge family-----	0-34	.28	.28	1	5	56
		34-55	.28	.28			
		55-60	.24	.24			
488326: Winnedumah-----	0-14	.24	.24	5	2	134	
	14-29	.32	.32				
	29-38	.28	.28				
	38-60	.24	.24				

Soil Survey of Manzanar National Historic Site, California

Table 19.—Total Soil Carbon

(This table displays soil organic carbon (SOC) and soil inorganic carbon (SIC) in kilograms per square meter to a depth of 2 meters or to the representative top depth of any kind of bedrock or any cemented soil horizon. SOC and SIC are reported on a volumetric whole soil basis, corrected for representative rock fragments indicated in the database. SOC is converted from horizon soil organic matter of the fraction of the soil less than 2 mm in diameter. If soil organic matter indicated in the database is NULL, SOC is assumed to be zero. SIC is converted from horizon calcium carbonate content fraction of the soil less than 2 mm in diameter. If horizon calcium carbonate indicated in the database is NULL, SIC is assumed to be zero. A weighted average of all horizons is used in the calculations. Only major components of a map unit are displayed in this table)

Map unit symbol, component name, and component percent	SOC	SIC
	kg/m ²	kg/m ²
488047: Cartago (85%)-----	3	0
488048: Cartago (85%)-----	3	0
488091: Goodale (60%)-----	1	0
Cartago (25%)-----	2	0
488173: Lubkin (85%)-----	2	0
488211: Morey family (40%)-----	18	4
Winnedumah (25%)-----	0	0
Rindge family (20%)-----	59	1
488326: Winnedumah (85%)-----	14	8

Soil Survey of Manzanar National Historic Site, California

Table 20.—Chemical Soil Properties

Map unit symbol and soil name	Depth	Cation- exchange capacity	Soil reaction	Calcium carbon- ate	Salinity	Sodium adsorp- tion ratio
	In	meq/100 g	pH	Pct	mmhos/cm	
488047:						
Cartago-----	0-14	1.0-5.0	6.6-7.8	0	0	0
	14-33	1.0-5.0	6.6-7.8	0	0	0
	33-53	5.0-10.0	6.6-7.8	0	0	0
	53-60	1.0-5.0	6.6-7.8	0	0	0
488048:						
Cartago-----	0-14	1.0-5.0	6.6-7.8	0	0	0
	14-33	1.0-5.0	6.6-7.8	0	0	0
	33-53	5.0-10.0	6.6-7.8	0	0	0
	53-60	1.0-5.0	6.6-7.8	0	0	0
488091:						
Goodale-----	0-2	5.0-10.0	6.6-7.8	0	0	0
	2-16	5.0-10.0	6.6-7.8	0	0	0
	16-22	5.0-10.0	6.6-7.8	0	0	0
	22-60	5.0-10.0	6.6-7.8	0	0	0
Cartago-----	0-10	5.0-10.0	6.6-7.8	0	0	0
	10-60	5.0-10.0	6.6-7.8	0	0	0
488173:						
Lubkin-----	0-5	5.0-10.0	6.6-7.8	0	0	0
	5-26	5.0-15.0	6.6-7.8	0	0	0
	26-46	5.0-10.0	6.6-7.8	0	0	0
	46-60	5.0-10.0	6.6-7.8	0	0	0
488211:						
Morey family-----	0-5	20.0-40.0	6.6-8.4	0-1	2.0-8.0	1-5
	5-18	20.0-40.0	6.6-7.8	1-2	0.0-2.0	0-5
	18-48	20.0-40.0	6.6-7.8	1-2	0.0-2.0	0-5
	48-60	15.0-40.0	5.6-7.3	0	0.0-2.0	0
Winnedumah-----	0-3	10.0-15.0	7.9-9.0	1-5	2.0-4.0	1-13
	3-44	10.0-25.0	7.4-8.4	1-5	0.0-2.0	1-5
	44-60	10.0-35.0	6.6-7.8	1-5	0.0-2.0	1-5
Rindge family-----	0-34	50.0-70.0	8.5-9.0	1-5	2.0-16.0	1-13
	34-55	15.0-45.0	7.4-7.8	0	0.0-2.0	0
	55-60	10.0-35.0	6.6-7.3	0	0.0-2.0	0
488326:						
Winnedumah-----	0-14	5.0-15.0	8.5-9.6	1-5	8.0-16.0	13-100
	14-29	5.0-20.0	7.9-9.0	1-5	8.0-16.0	13-40
	29-38	10.0-35.0	7.9-8.4	1-5	4.0-16.0	13-35
	38-60	5.0-15.0	6.6-7.8	1-5	0.0-4.0	1-13

Table 21.—Water Features

(See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated)

Map unit symbol and soil name	Hydro- logic group	Months	Water table			Ponding		Flooding	
			Upper limit Ft	Lower limit Ft	Surface water depth Ft	Duration	Frequency	Duration	Frequency
488047: Cartago-----	A	January	---	---	---	---	None	Very brief	Rare
		February	---	---	---	---	None	Very brief	Rare
		March	---	---	---	---	None	Very brief	Rare
		April	---	---	---	---	None	Very brief	Rare
		May	---	---	---	---	None	Very brief	Rare
		June	---	---	---	---	None	Very brief	Rare
		July	---	---	---	---	None	Very brief	Rare
		August	---	---	---	---	None	Very brief	Rare
		September	---	---	---	---	None	Very brief	Rare
		October	---	---	---	---	None	Very brief	Rare
		November	---	---	---	---	None	Very brief	Rare
		December	---	---	---	---	None	Very brief	Rare
488048: Cartago-----	A	January	---	---	---	---	None	Very brief	Rare
		February	---	---	---	---	None	Very brief	Rare
		March	---	---	---	---	None	Very brief	Rare
		April	---	---	---	---	None	Very brief	Rare
		May	---	---	---	---	None	Very brief	Rare
		June	---	---	---	---	None	Very brief	Rare
		July	---	---	---	---	None	Very brief	Rare
		August	---	---	---	---	None	Very brief	Rare
		September	---	---	---	---	None	Very brief	Rare
		October	---	---	---	---	None	Very brief	Rare
		November	---	---	---	---	None	Very brief	Rare
		December	---	---	---	---	None	Very brief	Rare

Table 21.--Water Features--Continued

Map unit symbol and soil name	Hydro- logic group	Months	Water table			Ponding		Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
488091: Goodale-----	A	January	---	---	---	---	None	Very brief	Rare
		February	---	---	---	---	None	Very brief	Rare
		March	---	---	---	---	None	Very brief	Rare
		April	---	---	---	---	None	Very brief	Rare
		May	---	---	---	---	None	Very brief	Rare
		June	---	---	---	---	None	Very brief	Rare
		July	---	---	---	---	None	Very brief	Rare
		August	---	---	---	---	None	Very brief	Rare
		September	---	---	---	---	None	Very brief	Rare
		October	---	---	---	---	None	Very brief	Rare
		November	---	---	---	---	None	Very brief	Rare
		December	---	---	---	---	None	Very brief	Rare
Cartago-----	A	January	---	---	---	---	None	Very brief	Rare
		February	---	---	---	---	None	Very brief	Rare
		March	---	---	---	---	None	Very brief	Rare
		April	---	---	---	---	None	Very brief	Rare
		May	---	---	---	---	None	Very brief	Rare
		June	---	---	---	---	None	Very brief	Rare
		July	---	---	---	---	None	Very brief	Rare
		August	---	---	---	---	None	Very brief	Rare
		September	---	---	---	---	None	Very brief	Rare
		October	---	---	---	---	None	Very brief	Rare
		November	---	---	---	---	None	Very brief	Rare
		December	---	---	---	---	None	Very brief	Rare
488173: Lubkin-----	A	January	---	---	---	---	None	Very brief	Rare
		February	---	---	---	---	None	Very brief	Rare
		March	---	---	---	---	None	Very brief	Rare
		April	---	---	---	---	None	Very brief	Rare
		May	---	---	---	---	None	Very brief	Rare
		June	---	---	---	---	None	Very brief	Rare
		July	---	---	---	---	None	Very brief	Rare
		August	---	---	---	---	None	Very brief	Rare
		September	---	---	---	---	None	Very brief	Rare
		October	---	---	---	---	None	Very brief	Rare
		November	---	---	---	---	None	Very brief	Rare
		December	---	---	---	---	None	Very brief	Rare

Table 21.—Water Features—Continued

Map unit symbol and soil name	Hydro- logic group	Months	Water table		Surface water depth	Ponding		Flooding	
			Upper limit	Lower limit		Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
488211: Morey family-----	C	Jan-Dec	---	---	---	---	None	---	None
Winnedumah-----	B	Jan-Dec	---	---	---	---	None	---	None
Rindge family-----	C	January	2.3	>6.0	---	---	None	---	None
		February	2.3	>6.0	---	---	None	---	None
		March	2.3	>6.0	---	---	None	---	None
		April	2.3	>6.0	---	---	None	---	None
		May-Nov	---	---	---	---	None	---	None
		December	2.3	>6.0	---	---	None	---	None
488326: Winnedumah-----	B	Jan-Feb	---	---	---	---	None	---	None
		March	4.5	>6.0	---	---	None	---	None
		April	4.5	>6.0	---	---	None	---	None
		May	4.5	>6.0	---	---	None	---	None
		Jun-Dec	---	---	---	---	None	---	None

Table 22.—Soil Features

(See text for definitions of terms used in this table. Absence of an entry indicates that data were not estimated)

Map unit symbol and soil name	Restrictive layer			Potential for frost action	Risk of corrosion	
	Kind	Depth to top <u>In</u>	Hardness		Uncoated steel	Concrete
488047: Cartago-----	None	---	---	Low	Low	Low
488048: Cartago-----	None	---	---	Low	Low	Low
488091: Goodale-----	None	---	---	Low	Low	Low
Cartago-----	None	---	---	Low	Low	Low
488173: Lubkin-----	None	---	---	Low	Moderate	Low
488211: Morey family-----	None	---	---	Low	Low	Low
Winnedumah-----	None	---	---	Moderate	High	Low
Rindge family-----	None	---	---	Low	High	Not rated
488326: Winnedumah-----	None	---	---	Moderate	High	Low

Soil Survey of Manzanar National Historic Site, California

Table 23.—Taxonomic Classification of the Soils

Soil name	Family or higher taxonomic class
Cartago-----	Sandy, mixed, thermic Xeric Torriorthents
Goodale-----	Sandy-skeletal, mixed, thermic Xeric Torriorthents
Lubkin-----	Loamy-skeletal, mixed, superactive, thermic Xeric Haplargids
Morey family-----	Fine-silty, mixed, superactive, thermic Typic Argiaquolls
Rindge family-----	Euic, thermic Typic Medisaprists
Winnedumah-----	Fine-loamy, mixed, superactive, thermic Xeric Haplargids

Soil Survey of Manzanar National Historic Site, California

Table 24.—Soil Classification Key

(An asterisk in the first column indicates a taxadjunct to the series. See text for a description of those characteristics that are outside the range of the series)

ORDER	
Suborder	
Great Group	
Subgroup	
Series or Higher Category	

ARIDISOLS	
Argids	
Haplargids	
Xeric Haplargids	
Winnedumah-----	Fine-loamy, mixed, superactive, thermic Xeric Haplargids
Lubkin-----	Loamy-skeletal, mixed, superactive, thermic Xeric Haplargids
ENTISOLS	
Orthents	
Torriorthents	
Xeric Torriorthents	
Cartago-----	Sandy, mixed, thermic Xeric Torriorthents
Goodale-----	Sandy-skeletal, mixed, thermic Xeric Torriorthents
HISTOSOLS	
Saprists	
Medisaprists	
Typic Medisaprists	
Rindge family-----	Euic, thermic Typic Medisaprists
MOLLISOLS	
Aquolls	
Argiaquolls	
Typic Argiaquolls	
Morey family-----	Fine-silty, mixed, superactive, thermic Typic Argiaquolls

Appendix

**UNITED STATES DEPARTMENT OF AGRICULTURE
NATURAL RESOURCES CONSERVATION SERVICE**

RANGE/ECOLOGICAL SITE DESCRIPTION

ECOLOGICAL SITE CHARACTERISTICS

Site Type: Rangeland

Site Name: Gravelly Loamy Sand

Site ID: 029XG018CA

Major Land Resource Area: 029G Southern Nevada Basin and Range

Physiographic Features:

This site occurs on alluvial fans and fan terraces. Elevations are 3700 to 4500 feet. Slopes range from 0 to 15 percent but slopes from 2 to 5 percent are most typical.

Landform: alluvial fans and fan terraces

Aspect:

	<u>Minimum</u>	<u>Maximum</u>
<u>Site elevation (feet):</u>	3700	4500
<u>Slope (percent):</u>	0	15 (mostly 2 to 5%)
<u>Water table depth (inches):</u>		

Flooding

Frequency:

Duration:

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Influencing Water Features

Cowardin wetland classification system: System(s) Subsystem(s) Class(es)

Rosgen stream code:

More than one stream type may be associated with the site.

Representative Soil Features

The soils that characterize this site are very deep and well-drained to somewhat excessively drained. They are formed in granitic alluvium. Surface textures are gravelly loamy sands. Available water capacities range from low to moderate and the hazard of water erosion is slight to moderate. Wind erosion hazard is slight to severe. Effective rooting depth is 4 to 60 inches.

Benton – Owens Valley Soil Survey

SSA	Map Unit Symbol	Component Name
802	413	Goodale – Cartago complex, 2 – 5% slopes
	461	Goodale bouldery – Cartago complex, 2 – 5% slopes
	482	Cartago gravelly loamy sand, 2 – 5% slopes
	490A	Lubkin – Tinemaha complex, 2 – 5% slopes
	492	Lubkin gravelly loamy sand, 5 – 15% slopes
	510	Inyo gravelly loamy coarse sand, 0 – 5% slopes
	512	Inyo sand, 5 – 15% slopes
	520	Dehy – Typic Argiaquolls – Lubkin association, 2 – 9% slopes

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Representative soil features include:

Predominant Parent materials:

Kind: granitic alluvium

Origin:

Surface texture: sand

Subsurface Texture Group:

	<u>Minimum</u>	<u>Maximum</u>
<u>Surface fragments <=3" (% Cover):</u>		
<u>Surface fragments >3" (% Cover):</u>		
<u>Subsurface fragments <=3" (% Volume):</u>		
<u>Subsurface fragments >3" (% Volume):</u>		
<u>Drainage class:</u>	well	somewhat excessively
<u>Permeability Class:</u>		
<u>Depth (inches):</u>	4	60+
<u>Electrical conductivity (mmhos/cm):</u>		
<u>Sodium adsorption ratio:</u>		
<u>Calcium Carbonate Equivalent (percent):</u>		
<u>Soil reaction (1:1 Water):</u>		
<u>Soil reaction (0.01M CaCl₂):</u>		
<u>Available Water Holding Capacity:</u>	low	moderate
<u>(Total inches for entire profile):</u>		

PLANT COMMUNITIES

Ecological dynamics of the site

Overgrazing would result in a decrease in the perennial grasses, bud sagebrush, spiny hopsage, and Winterfat. Shadscale and white bursage would increase. Russian thistle and white burrobrush are invaders of this site. Areas along drainages which receive additional moisture are dominated by basin big sagebrush.

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Plant communities

The interpretative plant community is the historic climax plant community. The plant community is dominated by Shadscale, white bursage, and desert needlegrass. Potential vegetation composition is about 25 percent grasses, 10 percent forbs, and 65 percent shrubs.

Plant Community Composition

The following table lists the major plant species and percentages by weight, air-dry, of the total plant community that each contributes in an average production year.

Group	Common Name	Scientific Name	% by Weight	Group Allowable		Annual Production	
				Low	High	Low	High
Grasses and Grasslike Plants 25%							
1	Annual Grasses						
2	Perennial Grasses						
	Desert needlegrass	<i>Achnatherum speciosum</i>	10 – 20				
	Other perennial grasses *			2	5		
	Indian ricegrass	<i>Oryzopsis hymenoides</i>	*				
	Sedges and Rushes						
Forbs 10%							
3	Forbs						
	Other perennial forbs *			2	5		
	Desert trumpet	<i>Eriogonum inflatum</i>	*				
	Indian paintbrush	<i>Castilleja spp.</i>	*				
	Other annual forbs		T – 5				

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Shrubs 65%							
4	Shrubs						
	Shadscale saltbush	<i>Atriplex confertifolia</i>	25 – 35				
	White bursage (Burrobrush)	<i>Ambrosia dumosa</i>	5 – 15				
	Bud sagebrush	<i>Picrothamnus desertorum</i>	5 – 15				
	Winterfat	<i>Krascheninnikovia lanata</i>	5 – 10				
	Allscale (Cattle) saltbush	<i>Atriplex polycarpa</i>	2 – 8				
	Other shrubs **			5	15		
	Spiny hopsage	<i>Graya spinosa</i>	**				
	Nevada ephedra	<i>Ephedra nevadensis</i>	**				
	Desert alysum (Desert pepperweed)	<i>Lepidium fremontii</i>	**				
	Anderson's wolfberry (Water jacket)	<i>Lycium andersonii</i>	**				
	Basin Big Sagebrush	<i>Artemisia tridentata tridentata</i>	**				
Trees							
5	Trees						

* Allow no more than 2% of each species of this group and no more than 8% in aggregate.

** Allow no more than 5% of each species of this group and no more than 15% in aggregate.

Ground cover and structure

Approximate ground cover (basal and crown) is 10 to 25 percent.

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Soil Surface Cover

Vegetative Cover						Non-Vegetative Cover					
Basal Cover				Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
Grass/Grasslike	Forb	Shrub/Vine	Tree								
to	to	to	to	to	to	to	To	to	to	to	to

Ground Cover

Vegetative Cover						Non-Vegetative Cover					
Grass/Grasslike	Forb	Shrub/Vine	Tree	Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
to	to	to	to	to	to	to	To	to	to	to	to

Structure of Canopy Cover

	Grasses/Grasslike	Forbs	Shrubs/Vines	Trees
0.5 Feet	to	to	to	to
0.5 - 1 Foot	to	to	to	to
1.0 - 2 Feet	to	to	to	to
2.0 - 4.5 Feet	to	to	to	to
4.5 - 13 Feet	to	to	to	to
13 - 40 Feet	to	to	to	to

Annual Production by Plant Type (historic climax plant community)

Annual Production (lbs/acre)			
Plant Type	Low	Representative Value	High
Grasses/Grasslike			

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Forbs			
Shrubs/Vines			
Trees			
TOTAL	150	300	500

Plant community growth curves

Growth Curve Name:

Growth Curve Number:

Growth Curve Description:

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

ECOLOGICAL SITE INTERPRETATIONS

Animal community

Wildlife interpretations

Tule elk use this site in the spring; there is occasional winter use of this site by mule deer.

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Livestock Interpretations

Season of Use – Other Management Considerations

This site is primarily used for spring grazing especially in favorable precipitation years when annual forbs are abundant.

General guide to initial stocking rate. Before making specific recommendations, an on-site evaluation must be made.

Production Year	lbs/acre (air-dry)	AUM/acre	Acres/AUM
Favorable Years	500		
Normal Years	300		
Unfavorable Years	150		

Plant community growth curves

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Growth Curve Name:
Growth Curve Number:
Growth Curve Description:

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Hydrologic functions

Runoff is slow to moderate; the water erosion hazard is slight to moderate.

Soil Series	Hydrologic Group	Hydrologic Conditions and Runoff Curves		
		Good	Fair	Poor
Cartago	A	41	46	61
Goodale	A	41	46	61
Inyo	A	41	46	61
Lubkin	B	63	66	76
Tinemaha	B	63	66	76

Recreational uses

This site is primarily located on the City of Los Angeles, Department of Water and Power (LADWP) property and is open to the public.

Wood products

Other products

Other information

Reduce the hazard of wind erosion by avoiding overgrazing, preserving the existing plant cover, and limiting operation of off-road vehicles.

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Threatened and endangered plants and animals: Planners will refer to state and federal lists of endangered species. Management recommendations will address impacts on endangered species and their critical habitats (refer to Endangered Species Policy and California Endangered Species Handbook).

Fire is very infrequent on this site.

SUPPORTING INFORMATION

Associated sites

Principle sites that commonly occur in association with this potential plant community include:
Bouldery Fan (029XG010CA)

Similar sites

Competing sites (and their differentiae) that are similar to this potential plant community:
Loamy (029XG017CA): white bursage is only a minor occurrence; it is also a more productive site.
Sandy Fan (029XG012CA): Indian ricegrass is the dominant grass and the site is less productive than Gravelly Loamy Sand

Inventory data references

State correlation

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Type locality

Northwest ¼ of Section 16, Township 14 South, Range 35 East
Approximately five miles south of Independence, Inyo County, California

Relationship to other established classification systems

Plant Symbol: ATCO – AMDU2/STSP3
Society of American Foresters Cover Type:
Kuchler:
CalVeg:
Wildlife Habitat Relationship:
Woodland Suitability Group:
Soil Survey Area: 802 Benton – Owens Valley Soil Survey

Developed in cooperation with the Los Angeles Department of Water and Power

Other references

SITE DESCRIPTION APPROVAL

Author	Agency	Date	Approval	Date
Patti J. Novak	SCS	4/1992	Leonard Jolley	

Soil Survey of Manzanar National Historic Site, California

Gravelly Loamy Sand
029XG018CA

Authorship

Original authors' names and date: Patti J. Novak, April 1992
Revision authors' names and revision date:

Site approval

Leonard Jolley, State Range Conservationist, SCS, California

**UNITED STATES DEPARTMENT OF AGRICULTURE
NATURAL RESOURCES CONSERVATION SERVICE**

RANGE/ECOLOGICAL SITE DESCRIPTION

ECOLOGICAL SITE CHARACTERISTICS

Site Type: Rangeland

Site Name: Sandy

Site ID: 029XG011CA

Major Land Resource Area: 029G Southern Nevada Basin and Range

Physiographic Features:

This site occurs on toe slopes of alluvial fans. Elevations are 3800 to 4400 feet. Slopes range from 0 to 5 percent.

Landform: toe slopes of alluvial fans

Aspect:

	<u>Minimum</u>	<u>Maximum</u>
<u>Site elevation (feet):</u>	3800	4400
<u>Slope (percent):</u>	0	5

Water table depth (inches):

Flooding

Frequency:

Duration:

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

Ponding:

Depth (inches):

Frequency:

Duration (days):

Runoff class: slow

Climatic Features

The climate on this site is characterized by mild, cool winters (30° to 60° F) and hot, dry summers (70° to 100° F). The average annual precipitation ranges from 4 to 8 inches with most falling as rain from November to March. The average frost-free period is 150 to 200 days.

	<u>Minimum</u>	<u>Maximum</u>
<u>Frost-free period (days):</u>	150	200
<u>Freeze-free period (days):</u>		
<u>Mean annual precipitation (inches):</u>	4	8

Monthly precipitation and temperature distribution (°F)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Min.												
Precip. Max.												
Temp. Min.												
Temp. Max.												

Location of climate stations:

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

Influencing Water Features

System(s) Subsystem(s) Class(es)

Cowardin wetland classification system:

Rosgen stream code:

More than one stream type may be associated with the site.

Representative Soil Features

Benton – Owens Valley Soil Survey

SSA	Map Unit Symbol	Component Name
802	480	Cartago gravelly loamy sand, 0 – 2% slopes
	698	Herperia – Cartago complex, 0 – 5% slopes

Representative soil features include:

The soils that characterize this site are very deep and well-drained to somewhat excessively drained. They are formed in granitic alluvium. Surface textures are loamy sands. Available water capacity is low and the hazard of erosion is slight. Wind erosion hazard is moderate to severe. Effective rooting depth is 60 inches or more.

Predominant Parent materials:
Kind: granitic alluvium
Origin;
Surface texture: loamy sands
Subsurface Texture Group:

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

	<u>Minimum</u>	<u>Maximum</u>
<u>Surface fragments <=3" (% Cover):</u>		
<u>Surface fragments >3" (% Cover):</u>		
<u>Subsurface fragments <=3" (% Volume):</u>		
<u>Subsurface fragments >3" (% Volume):</u>		
<u>Drainage class:</u>	well	somewhat excessively drained
<u>Permeability Class:</u>		
<u>Depth (inches):</u>	60+	
<u>Electrical conductivity (mmhos/cm):</u>		
<u>Sodium adsorption ratio:</u>		
<u>Calcium Carbonate Equivalent (percent):</u>		
<u>Soil reaction (1:1 Water):</u>		
<u>Soil reaction (0.01M CaCl₂):</u>		
<u>Available Water Holding Capacity:</u>	low	
<u>(Total inches for entire profile):</u>		

PLANT COMMUNITIES

Ecological dynamics of the site

Overgrazing this site will cause a decrease in the perennial grasses and the more palatable shrubs. Species likely to invade this site include Russian thistle (*Salsola kali*) and white bursage (burrobrush) (*Ambrosia dumosa*). Abandoned agricultural land and other disturbed areas are initially dominated by basin big sagebrush (*Artemisia tridentata tridentata*) and rubber rabbitbrush (*Ericameria nauseosa*). Areas receiving additional moisture (drainages) are dominated by basin big sagebrush.

Plant communities

The interpretative plant community is the historic climax plant community. The plant community is dominated by four-wing saltbush, Fremont dalea, and Indian ricegrass. Potential vegetation composition is about 30% grasses, 15% forbs, and 55% shrubs.

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

Plant Community Composition

The following table lists the major plant species and percentages by weight, air-dry, of the total plant community that each contributes in an average production year.

Group	Common Name	Scientific Name	% by Weight	Group Allowable		Annual Production	
				Low	High	Low	High
Grasses and Grasslike Plants 30%							
1	Annual Grasses						
2	Perennial Grasses						
	Indian ricegrass	<i>Oryzopsis hymenoides</i>	10 – 15				
	Other perennial grasses *			5	15		
	Bottlebrush squirreltail	<i>Sitanion hystrix</i>	*				
	Desert needlegrass	<i>Achnatherum speciosum</i>	*				
	Pine bluegrass	<i>Poa scabrella</i>	*				
	Sedges and Rushes						
Forbs 15%							
3	Forbs						
	Other perennial forbs **			2	8		
	Other annual forbs **			2	8		
Shrubs 55%							
4	Shrubs						
	Four-wing saltbush	<i>Atriplex canescens</i>	15 – 25				
	Fremont's dalea	<i>Psoralea fremontii</i>	5 – 15				
	Nevada ephedra	<i>Ephedra nevadensis</i>	5 – 10				

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

	Spiny hopsage	<i>Graya spinosa</i>	5 – 10				
	Winterfat	<i>Krascheninnikovia lanata</i>	5 – 10				
	Basin Big Sagebrush	<i>Artemisia tridentata tridentata</i>	2 - 5				
	Other shrubs ***			5	10		
	Longspine horsebrush	<i>Tetradymia axillaris</i>	***				
	Nevada dalea	<i>Psoralea polydenius</i>	***				
	Rubber rabbitbrush	<i>Ericameria nauseosa</i>	***				
	Shadscale saltbush	<i>Atriplex confertifolia</i>	***				
Trees							
5	Trees						

- * Allow no more than 5% of each species of this group and no more than 15% in aggregate.
- ** Allow no more than 2% of each species of this group and no more than 8% in aggregate.
- *** Allow no more than 5% of each species of this group and no more than 10% in aggregate.

Ground cover and structure

Approximate ground cover (basal and crown) is 20 to 35 percent.

Soil Surface Cover

Vegetative Cover				Non-Vegetative Cover							
Basal Cover				Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 – 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
Grass/ Grasslike	Forb	Shrub/ Vine	Tree								
to	to	to	to	to	to	to	to	to	to	to	to

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

Ground Cover

Vegetative Cover						Non-Vegetative Cover					
Grass/ Grasslike	Forb	Shrub/ Vine	Tree	Non- Vascular Plants	Biological Crust	Litter	Surface Fragments >¼ - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
to	to	to	to	to	to	to	to	to	to	to	to

Structure of Canopy Cover

	Grasses/Grasslike	Forbs	Shrubs/Vines	Trees
0.5 Feet	to	to	To	to
0.5 – 1 Foot	to	to	To	to
1.0 – 2 Feet	to	to	To	to
2.0 – 4.5 Feet	to	to	To	to
4.5 – 13 Feet	to	to	To	to
13 – 40 Feet	to	to	To	to

Annual Production by Plant Type (historic climax plant community)

Annual Production (lbs/acre)			
Plant Type	Low	Representative Value	High
Grasses/Grasslike			
Forbs			
Shrubs/Vines			
Trees			
TOTAL	300	550	700

Plant community growth curves

Growth Curve Name:
Growth Curve Number:
Growth Curve Description:

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

ECOLOGICAL SITE INTERPRETATIONS

Animal community

Wildlife interpretations

Occasional use by Tule elk.

Livestock Interpretations

Season of Use – Other Management Considerations

This site is primarily used for spring grazing when annual forbs are abundant.

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

General guide to initial stocking rate. Before making specific recommendations, an on-site evaluation must be made.

Production Year	lbs/acre (air-dry)	AUM/acre	Acres/AUM
Favorable Years	700		
Normal Years	550		
Unfavorable Years	300		

Plant community growth curves

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Growth Curve Name:
Growth Curve Number:
Growth Curve Description:

Hydrologic functions

Runoff is slow. The erosion hazard is slight.

Soil Series	Hydrologic Group	Hydrologic Conditions and Runoff Curves		
		Good	Fair	Poor
Cartago	A	41	46	61

Recreational uses

This site is primarily located on the City of Los Angeles, Department of Water and Power (LADWP) property and is open to the public.

Wood products

Other products

Other information

Reduce the hazard of wind erosion by avoiding overgrazing, preserving the existing plant cover, and limiting operation of off-road vehicles.

Threatened and endangered plants and animals: Planners will refer to state and federal lists of endangered species. Management recommendations will address impacts on endangered species and their critical habitats (refer to Endangered Species Policy and California Endangered Species Handbook).

Fire is very infrequent at this site.

SUPPORTING INFORMATION

Associated sites

Principle sites that commonly occur in association with this potential plant community include:
Bouldery Fan (029XG010CA)

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

Lava Flow (029XG004)

Similar sites

Competing sites (and their differentiae) that are similar to this potential plant community: None.

Inventory data references

State correlation

Type locality

Southeast ¼ of Section 21, Township 11 South, Range 34 East.
Approximately ½ mile west of Aberdeen, Inyo County, California.

Relationship to other established classification systems

Plant Symbol: ATCA2 – PSARM/ORHY
Society of American Foresters Cover Type:
Kuchler:
CalVeg:
Wildlife Habitat Relationship:
Woodland Suitability Group:
Soil Survey Area: 802 Benton – Owens Valley Soil Survey

Developed in cooperation with the Los Angeles Department of Water and Power

Soil Survey of Manzanar National Historic Site, California

Sandy
029XG011CA

Other references

SITE DESCRIPTION APPROVAL

Author	Agency	Date	Approval	Date
Patti J. Novak	SCS	3/1992	Leonard Jolley	

Authorship

Original authors' names and date: Patti J. Novak, March 1992
Revision authors' names and revision date:

Site approval

Leonard Jolley, State Range Conservationist, SCS, California

**UNITED STATES DEPARTMENT OF AGRICULTURE
NATURAL RESOURCES CONSERVATION SERVICE**

RANGE/ECOLOGICAL SITE DESCRIPTION

ECOLOGICAL SITE CHARACTERISTICS

Site Type: Rangeland

Site Name: Sodic Fan

Site ID: 029XG005CA

Major Land Resource Area: 029G Southern Nevada Basin and Range

Physiographic Features:

This site occurs on stream and lacustrine terraces. Elevations are 3650 to 4000 feet. Slopes range from 0 to 2 percent.

Landform: lacustrine terraces

Aspect:

	<u>Minimum</u>	<u>Maximum</u>
<u>Site elevation (feet):</u>	3650	4000
<u>Slope (percent):</u>	0	2
<u>Water table depth (inches):</u>	40	80

Flooding

Frequency:

Duration:

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Ponding:

Depth (inches):

Frequency:

Duration (days):

Runoff class: slow

Climatic Features

The climate on this site is characterized by mild, cool winters (30° to 60° F) and hot, dry summers (70° to 100° F). The average annual precipitation ranges from 4 to 6 inches with most falling as rain from November to March. The average frost-free period is 160 to 225 days.

	<u>Minimum</u>	<u>Maximum</u>
<u>Frost-free period (days):</u>	160	225
<u>Freeze-free period (days):</u>		
<u>Mean annual precipitation (inches):</u>	4	6

Monthly precipitation and temperature distribution (°F)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Min.												
Precip. Max.												
Temp. Min.												
Temp. Max.												

Location of climate stations:

Influencing Water Features

Cowardin wetland classification system: System(s) Subsystem(s) Class(es)

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Rosgen stream code:

More than one stream type may be associated with the site.

Representative Soil Features

The soils that characterize this site are very deep and somewhat poorly drained. They are formed in mixed alluvium and lacustrine sediments. Surface textures are silt loams and fine sandy loams. Available water capacity is moderate to high and the hazard of water erosion is slight. Wind erosion hazard is severe. Effective rooting depth is 60 inches or more. Water tables are 40 to 80 inches from March 1 to May 15. The Winnedumah soil is calcareous and saline/sodic.

Benton – Owens Valley Soil Survey

SSA	Map Unit Symbol	Component Name
802	450	Morey family – Winnedumah, drained – Rindge family complex, 0 – 2% slopes
	630	Winnedumah silt loam, saline - sodic, 0 – 2 % slopes
	631	Winnedumah fine sandy loam, saline – sodic, 0 – 2% slopes
	636	Manzanar – Winnedumah association, 0 – 2 % slopes

Representative soil features include:

Predominant Parent materials:

Kind: mixed alluvium and lacustrine sediments

Origin;

Surface texture: silt loams and fine sandy loams

Subsurface Texture Group:

Minimum Maximum

Surface fragments <=3" (% Cover):

Surface fragments >3" (% Cover):

Subsurface fragments <=3" (% Volume):

Subsurface fragments >3" (% Volume):

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Drainage class: somewhat poorly drained
Permeability Class:
Depth (inches): 60+”
Electrical conductivity (mmhos/cm):
Sodium adsorption ratio:
Calcium Carbonate Equivalent (percent):
Soil reaction (1:1 Water):
Soil reaction (0.01M CaCl₂):
Available Water Holding Capacity: moderate high
(Total inches for entire profile):

PLANT COMMUNITIES

Ecological dynamics of the site

As ecological condition deteriorates, the perennial grasses will decrease and watersage saltbush will become solely dominant.

Plant communities

The interpretative plant community is the historic climax plant community. The plant community is dominated by Nevada saltbush and alkali sacaton. Potential vegetation composition is about 25% grasses, 5% forbs, and 70% shrubs.

Plant Communities

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Plant Community Composition

The following table lists the major plant species and percentages by weight, air-dry, of the total plant community that each contributes in an average production year.

Group	Common Name	Scientific Name	% by Weight	Group Allowable		Annual Production	
				Low	High	Low	High
Grasses and Grasslike Plants							
1	Annual Grasses						
2	Perennial Grasses						
	Alkali sacaton	<i>Sporobolus airoides</i>	10 – 20				
	Inland saltgrass	<i>Distichlis spicata</i>	5 – 10				
	Basin wildrye	<i>Leymus cinereus</i>	5 – 10				
	Other perennial grasses *			2	5		
	Sedges and Rushes						
	Baltic rush*	<i>Juncus balticus</i>					
Forbs							
3	Forbs						
	Other perennial forbs *			2	5		
	American licorice *	<i>Glycyrrhiza lepidota</i>					
Shrubs							
4	Shrubs						
	Watersage (Torrey's) saltbush	<i>Atriplex torreyi</i>	40 -55				
	Black greasewood	<i>Sarcobatus vermiculatus</i>	5 – 15				
	Shadscale saltbush	<i>Atriplex confertifolia</i>	2 – 5				
	Other shrubs**			5	10		

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

	Rubber rabbitbrush **	<i>Ericameria nauseosa</i>					
	Basin Big Sagebrush **	<i>Artemisia tridentata tridentata</i>					
	Mojave seablight **	<i>Suaeda moquinii</i>					
Trees							
5	Trees						

* Allow no more than 2% of each species of this group and no more than 5% in aggregate.

** Allow no more than 5% of each species of this group and no more than 10% in aggregate.

Ground cover and structure

Approximate ground cover (basal and crown) is 20 to 40 percent.

Soil Surface Cover

Vegetative Cover						Non-Vegetative Cover					
Basal Cover				Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
Grass/ Grasslike	Forb	Shrub/ Vine	Tree								
to	to	to	To	To	to	to	to	to	to	to	to

Ground Cover

Vegetative Cover						Non-Vegetative Cover					
Grass/ Grasslike	Forb	Shrub/ Vine	Tree	Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
to	to	to	to	To	to	to	to	to	to	to	to

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Structure of Canopy Cover

	Grasses/Grasslike	Forbs	Shrubs/Vines	Trees
0.5 Feet	To	to	to	to
0.5 – 1 Foot	To	to	to	to
1.0 – 2 Feet	To	to	to	to
2.0 – 4.5 Feet	To	to	to	to
4.5 – 13 Feet	To	to	to	to
13 – 40 Feet	To	to	to	to

Annual Production by Plant Type (historic climax plant community)

Annual Production (lbs/acre)			
Plant Type	Low	Representative Value	High
Grasses/Grasslike			
Forbs			
Shrubs/Vines			
Trees			
TOTAL	700	1200	1500

Plant community growth curves

Growth Curve Name:

Growth Curve Number:

Growth Curve Description:

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

ECOLOGICAL SITE INTERPRETATIONS

Animal community

Wildlife interpretations

Occasional use by Tule elk.

Livestock Interpretations

Season of Use: Other management considerations.

This site is primarily used for winter and spring grazing. Continuous year-round grazing would result in a deteriorated plant community that has low value as forage. Grazing when soil is wet will result in trampling and compaction of the surface soil layer.

General guide to initial stocking rate. Before making specific recommendations, an on-site evaluation must be made.

Production Year	lbs/acre (air-dry)	AUM/acre	Acres/AUM
Favorable Years	1500		
Normal Years	1200		2 – 4
Unfavorable Years	700		

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Plant community growth curves

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Growth Curve Name:
Growth Curve Number:
Growth Curve Description:

Hydrologic functions

Runoff is slow. The erosion hazard is slight.

Soil Series	Hydrologic Group	Hydrologic Conditions and Runoff Curves		
		Good	Fair	Poor
Winnedumah	B	58	70	77

Recreational uses

This site is primarily located on the City of Los Angeles, Department of Water and Power (LADWP) property and is open to the public.

Wood products

Other products

Other information

Reduce the hazard of wind erosion by avoiding overgrazing, preserving the existing plant cover, and limiting operation of off-road vehicles.

Threatened and endangered plants and animals: Planners will refer to state and federal lists of endangered species. Management recommendations will address impacts on endangered species and their critical habitats (refer to Endangered Species Policy and California Endangered Species Handbook).

Fire is infrequent but can be used as a range management tool to decrease brush species and increase grass cover.

SUPPORTING INFORMATION

Associated sites

Principle sites that commonly occur in association with this potential plant community include:
Saline Meadow (29XG002CA)

Similar sites

Inventory data references

State correlation

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Type locality

Northeast ¼ of Section 15, Township 13 South, Range 35 South.
2 ½ miles east of Independence, Inyo County, California

Relationship to other established classification systems

Plant Symbol: ATTO - SPAI
Society of American Foresters Cover Type:
Kuchler:
CalVeg:
Wildlife Habitat Relationship:
Woodland Suitability Group:
Soil Survey Area: 802 Benton – Owens Valley Soil Survey

Developed in cooperation with the Los Angeles Department of Water and Power

Other references

SITE DESCRIPTION APPROVAL

Author	Agency	Date	Approval	Date
Patti J Novak	SCS	3/1992	Leonard Jolley	

Soil Survey of Manzanar National Historic Site, California

Sodic Fan
029XG005CA

Authorship

Original authors' names and date: Patti J. Novak, March 1992
Revision authors' names and revision date:

Site approval

Leonard Jolley

UNITED STATES DEPARTMENT OF AGRICULTURE
NATURAL RESOURCES CONSERVATION SERVICE

RANGE/ECOLOGICAL SITE DESCRIPTION

ECOLOGICAL SITE CHARACTERISTICS

Site Type: Rangeland

Site Name: Wetland

Site ID: 029XG003CA

Major Land Resource Area: 029G Southern Nevada Basin and Range

Physiographic Features:

This site occurs on fan and stream terraces. Elevations are 3700 to 4700 feet. Slopes range from 0 to 15 percent.

Landform: fan and stream terraces

Aspect:

	<u>Minimum</u>	<u>Maximum</u>
<u>Site elevation (feet):</u>	3700	4700
<u>Slope (percent):</u>	0	15
<u>Water table depth (inches):</u>	1	40
<u>Flooding</u>		
<u>Frequency:</u>	Frequent	
<u>Duration:</u>		

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

Ponding:

Depth (inches):

Frequency:

Duration (days):

during most of the year

Runoff class:

slow

Climatic Features

The climate on this site is characterized by mild, cool winters (30° to 60° F) and hot, dry summers (70° to 100° F). The average annual precipitation ranges from 5 to 8 inches with most falling as rain from November to March. The average frost-free period is 140 to 225 days.

	<u>Minimum</u>	<u>Maximum</u>
<u>Frost-free period (days):</u>	140	225
<u>Freeze-free period (days):</u>		
<u>Mean annual precipitation (inches):</u>	5	8

Monthly precipitation and temperature distribution (°F)

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Min.												
Precip. Max.												
Temp. Min.												
Temp. Max.												

Location of climate stations:

Influencing Water Features

Cowardin wetland classification system: System(s) Subsystem(s) Class(es)

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

Rosgen stream code:

More than one stream type may be associated with the site.

Representative Soil Features

The soils that characterize this site are very deep and poorly drained to very poorly drained. They are formed in organic matter and mixed alluvium. Surface textures are muck, mucky loams, and peaty muck. Available water capacity is high to very high and the hazard of water erosion is slight. Wind erosion hazard is slight to severe. Effective rooting depth is 60 inches or more. Water tables are from 1 to 40 inches. These soils are subject to frequent flooding and ponded water is present of the site during most of the year. This site is associated with springs, seeps, oxbows, and along fault escarpments.

802 Benton - Owns Valley Soil Survey

SSA	Map Unit Symbol	Component Name
802	450	Morey family – Winnedumah, drained - Rindge family complex, 0 – 2% slopes
802	604	Mountom – Artray family – Rounval complex, 0 – 2% slopes
802	614	Argids – Endoaquolls, 0 – 15% slopes

Representative soil features include:

Predominant Parent materials:

Kind: organic matter and mixed alluvium

Origin:

Surface texture: muck, mucky loams, and peaty muck

Subsurface Texture Group:

Minimum Maximum

Surface fragments <=3" (% Cover):

Surface fragments >3" (% Cover):

Subsurface fragments <=3" (% Volume):

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

<u>Subsurface fragments >3" (% Volume):</u>		
<u>Drainage class:</u>	poorly drained	very poorly drained
<u>Permeability Class:</u>		
<u>Depth (inches):</u>		60+"
<u>Electrical conductivity (mmhos/cm):</u>		
<u>Sodium adsorption ratio:</u>		
<u>Calcium Carbonate Equivalent (percent):</u>		
<u>Soil reaction (1:1 Water):</u>		
<u>Soil reaction (0.01M CaCl₂):</u>		
<u>Available Water Holding Capacity:</u>	high	very high
<u>(Total inches for entire profile):</u>		

PLANT COMMUNITIES

Ecological dynamics of the site

As ecological condition deteriorates, the more palatable grasses decrease and rose and willow increase. If the water table is lowered, a plant community dominated by watersage saltbush and rubber rabbitbrush will become the potential.

Plant communities

The interpretative plant community is the historic climax plant community. The plant community is dominated by bulrush and cattail. Potential vegetation composition is about 80% grasses and grass-like plants, 5% forbs, 10% shrubs, and 5% trees.

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

Plant community composition

The following table lists the major plant species and percentages by weight, air-dry, of the total plant community that each contributes in an average production year.

Group	Common Name	Scientific Name	% by Weight	Group Allowable		Annual Production	
				Low	High	Low	High
Grasses and Grasslike Plants							
1	Annual Grasses						
2	Perennial Grasses			5	15		
	Bluegrass	<i>Poa spp.</i>			2*		
	Common reed	<i>Phragmites australis</i>		5	10		
	Knot grass	<i>Paspalum distichum</i>			2*		
	Sedges and Rushes						
	Bulrush	<i>Scirpus spp.</i>	20 – 35				
	Spikerush	<i>Eleocharis spp.</i>	15 – 20				
	Baltic rush	<i>Juncus balticus</i>			2*		
	Sedge	<i>Carex spp.</i>	2 – 5				
Forbs							
3	Forbs			5	15		
	Broadleaf (Common) cattail	<i>Typha latifolia</i>	20 – 35				
	Yerba mansa	<i>Anemopsis californica</i>			2*		
	Clover	<i>Trifolium spp.</i>			2*		
	Buttercup	<i>Ranunculus spp.</i>			2*		
	Annual sunflower	<i>Helianthus anomalus</i>	T – 1				
Shrubs							

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

4	Shrubs			5	10**		
	Willow	<i>Salix spp.</i>			2		
	Woods rose	<i>Rosa woodsii</i>			2		
Trees							
5	Trees						
	Willow	<i>Salix spp.</i>	2 – 5				
	Cottonwood	<i>Populus fremontii</i>	T – 5				
* Allow no more than 2% of each species and no more than 5% in aggregate.							
** Allow no more than 2% of each species and no more than 10% in aggregate.							

Estimated total annual production – air-dry weight (lbs/ac):

Production Year	lbs/acre (air-dry)	AUM/acre	Acres/AUM
Favorable Years	5000		
Normal Years	3500		
Unfavorable Years	2500		

Ground cover and structure

Approximate ground cover (basal and crown) is 70 to 90 percent.

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

Soil Surface Cover

Vegetative Cover						Non-Vegetative Cover					
Basal Cover				Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
Grass/Grasslike	Forb	Shrub/Vine	Tree								
to	to	to	to	to	To	To	to	to	to	to	to

Ground Cover

Vegetative Cover						Non-Vegetative Cover					
Grass/Grasslike	Forb	Shrub/Vine	Tree	Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
to	to	to	to	to	To	To	to	to	to	to	to

Structure of Canopy Cover

	Grasses/Grasslike	Forbs	Shrubs/Vines	Trees
0.5 Feet	to	To	to	to
0.5 - 1 Foot	to	To	to	to
1.0 - 2 Feet	to	To	to	to
2.0 - 4.5 Feet	to	To	to	to
4.5 - 13 Feet	to	To	to	to
13 - 40 Feet	to	To	to	to

Annual Production by Plant Type (historic climax plant community)

Annual Production (lbs/acre)			
Plant Type	Low	Representative Value	High
Grasses/Grasslike			
Forbs			

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

Shrubs/Vines			
Trees			
TOTAL			

Plant community growth curves

Growth Curve Name:
Growth Curve Number:
Growth Curve Description:

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

ECOLOGICAL SITE INTERPRETATIONS

Animal community

Wildlife interpretations

Diverse and abundant wildlife due to the presence of water in a xeric environment: great blue herons, egrets, American bitterns, waterfowl, passerines, raptors, rodents, elk, mule deer, weasels, water snakes, and bullfrogs.

Livestock Interpretations

Season of Use: Other management considerations.

This site is primarily used for spring and summer grazing. Continuous grazing would result in a deteriorated plant community that has low value as forage.

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

General guide to initial stocking rate. Before making specific recommendations, an on-site evaluation must be made.

Production Year	lbs/acre (air-dry)	AUM/acre	Acres/AUM
Favorable Years	5000		
Normal Years	3500	1.0 – 1.5	
Unfavorable Years	2500		

Plant community growth curves

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Growth Curve Name:
Growth Curve Number:
Growth Curve Description:

Hydrologic functions

Runoff is slow. The erosion hazard is slight.

Soil Series	Hydrologic Group	Hydrologic Conditions and Runoff Curves		
		Good	Fair	Poor
Endoaquolls				
Rindge family	B	63	66	76
Mountom	D	81	83	88

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

Recreational uses

This site is primarily located on the City of Los Angeles, Department of Water and Power (LADWP) property and is open to the public.

Wood products

Other products

Other information

Reduce the hazard of wind erosion by avoiding overgrazing and preserving the existing plant cover.

Threatened and endangered plants and animals: Planners will refer to state and federal lists of endangered species. Management recommendations will address impacts on endangered species and their critical habitats (refer to Endangered Species Policy and California Endangered Species Handbook).

Fire: Fire is infrequent but it can be used as a range management tool to decrease the brush species and to increase grass cover.

SUPPORTING INFORMATION

Associated sites

Principle sites that commonly occur in association with this potential plant community include:

Saline Meadow	029XG002CA
Wet Meadow	029XG001CA

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

Similar sites

Competing sites and their differentiae that are similar to this potential plant community:
Wet Meadow 029XG001CA (ACAREX – JUBA dominant species)

Inventory data references

State correlation

Type locality

Northeast $\frac{1}{4}$ of Section 21 of Township 6 South and Range 31 East. Twelve miles west of Bishop, Round Valley, Inyo County, California.

Relationship to other established classification systems

Plant Symbol: SCIRP - TYPHA
Society of American Foresters Cover Type:
Kuchler:
CalVeg:
Wildlife Habitat Relationship:
Woodland Suitability Group:
Soil Survey Area: 802 Benton – Owens Valley Soil Survey

Developed in cooperation with the Los Angeles Department of Water and Power

Other references

Soil Survey of Manzanar National Historic Site, California

Wetland
029XG003CA

SITE DESCRIPTION APPROVAL

Author	Agency	Date	Approval	Date
Patti J. Novak	NRCS	1/1993	Leonard Jolley	

Authorship

Original author's names and date: Patti J. Novak, January 1993.

Revision authors' names and revision date:

Site approval

Leonard Jolley, State Range Conservationist

**UNITED STATES DEPARTMENT OF AGRICULTURE
NATURAL RESOURCES CONSERVATION SERVICE**

RANGE/ECOLOGICAL SITE DESCRIPTION

ECOLOGICAL SITE CHARACTERISTICS

Site Type: Rangeland

Site Name: Saline Meadow

Site ID: 029XG002CA

Major Land Resource Area: 029G Southern Nevada Basin and Range

Physiographic Features:

This site occurs on fan, stream, and lacustrine terraces and axial stream floodplains. Elevations are 3580 to 6000 feet. Slopes range from 0 to 9 percent but slope gradients of 0 to 2 percent are most typical.

Landform: fan, stream, and lacustrine terraces and axial stream floodplains

Aspect:

	<u>Minimum</u>	<u>Maximum</u>
<u>Site elevation (feet):</u>	3580	6000
<u>Slope (percent):</u>	0	9 (mostly 0 – 2%)
<u>Water table depth (inches):</u>	18	60 (from February 1 to May 15)

Flooding

Frequency:

Duration:

Soil Survey of Manzanar National Historic Site, California

Wet Meadow
029XG001CA

More than one stream type may be associated with the site.

Representative Soil Features

The soils that characterize this site are very deep and range from well drained to poorly drained. They are formed in mixed alluvium and lacustrine sediments. Surface textures are loam sands, sandy loams, loams, and silt loams. Available water capacity is moderate to high and the hazard of water erosion is slight. Wind erosion hazard is slight to severe. Effective rooting depth is 60 inches or more. Water tables are from 18 to 60 inches between February 1 and May 15. Capillary rise of the water table in spring enhances soil moisture during the growing season. These soils are slightly to strongly salt and sodium affected in the upper profile but decreasing with depth.

Benton – Owens Valley Soil Survey (802)

Map Unit Symbol	Component Name
450	Morey Family – Winnedumah, drained – Rindge Family, 0 – 2% slopes
520	Dehy – Typic Argiaquolls – Lubkin association, 0 – 2% slopes
531	Shabbell – Shondow – Fluvents association, 0 – 2% slopes
626	Torrifluvents, 0 – 2% slopes
635	Manzanar silt loam, saline – sodic, 0 – 2% slopes
636	Manzanar – Winnedumah association, 0 – 2% slopes
638	Shondow loam, 0 – 2% slopes
653	Manzanar – Westguard association, 0 – 2% slopes
660	Manzanar – Division association, 0 – 2% slopes
670	Dehy loam, 0 – 2% slopes
672	Dehy sandy loam, 0 – 2% slopes
674	Dehy - Dehy saline complex, 0 – 2% slopes
678	Dehy - gravelly loam sand, saline, 0 – 2% slopes, cool
701	Shondow loam, drained, 0 – 2% slopes
704	Shondow, drained-Hessica, sandy sub-irrigated association, 0 – 2% slopes
740	Dehy complex, 0 – 2% slopes

Soil Survey of Manzanar National Historic Site, California

Wet Meadow
029XG001CA

Representative soil features include:

Predominant Parent materials:

Kind: mixed alluvium and lacustrine sediments

Origin:

Surface texture: loamy sands, sandy loams, loams, and silt loams

Subsurface Texture Group:

	<u>Minimum</u>	<u>Maximum</u>
<u>Surface fragments <=3" (% Cover):</u>		
<u>Surface fragments >3" (% Cover):</u>		
<u>Subsurface fragments <=3" (% Volume):</u>		
<u>Subsurface fragments >3" (% Volume):</u>		
<u>Drainage class:</u>	well drained	poorly drained
<u>Permeability Class:</u>		
<u>Depth (inches):</u>	18	60+
<u>Electrical conductivity (mmhos/cm):</u>		
<u>Sodium adsorption ratio:</u>		
<u>Calcium Carbonate Equivalent (percent):</u>		
<u>Soil reaction (1:1 Water):</u>		
<u>Soil reaction (0.01M CaCl₂):</u>		
<u>Available Water Holding Capacity:</u>	moderate	high
(Total inches for entire profile):		

PLANT COMMUNITIES

The interpretive plant community contains plants that existed both in the historic climax plant community as well as plants from the naturalized plant community. The plant community is dominated by sedges and rushes. Potential vegetation composition is about 80% grasses, 15% forbs, and 5% shrubs. The following table lists the major plant species and percentages by weight (air-dry) of the total naturalized plant community that each contributes in an average production year. It illustrates the normal changes which occur in the plant community.

Soil Survey of Manzanar National Historic Site, California

Wet Meadow
029XG001CA

PLANT COMMUNITY DYNAMICS:

Overgrazing will result in a decrease in the palatable sedges and grasses and an increase in Baltic rush, wild iris, rose, and willows. Species likely to invade this site include Kentucky bluegrass, rabbitsfoot grass, and thistles, especially Canada thistle.

Group	Common Name	Scientific Name	% by Weight	Group Allowable		Annual Production	
				Low	High	Low	High
Grasses and Grasslike Plants							
1	Annual Grasses						
2	Perennial Grasses						
	Alkali sacaton	<i>Sporobolus airoides</i>	25 - 50				
	Inland saltgrass	<i>Distichlis spicata</i>	15 - 25				
	Beardless wildrye	<i>Leymus triticoides</i>	5 - 10		3		
	Other perennial grasses			2	8		
	Common reed	<i>Phragmites australis</i>			2		
	Nevada bluegrass	<i>Poa nevadensis</i>			2		
	Basin wildrye	<i>Leymus cinereus</i>			2		
	Sedges and Rushes						
	Baltic rush	<i>Juncus balticus</i>	5 - 15				
	Sedge	<i>Carex spp.</i>			2		
Forbs							
3	Forbs			5	10		
	Yerba mansa	<i>Anemopsis californica</i>			3		
	Wyoming Indian paintbrush	<i>Castilleja linariifolia</i>			3		
	Clustered goldenweed	<i>Pyrrocoma racemosa</i>			3		
	Other annual forbs		T - 2				

Soil Survey of Manzanar National Historic Site, California

Wet Meadow
029XG001CA

Shrubs							
4	Shrubs			5	10		
	Watersage (Torrey's) saltbush	<i>Atriplex torreyi</i>			3		
	Rubber rabbitbrush	<i>Ericameria nauseosa</i>			3		
	Willow	<i>Salix spp.</i>			3		
	Woods rose	<i>Rosa woodsii</i>			3		
	Black greasewood	<i>Sarcobatus vermiculatus</i>			3		
Trees							
5	Trees		0 – 5				
	Fremont cottonwood	<i>Populus fremontii</i>					
	Red willow	<i>Salix laevigata</i>					

Estimated total annual production – air-dry weight (lbs/ac):

Production Year	lbs/acre (air-dry)	AUM/acre	Acres/AUM
Favorable Years	3300		
Normal Years	2200		
Unfavorable Years	1000		

Stocking rate derived from total production minus 700 pounds per acre for mulch. Before making specific recommendations for initial stocking rates, an on-site evaluation must be made.

Ecological dynamics of the site

As ecological condition deteriorates, the perennial grasses will decline and the shrub species, especially watersage saltbush, will become dominant. Yerba mansa will also increase with deterioration of the community. Waterspreading

Soil Survey of Manzanar National Historic Site, California

Wet Meadow
029XG001CA

activities in high runoff years will alter the community by causing an increase in Baltic rush, sedge, and common reed, and an invasion of cattails, tules, and saltcedar.

Plant communities

Soil Surface Cover

Vegetative Cover						Non-Vegetative Cover					
Basal Cover				Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
Grass/ Grasslike	Forb	Shrub/ Vine	Tree								
To	to	to	to	To	To	to	to	to	to	to	to

Approximate ground cover (basal and crown) is 40 to 80 percent.

Ground Cover

Vegetative Cover						Non-Vegetative Cover					
Grass/ Grasslike	Forb	Shrub/ Vine	Tree	Non-Vascular Plants	Biological Crust	Litter	Surface Fragments >1/4 - 3"	Surface Fragments >3"	Bedrock	Water	Bare Ground
To	to	to	to	To	to	to	to	to	to	to	to

Structure of Canopy Cover

	Grasses/Grasslike	Forbs	Shrubs/Vines	Trees
0.5 Feet	To	to	to	to
0.5 - 1 Foot	To	to	to	to
1.0 - 2 Feet	To	to	to	to
2.0 - 4.5 Feet	To	to	to	to
4.5 - 13 Feet	To	to	to	to
13 - 40 Feet	To	to	to	to

Annual Production by Plant Type (historic climax plant community)

Annual Production (lbs/acre)			
Plant Type	Low	Representative Value	High
Grasses/Grasslike			
Forbs			
Shrubs/Vines			
Trees			
TOTAL			

Plant community growth curves

Growth Curve Name:
Growth Curve Number:
Growth Curve Description:

Percent Production by Month											
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

ECOLOGICAL SITE INTERPRETATIONS

Animal community

Wildlife interpretations

Year-round range for Tule Elk.
Productive passerine/flycatcher habitat
Habitat for small rodents and predatory raptors
Winter feeding areas for geese

Soil Survey of Manzanar National Historic Site, California

**Wet Meadow
029XG001CA**

Upland game habitat along brushy edges

Maintain tule elk numbers at levels compatible with existing land uses, such as livestock grazing. Irrigation water would enhance the area for existing wildlife populations.

Livestock Interpretations

Season of use: other management considerations

This site is primarily used for winter and early spring grazing. Continuous year-round grazing would result in a deteriorated plant community that has low value as forage.

Estimated total annual production – air-dry weight (lbs/ac):

Production Year	lbs/acre (air-dry)	AUM/acre	Acres/AUM
Favorable Years	3300		
Normal Years	2200	1.0	
Unfavorable Years	1000		

Before making specific recommendations, an on-site evaluation must be made.

Hydrologic functions

Soil Series	Hydrologic Group	Hydrologic Conditions and Runoff Curves		
		Good	Fair	Poor
Dehy	C	72	78	84
Manzanar	C	72	78	84
Morey family	C	72	78	84
Shondow	B	58	70	77
Torrifluvents				

Runoff is slow. The erosion hazard is slight.

Recreational uses

This site is primarily located on the City of Los Angeles, Department of Water and Power (LADWP) property and is open to the public.

Wood products

Other products

Other information

Major poisonous plants: Specklepod loco milkvetch (*Astragalus spp.*)

Threatened and endangered plants and animals: Planners will refer to state and federal lists of endangered species. Management recommendations will address impacts on endangered species and their critical habitats (refer to Endangered Species Policy and California Endangered Species Handbook).

Alkali Mariposa lily (*Calochortus excavatus*) – Federal candidate

Owens Valley Checker Mallow (*Sidalcea covillei*) – State endangered and Federal candidate

Fire: Fire is infrequent but it can be used as a range management tool to decrease the brush species and to increase grass cover.

SUPPORTING INFORMATION

Associated sites

Principle site(s) that commonly occur in association with this potential plant community include:

Wetland 029XG003CA,
Sodic Fan (029XG005CA)

Similar sites

Competing sites (and their differentiae) that are similar to this potential plant community:

Wet Meadow (029XG001CA)	(CAREX – JUBA dominant species)
Sodic Fan (029XG005CA)	(ATTO co-dominant with SPAI; less productive site)

Inventory data references

State correlation

Type locality

Southeast of Independence, Inyo County, California: SW1/4 of Section 35, T13S, R35E.

Relationship to other established classification systems

Plant Symbol: SPAI – DISPS2
Society of American Foresters Cover Type:
Kuchler:
CalVeg:
Wildlife Habitat Relationship:

Soil Survey of Manzanar National Historic Site, California

Wet Meadow
029XG001CA

Woodland Suitability Group:
Soil Survey Area: 802 Benton – Owens Valley Soil Survey

Developed in cooperation with the Los Angeles Department of Water and Power

Other references

SITE DESCRIPTION APPROVAL

Author	Agency	Date	Approval	Date
Original:	SCS	2/1992	Leonard Jolley	3/1992
Revised:	SCS	12/1993	Leonard Jolley	1/1994

Authorship

Patti J. Novak, Range Conservationist

Site approval

Leonard Jolley, State Range Conservationist
SCS, California

NRCS Accessibility Statement

The Natural Resources Conservation Service (NRCS) is committed to making its information accessible to all of its customers and employees. If you are experiencing accessibility issues and need assistance, please contact our Helpdesk by phone at 1-800-457-3642 or by e-mail at ServiceDesk-FTC@ftc.usda.gov. For assistance with publications that include maps, graphs, or similar forms of information, you may also wish to contact our State or local office. You can locate the correct office and phone number at <http://offices.sc.egov.usda.gov/locator/app>.