

NRCS Accessibility Statement

The Natural Resources Conservation Service (NRCS) is committed to making its information accessible to all of its customers and employees. If you are experiencing accessibility issues and need assistance, please contact our Helpdesk by phone at 1-800-457-3642 or by e-mail at ServiceDesk-FTC@ftc.usda.gov. For assistance with publications that include maps, graphs, or similar forms of information, you may also wish to contact our State or local office. You can locate the correct office and phone number at <http://offices.sc.egov.usda.gov/locator/app>.



United States
Department of
Agriculture



NRCS

Natural
Resources
Conservation
Service

In cooperation with
The Connecticut
Agricultural Experiment
Station, The Storrs
Agricultural Experiment
Station, and Connecticut
Department of
Environmental Protection

Soil Survey of the State of Connecticut



How To Use This Soil Survey

General Soil Map

The general soil map associated with this publication is the Digital General Soil Map of the United States, formerly known as STATSGO. This map shows broad areas with a distinctive pattern of soils, relief, and drainage. The map is available for download from the Soil Data Mart of the Natural Resources Conservation Service, accessed at <http://soils.usda.gov>.

To find information about your area of interest, locate that area on the map, identify the name of the map unit in the area, then refer to the section **General Soil Map Units** for a general description of the soils in your area.


Detailed Soil Maps

The detailed soil maps can be useful in planning the use and management of small areas.

To find information about your area of interest, locate that area on the **Index to Map Sheets**. Note the number of the map sheet and turn to that sheet.

Locate your area of interest on the map sheet. Note the map unit symbols that are in that area. Turn to the **Contents**, which lists the map units by symbol and name and shows the page where each map unit is described.

The **Contents** shows which table has data on a specific land use for each detailed soil map unit. Also see the **Contents** for sections of this publication that may address your specific needs.



NOTE: Map unit symbols in a soil survey may consist only of numbers or letters, or they may be a combination of numbers and letters.

This soil survey is a publication of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (formerly the Soil Conservation Service) has leadership for the Federal part of the National Cooperative Soil Survey.

Earlier soil surveys of the eight Connecticut counties (Fairfield, Hartford, Litchfield, Middlesex, New Haven, New London, Tolland, and Windham) were published by the United States Department of Agriculture Soil Conservation Service between 1958 and 1983. This survey supercedes the earlier ones and provides additional information and maps that show the soil in greater detail. Major fieldwork for this soil survey was completed in 2002. Soil names and descriptions were approved in 2003. Unless otherwise indicated, statements in this publication refer to conditions in the survey area in 2003. This survey was made cooperatively by the Natural Resources Conservation Service and the Connecticut Agricultural Experiment Station, the Storrs Agricultural Experiment Station, and the Connecticut Department of Environmental Protection. Partial funding for this survey was provided by the United States Environmental Protection Agency and the Connecticut Department of Transportation.

Soil maps in this survey may be copied without permission. Enlargement of these maps, however, could cause misunderstanding of the detail of mapping. If enlarged, maps do not show the small areas of contrasting soils that could have been shown at a larger scale.

The United States Department of Agriculture (USDA) prohibits discrimination in all of its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice or TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue SW, Washington, DC 20250-9410, or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Cover: This stone wall, located in Woodstock, Connecticut, is in an area of Woodbridge fine sandy loam. Old stone walls in the woods indicate that the land was cleared for agriculture in the past.

Additional information about the Nation's natural resources is available online from the Natural Resources Conservation Service at <http://www.nrcs.usda.gov>.

Contents

How To Use This Soil Survey	i
Foreword	xv
General Nature of the Survey Area	1
How This Survey Was Made	7
General Soil Map Units	9
S1438. Charlton-Canton-Paxton	9
S1439. Charlton-Chatfield-Hollis	10
S1440. Woodbridge-Paxton-Ridgebury	10
S1441. Hinckley-Agawam-Merrimac	11
S1442. Catden-Merrimac-Saco	11
S1443. Urban land-Udorthents-Cheshire	12
S1444. Stockbridge-Charlton-Mudgepond	12
S1445. Copake-Groton-Hero	13
S1446. Windsor-Merrimac-Hinckley	13
S1447. Windsor-Agawam-Haven	14
S1448. Brancroft-Scitico-Shaker	14
S1449. Brookfield-Brimfield-Leicester	15
S1451. Holyoke-Rock outcrop-Yalesville	15
S1452. Cheshire-Yalesville-Wethersfield	16
S1453. Narragansett-Cheshire-Wapping	16
S1454. Wethersfield-Ludlow-Wilbraham	17
S1455. Manchester-Branford-Rippowam	17
S1456. Manchester-Penwood-Hartford	18
S1457. Wethersfield-Berlin-Belgrade	18
S1458. Winooski-Hadley-Occum	18
S3111. Macomber-Taconic-Lanesboro	19
S3114. Copake-Hero-Halsey	20
S3115. Hinckley-Windsor-Merrimac	21
S3120. Wethersfield-Meckesville-Scarboro	21
S3121. Brookfield-Brimfield-Paxton	22
S3122. Paxton-Woodbridge-Hollis	22
S3136. Hollis-Chatfield-Rock outcrop	23
S6623. Charlton-Canton-Sutton	24
S6625. Woodbridge-Paxton-Ridgebury	24
S6627. Carlisle-Adrian-Scarboro	25
S9569. Bice-Shelburne-Ashfield	25
S9570. Bice-Millsite-Westminster	26
S9571. Ashfield-Shelburne-Loonmeadow	27
Detailed Soil Map Units	29
2—Ridgebury fine sandy loam	30
3—Ridgebury, Leicester, and Whitman soils, extremely stony	31
4—Leicester fine sandy loam	33
5—Wilbraham silt loam	34
6—Wilbraham and Menlo soils, extremely stony	35
7—Mudgepond silt loam	37

8—Mudgepond and Alden soils, extremely stony	38
9—Scitico, Shaker, and Maybid soils	39
10—Raynham silt loam	41
12—Raypol silt loam	42
13—Walpole sandy loam	44
14—Fredon silt loam	45
15—Scarboro muck	46
16—Halsey silt loam	47
17—Timakwa and Natchaug soils	48
18—Catden and Freetown soils	50
20A—Ellington silt loam, 0 to 5 percent slopes	52
21A—Ninigret and Tisbury soils, 0 to 5 percent slopes	53
22A—Hero gravelly loam, 0 to 3 percent slopes	55
22B—Hero gravelly loam, 3 to 8 percent slopes	56
23A—Sudbury sandy loam, 0 to 5 percent slopes	57
24A—Deerfield loamy fine sand, 0 to 3 percent slopes	58
25A—Brancroft silt loam, 0 to 3 percent slopes	59
25B—Brancroft silt loam, 3 to 8 percent slopes	60
25C—Brancroft silt loam, 8 to 15 percent slopes	62
26A—Berlin silt loam, 0 to 3 percent slopes	63
26B—Berlin silt loam, 3 to 8 percent slopes	64
27A—Belgrade silt loam, 0 to 5 percent slopes	65
28A—Elmridge fine sandy loam, 0 to 3 percent slopes	66
28B—Elmridge fine sandy loam, 3 to 8 percent slopes	67
29A—Agawam fine sandy loam, 0 to 3 percent slopes	68
29B—Agawam fine sandy loam, 3 to 8 percent slopes	69
29C—Agawam fine sandy loam, 8 to 15 percent slopes	70
30A—Branford silt loam, 0 to 3 percent slopes	72
30B—Branford silt loam, 3 to 8 percent slopes	73
30C—Branford silt loam, 8 to 15 percent slopes	74
31A—Copake fine sandy loam, 0 to 3 percent slopes	75
31B—Copake fine sandy loam, 3 to 8 percent slopes	76
31C—Copake gravelly loam, 8 to 15 percent slopes	77
32A—Haven and Enfield soils, 0 to 3 percent slopes	78
32B—Haven and Enfield soils, 3 to 8 percent slopes	80
32C—Haven and Enfield soils, 8 to 15 percent slopes	81
33A—Hartford sandy loam, 0 to 3 percent slopes	83
33B—Hartford sandy loam, 3 to 8 percent slopes	85
34A—Merrimac sandy loam, 0 to 3 percent slopes	86
34B—Merrimac sandy loam, 3 to 8 percent slopes	87
34C—Merrimac sandy loam, 8 to 15 percent slopes	88
35A—Penwood loamy sand, 0 to 3 percent slopes	89
35B—Penwood loamy sand, 3 to 8 percent slopes	90
36A—Windsor loamy sand, 0 to 3 percent slopes	91
36B—Windsor loamy sand, 3 to 8 percent slopes	92
36C—Windsor loamy sand, 8 to 15 percent slopes	93
37A—Manchester gravelly sandy loam, 0 to 3 percent slopes	94
37C—Manchester gravelly sandy loam, 3 to 15 percent slopes	96
37E—Manchester gravelly sandy loam, 15 to 45 percent slopes	97
38A—Hinckley gravelly sandy loam, 0 to 3 percent slopes	98
38C—Hinckley gravelly sandy loam, 3 to 15 percent slopes	99
38E—Hinckley gravelly sandy loam, 15 to 45 percent slopes	100
39A—Groton gravelly sandy loam, 0 to 3 percent slopes	102
39C—Groton gravelly sandy loam, 3 to 15 percent slopes	103

39E—Groton gravelly sandy loam, 15 to 45 percent slopes	104
40A—Ludlow silt loam, 0 to 3 percent slopes	105
40B—Ludlow silt loam, 3 to 8 percent slopes	106
41B—Ludlow silt loam, 2 to 8 percent slopes, very stony	107
42C—Ludlow silt loam, 2 to 15 percent slopes, extremely stony	109
43A—Rainbow silt loam, 0 to 3 percent slopes	110
43B—Rainbow silt loam, 3 to 8 percent slopes	111
44B—Rainbow silt loam, 2 to 8 percent slopes, very stony	112
45A—Woodbridge fine sandy loam, 0 to 3 percent slopes	113
45B—Woodbridge fine sandy loam, 3 to 8 percent slopes	114
45C—Woodbridge fine sandy loam, 8 to 15 percent slopes	116
46B—Woodbridge fine sandy loam, 2 to 8 percent slopes, very stony	117
46C—Woodbridge fine sandy loam, 8 to 15 percent slopes, very stony	118
47C—Woodbridge fine sandy loam, 2 to 15 percent slopes, extremely stony	119
48B—Georgia and Amenia silt loams, 2 to 8 percent slopes	121
48C—Georgia and Amenia silt loams, 8 to 15 percent slopes	123
49B—Georgia and Amenia silt loams, 3 to 8 percent slopes, very stony	124
49C—Georgia and Amenia silt loams, 8 to 15 percent slopes, very stony	126
50A—Sutton fine sandy loam, 0 to 3 percent slopes	128
50B—Sutton fine sandy loam, 3 to 8 percent slopes	129
51B—Sutton fine sandy loam, 2 to 8 percent slopes, very stony	130
52C—Sutton fine sandy loam, 2 to 15 percent slopes, extremely stony	131
53A—Wapping very fine sandy loam, 0 to 3 percent slopes	132
53B—Wapping very fine sandy loam, 3 to 8 percent slopes	134
54B—Wapping very fine sandy loam, 2 to 8 percent slopes, very stony	135
55A—Watchaug fine sandy loam, 0 to 3 percent slopes	136
55B—Watchaug fine sandy loam, 3 to 8 percent slopes	137
56B—Watchaug fine sandy loam, 2 to 8 percent slopes, very stony	138
57B—Gloucester gravelly sandy loam, 3 to 8 percent slopes	139
57C—Gloucester gravelly sandy loam, 8 to 15 percent slopes	140
57D—Gloucester gravelly sandy loam, 15 to 25 percent slopes	142
58B—Gloucester gravelly sandy loam, 3 to 8 percent slopes, very stony	143
58C—Gloucester gravelly sandy loam, 8 to 15 percent slopes, very stony	144
59C—Gloucester gravelly sandy loam, 3 to 15 percent slopes, extremely stony	145
59D—Gloucester gravelly sandy loam, 15 to 35 percent slopes, extremely stony	146
60B—Canton and Charlton soils, 3 to 8 percent slopes	147
60C—Canton and Charlton soils, 8 to 15 percent slopes	149
60D—Canton and Charlton soils, 15 to 25 percent slopes	151
61B—Canton and Charlton soils, 3 to 8 percent slopes, very stony	152
61C—Canton and Charlton soils, 8 to 15 percent slopes, very stony	154
62C—Canton and Charlton soils, 3 to 15 percent slopes, extremely stony	156
62D—Canton and Charlton soils, 15 to 35 percent slopes, extremely stony	157
63B—Cheshire fine sandy loam, 3 to 8 percent slopes	159
63C—Cheshire fine sandy loam, 8 to 15 percent slopes	160
63D—Cheshire fine sandy loam, 15 to 25 percent slopes	161
64B—Cheshire fine sandy loam, 3 to 8 percent slopes, very stony	162
64C—Cheshire fine sandy loam, 8 to 15 percent slopes, very stony	163
65C—Cheshire fine sandy loam, 3 to 15 percent slopes, extremely stony	164
65D—Cheshire fine sandy loam, 15 to 35 percent slopes, extremely stony	165
66B—Narragansett silt loam, 2 to 8 percent slopes	166
66C—Narragansett silt loam, 8 to 15 percent slopes	168
67B—Narragansett silt loam, 3 to 8 percent slopes, very stony	169

67C—Narragansett silt loam, 8 to 15 percent slopes, very stony	170
68C—Narragansett silt loam, 3 to 15 percent slopes, extremely stony	171
68D—Narragansett silt loam, 15 to 25 percent slopes, extremely stony	172
69B—Yalesville fine sandy loam, 3 to 8 percent slopes	174
69C—Yalesville fine sandy loam, 8 to 15 percent slopes	175
70C—Branford–Holyoke complex, 3 to 15 percent slopes, very rocky	176
71C—Brookfield–Brimfield–Rock outcrop complex, 3 to 15 percent slopes	178
71E—Brookfield–Brimfield–Rock outcrop complex, 15 to 45 percent slopes	179
73C—Charlton–Chatfield complex, 3 to 15 percent slopes, very rocky	181
73E—Charlton–Chatfield complex, 15 to 45 percent slopes, very rocky	183
74C—Narragansett–Hollis complex, 3 to 15 percent slopes, very rocky	185
75C—Hollis–Chatfield–Rock outcrop complex, 3 to 15 percent slopes	187
75E—Hollis–Chatfield–Rock outcrop complex, 15 to 45 percent slopes	189
76E—Rock outcrop–Hollis complex, 3 to 45 percent slopes	191
76F—Rock outcrop–Hollis complex, 45 to 60 percent slopes	192
77C—Cheshire–Holyoke complex, 3 to 15 percent slopes, very rocky	193
77D—Cheshire–Holyoke complex, 15 to 35 percent slopes, very rocky	195
78C—Holyoke–Rock outcrop complex, 3 to 15 percent slopes	197
78E—Holyoke–Rock outcrop complex, 15 to 45 percent slopes	198
79E—Rock outcrop–Holyoke complex, 3 to 45 percent slopes	199
80B—Bernardston silt loam, 3 to 8 percent slopes	201
80C—Bernardston silt loam, 8 to 15 percent slopes	202
81C—Bernardston silt loam, 3 to 15 percent slopes, extremely stony	203
81D—Bernardston silt loam, 15 to 25 percent slopes, extremely stony	204
82B—Broadbrook silt loam, 3 to 8 percent slopes	205
82C—Broadbrook silt loam, 8 to 15 percent slopes	206
82D—Broadbrook silt loam, 15 to 25 percent slopes	207
83B—Broadbrook silt loam, 3 to 8 percent slopes, very stony	209
83C—Broadbrook silt loam, 8 to 15 percent slopes, very stony	210
84B—Paxton and Montauk fine sandy loams, 3 to 8 percent slopes	211
84C—Paxton and Montauk fine sandy loams, 8 to 15 percent slopes	213
84D—Paxton and Montauk fine sandy loams, 15 to 25 percent slopes	215
85B—Paxton and Montauk fine sandy loams, 3 to 8 percent slopes, very stony	216
85C—Paxton and Montauk fine sandy loams, 8 to 15 percent slopes, very stony	218
86C—Paxton and Montauk fine sandy loams, 3 to 15 percent slopes, extremely stony	220
86D—Paxton and Montauk fine sandy loams, 15 to 35 percent slopes, extremely stony	222
87B—Wethersfield loam, 3 to 8 percent slopes	224
87C—Wethersfield loam, 8 to 15 percent slopes	225
87D—Wethersfield loam, 15 to 25 percent slopes	226
88B—Wethersfield loam, 3 to 8 percent slopes, very stony	228
88C—Wethersfield loam, 8 to 15 percent slopes, very stony	229
89C—Wethersfield loam, 3 to 15 percent slopes, extremely stony	230
89D—Wethersfield loam, 15 to 35 percent slopes, extremely stony	231
90B—Stockbridge loam, 3 to 8 percent slopes	233
90C—Stockbridge loam, 8 to 15 percent slopes	234
90D—Stockbridge loam, 15 to 25 percent slopes	235
91B—Stockbridge loam, 3 to 8 percent slopes, very stony	237
91C—Stockbridge loam, 8 to 15 percent slopes, very stony	238
91D—Stockbridge loam, 15 to 35 percent slopes, very stony	239
92B—Nellis fine sandy loam, 3 to 8 percent slopes	240

92C—Nellis fine sandy loam, 8 to 15 percent slopes	241
92D—Nellis fine sandy loam, 15 to 25 percent slopes	242
93C—Nellis fine sandy loam, 3 to 15 percent slopes, very stony	244
94C—Farmington-Nellis complex, 3 to 15 percent slopes, very rocky	245
94E—Farmington-Nellis complex, 15 to 35 percent slopes, very rocky	247
95C—Farmington-Rock outcrop complex, 3 to 15 percent slopes	249
95E—Farmington-Rock outcrop complex, 15 to 45 percent slopes	250
96—Ipswich mucky peat	251
97—Pawcatuck mucky peat	253
98—Westbrook mucky peat	254
99—Westbrook mucky peat, low salt	255
100—Suncook loamy fine sand	256
101—Occum fine sandy loam	257
102—Pootatuck fine sandy loam	258
103—Rippowam fine sandy loam	260
104—Bash silt loam	261
105—Hadley silt loam	262
106—Winooski silt loam	263
107—Limerick and Lim soils	264
108—Saco silt loam	265
109—Fluvaquents-Udifluvents complex, frequently flooded	266
221A—Ninigret-Urban land complex, 0 to 5 percent slopes	268
224A—Deerfield-Urban land complex, 0 to 3 percent slopes	269
225B—Brancroft-Urban land complex, 0 to 8 percent slopes	271
226B—Berlin-Urban land complex, 0 to 8 percent slopes	272
228B—Elmridge-Urban land complex, 0 to 8 percent slopes	273
229B—Agawam-Urban land complex, 0 to 8 percent slopes	275
229C—Agawam-Urban land complex, 8 to 15 percent slopes	276
230B—Branford-Urban land complex, 0 to 8 percent slopes	277
230C—Branford-Urban land complex, 8 to 15 percent slopes	279
232B—Haven-Urban land complex, 0 to 8 percent slopes	280
234B—Merrimac-Urban land complex, 0 to 8 percent slopes	281
235B—Penwood-Urban land complex, 0 to 8 percent slopes	282
236B—Windsor-Urban land complex, 0 to 8 percent slopes	284
237A—Manchester-Urban land complex, 0 to 3 percent slopes	285
237C—Manchester-Urban land complex, 3 to 15 percent slopes	286
238A—Hinckley-Urban land complex, 0 to 3 percent slopes	288
238C—Hinckley-Urban land complex, 3 to 15 percent slopes	289
240B—Ludlow-Urban land complex, 0 to 8 percent slopes	290
243B—Rainbow-Urban land complex, 0 to 8 percent slopes	292
245B—Woodbridge-Urban land complex, 0 to 8 percent slopes	293
245C—Woodbridge-Urban land complex, 8 To 15 percent slopes	294
248B—Georgia-Urban land complex, 2 to 8 percent slopes	296
250B—Sutton-Urban land complex, 0 to 8 percent slopes	297
253B—Wapping-Urban land complex, 0 to 8 percent slopes	298
255B—Watchaug-Urban land complex, 0 to 8 percent slopes	300
260B—Charlton-Urban land complex, 3 to 8 percent slopes	301
260C—Charlton-Urban land complex, 8 to 15 percent slopes	302
260D—Charlton-Urban land complex, 15 to 25 percent slopes	304
263B—Cheshire-Urban land complex, 3 to 8 percent slopes	305
263C—Cheshire-Urban land complex, 8 to 15 percent slopes	306
266B—Narragansett-Urban land complex, 3 to 8 percent slopes	307
269B—Yalesville-Urban land complex, 3 to 8 percent slopes	309
269C—Yalesville-Urban land complex, 8 to 15 percent slopes	310

273C—Urban land-Charlton-Chatfield complex, rocky, 3 to 15 percent slopes ..	311
273E—Urban land-Charlton-Chatfield complex, rocky, 15 to 45 percent slopes	313
275C—Urban land-Chatfield complex, rocky, 3 to 15 percent slopes	315
275E—Urban land-Chatfield-Rock outcrop complex, 15 to 45 percent slopes ...	317
282B—Broadbrook-Urban land complex, 3 to 8 percent slopes	318
284B—Paxton-Urban land complex, 3 to 8 percent slopes	320
284C—Paxton-Urban land complex, 8 to 15 percent slopes	321
284D—Paxton-Urban land complex, 15 to 25 percent slopes	322
287B—Wethersfield-Urban land complex, 3 to 8 percent slopes	324
287C—Wethersfield-Urban land complex, 8 to 15 percent slopes	325
287D—Wethersfield-Urban land complex, 15 to 25 percent slopes	326
290B—Stockbridge-Urban land complex, 3 to 8 percent slopes	328
290C—Stockbridge-Urban land complex, 8 to 15 percent slopes	329
290D—Stockbridge-Urban land complex, 15 to 25 percent slopes	330
301—Beaches-Udipsamments complex, coastal	332
302—Dumps	333
303—Pits, Quarries	334
304—Udorthents, loamy, very steep	334
305—Udorthents-Pits complex, gravelly	335
306—Udorthents-Urban land complex	336
307—Urban Land	337
308—Udorthents, smoothed	338
309—Udorthents, flood control	339
310—Udorthents, periodically flooded	340
401C—Macomber-Taconic complex, 3 to 15 percent slopes, very rocky	341
402D—Taconic-Macomber-Rock outcrop complex, 15 to 25 percent slopes	342
403C—Taconic-Rock outcrop complex, 3 to 15 percent slopes	344
403E—Taconic-Rock outcrop complex, 15 to 45 percent slopes	345
403F—Taconic-Rock outcrop complex, 45 to 70 percent slopes	346
405C—Dummerston gravelly loam, 3 to 15 percent slopes, very stony	348
405E—Dummerston gravelly loam, 15 to 45 percent slopes, very stony	349
407C—Lanesboro loam, 3 to 15 percent slopes, very stony	350
407E—Lanesboro loam, 15 to 45 percent slopes, very stony	351
408C—Fullam silt loam, 3 to 15 percent slopes, very stony	352
409B—Brayton mucky silt loam, 0 to 8 percent slopes, very stony	354
412B—Bice fine sandy loam, 3 to 8 percent slopes	355
412C—Bice fine sandy loam, 8 to 15 percent slopes	356
412D—Bice fine sandy loam, 15 to 25 percent slopes	357
413C—Bice-Millsite complex, 3 to 15 percent slopes, very rocky	358
413E—Bice-Millsite complex, 15 to 45 percent slopes, very rocky	360
414—Fredon silt loam, cold	361
415C—Westminster-Millsite-Rock outcrop complex, 3 to 15 percent slopes	362
415E—Westminster-Millsite-Rock outcrop complex, 15 to 45 percent slopes	364
416E—Rock outcrop-Westminster complex, 8 to 45 percent slopes	366
416F—Rock outcrop-Westminster complex, 45 to 70 percent slopes	367
417B—Bice fine sandy loam, 3 to 8 percent slopes, very stony	368
417C—Bice fine sandy loam, 8 to 15 percent slopes, very stony	369
417D—Bice fine sandy loam, 15 to 25 percent slopes, very stony	370
418C—Schroon fine sandy loam, 2 to 15 percent slopes, very stony	371
420A—Schroon fine sandy loam, 0 to 3 percent slopes	373
420B—Schroon fine sandy loam, 3 to 8 percent slopes	374
421A—Ninigret fine sandy loam, cold, 0 to 3 percent slopes	375
423A—Sudbury sandy loam, cold, 0 to 3 percent slopes	376

424B—Shelburne fine sandy loam, 3 to 8 percent slopes	377
424C—Shelburne fine sandy loam, 8 to 15 percent slopes	379
424D—Shelburne fine sandy loam, 15 to 25 percent slopes	380
425B—Shelburne fine sandy loam, 3 to 8 percent slopes, very stony	381
425C—Shelburne fine sandy loam, 8 to 15 percent slopes, very stony	382
426D—Shelburne fine sandy loam, 15 to 35 percent slopes, extremely stony ...	384
427B—Ashfield fine sandy loam, 2 to 8 percent slopes, very stony	385
427C—Ashfield fine sandy loam, 8 to 15 percent slopes, very stony	386
428A—Ashfield fine sandy loam, 0 to 3 percent slopes	388
428B—Ashfield fine sandy loam, 3 to 8 percent slopes	389
428C—Ashfield fine sandy loam, 8 to 15 percent slopes	390
429A—Agawam fine sandy loam, cold, 0 to 3 percent slopes	391
429B—Agawam fine sandy loam, cold, 3 to 8 percent slopes	393
429C—Agawam fine sandy loam, cold, 8 to 15 percent slopes	394
433—Moosilauke sandy loam	395
434A—Merrimac sandy loam, cold, 0 to 3 percent slopes	396
434B—Merrimac sandy loam, cold, 3 to 8 percent slopes	397
434C—Merrimac sandy loam, cold, 8 to 15 percent slopes	398
435—Scarboro muck, cold	399
436—Halsey silt loam, cold	400
437—Wonsqueak mucky peat	401
438—Bucksport muck	402
440A—Boscawen gravelly sandy loam, 0 to 3 percent slopes	403
440C—Boscawen gravelly sandy loam, 3 to 15 percent slopes	404
440E—Boscawen gravelly sandy loam, 15 to 45 percent slopes	406
442—Brayton loam	407
443—Brayton-Loonmeadow complex, extremely stony	408
448B—Hogansburg loam, 3 to 8 percent slopes	410
449B—Hogansburg loam, 3 to 8 percent slopes, very stony	411
449C—Hogansburg loam, 8 to 15 percent slopes, very stony	412
450B—Pyrities loam, 3 to 8 percent slopes	413
450C—Pyrities loam, 8 to 15 percent slopes	414
450D—Pyrities loam, 15 to 25 percent slopes	415
451B—Pyrities loam, 3 to 8 percent slopes, very stony	416
451C—Pyrities loam, 8 to 15 percent slopes, very stony	417
451D—Pyrities loam, 15 to 25 percent slopes, very stony	419
457—Mudgepond silt loam, cold	420
458—Mudgepond and Alden soils, extremely stony, cold	421
501—Ondawa fine sandy loam	422
503—Rumney fine sandy loam	423
508—Medomak silt loam	424
Prime Farmland	427
Use and Management of the Soils	429
Interpretive Ratings	429
Rating Class Terms	429
Numerical Ratings	429
Crops and Pasture	430
Yields per Acre	431
Land Capability Classification	432
Forest Productivity and Management	433
Forest Productivity	434
Forest Management	434
Conservation and Environmental Plantings	436
Recreation	436

Wildlife Habitat	438
Hydric Soils	440
Connecticut Inland Wetland Soils	442
Engineering	442
Building Site Development	443
Sanitary Facilities	444
Construction Materials	446
Water Management	447
Soil Properties	449
Engineering Index Properties	449
Physical Properties	450
Chemical Properties	452
Water Features	452
Soil Features	454
Storm Water Runoff Management Systems	455
Classification of the Soils	457
Soil Series and Their Morphology	457
Agawam Series	458
Alden Series	459
Amenia Series	461
Ashfield Series	462
Bash Series	464
Belgrade Series	465
Berlin Series	466
Bernardston Series	468
Bice Series	469
Boscawen Series	471
Brancroft Series	472
Branford Series	474
Brayton Series Taxadjunct	475
Brayton Series Taxadjunct	477
Brimfield Series	479
Broadbrook Series	480
Brookfield Series	481
Bucksport Series	482
Canton Series	483
Catden Series	485
Charlton Series	486
Chatfield Series	487
Cheshire Series	489
Copake Series	490
Deerfield Series	491
Dummerston Series	493
Ellington Series	494
Elmridge Series	496
Enfield Series	497
Farmington Series	499
Fluvaquents	500
Fredon Series	501
Freetown Series	502
Fullam Series	503
Georgia Series	505
Gloucester Series	506
Groton Series	507

Hadley Series	509
Halsey Series.....	510
Hartford Series	511
Haven Series	513
Hero Series.....	514
Hinckley Series	515
Hogansburg Series	517
Hollis Series.....	518
Holyoke Series.....	519
Ipswich Series.....	520
Lanesboro Series	521
Leicester Series	523
Lim Series	525
Limerick Series	526
Loonmeadow series	528
Ludlow Series	529
Macomber Series	530
Manchester Series.....	531
Maybid Series	532
Medomak Series Taxadjunct.....	534
Menlo Series.....	535
Merrimac Series	537
Millsite Series	538
Montauk Series.....	539
Moosilauke Series	541
Mudgepond Series	542
Narragansett Series.....	544
Natchaug Series	545
Nellis Series.....	547
Ninigret Series	548
Occum Series	550
Ondawa Series	551
Pawcatuck Series	552
Paxton Series	553
Penwood Series.....	555
Pootatuck Series.....	556
Pyrities Series.....	557
Rainbow Series.....	559
Raynham Series	560
Raypol Series	562
Ridgebury Series Taxadjunct	563
Rippowam Series.....	565
Rumney Series	566
Saco Series	568
Scarboro Series	569
Schroon Series	570
Scitico Series	572
Shaker Series	573
Shelburne Series	575
Stockbridge Series	576
Sudbury Series	578
Suncook Series	579
Sutton Series	580
Taconic Series	582

Timakwa Series	583
Tisbury Series.....	584
Udifluvents	586
Udipsamments	587
Udorthents	588
Walpole Series.....	589
Wapping Series	590
Watchaug Series	591
Westbrook Series	593
Westminster Series.....	594
Wethersfield Series.....	595
Whitman Series	596
Wilbraham Series	598
Windsor Series	599
Winooski Series	600
Wonsqueak Series	602
Woodbridge Series	603
Yalesville Series.....	604
Formation of the Soils	607
Factors of Soil Formation.....	607
Processes of Soil Formation	609
References	611
Glossary	615
Tables	635
Table 1.—Temperature and Precipitation	636
Table 2.—Freeze Dates in Spring and Fall	646
Table 3.—Growing Season	656
Table 4.—Acreage and Proportionate Extent of the Soils	661
Table 5.—Prime and other Important Farmland	666
Table 6.—Non-Irrigated Yields by Map Unit Component	668
Table 7.—Acreage by Capability Class and Subclass	687
Table 8.—Forestland Productivity	688
Table 9.—Forestland Management.....	727
Table 10.—Hazard of Erosion and Suitability for Roads on Forestland	765
Table 11.—Damage by Fire and Seedling Mortality on Forestland	798
Table 12.—Conservation and Environmental Plantings	835
Table 13.—Recreation (Part 1)	886
Table 14.—Recreation (Part 2)	933
Table 15.—Wildlife Habitat	969
Table 16.—Connecticut Inland Wetlands	994
Table 17.—Building Site Development (Part 1)	995
Table 18.—Building Site Development (Part 2)	1029
Table 19.—Sewage Disposal	1073
Table 20.—Source of Sand and Gravel	1118
Table 21.—Construction Materials.....	1151
Table 22.—Ponds and Embankments	1198
Table 23.—Engineering Properties.....	1235
Table 24.—Physical Properties of the Soils.....	1418
Table 25.—Chemical Properties of the Soils	1479
Table 26.—Water Features	1525
Table 27.—Soil Features	1582
Table 28.—Storm Water Runoff Systems	1607

Table 29.—Taxonomic Classification of the Soils	1700
Table 30.—Relationships Among Parent Material, Dominant Texture, and Drainage of the Soils	1703

Issued 2008

Foreword

This soil survey contains information that affects land use planning in this survey area. It contains predictions of soil behavior for selected land uses. The survey also highlights soil limitations, improvements needed to overcome the limitations, and the impact of selected land uses on the environment.

This soil survey is designed for many different users. Farmers, foresters, and agronomists can use it to evaluate the potential of the soil and the management needed for maximum food and fiber production. Planners, community officials, engineers, developers, builders, and home buyers can use the survey to plan land use, select sites for construction, and identify special practices needed to ensure proper performance. Conservationists, teachers, students, and specialists in recreation, wildlife management, waste disposal, and pollution control can use the survey to help them understand, protect, and enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. The information in this report is intended to identify soil properties that are used in making various land use or land treatment decisions. Statements made in this report are intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are shallow to bedrock. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

These and many other soil properties that affect land use are described in this soil survey. Broad areas of soils are shown on the general soil map. The location of each soil is shown on the detailed soil maps. Each soil in the survey area is described. Information on specific uses is given for each soil. Help in using this publication and additional information are available at the local offices of the Natural Resources Conservation Service or the Cooperative Extension Service.

Margo Wallace
State Conservationist
Natural Resources Conservation Service

Soil Survey of The State of Connecticut

By Marjorie Faber, Natural Resources Conservation Service

Fieldwork by Barbara Alexander, Mark Beroz, Christine Clarke, Marjorie Faber, Nancy Gottung, Kipen Kolesinskas, Shawn McVey, Donald Parizek, Alfred Roberts, Roy Shook, Glenn Stanisewski, and Deborah Surabian

United States Department of Agriculture, Natural Resources Conservation Service, in cooperation with
The Connecticut Agricultural Experiment Station, the Storrs Agricultural Experiment Station, and the Connecticut Department of Environmental Protection

General Nature of the Survey Area

Connecticut is the southernmost state in New England. The total area of the state is 3,194,700 acres, or 4,992 square miles. Connecticut is bordered by New York State on the west, Rhode Island on the east, Massachusetts on the north, and Long Island Sound on the south ([fig. 1](#)). The three major waterways draining the state are the Housatonic River, the Connecticut River, and the Thames River. These rivers drain into Long Island Sound.

The coastal slope and central valley are mostly flat and contain most of the population of the state. The eastern uplands and western uplands are mostly in farmland, open space, and forests. Agriculture is no longer as prominent as it once was in Connecticut because of the displacement of agricultural lands by expanding suburbs, new ex-urban areas (development not connected to older urban areas), and industrial growth.

Below is general information about the climate, physiography, relief, and drainage, water supply, agriculture, recreation, history, and industry and transportation of Connecticut.

Climate

Prepared by the Natural Resources Conservation Service National Water and Climate Center, Portland, Oregon.

Climate tables are created from climate stations in Bridgeport, Cockaponset Ranger Station, Falls Village, Groton, Hartford Brainard Field, Mt. Carmel, Norfolk 2 SW, Shepaug Dam, Storrs, and West Thompson Lake, Connecticut.

Thunderstorm days, relative humidity, percent sunshine, and wind information are estimated from First Order stations Bridgeport and Hartford, Connecticut.

[Table 1](#) gives data on temperature and precipitation for the survey area as recorded at these stations in the period 1961 to 1990, except 1966 to 1990 at West Thompson Lake. [Table 2](#) shows probable dates of the first freeze in fall and the last freeze in spring. [Table 3](#) provides data on the length of the growing season. In the narrative below, normals are for these averaging periods, while extremes are for the full period of record for each station. These generally extend from 1948 to 1998,



Figure 1.—The location of the State of Connecticut.

except records at Storrs begin in 1900, in 1920 at Hartford, in 1936 at Mt. Carmel, and in 1943 at Norfolk.

In winter, average temperatures range from 21.5 degrees F at Norfolk and the higher hills in the northwest, to 31.2 degrees at Bridgeport. Most of the central interior averages around 27 degrees, and the coastal areas generally average around 30. Average daily minimum temperatures in winter range from around 13 degrees in Norfolk and most of the higher northwest region, to 24 degrees at Bridgeport. Most of the central portion of the state averages around 20 degrees. The lowest temperatures on record include -25 at Norfolk on February 16, 1943; -24 at Hartford on the same date; -20 at Storrs on February 9, 1934; -30 at Falls Village on January 22, 1961; and -7 at Bridgeport on January 22, 1984.

In summer, average temperatures range from around 65 degrees at Norfolk to around 70 at Hartford and 71 at Bridgeport. Average daily maximum temperatures in summer range from 75 at Norfolk to around 78 along most of the coast, and around 81 at Hartford, Cockaponset, Falls Village and most of the interior. Highest temperatures ever recorded include: 101 at Falls Village on September 2, 1953; 101 at Storrs on July 6, 1919; 102 at Hartford on August 3, 1975; 103 at Bridgeport on July 22, 1957; and 93 at Norfolk on August 6, 1955.

Growing degree days are shown in [table 1](#). They are equivalent to “heat units”. During the month, growing degree days accumulate by the amount that the average temperature each day exceeds a base temperature (40 degrees F). The normal monthly accumulation is used to schedule single or successive plantings of a crop between the last freeze in spring and the first freeze in fall.

Average annual precipitation is generally between 44 and 54 inches across the state. Lowest average totals are 42.65 inches at Hartford and 41.67 at Bridgeport (the central Connecticut River Valley, and along the southwest coast). Highest annual averages include 51.24 inches at Norfolk, 50.32 inches at Cockaponset, and 49.54 inches at West Thompson Lake, with most of the highlands north of Bridgeport and also northwest of Torrington receiving between 52 and 54 inches annually. Of these annual average totals, around 22 to 25 inches, or about 50 percent of the annual total, generally falls during the growing season, which is usually May to October. The heaviest 1-day precipitation events during the periods of record included: 9.50 inches at Falls Village on August 19, 1955; 10.47 inches at Cockaponset on June 6, 1982; 10.67 inches at Norfolk on August 19, 1955; 6.10 inches at Hartford on September 20, 1938; 6.21 inches at Storrs, also on September 21, 1938; and 7.43 inches at

Groton on September 21, 1961. Thunderstorms occur on about 21 days each year, and most occur in between May and August.

Average seasonal snowfall is quite variable across the state, with lowest average totals along the coast where many winter events fall as rain or freezing rain. This includes 23 inches annually at Groton, and 26 inches at Bridgeport. Inland, annual average snowfall is greater, including 32 inches at Mt. Carmel, 36 inches at Cockaponset, 28 inches at Hartford, 38 inches at Storrs, and 34 inches in the northeast at West Thompson Lake. The snowiest area is the northwest, where annual averages include 40 inches at Falls Village, 54 inches at Shepaug Dam, and 99 inches at Norfolk. These totals are also aligned with the average number of days per year that have at least 1 inch of snow on the ground, which ranges from 26 at Groton and 29 at Bridgeport; to 39 at Hartford and 33 at Cockaponset; and to 110 days per year at Norfolk and most of the higher terrain in the northwest.

Greatest snow depths at any one time during the periods of record included: 20 inches at Bridgeport on January 10, 1996; 19 inches at Groton on February 4, 1961; 23 inches at Cockaponset on February 5, 1961; 45 inches at Falls Village on February 10, 1969; 30 inches at West Thompson Lake on February 8, 1978; and 55 inches at Norfolk on February 5, 1961.

Heaviest 1-day snowfall records include: 25.7 inches at Norfolk on February 10, 1969; 17.7 inches at Hartford on December 29, 1945; 19.8 inches at Cockaponset on February 8, 1978; 24.0 inches at Falls Village on January 20, 1961; 20.0 inches at Shepaug Dam on February 12, 1983; and 16.0 inches at Bridgeport on December 19, 1948.

The average relative humidity in mid-afternoon is generally around 60 percent along the coast and 50 to 55 percent inland. Humidity is higher at night, and the average at dawn is generally around 70 percent, except around 80 to 85 percent in the late summer and fall. The sun shines about 60 percent of the time in summer and about 50 to 55 percent in winter. The prevailing wind is from the northwest from November to March, and from the south or southwest for much of the rest of the year. Northeast winds predominate in September and much of October. Wind direction is quite variable; although, can be influenced by local topography. Average wind speeds are highest, generally around 13 miles per hour along the coast and around 10 mph in inland valleys, in the winter and early spring.

Physiography, Relief, and Drainage

Connecticut lies within the New England physiographic province. This area is divided into four physiographic regions: Western Uplands, Central Valley, Eastern Uplands, and Coastal Slope. These physiographic regions are each characterized by different landscapes and geology. They may be categorized in still further detail (Bell, 1985).

Combining physiography with vegetation and other factors, an ecoregional classification has been developed for the state. The ecoregions are characterized by a distinctive pattern of landscapes and regional climate as expressed by the vegetation composition and pattern and the presence or absence of certain indicator species and species groups (Dowhan and Craig, 1976). Additional work by the USDA Natural Resources Conservation Service on describing soils and ecoregions is anticipated.

The Central Valley, a north-south trending physiographic region, lies between the Western and Eastern Uplands. It is a broad, flat valley developed on fairly weak, tilted, stratified rocks. The rocks are Triassic in age, from which more resistant Triassic lava flows of basalt, known as traprock, are exposed primarily at high ridges such as Vexation Hill, Cedar Mountain, and Talcott Mountain. In this region, most soils formed

in glacial till, glaciofluvial, and glaciolacustrine deposits of sandstone, shale, conglomerate, and basalt.

Dividing the Central Valley longitudinally is the Talcott Mountain Range. The area east of the mountain range is drained by the Connecticut River and the smaller area west of the range is drained by the Farmington River (a tributary of the Connecticut River). The Connecticut Valley consists of flood plains along the Connecticut and Farmington rivers, with nearly level to sloping terraces, low glacial upland hills, and narrow ridges of basalt. Elevations above sea level in this region range from about 10 feet on the flood plain of the Connecticut River, to about 500 to 950 feet on the highest basalt ridges.

Flanking the Central Lowlands to the west are the Western Uplands which include the Northwest Highlands. To the east are the Eastern Highlands. The Western and Eastern Uplands consist mainly of till plains and drumlins dissected by narrow valleys that are underlain by metamorphic rocks of early Paleozoic age. Some areas of the Western Uplands are dominated by soils formed in glacial till and glaciofluvial deposits derived from limestone or dolomite. The Eastern Uplands are dominated by soils formed in glacial till and glaciofluvial deposits derived from granite, gneiss, or schist.

Elevations above sea level in the Western Uplands range from 350 feet on the lowest valley floor, to about 2,350 feet on the highest peak in the northwest corner of the state. The area is sloping to steep, and drainage generally runs south and southeast into the Farmington River or Housatonic River.

Elevations above sea level in the Eastern Uplands range from about 300 feet on the lowest valley floor, to over 1,000 feet on the highest hills. The area is sloping to hilly and drainage generally flows to the west-southwest into the Connecticut River or south into the Thames River.

The Coastal Slope includes the areas significantly affected by ocean processes. This area is rather rocky, where hills extend out into the water and coves lie in between. It is a sheltered coastline with an abundance of calm, secure harbors. The Coastal Slope also includes a scattering of small islands along the coastline formed by a recessional moraine. The two small, rocky recessional moraines, the Madison and the Old Saybrook, each mark a short halt in the retreat of glaciation from New England. Some areas of the Coastal Slope are affected by tidal water and are dominated by soils formed in organic material.

Within 10 miles of Long Island Sound, elevations in the Coastal Slope range from sea level to a few hilltops approximately 400 feet above sea level. The area is level and drainage flows into the Long Island Sound. Many Connecticut rivers that flow into the Long Island Sound are tidal rivers whose water levels respond to the rise and fall of tide. Examples include the Niantic, Connecticut, Thames, and Housatonic rivers.

Water Supply

Connecticut's ground water resources are the source of drinking water for approximately one million people, which is about one-third of the state's population. In addition, it is also the baseflow for many rivers, streams, and wetlands. Thus, the quality of ground water plays an important role in the quality of surface water resources.

Ground water is withdrawn through 1,200 community wells (public supply), more than 3,000 non-community wells (schools, hospitals, etc.), and about 250,000 individual private home wells. In all, about 83 percent of the state's population receives its water from public water utilities while the remainder relies on individual private wells.

Connecticut has two major types of aquifers, glaciofluvial and bedrock. Glaciofluvial aquifers are composed of unconsolidated, stratified sand and gravel of

glacial origin. These aquifers line the larger river valleys, are the most productive, and are the primary source of ground water for water utilities that serve populations of greater than 1,000 people. Bedrock aquifers are composed of sedimentary, crystalline or carbonate-rock. These aquifers underlie most of the state and are the primary source of ground water for non-community water supplies and private wells.

Connecticut's aquifers are shallow, typically less than 300 feet deep with the water table within 50 feet of the land surface, and are therefore susceptible to contamination. Some of the sources of contamination include historical industrial activities, underground storage tanks, landfills, salt storage facilities, road salt application, pesticide and fertilizer application, and numerous accidental spills of chemicals. On average, 75 to 100 contaminated drinking water wells are identified every year. Thus, all sources of water must be continuously protected from intensive development and potentially deleterious land uses.

Agriculture

About 10 percent of the land of Connecticut is in cropland or pastureland and about 55 percent is in forests (1997 NRI). Important agriculture and aquaculture products of the state include: oysters, milk, poultry, beef, greenhouse and nursery crops, fruit, tobacco, maple syrup, Christmas trees, and mushrooms.

Recreation

With nearly two-thirds of the state classified as open land, Connecticut has many areas of scenic, geologic, and historical interest. There are more than 100 state parks and forests that provide a variety of recreational uses including boating, fishing, hunting, camping, horseback riding, snowmobiling, hiking, and picnicking. Public lands available for recreation include Patchaug State Forest, Hammonasset Beach State Park, West Rock Ridge State Park, Dinosaur Park, and Kent Falls State Park.

The Appalachian Trail, stretching from Springer Mountain, Georgia, to Mt. Katahdin, Maine, passes through 14 states, including Connecticut. The trail, started in 1921 and completed in 1937, is known as the longest national park in the world. It is 2,150 miles long, with over 50 miles and nearly 7,000 acres of National Park land along the Appalachian Trail corridor in Connecticut.

Many soils in the state are well suited to the development of recreational areas. Wooded slopes, rolling topography, exposed rock formations, and many streams provide a variety of possibilities for recreation.

History

The native Americans who settled in what was then known as Quinnehtukqut (a Mohegan word meaning beside the Long Tidal River), migrated in series. There were four distinct groups of Algonkians, followed by the Delaware Indians. Over time, the tribes from earlier migrations formed affiliations. This led to further localization of smaller, scattered tribes.

The Pequots were the last to migrate, settling in 1600. The northeastern portion of the state (along with part of Massachusetts) was occupied by the Nipmuck Tribe; the southeastern section by the Mohegan and Pequot Tribes; the River Valley by the River Tribes; and the western section by two groups—the Mahicans and the Mattabesec-Wappinger Confederacy. The population remained constant for almost 30 years.

In late 1629, a powerful and popular preacher named Thomas Hooker began looking for relief from England's religious persecution. Hooker, who later became known as the Father of Connecticut, eventually fled to the New World. He, along with several other leaders, arrived in Boston in 1633. Cramped living conditions made

Hooker and his congregation to decide to move to the rich lands of the Connecticut River Valley. In June 1636, Thomas Hooker, 100 people, 160 head of cattle, and a number of swine and goats left Newtown, Massachusetts for Connecticut. They settled in an area they called Newtown, now Hartford.

In the following years, Connecticut was ruled by a governor and eight magistrates. In 1638, Hooker delivered a sermon that put forth the idea it was the people's God-given right to select magistrates, and that they also had a right to limit the elected magistrate's powers. Written and adopted in 1639, the Fundamental Orders became the beginning of government in Connecticut. It was the first constitution adopted in North America, establishing representative government, thus giving Connecticut the nickname The Constitution State.

As settlements grew, towns were formed and town greens set aside. The greens were generally set aside for public uses such as grazing cattle or sheep, military parades, marketplaces, and as general meeting places.

On January 9, 1788, Connecticut became the fifth territory to receive statehood and in that same year built the first State House in America. Currently, Connecticut consists of 169 towns, and eight counties—Fairfield, Hartford, Middlesex, New Haven, New London, Litchfield, Tolland, and Windham. As of the 2000 Census, the state's population has swelled to over 3.4 million people, Fairfield County being the most populous, and Windham the least.

Connecticut has a population of over 9,000 American Indians. The state recognizes the Mashantucket (Western Pequot), Paucatuck (Eastern Pequot), Mohegan, Golden Hill Paugussett (who boast the oldest Indian reservation in the country), Schaghticoke, and two Mattabesic tribes.

Industry and Transportation

Industry and transportation systems have been strongly linked in Connecticut for several centuries. The types of industries, their locations relative to geographic features (e.g. rivers and ports), and the scales at which industry took place have changed in the last three centuries as have the modes and networks of transportation.

Many early roads in Connecticut were well-worn pathways of American Indians. In 1633, Connecticut's first primitive road was established and by 1635, the now named Boston Post Road began to establish settlements and connect cities such as Hartford and Windsor. By 1671, roads were built connecting Hartford to Boston, New York, and Providence.

Connecticut's waterways also provided a primary means for transporting goods and people. By the early 1800's, the state had developed several canals and locks. The now Windsor Locks in Enfield was built in 1824 and the Farmington Canal, connecting New Haven and Northampton, Massachusetts, was completed in 1835.

In the early 1800s industrialization began to transform classic New England towns. Towns soon became centered around factories located in river valleys beside streams rather than a central town green bordered by colonial homes and a church. In 1806, the Town of Seymour became the first planned and established factory-town in America.

Much of Connecticut's industry was small in scale and often run by families. In the 19th century, some of the products local industries produced included woolens, sewing machines and spools, arms and hardware, agricultural implements, greeting cards, glass, textiles, silk velvet, and clocks. Beginning in the 1830s, railroads provided a flexible and effective way to transport manufactured goods. Shipping between railroads and steamship lines were important to port cities such as Bridgeport, New Haven, New London, and Norwich.

Journeying by railway and steamboat was popular, but more hazardous than by stagecoach or foot. In 1810, the nation's first insurance company, ITT Hartford Group, Inc., was opened enabling people to get insurance covering loss of life or personal injury incurred while traveling.

In the 20th century, highways and roadways were improved with the completion of the interstate highway system in 1958. Suddenly, trucks and automobiles became the primary means of transportation of goods and individuals. The automobile also encouraged individuals to move from the city center to suburban areas.

As technology advanced, a concentration of companies and industries settled into Connecticut. The boom was due to the interconnection of markets served and type of products produced, as well as the presence of suppliers, trade associations, and educational institutions. Through the years, Connecticut's industrial genius gave rise to inventions such as the first helicopter—designed by Igor Sikorsky in 1939; the first nuclear submarine—launched in New London in 1954; and the first artificial heart—invented by Dr. Robert K. Jarvik in 1982.

In the 21st century, some of the industries key to Connecticut's economic competitiveness in the global economy include manufacturing, transportation equipment, financial services, telecommunications and information, health care services, high technology, and tourism. Only a few visible signs of industrial activities of the past such as buried foundations, waterworks, and abandoned machinery and products remain.

Today, Connecticut has an interconnecting transportation system that includes interstate highways, aviation facilities, rail and bus service, ports, and ferries. A network of interstate highway systems provides major routes for travelers and trucking. The Merritt Parkway, one of the oldest in America, extends from Meriden to the New York state line.

The state boasts 123 public and private aviation facilities in addition to six state-owned airports, the largest being Bradley International Airport in Windsor Locks. Bradley is New England's second largest airport, having 21 carriers with over 250 flights daily.

Rail passenger service connects Metro-North, Amtrak, and Shore Line East, and now includes a high-speed rail service from Washington to Boston. Freight rail services are also available, along with local bus service in eighteen transit areas.

Connecticut has three major port operations in Bridgeport, New Haven, and New London that handle cargo such as food, steel, tin, paper, woodpulp, lumber, and automobiles. The state also has two seasonal river ferries, one in the Rocky Hill-Glastonbury area, and the other in Chester-Hadlyme. The Rocky Hill-Glastonbury ferry is the nation's oldest continuously operating commuter boat, providing service since 1655.

How This Survey Was Made

This survey was made to provide information about the soils and miscellaneous areas in the survey area. The information includes a description of the soils and miscellaneous areas and their location and a discussion of their suitability, limitations, and management for specified uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They dug many holes to study the soil profile, which is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

The soils and miscellaneous areas in the survey area are in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area.

Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept or model of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

The descriptions, names, and delineations of the soils in this survey area do not fully agree with those of the soils in adjacent survey areas. Differences are the result of a better knowledge of soils, modifications in series concepts, or variations in the intensity of mapping or in the extent of the soils in the survey area.

General Soil Map Units

The general soil map associated with this publication is the Digital General Soil Map of the United States, formerly known as STATSGO. This map shows broad areas with a distinctive pattern of soils, relief, and drainage. The digital soil survey area maps and attribute tables are available for download from the Soil Data Mart of the Natural Resources Conservation Service, accessed at <http://soils.usda.gov>. Each map unit on the Digital General Soil Map of the U.S. is a unique natural landscape. Typically, a map unit is multi-state in extent and consists of one or more major soils and some minor soils. It is named for the major soils. The soils making up one map unit can occur in other units but in a different pattern. Some of the soil names used in these map units are common to and dominant in adjacent states only, and were not mapped in the detailed *Soil Survey of the State of Connecticut*.

The Digital General Soil Map of the U.S. can be used to compare the suitability of large areas for general land uses. Areas of suitable soils can be identified on the map. Likewise, areas where the soils are not suitable can be identified.

Because of its small scale (1:250,000), the map is not suitable for planning the management of a farm or field or for selecting a site for a road or building or other structure. The soils in any one map unit differ from place to place in slope, depth, drainage, and other characteristics that affect management.

The 33 Digital General Soil Map of the U.S. soil map units in Connecticut are described in the following pages.

S1438. Charlton-Canton-Paxton

Gently sloping to steep, well drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 34 percent of the state. The landscape is predominantly upland hill landforms. Charlton and Canton soils formed in melt-out till and Paxton soils formed in lodgement till. Stones and boulders are common on the surface in many places.

Charlton soils make up about 23 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Charlton soils are gently sloping to steep and are on hills.

Canton soils make up about 15 percent of this map unit. They are very deep, well drained, loamy over sandy and gravelly soils with rapid permeability in the substratum. Canton soils are gently sloping to steep and are on hills.

Paxton soils make up about 14 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Paxton soils are gently sloping to steep and are on hills and drumlins.

Soils of minor extent make up about 48 percent of this map unit. They are mainly Leicester, Woodbridge, Sutton, Chatfield, and Hollis soils. Most areas of this map unit are in woodland or cropland. Some areas are in community development. The soils of this map unit are suited to forestry, cropland, community development, and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in Paxton soils and the rapid permeability of the substratum in the

Canton soils. Stones and boulders need to be removed from the surface in some areas.

S1439. Charlton-Chatfield-Hollis

Gently sloping to very steep, well drained and somewhat excessively drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 10 percent of the state. The landscape is predominantly upland hill and ridges landscapes. The soils formed in melt-out till. Stones and boulders are common on the surface in most places, and many areas have outcrops of bedrock.

Charlton soils make up about 31 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Charlton soils are gently sloping to steep and are on hills.

Chatfield soils make up about 18 percent of this map unit. They are moderately deep to bedrock, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Chatfield soils are gently sloping to steep and are on bedrock-controlled hills and ridges.

Hollis soils make up about 17 percent of this map unit. They are shallow to bedrock, somewhat excessively drained, loamy soils with moderate or moderately rapid permeability in the substratum. Hollis soils are gently sloping to very steep and are on bedrock-controlled hills and ridges.

Soils of minor extent make up about 34 percent of this map unit. They are mainly Leicester, Rock outcrop, Canton, Catden, Sutton, Hinckley, and Paxton soils. Most areas of this map unit are in woodland and some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factor for community development is the depth to bedrock in Chatfield and Hollis soils. Stones and boulders need to be removed from the surface in some areas.

S1440. Woodbridge-Paxton-Ridgebury

Nearly level to steep, poorly drained to well drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 16 percent of the state. The landscape is predominantly upland hill and drumlin landforms with a north-south orientation. The soils formed in lodgement till. Stones and boulders are common on the surface in many places.

Woodbridge soils make up about 32 percent of this map unit. They are very deep, moderately well drained, loamy soils with slow or very slow permeability in the substratum. Woodbridge soils are nearly level to strongly sloping and are on hills and drumlins.

Paxton soils make up about 28 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Paxton soils are gently sloping to steep and are on hills and drumlins.

Ridgebury soils make up about 11 percent of this map unit. They are very deep, poorly drained, loamy soils with slow or very slow permeability in the substratum. Ridgebury soils are nearly level to gently sloping and are in depressions and drainageways.

Soils of minor extent make up about 29 percent of this map unit. They are mainly Canton, Charlton, Catden, Chatfield, Hollis, and Merrimac soils. Most areas of this map unit are in woodland or cropland. Some areas are in community development. The soils of this map unit are suited to forestry, cropland, and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in the substratum and the seasonal high water table in Woodbridge and

Ridgebury soils. Stones and boulders need to be removed from the surface in some areas.

S1441. Hinckley-Agawam-Merrimac

Nearly level to steep, well drained to excessively drained, sandy to loamy soils; on outwash plains and terraces

The soils in this map unit make up about 11 percent of the state. The landscape is predominantly outwash plains, stream terraces, kames, and eskers in valleys between glacial till uplands. The soils formed in glacial outwash.

Hinckley soils make up about 34 percent of this map unit. They are very deep, excessively drained, soils with very rapid permeability in the substratum. Hinckley soils are nearly level to steep and are on kames, eskers, and outwash plains and terraces.

Agawam soils make up about 15 percent of this map unit. They are very deep, well drained, loamy soils with rapid or very rapid permeability in the substratum. Agawam soils are nearly level to strongly sloping and are on outwash plain and stream terraces.

Merrimac soils make up about 13 percent of this map unit. They are very deep, somewhat excessively drained, sandy soils with rapid or very rapid permeability in the substratum. Merrimac soils are nearly level to strongly sloping and are on outwash plains, stream terraces, and kames.

Soils of minor extent make up about 38 percent of this map unit. They are mainly Ninigret, Paxton, Rippowam, Timakwa, Walpole, Occum, Charlton, and Chatfield soils. Most areas of this map unit are in cropland or community development. The soils of this map unit are suited to cropland and community development, and are also a source of sand and gravel.

The major limiting factors for community development are rapid and very rapid permeability in the substratum.

The major limiting factors for cropland are droughtiness in Hinckley soils and steep slopes in some areas limiting the use of farming equipment.

S1442. Catden-Merrimac-Saco

Nearly level to strongly sloping, very poorly drained to somewhat excessively drained, organic, sandy, and silty soils; on drainageways, stream terraces, and outwash plains

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly stream terrace landforms.

Catden soils make up about 31 percent of this map unit. They are very deep, very poorly drained, woody organic soils with variable mineral substrata below 51 inches. Catden soils are nearly level and are in depressions.

Merrimac soils make up about 15 percent of this map unit. They are very deep, somewhat excessively drained, sandy soils with rapid or very permeability in the substratum. Merrimac soils are nearly level to strongly sloping and are on outwash plains, stream terraces, and kames.

Saco soils make up about 8 percent of this map unit. They are very deep, very poorly drained, silty soils formed in alluvium. Saco soils are nearly level and are in depressions and drainageways.

Soils of minor extent make up about 46 percent of this map unit. They are mainly Lim, Natchaug, Timakwa, Paxton, Walpole, Charlton, Deerfield, and Sutton soils. Most areas of this map unit are in woodland and wildlife habitat. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for forestry are the seasonal high water table in Catden and Saco soils and the low load bearing strength of these soils.

S1443. Urban land-Udorthents-Cheshire

Nearly level to very steep, variably drained, sandy to loamy soils; on mostly human influenced landforms

The soils in this map unit make up about 4 percent of the state. The natural landscapes have been altered by human activity.

Urban lands make up about 40 percent of this map unit. Urban land is land mostly covered by buildings, streets, parking lots or other impervious surfaces.

Udorthents soils make up about 30 percent of this map unit. They occur in cut and fill areas, road and railroad beds, and on spoil piles with a wide range of soil textures and permeability. The substratum often resembles the original, preconstruction soil existing in the area. Udorthents soils are nearly level to very steep.

Cheshire soils make up about 5 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Cheshire soils are gently sloping to steep and are on upland hills and till plains.

Soils of minor extent make up about 25 percent of this map unit. They are mainly Charlton, Penwood, Windsor, Leicester, Walpole, Wethersfield, and Rippowam soils. Most areas of this map unit are in community development and the soils have been modified to be suited to community development.

S1444. Stockbridge-Charlton-Mudgepond

Gently sloping to steep, poorly drained to well drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly upland hill and drainageway landforms in the Northwestern part of Connecticut. The soils formed in melt-out till (fig. 2).

Stockbridge soils make up about 37 percent of this map unit. They are very deep, well drained, loamy soils with moderately slow permeability in the substratum. Stockbridge soils are gently sloping to steep and are on hills.




Figure 2.—Typical pattern of soils in the Stockbridge-Charlton-Mudgepond general soil map unit.

Charlton soils make up about 11 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Charlton soils are gently sloping to steep and are on hills.

Mudgepond soils make up about 10 percent of this map unit. They are very deep, very poorly drained, loamy soils with moderate permeability in the substratum. Mudgepond soils are nearly level to gently sloping and are in depressions and drainageways.

Soils of minor extent make up about 42 percent of this map unit. They are mainly Charlton, Georgia, Amenia, Copake, Catden, Nellis, Farmington, Chatfield, and Hollis soils. Most areas of this map unit are in cropland or woodland. Some areas are in community development. The soils of this map unit are suited to these uses.

The major limiting factor for community development is the seasonal high water table in Mudgepond soils.

S1445. Copake-Groton-Hero

Nearly level to steep, moderately well drained to excessively drained, loamy to sandy soils; on outwash valley uplands

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly outwash plain, terrace, and kame landforms in upland valleys of Northwestern Connecticut. The soils formed in outwash.

Copake soils make up about 33 percent of this map unit. They are very deep, well drained, loamy soils with rapid or very permeability in the substratum. Copake soils are nearly level to strongly sloping and are on kame, terrace, and plain landforms.

Groton soils make up about 14 percent of this map unit. They are very deep, excessively drained, sandy soils with rapid or very permeability in the substratum. Groton soils are nearly level to steep and are on kame, esker, terrace and plain landforms.

Hero soils make up about 12 percent of this map unit. They are very deep, moderately well drained, loamy soils with rapid or very rapid permeability in the substratum. Hero soils are nearly level to gently sloping and are on terrace and plain landforms.

Soils of minor extent make up about 41 percent of this map unit. They are mainly Fredon, Saco, Charlton, Catden, Lim, Stockbridge, Chatfield, Hollis, and Nellis soils. Most areas of this map unit are in cropland or community development. Some areas are in woodland. The soils of this map unit are suited to these uses and are also a source of sand and gravel.

The major limiting factors for community development are the rapid or very rapid permeability of these soils and the seasonal high water table in Hero soils.

S1446. Windsor-Merrimac-Hinckley

Nearly level to steep, somewhat excessively drained to excessively drained, sandy and gravelly soils; on outwash plains, terrace, kames, and eskers

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly outwash plain, terrace, esker, and kame landforms. The soils formed in outwash.

Windsor soils make up about 46 percent of this map unit. They are very deep, excessively drained, sandy soils with rapid or very permeability in the substratum. Windsor soils are nearly level to strongly sloping and are on kames, terraces and plains.

Merrimac soils make up about 10 percent of this map unit. They are very deep, somewhat excessively drained, sandy soils with rapid or very rapid permeability in the substratum. Merrimac soils are nearly level to strongly sloping and are on kames, terraces and plains.

Hinckley soils make up about 10 percent of this map unit. They are very deep, excessively drained, sandy and gravelly soils with very rapid permeability in the substratum. Hinckley soils are nearly level to steep on kames, terraces, plains, and eskers.

Soils of minor extent make up about 34 percent of this map unit. They are mainly Agawam, Raypol, Elmridge, Occum, Saco, Cheshire, Lim, Scarboro, Shaker, and Winooski soils. Most areas of this map unit are in woodland or community development. Some small areas are in cropland. The soils of this map unit are suited to woodland and community development and also are a source of sand and gravel.

These soils are droughty for cropland, unless irrigation is provided. Steep slopes in some areas limit the use of farming equipment.

The major limiting factor for community development is the rapid or very rapid permeability of Windsor and Hinckley soils.

S1447. Windsor-Agawam-Haven

Nearly level to strongly sloping, well drained to excessively drained, sandy and loamy soils; on outwash plains, terraces, and kames

The soils in this map unit make up about 2 percent of the state. The landscape is predominantly upland hill, drumlin, and drainageway landforms with a north-south orientation. The landscape is predominantly outwash plain, terrace, and kame landforms. The soils formed in outwash.

Windsor soils make up about 22 percent of this map unit. They are very deep, excessively drained, sandy soils with rapid or very rapid permeability in the substratum. Windsor soils are nearly level to strongly sloping and are on kames, terraces and plains.

Agawam soils make up about 15 percent of this map unit. They are very deep, well drained, loamy soils with rapid or very rapid permeability in the substratum. Agawam soils are nearly level to strongly sloping and are on outwash plains and terraces.

Haven soils make up about 15 percent of this map unit. They are very deep, well drained, loamy soils with very rapid permeability in the substratum. Haven soils are nearly level to strongly sloping and are on outwash plains and terraces.

Soils of minor extent make up about 48 percent of this map unit. They are mainly Manchester, Walpole, Ninigret, Elmridge, Typic Udorthents, Rippowam, Timakwa, Wapping, and Saco soils. Most areas of this map unit are in cropland or community development. Some areas are in woodland. The soils of this map unit are suited to these uses, although Windsor soil can be droughty for cropland unless irrigation is provided.

The major limiting factor for community development is the rapid or very rapid permeability of Windsor and Agawam soils.

S1448. Brancroft-Scitico-Shaker

Nearly level to strongly sloping, poorly drained to moderately well drained, silty soils; on lake plain, terrace, and drainageway landforms

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly lake plain landforms. The soils formed in silty and clayey glaciolacustrine materials.

Brancroft soils make up about 24 percent of this map unit. They are very deep, moderately well drained, silty and clayey soils with very slow permeability in the substratum. Brancroft soils are nearly level to sloping and are on terraces.

Scitico soils make up about 17 percent of this map unit. They are very deep, poorly drained, clayey soils with very slow permeability in the substratum. Scitico soils are nearly level and are in depressions and drainageways.

Shaker soils make up about 14 percent of this map unit. They are very deep, poorly drained, clayey soils with slow or very slow permeability in the substratum. Shaker soils are nearly level and are in depressions and drainageways.

Soils of minor extent make up about 45 percent of this map unit. They are mainly Elmridge, Maybid, Scarboro, Broadbrook, Belgrade, Lim, Ludlow, Ninigret, Wilbraham, and Timakwa soils. Most areas of this map unit are in woodland, hayland, or pasture. Some areas are in community development. The soils of this map unit are suited to forestry, hayland, and pastureland.

The major limiting factors for community development are the slow to very slow permeability of these soils and the seasonal high water table in Scitico and Shaker soils.

S1449. Brookfield-Brimfield-Leicester

Nearly level to steep, poorly drained to somewhat excessively drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly upland hill and ridge landforms in Northeastern Connecticut. The soils formed in melt-out till. Stones and boulders are common on the surface in most places, and many areas have outcrops of bedrock.

Brookfield soils make up about 40 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Brookfield soils are gently sloping to steep and are on bedrock-controlled hills and ridges.

Brimfield soils make up about 21 percent of this map unit. They are shallow to bedrock, somewhat excessively drained, loamy soils with moderate or moderately rapid permeability in the substratum. Brimfield soils are gently sloping to steep and are on bedrock-controlled hills and ridges.

Leicester soils make up about 12 percent of this map unit. They are very deep, poorly drained, loamy soils with moderate to rapid permeability in the substratum. Leicester soils are nearly level to gently sloping and are in depressions and drainageways.

Soils of minor extent make up about 27 percent of this map unit. They are mainly Sutton, Paxton, Catden, Hinckley, and Rippowam soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to these uses.

The major limiting factor for community development is the shallow to bedrock in Brimfield soils. Stones and boulders need to be removed from the surface in some areas.

S1451. Holyoke-Rock outcrop-Yalesville

Gently sloping to very steep, well drained, silty to loamy soils; on glacial till uplands

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly bedrock-controlled upland hill and ridge landforms in the Connecticut Valley with a north-south orientation. The soils formed in melt-out till. Stones and boulders are common on the surface in most places, and many areas have outcrops of bedrock.

Holyoke soils make up about 35 percent of this map unit. They are shallow to bedrock, well drained, silty soils with moderate permeability in the substratum. Holyoke soils are gently sloping to steep and are on bedrock-controlled hills and ridges.

Rock outcrop makes up about 18 percent of this map unit and occurs on bedrock-controlled landforms with slopes that are sometimes very steep.

Yalesville soils make up about 14 percent of this map unit. They are moderately deep to bedrock, well drained, loamy soils with moderately rapid permeability in the substratum. Yalesville soils are gently sloping to strongly sloping and are on bedrock-controlled hills and ridges.

Soils of minor extent make up about 33 percent of this map unit. They are mainly Wilbraham, Cheshire, Wethersfield, and Catden soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the depth to bedrock in Holyoke and Yalesville soils, the steep slopes, and rock outcrops. Stones and boulders need to be removed from the surface in some areas.

S1452. Cheshire-Yalesville-Wethersfield

Nearly level to steep, well drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 2 percent of the state. The landscape is predominantly upland hill, drumlin, and plain landforms in the Connecticut Valley with a north-south orientation. The soils formed in melt-out or lodgement till. Stones and boulders are common on the surface in many places.

Cheshire soils make up about 26 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Cheshire soils are gently sloping to steep and are on hills and plains.

Yalesville soils make up about 21 percent of this map unit. They are moderately deep to bedrock, well drained, loamy soils with moderately rapid permeability in the substratum. Yalesville soils are gently sloping to strongly sloping and are on bedrock-controlled hills and ridges.

Wethersfield soils make up about 11 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Wethersfield soils are nearly level to steep and are on hills and drumlins.

Soils of minor extent make up about 42 percent of this map unit. They are mainly Wilbraham, Manchester, Raypol, Bash, Ludlow, and Catden soils. Most areas of this map unit are in cropland or community development. Some small areas are in woodland. The soils of this map unit are suited to these uses.

The major limiting factors for community development are the slow or very slow permeability in Wethersfield soils and the depth to bedrock in Yalesville soils. Stones and boulders need to be removed from the surface in some areas.

S1453. Narragansett-Cheshire-Wapping

Nearly level to steep, moderately well drained to well drained, silty to loamy soils; on glacial till uplands

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly upland hill and plain landforms in the Connecticut Valley. The soils formed in melt-out till. Stones and boulders are common on the surface in many places.

Narragansett soils make up about 32 percent of this map unit. They are very deep, well drained, silty soils with moderate or moderately rapid permeability in the substratum. Narragansett soils are nearly level to moderately steep and are on hills and plains.

Cheshire soils make up about 29 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately permeability in the substratum. Cheshire soils are gently sloping to steep and are on hills and plains.

Wapping soils make up about 9 percent of this map unit. They are very deep, moderately well drained, silty soils with moderately rapid or rapid permeability in the substratum. Wapping soils are nearly level to gently sloping and are on hills and plains.

Soils of minor extent make up about 30 percent of this map unit. They are mainly Wilbraham, Haven, Broadbrook, Manchester, Rippowam, Watchaug, and Catden soils. Most areas of this map unit are in cropland or woodland. Some areas are in community development. The soils of this map unit are suited to these uses.

Stones and boulders need to be removed from the surface in some areas.

S1454. Wethersfield-Ludlow-Wilbraham

Nearly level to steep, poorly drained to well drained, loamy and silty soils; on glacial till uplands

The soils in this map unit make up about 5 percent of the state. The landscape is predominantly upland hill, drumlin, and drainageway landforms in the Connecticut Valley with a north-south orientation. The soils formed in lodgement till.

Wethersfield soils make up about 37 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Wethersfield soils are nearly level to steep and are on hills and drumlins.

Ludlow soils make up about 14 percent of this map unit. They are very deep, moderately well drained, loamy soils with slow or very slow permeability in the substratum. Ludlow soils are nearly level to strongly sloping and are on hills and drumlins.

Wilbraham soils make up about 11 percent of this map unit. They are very deep, poorly drained, silty soils with slow or very slow permeability in the substratum. Wilbraham soils are nearly level and are in depressions and drainageways.

Soils of minor extent make up about 38 percent of this map unit. They are mainly Branford, Cheshire, Holyoke, Yalesville, and Manchester soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in these soils and the seasonal high water table in Wilbraham soils.

S1455. Manchester-Branford-Rippowam

Nearly level to steep, poorly drained to excessively drained, sandy and gravelly to loamy soils; on glacial outwash plains, kames, terraces, and drainageways

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly outwash plain and drainageway landforms in the Connecticut Valley. The soils formed in outwash and silty alluvium.

Manchester soils make up about 31 percent of this map unit. They are very deep, excessively drained, sandy and gravelly soils with rapid or very permeability in the substratum. Manchester soils are nearly level to steep sloping and are on plains, kames, eskers, and terraces.

Branford soils make up about 31 percent of this map unit. They are very deep, well drained, loamy soils with rapid or very rapid permeability in the substratum. Branford soils are nearly level to strongly sloping and are on plains and terraces.

Rippowam soils make up about 6 percent of this map unit. They are very deep, poorly drained, silty soils with rapid or very rapid permeability in the substratum. Rippowam soils are nearly level and are in depressions on flood plains.

Soils of minor extent make up about 32 percent of this map unit. They are mainly Sudbury, Udorthents, Raypol, Penwood, Timakwa, Wethersfield, Cheshire, and Holyoke soils. Most areas of this map unit are in cropland, woodland or community development. The soils of this map unit are suited to these uses.

The major limiting factors for community development are the rapid or very rapid permeability in Manchester and Branford soils and the seasonal high water table in Rippowam soils.

S1456. Manchester-Penwood-Hartford

Nearly level to steep, somewhat excessively drained to excessively drained, sandy and gravelly soils; on outwash plains

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly outwash plain, terrace, and kame landforms in the Connecticut Valley. The soils formed in outwash.

Manchester soils make up about 37 percent of this map unit. They are very deep, excessively drained, sandy and gravelly soils with rapid or very rapid permeability in the substratum. Manchester soils are nearly level to steep and are on plains, kames, eskers, and terraces.

Penwood soils make up about 14 percent of this map unit. They are very deep, excessively drained, sandy soils with rapid or very rapid permeability in the substratum. Penwood soils are nearly level to gently sloping and are on plains and terraces.

Hartford soils make up about 12 percent of this map unit. They are very deep, somewhat excessively drained, sandy soils with rapid or very rapid permeability in the substratum. Hartford soils are nearly level to gently sloping and are on plains and terraces.

Soils of minor extent make up about 37 percent of this map unit. They are mainly Walpole, Haven, Ellington, Sudbury, Wethersfield, Rippowam, Timakwa, and Cheshire soils. Most areas of this map unit are in community development. Some areas are in cropland. The soils of this map unit are suited to these uses, although these soils can be droughty unless irrigation is provided.

The major limiting factor for community development is the rapid or very rapid permeability in Manchester and Penwood soils.

S1457. Wethersfield-Berlin-Belgrade

Nearly level to steep, moderately well drained and well drained, loamy and silty soils; on glacial till uplands

The soils in this map unit make less than 1 percent of the state. The landscape is predominantly drumlin and lake plain and terrace landforms in the Connecticut Valley. The soils formed in lodgement till and glaciolacustrine materials. Stones and boulders are common on the surface in many places.

Wethersfield soils make up about 25 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Wethersfield soils are nearly level to steep and are on hills and drumlins.

Berlin soils make up about 24 percent of this map unit. They are very deep, moderately well drained, silty soils with very slow permeability in the substratum. Berlin soils are nearly level to gently sloping and are on plains and terraces.

Belgrade soils make up about 20 percent of this map unit. They are very deep, moderately well drained, silty soils with moderate permeability in the substratum. Belgrade soils are nearly level to gently sloping and are on plains and terraces.

Soils of minor extent make up about 31 percent of this map unit. They are mainly Scitico, Shaker, Branford, Bash, Ludlow, Manchester, Maybid, and Raynham soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factor for community development is the slow or very slow permeability in the Wethersfield and Berlin soils.

S1458. Winooski-Hadley-Occum

Nearly level, moderately well drained and well drained, silty and loamy soils; on flood plains

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly flood plain landforms (fig. 3). The soils formed in alluvium.




Figure 3.—Relationship of soils, landscapes, and parent material along the Connecticut River.

Winooski soils make up about 31 percent of this map unit. They are very deep, moderately well drained, silty soils with moderate or moderately rapid permeability in the substratum. Winooski soils are nearly level and are on flood plains subject to frequent flooding.

Hadley soils make up about 26 percent of this map unit. They are very deep, well drained, silty soils with moderate to rapid permeability in the substratum. Hadley soils are nearly level and are on flood plains subject to occasional flooding.

Occum soils make up about 19 percent of this map unit. They are very deep, well drained, loamy soils with rapid or very rapid permeability in the substratum. Occum soils are nearly level and are on flood plains subject to occasional flooding.

Soils of minor extent make up about 24 percent of this map unit. They are mainly Limerick, Pootatuck, Lim, Saco, and Suncook soils. Most areas of this map unit are in cropland or woodland. The soils of this map unit are suited to these uses and are some of the most productive soils in Connecticut.

The major limiting factors for cropland are the occasional to frequent flooding hazards associated with these soils.

S3111. Macomber-Taconic-Lanesboro

Gently sloping to very steep, well drained to somewhat excessively drained, loamy soils; on glacial till uplands

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly bedrock-controlled upland landforms in Northwestern Connecticut (fig.4). The soils formed in melt-out till and lodgement till. Stones and boulders are common on the surface in many places.

Macomber soils make up about 32 percent of this map unit. They are moderately deep to bedrock, well drained, loamy soils with moderate permeability in the




Figure 4.—Typical pattern of soils in the Macomber-Taconic-Lanesboro general soil map unit.

substratum. Macomber soils are gently sloping to moderately steep and are on bedrock-controlled hills and ridges.

Taconic soils make up about 32 percent of this map unit. They are shallow to bedrock, somewhat excessively drained, loamy soils with moderate or moderately rapid permeability in the substratum. Taconic soils are gently sloping to very steep and are on bedrock-controlled hills and ridges.

Lanesboro soils make up about 11 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Lanesboro soils are gently sloping to steep and are on hills.

Soils of minor extent make up about 25 percent of this map unit. They are mainly Fullam, Rock outcrop, Dummerston, Brayton, Farmington, Fluvaquents, and Hoosic soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in Lanesboro soils and the depth to bedrock in Macomber and Taconic soils. Stones and boulders need to be removed from the surface in some areas.

S3114. Copake-Hero-Halsey

Nearly level to strongly sloping, very poorly drained to well drained, loamy soils; on upland valley outwash plains

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly upland valley landforms in Northwestern Connecticut. The soils formed in outwash.

Copake soils make up about 38 percent of this map unit. They are very deep, well drained, loamy soils with rapid or very rapid permeability in the substratum. Copake soils are nearly level to strongly sloping and are on kames, plains, and terraces.

Hero soils make up about 9 percent of this map unit. They are very deep, moderately well drained, loamy soils with rapid or very rapid permeability in the substratum. Hero soils are nearly level to gently sloping and are on plains and terraces.

Halsey soils make up about 8 percent of this map unit. They are very deep, very poorly drained, loamy soils with rapid or very rapid permeability in the substratum. Halsey soils are nearly level and are on in depressions and drainageways.

Soils of minor extent make up about 45 percent of this map unit. They are mainly Fredon, Natchaug, Deerfield, Groton, Farmington, Catden, Hinckley, Oakville, and Winooski soils. Most areas of this map unit are in woodland or cropland. The soils of this map unit are suited to these uses.

The major limiting factors for cropland are steep slopes in some areas and the seasonal high water table in Halsey soils.

S3115. Hinckley-Windsor-Merrimac

Nearly level to steep, somewhat excessively drained and excessively drained, loamy soils; on outwash plains

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly outwash plain, terrace, kame and esker landforms. The soils formed in outwash.

Hinckley soils make up about 26 percent of this map unit. They are very deep, excessively drained, soils with very rapid permeability in the substratum. Hinckley soils are nearly level to steep and are on kames, eskers, and outwash plains and terraces.

Windsor soils make up about 20 percent of this map unit. They are very deep, excessively drained, sandy soils with rapid or very rapid permeability in the substratum. Windsor soils are nearly level to strongly sloping and are on kames, terraces and plains.

Merrimac soils make up about 18 percent of this map unit. They are very deep, somewhat excessively drained, sandy soils with rapid or very rapid permeability in the substratum. Merrimac soils are nearly level to strongly sloping and are on outwash plains, terraces, and kames.

Soils of minor extent make up about 36 percent of this map unit. They are mainly Freetown, Deerfield, Urban land, Pits, Belgrade, Scarboro, Wareham, Saco, and Sudbury soils. Most areas of this map unit are in woodland, cropland, or community development. The soils of this map unit are suited to these uses and also are a source of sand and gravel.

The major limiting factors for community development are steep slopes in some areas and the rapid to very rapid permeability.

S3120. Wethersfield-Meckesville-Scarboro

Nearly level to steep, very poorly drained to well drained, loamy soils; on glacial till uplands

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly upland hill, drumlin, and drainageway landforms with a north-south orientation. The soils formed in lodgement till. Stones and boulders are common on the surface in many places.

Wethersfield soils make up about 47 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Wethersfield soils are nearly level to steep and are on hills and drumlins.

Meckesville soils make up about 15 percent of this map unit in the U.S.; however the Meckesville soils do not occur in Connecticut. They are very deep, well drained, loamy soils with very slow permeability in the substratum. Meckesville soils are nearly level to steep and are on hills and ridges.

Scarboro soils make up about 12 percent of this map unit. They are very deep, very poorly drained, loamy soils with rapid or very rapid permeability in the substratum. Scarboro soils are nearly level and are in depressions and drainageways.

Soils of minor extent make up about 26 percent of this map unit. They are mainly Windsor, Ridgebury, Wareham, Agawam, Holyoke, and Swansea. Most areas of this map unit are in woodland or community development. The soils of this map unit are suited to these uses.

The major limiting factors for community development are the slow or very slow permeability in Wethersfield and Meckesville soils and the seasonal high water table in Scarboro soils. Stones and boulders need to be removed from the surface in some areas.

S3121. Brookfield-Brimfield-Paxton

Gently sloping to steep, well drained and somewhat excessively drained, loamy soils; on glacial till uplands

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly upland hill, drumlin, and ridge landforms in Northeastern Connecticut. The soils formed in melt-out till and lodgement till. Stones and boulders are common on the surface in many places.

Brookfield soils make up about 30 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Brookfield soils are gently sloping to steep and are on bedrock-controlled hills and ridges.

Brimfield soils make up about 19 percent of this map unit. They are shallow to bedrock, somewhat excessively drained, loamy soils with moderate or moderately rapid permeability in the substratum. Brimfield soils are gently sloping to steep and are on bedrock-controlled hill tops and ridges.

Paxton soils make up about 12 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Paxton soils are gently sloping to steep and are on hills and drumlins.

Soils of minor extent make up about 39 percent of this map unit. They are mainly Rock outcrop, Freetown, Hinckley, Ridgebury, and Woodbridge soils. Most areas of this map unit are in woodland. Some areas are in cropland and community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in Paxton soils and depth to bedrock in Brimfield soils. Stones and boulders need to be removed from the surface in some areas.

S3122. Paxton-Woodbridge-Hollis

Nearly level to very steep, moderately well drained to somewhat excessively drained, loamy soils; on glacial till uplands

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly upland hill, drumlin, and ridge landforms with a north-south orientation (fig. 5). The soils formed in lodgement till. Stones and boulders are common on the surface in many places.

Paxton soils make up about 35 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Paxton soils are nearly level to gently sloping to steep and are on hills and drumlins.




Figure 5.—Relationship of soils, landscapes, and parent material in the Paxton-Woodbridge-Hollis general soil map unit.

Woodbridge soils make up about 22 percent of this map unit. They are very deep, moderately well drained, loamy soils with slow or very slow permeability in the substratum. Woodbridge soils are nearly level to strongly sloping and are on hills and drumlins.

Hollis soils make up about 11 percent of this map unit. They are shallow to bedrock, somewhat excessively drained, loamy soils with moderate or moderately rapid permeability in the substratum. Hollis soils are gently sloping to very steep and are on bedrock-controlled hills and ridges.

Soils of minor extent make up about 32 percent of this map unit. They are mainly Freetown, Whitman, Charlton, Ridgebury, and Rock outcrop. Most areas of this map unit are in woodland, cropland, or community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in Paxton and Woodbridge soils and the depth to bedrock in Hollis soils. Stones and boulders need to be removed from the surface in some areas.

S3136. Hollis-Chatfield-Rock outcrop

Nearly level to very steep, well drained to somewhat excessively drained, loamy soils; on bedrock-controlled glacial till uplands

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly bedrock-controlled upland hill and ridge landforms. The soils formed in melt-out till. Stones and boulders, as well as exposed areas of Rock outcrop are common on the surface in many places.

Hollis soils make up about 20 percent of this map unit. They are shallow to bedrock, somewhat excessively drained, loamy soils with moderate or moderately

rapid permeability in the substratum. Hollis soils are gently sloping to very steep and are on bedrock-controlled hills and drumlins.

Chatfield soils make up about 19 percent of this map unit. They are moderately deep to bedrock, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Chatfield soils are gently sloping to steep and are on bedrock-controlled hills and ridges.

Rock outcrop makes up about 17 percent of this map unit and occurs on bedrock-controlled landforms with slopes that are sometimes very steep.

Soils of minor extent make up about 44 percent of this map unit. They are mainly Charlton, Woodbridge, Paxton, Ridgebury, Whitman, Scarborough, Freetown, Urban land, and Udorthents soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the depth to bedrock in and steep slopes in some areas. Stones and boulders need to be removed from the surface in some areas.

S6623. Charlton-Canton-Sutton

Nearly level to steep, moderately well drained and well drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 2 percent of the state. The landscape is predominantly upland hill landforms. The soils formed in melt-out till. Stones and boulders are common on the surface in many places.

Charlton soils make up about 24 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Charlton soils are gently sloping to steep and are on hills.

Canton soils make up about 24 percent of this map unit. They are very deep, well drained, loamy over sandy and gravelly soils with rapid permeability in the substratum. Canton soils are gently sloping to steep and are on hills.

Sutton soils make up about 12 percent of this map unit. They are very deep, moderately well drained, loamy soils with moderately rapid permeability in the substratum. Sutton soils are nearly level to strongly sloping and are on hills.

Soils of minor extent make up about 40 percent of this map unit. They are mainly Leicester, Paxton, Catden, Chatfield, Hollis, and Woodbridge soils. Most areas of this map unit are in cropland or woodland. Some areas are in community development. The soils of this map unit are suited to these uses.

Stones and boulders need to be removed from the surface in some areas.

S6625. Woodbridge-Paxton-Ridgebury

Nearly level to steep, poorly drained to well drained, loamy soils; on glacial till uplands

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly upland hill and drumlin landforms with a north-south orientation. The soils formed in lodgement till. Stones and boulders are common on the surface in many places.

Woodbridge soils make up about 37 percent of this map unit. They are very deep, moderately well drained, loamy soils with slow or very slow permeability in the substratum. Woodbridge soils are nearly level to strongly sloping and are on hills and drumlins.

Paxton soils make up about 23 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Paxton soils are gently sloping to steep and are on hills and drumlins.

Ridgebury soils make up about 20 percent of this map unit. They are very deep, poorly drained, loamy soils with slow or very slow permeability in the substratum.

Ridgebury soils are nearly level to gently sloping and are in depressions and drainageways.

Soils of minor extent make up about 20 percent of this map unit. They are mainly Canton, Catden, Charlton, Chatfield, and Hollis soils. Most areas of this map unit are in cropland, woodland, or community development. The soils of this map unit are suited to these uses.

The major limiting factors for community development are the slow or very slow permeability and the seasonal high water table in Woodbridge and Ridgebury soils. Stones and boulders need to be removed from the surface in some areas.

S6627. Carlisle-Adrian-Scarboro

Nearly level, very poorly drained, woody organic soils; on drainageways

The soils in this map unit make up less than 1 percent of the state. The landscape is predominantly depressions and drainageway landforms. The soils formed in organic materials or outwash.

Carlisle soils make up about 65 percent of this map unit in the U.S.; however Carlisle soils are not mapped in Connecticut. Catden soils are mapped in Connecticut and are similar to Carlisle soils. They are very deep, very poorly drained, organic soils with variable mineral substrata below 51 inches. Catden soils are nearly level and are in depressions.

Adrian soils make up about 13 percent of this map unit in the U.S.; however Adrian soils are not mapped in Connecticut. Timakwa soils are mapped in Connecticut and are similar to Adrian soils. They are very deep, very poorly drained, organic soils with sandy mineral substrata between 16 and 51 inches below the surface. Timakwa soils are nearly level and are in depressions.

Scarboro soils make up about 11 percent of this map unit. They are very deep, very poorly drained, loamy soils with rapid or very rapid permeability in the substratum. Scarboro soils are nearly level and are in depressions and drainageways.

Soils of minor extent make up about 11 percent of this map unit. They are mainly Merrimac, Woodbridge, and Hinckley soils.

Most areas of this map unit are in woodland and wildlife habitat, which these soils are well suited.

S9569. Bice-Shelburne-Ashfield

Nearly level to steep, moderately well drained and well drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 3 percent of the state. The landscape is predominantly upland hill and drumlin landforms. The soils formed in melt-out till and lodgement till. Stones and boulders are common on the surface in many places.

Bice soils make up about 30 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Bice soils are gently sloping to steep and are on hills.

Shelburne soils make up about 25 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Shelburne soils are gently to steep and are on hills and drumlins.

Ashfield soils make up about 20 percent of this map unit. They are very deep, moderately well drained, loamy soils with slow or very slow permeability in the substratum. Ashfield soils are nearly level to strongly sloping and are on hills and drumlins.

Soils of minor extent make up about 25 percent of this map unit. They are mainly Loonmeadow, Millsite, Schroon, and Westminster soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in Shelburne and Ashfield soils and the seasonal high water table in Ashfield soils. Stones and boulders need to be removed from the surface in some areas.

S9570. Bice-Millsite-Westminster

Gently sloping to very steep, well drained and somewhat excessively drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly upland hill and ridge landforms (fig. 6). The soils formed in melt-out till. Rock outcrop, stones and boulders are common on the surface in many places.

Bice soils make up about 30 percent of this map unit. They are very deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Bice soils are gently sloping to steep and are on hills.

Millsite soils make up about 25 percent of this map unit. They are moderately deep, well drained, loamy soils with moderate or moderately rapid permeability in the substratum. Millsite soils are gently sloping to steep and are on bedrock-controlled hills and ridges.

Westminster soils make up about 15 percent of this map unit. They are shallow to bedrock, somewhat excessively drained, loamy soils with moderately rapid permeability in the substratum. Westminster soils are gently sloping to very steep and are on bedrock-controlled hills and ridges.

Soils of minor extent make up about 30 percent of this map unit. They are mainly Schroon, Loonmeadow, and rock outcrop. Most areas of this map unit are in




Figure 6.—Typical pattern of soils in the Bice-Millsite-Westminster general soil map unit and the Ashfield-Shelburne-Loonmeadow general soil map unit.

woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factor for community development is the depth to bedrock in Millsite and Westminster soils. Stones and boulders need to be removed from the surface in some areas.

S9571. Ashfield-Shelburne-Loonmeadow

Nearly level to steep, very poorly drained to well drained, loamy soils; on glacial till uplands

The soils in this map unit make up about 1 percent of the state. The landscape is predominantly upland hill and drumlin landforms with a north-south orientation. The soils formed in lodgement till. Stones and boulders are common on the surface in many places.

Ashfield soils make up about 20 percent of this map unit. They are very deep, moderately well drained, loamy soils with slow or very slow permeability in the substratum. Ashfield soils are nearly level to strongly sloping and are on hills and drumlins.

Shelburne soils make up about 20 percent of this map unit. They are very deep, well drained, loamy soils with slow or very slow permeability in the substratum. Shelburne soils are gently sloping to steep and are on hills and drumlins.

Loonmeadow soils make up about 15 percent of this map unit. They are very deep, very poorly drained, loamy soils with moderately slow to rapid permeability in the substratum. Loonmeadow soils are nearly level and are in depressions and drainageways.

Soils of minor extent make up about 45 percent of this map unit. They are mainly Brayton, Bice, Wonsqueak, and Fullam soils. Most areas of this map unit are in woodland. Some areas are in community development. The soils of this map unit are suited to forestry and wildlife habitat.

The major limiting factors for community development are the slow or very slow permeability in Ashfield and Shelburne soils and the seasonal high water table in Ashfield and Loonmeadow soils. Stones and boulders need to be removed from the surface in some areas.

Detailed Soil Map Units

The map units delineated on the detailed soil maps in this survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this section, along with the maps, can be used to determine the suitability and potential of a unit for specific uses. They also can be used to plan the management needed for those uses.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. The contrasting components are mentioned in the map unit descriptions. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives the principal hazards and limitations to be considered in planning for specific uses.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown

on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Windsor loamy sand, 0 to 3 percent slopes is a phase of the Windsor series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Hollis-Chatfield-Rock outcrop complex, 15 to 45 percent slopes is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Mudgepond and Alden soils, extremely stony is an undifferentiated group in this survey area.

This survey includes *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Beaches, Dumps, Pits, Rock outcrop, and Urban land are examples. Miscellaneous areas have few, if any, significant soil properties and typically little information is mentioned in the map unit descriptions.

Table 4 gives the acreage and proportionate extent of each map unit. Other tables give properties of the soils and the limitations, capabilities, and potentials for many uses. The Glossary defines many of the terms used in describing the soils or miscellaneous areas.

2—Ridgebury fine sandy loam

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Size of map unit: Areas commonly range from 3 to 50 acres

Map Unit Composition

Ridgebury and similar soils: 80 percent

Minor components: 20 percent

Major Component

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; fine sandy loam

Bg1—5 to 14 inches; fine sandy loam

Bg2—14 to 21 inches; fine sandy loam

Cd—21 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 0 to 6 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of moderately well drained Sutton and Woodbridge soils on slightly higher areas of the landscape. Sutton soils lack the dense substratum that Woodbridge soils have. Poorly drained Leicester soils are included in areas that lack a dense substratum. Very poorly drained Whitman soils are in depressions and drainageways. Also included are very poorly drained Timakwa and Natchaug soils in marshy areas where muck is between 16 and 51 inches thick over mineral substratum. A few areas include a silt loam surface layer and subsoil, stones on the surface, or slopes steeper than 5 percent. Minor components make up about 20 percent of this map unit.

Use and Management

This soil is mostly in woodland. Some areas are in pasture, cropland, or community development.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil. The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

3—Ridgebury, Leicester, and Whitman soils, extremely stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Surface cover: 3 to 14 percent stones

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Ridgebury and similar soils: 40 percent

Leicester and similar soils: 35 percent

Whitman and similar soils: 15 percent

Minor components: 10 percent

Major Component

Ridgebury and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; fine sandy loam

Bg1—5 to 14 inches; fine sandy loam

Bg2—14 to 21 inches; fine sandy loam

Cd—21 to 60 inches; sandy loam

Leicester and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 7 inches; fine sandy loam

Bg1—7 to 10 inches; fine sandy loam

Bg2—10 to 18 inches; fine sandy loam

BC—18 to 24 inches; fine sandy loam

C1—24 to 43 inches; gravelly fine sandy loam

C2—43 to 65 inches; gravelly fine sandy loam

Whitman and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi —0 to 1 inch; slightly decomposed plant material

A—1 to 9 inches; fine sandy loam

Bg—9 to 16 inches; fine sandy loam

Cdg1—16 to 22 inches; fine sandy loam

Cdg2—22 to 60 inches; fine sandy loam

Major Component Properties and Qualities**Ridgebury and similar soils**

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 0 to 6 inches

Flooding: none

Leicester and similar soils

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 18 inches

Flooding: none

Whitman and similar soils

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: very low

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 12 to 20 inches to densic material

Ponding depth: 0 to 12 inches above surface

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Ridgebury and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Leicester and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Whitman and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of moderately well drained Sutton and Woodbridge soils that are slightly higher on the landscape. Sutton soils lack the dense substratum that Woodbridge soils have. Also included are a few non-stony surface soils, small areas of soils subject to flooding, small areas with steeper slopes, and areas with silt loam surface and subsoil textures. Minor components make up about 10 percent of the map unit.

Use and Management

This soil is mostly in woodland. Some areas are in pasture.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Ponding is a limitation in areas of Whitman soils. Slow percolation is also a limitation for septic tank absorption fields in areas of Ridgebury and Whitman soils. Large stones are a limitation for lawns and landscaping. A more suitable site should be selected for these uses in a drier inclusion or nearby soil. The seasonal high water table and frost action are the main limitations for local roads and streets. Ponding is also a limitation for local roads and streets in Whitman soils. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

4—Leicester fine sandy loam

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Leicester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; fine sandy loam

Bg1—7 to 10 inches; fine sandy loam

Bg2—10 to 18 inches; fine sandy loam

BC—18 to 24 inches; fine sandy loam

C1—24 to 43 inches; gravelly fine sandy loam

C2—43 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 18 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are some areas of moderately well drained Sutton and Woodbridge soils on slightly higher areas of the landscape. Sutton soils lack the dense substratum that Woodbridge soils have. Also included are poorly drained Ridgebury soils and very poorly drained Whitman soils that have a dense substratum and occur along drainageways and in depressions. Soils with stones on the surface are included in a few small areas. Minor components make up about 20 percent of this map unit.

Use and Management

This soil is mostly in woodland. Some areas are in pasture.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil. The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

5—Wilbraham silt loam

Map Unit Setting

Slope: nearly level

Landscape: drainageways on uplands, depressions on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Wilbraham and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; silt loam

Bw1—4 to 8 inches; silt loam

Bw2—8 to 20 inches; silt loam

Cd—20 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 36 inches to densic material

Depth to seasonal water table: 0 to 18 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of well drained Cheshire and Wethersfield soils that are higher on the landscape. Cheshire soils lack the very firm, dense substratum that Wethersfield soils have. Also included are moderately well drained Watchaug and Ludlow soils in slightly higher areas. Watchaug soils lack the dense substratum that Ludlow soils have. Very poorly drained Menlo soils are included in depressions and drainageways. Also included are small areas with slopes up to 8 percent, areas that lack a dense substratum, soils with a stony surface, and soils with a loam or fine sandy loam surface layer. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or brushland reverting to woodland.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

6—Wilbraham and Menlo soils, extremely stony

Map Unit Setting

Slope: nearly level

Landscape: drainageways on uplands, depressions on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres

Map Unit Composition

Wilbraham and similar soils: 60 percent

Menlo and similar soils: 25 percent

Minor components: 15 percent

Major Components

Wilbraham and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; silt loam
 Bw1—4 to 8 inches; silt loam
 Bw2—8 to 20 inches; silt loam
 Cd—20 to 65 inches; gravelly loam

Menlo and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 5 inches; highly decomposed plant material
 A—5 to 16 inches; mucky silt loam
 Bg1—16 to 22 inches; flaggy very fine sandy loam
 Bg2—22 to 27 inches; flaggy fine sandy loam
 Cd1—27 to 40 inches; fine sandy loam
 Cd2—40 to 60 inches; fine sandy loam

Major Component Properties and Qualities

Wilbraham and similar soils

Depth to bedrock: very deep
Drainage class: poorly drained
Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale
Permeability: very slow to moderate
Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 36 inches to densic material
Depth to seasonal water table: 0 to 18 inches
Flooding: none

Menlo and similar soils

Depth to bedrock: very deep
Drainage class: very poorly drained
Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale
Permeability: very slow to moderate
Available water capacity: moderate
Reaction: very strongly acid to moderately alkaline
Depth to restrictive feature: 20 to 36 inches to densic material
Ponding depth: 0 to 12 inches above surface
Depth to seasonal water table: 0 to 12 inches
Flooding: none

Interpretative Groups

Wilbraham and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Menlo and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of well drained Cheshire and Wethersfield soils that are higher on the landscape. Cheshire soils lack the very firm, dense substratum that Wethersfield soils have. Also included are moderately well drained Watchaug and Ludlow soils on slightly higher areas of the landscape. Watchaug soils lack the dense substratum that Ludlow soils have. Also included are

small areas with slopes up to 8 percent, areas that lack a dense substratum, and soils with a loam or fine sandy loam surface layer. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in brushland reverting to woodland.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a limitation for septic tank absorption fields. Ponding is a limitation in areas of Menlo soils for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Excess humus is also a limitation for lawns and landscaping. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

7—Mudgepond silt loam

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: drainageways on uplands, depressions on uplands

Size of map unit: Areas range from 3 to 50 acres

Map Unit Composition

Mudgepond and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 11 inches; silt loam

Bg—11 to 16 inches; loam

Bw1—16 to 26 inches; fine sandy loam

Bw2—26 to 35 inches; gravelly fine sandy loam

C —35 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderate to moderately rapid

Available water capacity: high

Reaction: neutral to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are moderately well drained Georgia and Amenia soils on slightly higher areas of the landscape. Georgia soils have carbonates between 40 and 80 inches below the surface. Amenia soils have carbonates above

40 inches. Also included are very poorly drained Alden soils in depressions. Minor components make up about 15 percent of this map unit.

Use and Management

This soil is mostly in woodland. Other areas are in pasture or cropland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised filled materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

8—Mudgepond and Alden soils, extremely stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas range from 3 to 50 acres

Map Unit Composition

Mudgepond and similar soils: 45 percent

Alden and similar soils: 35 percent

Minor components: 20 percent

Major Components

Mudgepond and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 11 inches; silt loam

Bg—11 to 16 inches; loam

Bw1—16 to 26 inches; fine sandy loam

Bw2—26 to 35 inches; gravelly fine sandy loam

C—35 to 65 inches; gravelly fine sandy loam

Alden and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A1—0 to 4 inches; mucky silt loam

A2—4 to 13 inches; silt loam

Bg1—13 to 23 inches; silt loam

Bg2—23 to 29 inches; silt loam

Cg1—29 to 43 inches; gravelly loam

Cg2—43 to 60 inches; loam

Major Component Properties and Qualities

Mudgepond and similar soils

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: neutral to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 0 to 12 inches
Flooding: none

Alden and similar soils

Depth to bedrock: very deep
Drainage class: very poorly drained
Parent material: fine-loamy till derived from limestone and dolomite and/or schist
Permeability: moderately slow or moderate
Available water capacity: high
Reaction: strongly acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Ponding depth: 0 to 6 inches above surface
Depth to seasonal water table: 0 to 12 inches
Flooding: none

Interpretative Groups

Mudgepond and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Alden and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Minor Components

Included with this soil in mapping are moderately well drained Georgia and Amenia soils. Georgia and Amenia soils are on slightly higher areas of the landscape. Also included are small areas of steeper slopes. Minor components make up about 15 percent of this map unit.

Use and Management

This soil is mostly in woodland. Some areas are in pasture.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Ponding is a limitation in areas of Alden soils. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

The seasonal high water table and frost action are the main limitations for local roads and streets. Ponding is also a limitation in areas of Alden soils. Construction on raised filled materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

9—Scitico, Shaker, and Maybid soils

Map Unit Setting

Slope: nearly level
Landscape: depressions on lake plains, drainageways on lake plains, terraces
Size of map unit: Areas range from 3 to 200 acres

Map Unit Composition

Scitico and similar soils: 40 percent

Shaker and similar soils: 30 percent

Maybid and similar soils: 15 percent

Minor components: 15 percent

Major Components

Scitico and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Eg—8 to 11 inches; silt loam

Bg1—11 to 18 inches; silty clay loam

Bg2—18 to 30 inches; silty clay loam

Bg3—30 to 38 inches; silty clay

Cg1—38 to 52 inches; silty clay loam

Cg2—52 to 65 inches; silty clay

Shaker and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 2 inches; moderately decomposed plant material

Ap—2 to 6 inches; fine sandy loam

Bg—6 to 20 inches; sandy loam

Bw—20 to 30 inches; sandy loam

2C—30 to 65 inches; silty clay

Maybid and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 9 inches; silt loam

Bg1—9 to 18 inches; silty clay loam

Bg2—18 to 26 inches; silty clay loam

Cg1—26 to 36 inches; silty clay loam

Cg2—36 to 60 inches; silty clay loam

Major Component Properties and Qualities

Scitico and similar soils

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: clayey glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Shaker and similar soils

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy eolian deposits over clayey glaciolacustrine deposits

Permeability: very slow to moderately rapid

Available water capacity: very high

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 18 inches

Flooding: none

Maybid and similar soils

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: clayey glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high

Reaction: strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 6 inches above surface

Depth to seasonal water table: 0 to 6 inches

Flooding: none

Interpretative Groups

Scitico and similar soils

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Shaker and similar soils

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Maybid and similar soils

Land capability classification (non-irrigated): 6w

Hydrologic group: D

Minor Components

Included with this unit in mapping are moderately well drained Elmridge and Brancroft soils. Elmridge soils are loamy over clayey and Brancroft soils are silty and clayey. Elmridge and Brancroft soils are on higher areas of the landscape. Also included are areas of sand and gravel at 2 to 4 foot depths and soils that are redder in color. Minor components make up 15 percent of this unit.

Use and Management

Most areas are in woodland. Some areas are cleared and in hayland, pasture or cultivated cropland. A few areas are in sod farming or community development.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Ponding is also a limitation in areas of Maybid soils. Slow percolation is also a limitation for septic tank adsorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

Seasonal high water table and frost action are the main limitations for local roads and streets. Ponding and low strength are also limitations in areas of Maybid soils. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

10—Raynham silt loam

Map Unit Setting

Slope: nearly level

Landscape: drainageways on lake plains, depressions on lake plains

Size of map unit: Areas range from 3 to 50 acres

Map Unit Composition

Raynham and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 10 inches; silt loam

Bg1—10 to 16 inches; silt loam

Bg2—16 to 26 inches; silt loam

Bw—26 to 34 inches; very fine sandy loam

Cg—34 to 47 inches; silt loam

C—47 to 60 inches; silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-silty glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are moderately well drained Belgrade soils. Belgrade soils are on higher areas of the landscape. Also included are Scitico soils and very poorly drained Maybid soils in the deeper depressions and along drainageways. In New Haven County, moderately well drained Ellington soils are included on higher areas and are underlain by sandy materials. Minor components make up 20 percent of this unit.

Use and Management

Most areas are in woodland. Some areas are drained and in cultivated cropland or pasture. Other areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Slow percolation is also a limitation for septic tank adsorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

Seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

12—Raypol silt loam

Map Unit Setting

Slope: nearly level

Landscape: depressions on outwash plains, drainageways on outwash plains

Size of map unit: Areas commonly range from 3 to 50 acres

Map Unit Composition

Raypol and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bg1—8 to 12 inches; very fine sandy loam

Bg2—12 to 20 inches; silt loam

Bw1—20 to 26 inches; silt loam

Bw2—26 to 29 inches; very fine sandy loam

2C1—29 to 52 inches; stratified very gravelly coarse sand to loamy fine sand

2C2—52 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are well drained Haven and Enfield soils, moderately well drained Ninigret and Tisbury soils, poorly drained Walpole soils, and very poorly drained Scarboro soils. Haven and Enfield soils are on higher areas of the landscape and Ninigret and Tisbury soils are on slightly higher areas. Walpole soils are sandy throughout. Scarboro soils are in depressions. Also included are small areas in Fairfield County with loamy material deeper than 40 inches. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Cleared areas are in cultivated cropland or pasture. Some areas are drained. Other areas are in residential development

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Poor filtering is also a limitation for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Frost action and a seasonal high water table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

13—Walpole sandy loam

Map Unit Setting

Slope: nearly level

Landscape: drainageways on terraces, depressions on terraces, drainageways on outwash plains, depressions on outwash plains

Size of map unit: Areas commonly range from 3 to 100 acres

Map Unit Composition

Walpole and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 7 inches; sandy loam

Bg—7 to 21 inches; sandy loam

Bw—21 to 25 inches; gravelly sandy loam

C1—25 to 41 inches; stratified very gravelly coarse sand to loamy fine sand

C2—41 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils, somewhat excessively drained Merrimac soils, and moderately well drained and somewhat poorly drained Sudbury Soils. Hinckley soils and Merrimac soils are on higher areas and Sudbury soils are on slightly higher areas of the landscape. Also included are moderately well drained Ninigret soils on slightly higher areas and the very poorly drained Scarboro soils in the depressions. Raypol soils are in areas that have a loamy surface and subsoil and Raynham soils which are silty throughout. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland. Cleared areas are in cultivated cropland or pasture. Some areas are drained. Other areas are in residential development

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Poor filtering is also a

limitation for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Frost action and seasonal high water table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

14—Fredon silt loam

Map Unit Setting

Slope: nearly level

Landscape: drainageways on outwash plains, terraces on outwash plains, depressions on outwash plains

Size of map unit: Areas commonly range from 3 to 50 acres

Map Unit Composition

Fredon and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bg—8 to 17 inches; fine sandy loam

Bw—17 to 24 inches; fine sandy loam

2Cg1—24 to 29 inches; stratified gravelly sand to loamy fine sand

2C—29 to 48 inches; stratified gravelly sand to loamy fine sand

2Cg2—48 to 60 inches; stratified gravelly sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic Group: D

Minor Components

Included with this soil in mapping are areas of well drained Copake and Groton soils that are higher on the landscape. Also included are moderately well drained Hero soils on slightly higher areas of the landscape and very poorly drained Halsey soils in depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Cleared areas are in pasture or cropland. Some cleared areas are drained. Other areas are in residential development.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Poor filtering is also a limitation for septic tank adsorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Frost action and seasonal high water table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

15—Scarboro muck

Map Unit Setting

Slope: nearly level

Landscape: terraces on outwash plains, drainageways on outwash plains, depressions on outwash plains

Size of map unit: Areas commonly range from 3 to 100 acres. Slopes range from 0 to 2 percent.

Map Unit Composition

Scarboro and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 12 inches; muck

A—12 to 17 inches; loamy sand

Cg1—17 to 31 inches; stratified sand to loamy fine sand

Cg2—31 to 72 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 6 inches above surface

Depth to seasonal water table: 0 to 6 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 5w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of excessively drained Windsor soils on the highest areas of the landscape. Also included are moderately well drained

Sudbury soils in slightly higher areas of the landscape. Poorly drained Walpole and Raypol soils are in slight depressions. Timakwa and Natchaug soils are included in areas of muck that is 16 to 51 inches thick over mineral soil and Catden and Freetown soils are in areas with more than 51 inches of muck. Soils with a silt loam surface are included in New London County and soils with a sandy loam surface are included in New Haven County. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are brushland reverting to woodland.

Ponding is the main limitation for dwellings with basements and lawns and landscaping. Ponding and poor filtering are the main limitations for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Ponding and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

16—Halsey silt loam

Map Unit Setting

Slope: nearly level

Landscape: terraces on outwash plains, depressions on outwash plains, drainageways on outwash plains

Size of map unit: Areas commonly range from 3 to 40 acres

Map Unit Composition

Halsey and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 8 inches; silt loam

Bg1—8 to 16 inches; silt loam

Bg2—16 to 28 inches; fine sandy loam

2Cg1—28 to 38 inches; loamy sand

2Cg2—38 to 60 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 6 inches above surface

Depth to seasonal water table: 0 to 6 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 5w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of poorly drained Fredon soils in shallow depressions and drainageways. Also included are areas of well drained Copake soils and excessively drained Groton soils that are higher on the landscape. Moderately well drained Hero soils are included in slightly higher areas of the landscape. Very poorly drained Timakwa soils are included in marshy areas where the muck is between 16 and 51 inches thick over the sandy mineral substratum. A few soils with silt loam subsoil are included. Minor components make up 20 percent of this map unit

Use and Management

Most areas are in woodland. Some areas are in pasture or water tolerant shrubs and herbaceous plants.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Poor filtering is also a limitation for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Frost action and seasonal high water table are the main limitations for local roads and streets. Constructing on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

17—Timakwa and Natchaug soils

Map Unit Setting

Slope: nearly level

Landscape: depressions

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Timakwa and similar soils: 45 percent

Natchaug and similar soils: 40 percent

Minor components: 15 percent

Major Components

Timakwa and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa1—0 to 10 inches; muck

Oa2—10 to 21 inches; muck

Oa3—21 to 24 inches; muck

Oa4—24 to 37 inches; muck

2Cg1—37 to 47 inches; very gravelly loamy coarse sand

2Cg2—47 to 60 inches; gravelly loamy very fine sand

Natchaug and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi1—0 to 2 inches; peat
 Oi2—2 to 4 inches; peat
 Oa1—4 to 6 inches; muck
 Oa2—6 to 11 inches; muck
 Oa3—11 to 18 inches; muck
 Oa4—18 to 24 inches; muck
 2Cg1—24 to 33 inches; fine sandy loam
 2Cg2—33 to 36 inches; fine sandy loam
 2Cg3—36 to 80 inches; loam

Major Component Properties and Qualities

Timakwa and similar soils

Depth to bedrock: very deep
Drainage class: very poorly drained
Parent material: woody organic material over sandy and gravelly glaciofluvial deposits
Permeability: moderate to very rapid
Available water capacity: very high
Reaction: ultra acid to neutral
Depth to restrictive feature: greater than 72 inches
Ponding depth: 0 to 12 inches above surface
Depth to seasonal water table: 0 to 12 inches
Flooding: rare

Natchaug and similar soils

Depth to bedrock: very deep
Drainage class: very poorly drained
Parent material: woody organic material over loamy alluvium and/or loamy glaciofluvial deposits and/or loamy till
Permeability: moderately slow to very rapid
Available water capacity: very high
Reaction: extremely acid to neutral
Depth to restrictive feature: greater than 72 inches
Ponding depth: 0 to 12 inches above surface
Depth to seasonal water table: 0 to 12 inches
Flooding: rare

Interpretative Groups

Timakwa and similar soils

Land capability classification (non-irrigated): 5w
Hydrologic group: D

Natchaug and similar soils

Land capability classification (non-irrigated): 5w
Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of very poorly drained Catden soils where the muck is more than 51 inches thick over mineral substratum. Also included are areas of very poorly drained Whitman, Menlo, Scarborough, Maybid, and Saco soils. Whitman and Menlo soils formed in loamy glacial till. Scarborough soils are sandy and Maybid soils are silty and clayey. Saco soils are on flood plains and are silty. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas of this soil are in woodland or wildlife habitat.

Ponding and subsidence are the main limitations for dwellings with basements, septic tank absorption fields, local roads, and streets. Excess humus is also a limitation for lawns and landscaping. Slow percolation is also a limitation for septic tank absorption fields. Frost action is also a limitation for local roads and streets. A more suitable site for all these uses should be selected on a drier soil.

18—Catden and Freetown soils

Map Unit Setting

Slope: nearly level

Landscape: depressions

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Catden and similar soils: 40 percent

Freetown and similar soils: 40 percent

Minor components: 20 percent

Major Components

Catden and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa1—0 to 2 inches; muck

Oa2—2 to 18 inches; muck

Oa3—18 to 47 inches; muck

Oa4—47 to 49 inches; muck

Oa5—49 to 61 inches; muck

Freetown and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi1—0 to 4 inches; peat

Oi2—4 to 10 inches; peat

Oa1—10 to 22 inches; muck

Oa2—22 to 35 inches; muck

Oa3—35 to 41 inches; muck

Oa4—41 to 55 inches; muck

Oa5—55 to 71 inches; muck

Oa6—71 to 91 inches; muck

Major Component Properties and Qualities

Catden and similar soils

Depth to bedrock: very deep

Drainage class: very poorly drained (fig. 7)

Parent material: not specified

Permeability: moderate or moderately rapid

Available water capacity: very high

Reaction: extremely acid to neutral

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 12 inches above surface

Depth to seasonal water table: 0 to 12 inches

Flooding: rare



Figure 7.—Wetland vegetation is common in areas of very poorly drained Catden and Freetown soils.

Freetown and similar soils

Depth to bedrock: very deep
Drainage class: very poorly drained
Parent material: woody organic material
Permeability: moderate or moderately rapid
Available water capacity: very high
Reaction: ultra acid to extremely acid
Depth to restrictive feature: greater than 72 inches
Ponding depth: 0 to 12 inches above surface
Depth to seasonal water table: 0 inches
Flooding: rare

Interpretative Groups

Catden and similar soils

Land capability classification (non-irrigated): 5w
Hydrologic group: D

Freetown and similar soils

Land capability classification (non-irrigated): 5w
Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of very poorly drained Timakwa and Natchaug soils. Timakwa soils have muck between 16 to 51 inches thick over sandy substratum; Natchaug soils have muck between 16 to 51 inches thick over loamy substratum. Also included are very poorly drained Whitman, Menlo, Scarboro, Maybid, and Saco soils. Whitman and Menlo soils formed in loamy glacial till.

Scarboro soils are sandy and Maybid soils are silty and clayey. Saco soils are on flood plains and are silty. Minor Components make up about 20 percent of this map unit.

Use and Management

Most areas of this soil are in woodland or wildlife habitat.

Ponding and subsidence are the main limitations for dwellings with basements, septic tank absorption fields, local roads and streets. Low strength is also a limitation for dwellings with basements. Excess humus is a limitation for lawns and landscaping. Slow percolation is also a limitation for septic tank absorption fields. Frost action is also a limitation for local roads and streets. A more suitable site for all these uses should be selected on a drier soil.

20A—Ellington silt loam, 0 to 5 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 25 acres.

Map Unit Composition

Ellington and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; silt loam

Bw2—18 to 26 inches; very fine sandy loam

2C—26 to 65 inches; stratified loamy fine sand to very gravelly coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Branford soils and poorly drained Raypol and Raynham soils. Branford soils are on higher areas of the landscape. Raypol and Raynham soils are in shallow depressions and along

drainageways. Raynham soils are silty throughout; Raypol soils are silty over sand and gravel. A few areas in Middlesex and New Haven counties include soils with a fine sandy loam surface layer. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland. Some areas are in community development, woodland, or pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

21A—Ninigret and Tisbury soils, 0 to 5 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ninigret and similar soils: 60 percent

Tisbury and similar soils: 25 percent

Minor components: 15 percent

Major Components

Ninigret and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

2C—26 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Tisbury and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; silt loam

Bw2—18 to 26 inches; silt loam

2C—26 to 60 inches; stratified very gravelly sand to loamy sand

Major Component Properties and Qualities

Ninigret and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Tisbury and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-silty eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Ninigret and similar soils

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Tisbury and similar soils

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of somewhat excessively drained Merrimac soils and well drained Agawam, Enfield, and Haven soils that are on higher areas of the landscape. Agawam soils are loamy over sand and gravel; Enfield and Haven soils are silty over sand and gravel. Also included are moderately well drained Sudbury soils that are sandy and gravelly throughout. Small areas poorly drained Walpole soils and Raypol soils are included in shallow depressions and drainageways. A few areas include soils with a red color in the central lowlands of the Connecticut River Valley. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are cleared and in cultivated crops, hay, pasture, or brushland. Some areas are in woodland or community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce the wetness.

Poor filtering and the seasonal high water table are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal. There is also a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

22A—Hero gravelly loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Hero and similar soils: 85 percent

Minor components: 15 percent

Major Components

Hero and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; gravelly loam

Bw1—9 to 18 inches; gravelly silt loam

Bw2—18 to 24 inches; gravelly silt loam

Bw3—24 to 27 inches; gravelly sandy loam

2C—27 to 60 inches; stratified extremely gravelly coarse sand to gravelly loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Groton soils and well drained Copake soils that are higher on the landscape. Also included are small areas of poorly drained Fredon soils and very poorly drained Halsey soils in depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Cleared areas are in pasture or cropland. Some areas are drained. Other areas are in residential development.

The seasonal high water table is the main limitation for dwelling with basements, lawns and landscaping, and septic tank absorption fields. Small stones are also a limitation for lawns and landscaping. Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly

permeable substratum does not adequately filter effluent. A more suitable site for these uses should be considered in a drier inclusion or nearby soil.

Frost action and the seasonal high water table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

22B—Hero gravelly loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Hero and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; gravelly loam

Bw1—9 to 18 inches; gravelly silt loam

Bw2—18 to 24 inches; gravelly silt loam

Bw3—24 to 27 inches; gravelly sandy loam

2C—27 to 60 inches; stratified extremely gravelly coarse sand to gravelly loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Groton soils and well drained Copake soils that are higher on the landscape. Also included are small areas of poorly drained Fredon soils and very poorly drained Halsey soils in depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Cleared areas are in pasture or cropland. Some areas are drained. Other areas are in residential development.

The seasonal high water table is the main limitation for dwelling with basements, lawns and landscaping, and septic tank absorption fields. Small stones are also a limitation for lawns and landscaping. Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site for these uses should be considered in a drier inclusion or nearby soil.

Frost action and the seasonal high water table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

23A—Sudbury sandy loam, 0 to 5 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas range from 3 to 40 acres.

Map Unit Composition

Sudbury and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 5 inches; sandy loam

Bw1—5 to 17 inches; gravelly sandy loam

Bw2—17 to 25 inches; sandy loam

2C—25 to 60 inches; stratified gravel to sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of somewhat excessively drained Merrimac soils and well drained Agawam soils that are higher on the landscape. Also included are moderately well drained Ninigret and Tisbury soils in areas with a finer surface texture. Small areas of poorly drained Walpole soils are included in drainageways and shallow depressions. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, hay, or pasture. Some areas are in woodland or community development.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Modifying a conventional septic system by extending the length of distribution lines and adding fill usually will allow on site sewage disposal. Specially designed septic systems are necessary in some areas of Sudbury soils.

The seasonal high water table and frost action are the main limitations for local roads and streets. Constructing roads on raised fill materials and installing a drainage system will reduce the wetness limitation. Providing a coarse grained subgrade to frost depth will reduce the limitation.

24A—Deerfield loamy fine sand, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Deerfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; loamy fine sand

Bw1—8 to 16 inches; loamy sand

Bw2—16 to 28 inches; loamy sand

C1—28 to 34 inches; fine sand

C2—34 to 60 inches; fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: sandy glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Windsor, Penwood, and Hinckley soils that are higher on the landscape. Windsor soils are sandy throughout; Penwood soils have red subsoil; Hinckley soils are sandy and gravelly. Also included are moderately well drained Ninigret soils in areas with a finer surface texture. Small areas of poorly drained Walpole soils and very poorly drained Scarboro soils are included in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland or residential development. Some areas are in cultivated cropland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Droughtiness can make establishment and maintenance of lawns difficult. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

25A—Brancroft silt loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 25 acres.

Map Unit Composition

Brancroft and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 17 inches; silt loam

Bw2—17 to 22 inches; silty clay loam

Bw3—22 to 32 inches; silt loam

C1—32 to 43 inches; silty clay loam

C2—43 to 66 inches; silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: fine-silty glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high
Reaction: very strongly acid to neutral
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w
Hydrologic group: C

Minor Components

Included with this soil in mapping are some areas of moderately well drained Elmridge and Berlin soils. Elmridge soils have a loamy over clayey substratum; Berlin soils are reddish brown in color. Also included are poorly drained Scitico soils in slight depressions on the landscape. Very poorly drained Maybid soils are included in deep depressions and drainageways (fig. 8). A few areas are underlain by sand and gravel or glacial till at 2 to 3 feet depths. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are wooded. A small acreage is in sod development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Low strength and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

25B—Brancroft silt loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping
Landscape: terraces on lake plains
Size of map unit: Areas commonly range from 3 to 25 acres.

Map Unit Composition

Brancroft and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam
 Bw1—6 to 17 inches; silt loam
 Bw2—17 to 22 inches; silty clay loam
 Bw3—22 to 32 inches; silt loam
 C1—32 to 43 inches; silty clay loam
 C2 43 to 66 inches; silt loam



Figure 8. —Small areas of very poorly drained Maybid soils are common in glaciolacustrine soils such as Brancroft silt loam.

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: fine-silty glaciolacustrine deposits
Permeability: very slow to moderate
Available water capacity: very high
Reaction: very strongly acid to neutral
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e
Hydrologic group: C

Minor Components

Included with this soil in mapping are some areas of moderately well drained Elmridge and Berlin soils. Elmridge soils have a loamy over clayey substratum; Berlin soils are reddish brown in color. Also included are poorly drained Scitico soils in slight depressions on the landscape. Very poorly drained Maybid soils are included in deep depressions and drainageways. A few areas are underlain by sand and gravel or glacial till at 2 to 3 feet depths. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are wooded. A small acreage is in sod development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Low strength and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

25C—Brancroft silt loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 25 acres.

Map Unit Composition

Brancroft and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 17 inches; silt loam

Bw2—17 to 22 inches; silty clay loam

Bw3—22 to 32 inches; silt loam

C1—32 to 43 inches; silty clay loam

C2—43 to 66 inches; silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: fine-silty glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Minor Components

Included with this soil in mapping are some areas of moderately well drained Elmridge and Berlin soils. Elmridge soils have a loamy over clayey substratum; Berlin soils are reddish brown in color. Also included are poorly drained Scitico soils in slight depressions on the landscape. Very poorly drained Maybid soils are included in deep depressions and drainageways. A few areas are underlain by sand and gravel or

glacial till at 2 to 3 feet depths. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are wooded. A small acreage is in sod development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Low strength and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

26A—Berlin silt loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Berlin and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 12 inches; silt loam

Bw2—12 to 20 inches; silty clay loam

Bw3—20 to 34 inches; silty clay loam

C1—34 to 48 inches; silty clay loam

C2—48 to 65 inches; silty clay loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: fine-silty glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 12 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Brancroft, Elmridge, and Belgrade soils. These soils are yellower in the subsoil and substratum. Also included are areas of moderately well drained Ludlow soils, which have a dense substratum. Well drained Wethersfield soils are included in areas that are higher on the landscape and have a dense substratum. Small areas of poorly drained Scitico soils and very poorly drained Maybid soils are included in depressions and along drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development. Some areas are in corn, vegetable or nursery crops, hay, or pasture. A few areas are in woodland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Low strength and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

26B—Berlin silt loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Berlin and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 12 inches; silt loam

Bw2—12 to 20 inches; silty clay loam

Bw3—20 to 34 inches; silty clay loam

C1—34 to 48 inches; silty clay loam

C2—48 to 65 inches; silty clay loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: fine-silty glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 12 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Brancroft, Elmridge, and Belgrade soils. These soils are yellower in the subsoil and substratum. Also included are areas of moderately well drained Ludlow soils, which have a dense substratum. Well drained Wethersfield soils are included in areas that are higher on the landscape and have a dense substratum. Small areas of poorly drained Scitico soils and very poorly drained Maybid soils are included in depressions and along drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development. Some areas are in corn, vegetable or nursery crops, hay, or pasture. A few areas are in woodland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Low strength and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

27A—Belgrade silt loam, 0 to 5 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Belgrade and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 16 inches; silt loam

Bw2—16 to 27 inches; silt loam

C1—27 to 45 inches; silt loam

C2—45 to 60 inches; silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: silty glaciolacustrine deposits

Permeability: moderate

Available water capacity: very high

Reaction: very strongly acid to neutral
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 42 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w
Hydrologic group: B

Minor Components

Included with this soil in mapping are some areas of moderately well drained Elmridge, Brancroft, and Berlin soils. Elmridge soils have a loamy over clayey substratum; Brancroft soils are silty and clayey; Berlin soils are reddish brown in color. Also included are poorly drained Scitico, Raynham, and Shaker soils, and very poorly drained Maybid soils in depressions and along drainageways. Included in some areas are soils red in color. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in vegetables and nursery stock. A few areas are in woodland or community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

28A—Elmridge fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level
Landscape: terraces on lake plains
Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Elmridge and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam
 Bw1—6 to 10 inches; fine sandy loam
 Bw2—10 to 18 inches; fine sandy loam
 Bw3—18 to 25 inches; sandy loam
 2C—25 to 65 inches; silty clay

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained

Parent material: coarse-loamy eolian sands over clayey glaciolacustrine deposits

Permeability: very slow to moderately rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Brancroft, Sudbury, Ninigret, Berlin, and Belgrade soils. Brancroft soils are silty and clayey; Sudbury soils are sandy and gravelly; Ninigret soils are loamy over sand and gravel; Berlin soils are redder; and Belgrade soils are silty throughout. Also included are small areas of poorly drained Shaker and Scitico soils in slight depressions on the landscape. Small areas of very poorly drained Maybid soils are also included in depressions and along drainageways. A few areas include soils from red parent materials. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in vegetables or nursery stock. A few areas are in woodland or community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

28B—Elmridge fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Elmridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

Bw1—6 to 10 inches; fine sandy loam

Bw2—10 to 18 inches; fine sandy loam

Bw3—18 to 25 inches; sandy loam

2C—25 to 65 inches; silty clay

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian sands over clayey glaciolacustrine deposits

Permeability: very slow to moderately rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Brancroft, Sudbury, Ninigret, Berlin, and Belgrade soils. Brancroft soils are silty and clayey; Sudbury soils are sandy and gravelly; Ninigret soils are loamy over sand and gravel; Berlin soils are redder; and Belgrade soils are silty throughout. Also included are small areas of poorly drained Shaker and Scitico soils in slight depressions on the landscape. Small areas of very poorly drained Maybid soils are also included in depressions and along drainageways. A few areas include soils from red parent materials. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in vegetables or nursery stock. A few areas are in woodland or community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

29A—Agawam fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Agawam and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam
 Bw1—8 to 14 inches; fine sandy loam
 Bw2—14 to 24 inches; fine sandy loam
 2C—24 to 60 inches; stratified very gravelly coarse sand to fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss
Permeability: moderately rapid to very rapid
Available water capacity: moderate
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 1
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are higher on the landscape; Hinckley and Merrimac soils are sandier in the subsoil. Also included are some moderately well drained Ninigret soils in slightly lower areas on the landscape. Poorly drained Walpole soils and very poorly drained Scarboro soils are included in depressions and drainageways. Scarboro soils have a mucky surface layer. A few areas in Hartford County include soils with reddish brown subsoil. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, community development, woodland, or are mined for sand and gravel. Some areas are in vegetables or nursery cropland.

This soil has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

This soil has few limitations for local roads and streets.

29B—Agawam fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping
Landscape: outwash plains on valleys, terraces on valleys
Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Agawam and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are higher on the landscape; Hinckley and Merrimac soils are sandier in the subsoil. Also included are some moderately well drained Ninigret soils in slightly lower areas on the landscape. Poorly drained Walpole soils and very poorly drained Scarboro soils are included in depressions and drainageways. Scarboro soils have a mucky surface layer. A few areas in Hartford County include soils with reddish brown subsoil. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, community development, woodland, or are mined for sand and gravel.

This soil has few limitations for dwellings with basements and lawns and landscaping. Droughtiness can make establishment and maintenance of lawns difficult.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

This soil has few limitations for local roads and streets.

29C—Agawam fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Agawam and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are higher on the landscape; Hinckley and Merrimac soils are sandier in the subsoil. Also included are some moderately well drained Ninigret soils in slightly lower areas on the landscape. Poorly drained Walpole soils and very poorly drained Scarboro soils are included in depressions and drainageways. Scarboro soils have a mucky surface layer. A few areas in Hartford County include soils with reddish brown subsoil. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, community development, woodland, or are mined for sand and gravel.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

30A—Branford silt loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Branford and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; loam

Bw2—18 to 24 inches; gravelly loam

2C—24 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Haven and Enfield soils. Enfield soils are coarse-silty over sand and gravel and Haven soils are coarse-loamy over sand and gravel. Also included are moderately well drained Ellington soils in slightly lower areas of the landscape. Excessively drained, sandy and gravelly Manchester soils and somewhat excessively drained, sandy Hartford soils are included in areas that are higher on the landscape. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland, nursery crops, or vegetable crops.

This soil has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

30B—Branford silt loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 80 acres.

Map Unit Composition

Branford and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; loam

Bw2—18 to 24 inches; gravelly loam

2C—24 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Haven and Enfield soils. Enfield soils are coarse-silty over sand and gravel and Haven soils are coarse-loamy over sand and gravel. Also included are moderately well drained Ellington soils in slightly lower areas of the landscape. Excessively drained, sandy and gravelly Manchester soils and somewhat excessively drained, sandy Hartford soils are included in areas that are higher on the landscape. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland.

This soil has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

30C—Branford silt loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Branford and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; loam

Bw2—18 to 24 inches; gravelly loam

2C—24 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Haven and Enfield soils. Enfield soils are coarse-silty over sand and gravel and Haven soils are coarse-loamy over sand and gravel. Also included are moderately well drained Ellington soils in slightly lower areas of the landscape. Excessively drained, sandy and gravelly Manchester soils and somewhat excessively drained, sandy Hartford soils are included in areas that are higher on the landscape. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland or pasture.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

31A—Copake fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on valleys, outwash plains on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Copake and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

AB—6 to 13 inches; gravelly fine sandy loam

Bw1—13 to 21 inches; gravelly fine sandy loam

Bw2—21 to 31 inches; gravelly fine sandy loam

2C1—31 to 56 inches; very gravelly coarse sand

2C2—56 to 65 inches; fine sand

2C3—65 to 75 inches; gravelly sand

2C4—75 to 80 inches; gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained, sandy and gravelly Groton soils that are higher on the landscape. Also included are areas of moderately well drained Hero soils in slightly lower areas of the landscape. Small areas of poorly drained Fredon soils and very poorly drained Halsey soils are included in depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in woodland, residential development, or are mined for sand and gravel.

This soil has few limitations for dwelling with basements, lawns and landscaping. Small stones are also a limitation for lawns and landscaping.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

Frost action is the main limitation for local roads and streets. Providing a coarser grained subgrade to frost depth will reduce this limitation.

31B—Copake fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: terraces on valleys, outwash plains on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Copake and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

AB—6 to 13 inches; gravelly fine sandy loam

Bw1—13 to 21 inches; gravelly fine sandy loam

Bw2—21 to 31 inches; gravelly fine sandy loam

2C1—31 to 56 inches; very gravelly coarse sand

2C2—56 to 65 inches; fine sand

2C3—65 to 75 inches; gravelly sand

2C4—75 to 80 inches; gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained, sandy and gravelly Groton soils that are higher on the landscape. Also included are areas of moderately well drained Hero soils in slightly lower areas on the landscape. Small areas of poorly drained Fredon soils and very poorly drained Halsey soils are included in depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in woodland, residential development, or are mined for sand and gravel.

This soil has few limitations for dwelling with basements, lawns and landscaping. Small stones are also a limitation for lawns and landscaping.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

Frost action is the main limitation for local roads and streets. Providing a coarser grained subgrade to frost depth will reduce this limitation.

31C—Copake gravelly loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: outwash plains on valleys, terraces on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Copake and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

AB—6 to 13 inches; gravelly fine sandy loam

Bw1—13 to 21 inches; gravelly fine sandy loam

Bw2—21 to 31 inches; gravelly fine sandy loam

2C1—31 to 56 inches; very gravelly coarse sand

2C2—56 to 65 inches; fine sand

2C3—65 to 75 inches; gravelly sand

2C4—75 to 80 inches; gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: high
Reaction: strongly acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained, sandy and gravelly Groton soils that are higher on the landscape. Also included are areas of moderately well drained Hero soils in slightly lower areas on the landscape. Small areas of poorly drained Fredon soils and very poorly drained Halsey soils are included in depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in woodland, residential development, or are mined for sand and gravel.

Slope is the main limitation for dwelling with basements, lawns and landscaping. Small stones are also a limitation for lawns and landscaping. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Erosion is a moderate hazard during construction.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

Frost action and slope are the main limitations for local roads and streets. Providing a coarser grained subgrade to frost depth and constructing roads on the contour will reduce these limitations.

32A—Haven and Enfield soils, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level
Landscape: outwash plains on valleys, terraces on valleys
Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Haven and similar soils: 60 percent
Enfield and similar soils: 25 percent
Minor components: 15 percent

Major Components

Haven and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; silt loam
 Bw1—7 to 14 inches; silt loam
 Bw2—14 to 20 inches; silt loam
 BC—20 to 24 inches; fine sandy loam
 2C—24 to 60 inches; stratified very gravelly sand to gravelly fine sand

Enfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 3 inches; slightly decomposed plant material

Oe—3 to 4 inches; moderately decomposed plant material

Ap—4 to 12 inches; silt loam

Bw1—12 to 20 inches; silt loam

Bw2—20 to 26 inches; silt loam

Bw3—26 to 30 inches; silt loam

2C—30 to 37 inches; stratified coarse sand to very gravelly loamy sand

3C—37 to 65 inches; stratified very gravelly coarse sand to loamy sand

Major Component Properties and Qualities**Haven and similar soils**

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Enfield and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-silty eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups**Enfield and similar soils**

Land capability classification (non-irrigated): 1

Hydrologic group: B

Haven and similar soils

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Branford and Agawam soils. Branford soils are silty over sand and gravel, and are red in color. Agawam soils are sandier in the surface layer and subsoil. Also included are moderately well drained Ninigret and Tisbury soils in slightly lower areas of the landscape. Poorly drained Raypol soils are included in depressions and drainageways. A few areas in New London County include soils with a gravelly surface layer. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, woodland, or community development.

This unit has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

32B—Haven and Enfield soils, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Haven and similar soils: 60 percent

Enfield and similar soils: 25 percent

Minor components: 15 percent

Major Components

Haven and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; silt loam

Bw1—7 to 14 inches; silt loam

Bw2—14 to 20 inches; silt loam

BC—20 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly sand to gravelly fine sand

Enfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 3 inches; slightly decomposed plant material

Oe—3 to 4 inches; moderately decomposed plant material

Ap—4 to 12 inches; silt loam

Bw1—12 to 20 inches; silt loam

Bw2—20 to 26 inches; silt loam

Bw3—26 to 30 inches; silt loam

2C—30 to 37 inches; stratified coarse sand to very gravelly loamy sand

3C—37 to 65 inches; stratified very gravelly coarse sand to loamy sand

Major Component Properties and Qualities

Haven and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Enfield and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-silty eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Haven and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Enfield and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Branford and Agawam soils. Branford soils are silty over sand and gravel, and are red in color. Agawam soils are sandier in the surface layer and subsoil. Also included are moderately well drained Ninigret and Tisbury soils in slightly lower areas of the landscape. Poorly drained Raypol soils are included in depressions and drainageways. A few areas in New London County include soils with a gravelly surface layer. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, woodland, or community development.

This unit has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

32C—Haven and Enfield soils, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Haven and similar soils: 60 percent

Enfield and similar soils: 25 percent

Minor components: 15 percent

Major Components

Haven and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; silt loam

Bw1—7 to 14 inches; silt loam

Bw2—14 to 20 inches; silt loam

BC—20 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly sand to gravelly fine sand

Enfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 3 inches; slightly decomposed plant material

Oe—3 to 4 inches; moderately decomposed plant material

Ap—4 to 12 inches; silt loam

Bw1—12 to 20 inches; silt loam

Bw2—20 to 26 inches; silt loam

Bw3—26 to 30 inches; silt loam

2C—30 to 37 inches; stratified coarse sand to very gravelly loamy sand

3C—37 to 65 inches; stratified very gravelly coarse sand to loamy sand

Major Component Properties and Qualities

Haven and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Enfield and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-silty eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Haven and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Enfield and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Branford and Agawam soils. Branford soils are silty over sand and gravel, and are red in color. Agawam soils are sandier in the surface layer and subsoil. Also included are moderately well drained Ninigret and Tisbury soils in slightly lower areas of the landscape. Poorly drained Raypol soils are included in depressions and drainageways. A few areas in New London County include soils with a gravelly surface layer. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, woodland, or community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the limitation.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

33A—Hartford sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Hartford and similar soils: 80 percent

Minor components: 20 percent

Major Components

Hartford and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; sandy loam ([fig. 9](#))

Bw1—8 to 20 inches; sandy loam

Bw2—20 to 26 inches; loamy sand

2C—26 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy glaciofluvial deposits derived from sandstone and/or basalt

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none



Figure 9.—The upper part of a typical profile of Hartford sandy loam.

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Manchester and Penwood soils that are higher on the landscape. Manchester soils are sandy and gravelly throughout and Penwood soils are sandier in the surface layer and subsoil. Also included are well drained Branford soils and moderately well drained Ellington soils. Branford soils are silty over sand and gravel and Ellington soils are in slightly lower areas and broad drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development, cultivated crops, or nursery stock. This soil has few limitations for dwelling with basements, lawns and landscaping, and local roads and streets.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

33B—Hartford sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Hartford and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; sandy loam

Bw1—8 to 20 inches; sandy loam

Bw2—20 to 26 inches; loamy sand

2C—26 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy glaciofluvial deposits derived from sandstone and/or basalt

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Manchester and Penwood soils that are higher on the landscape. Manchester soils are sandy and gravelly throughout and Penwood soils are sandier in the surface layer and subsoil. Also included are well drained Branford soils and moderately well drained Ellington soils. Branford soils are silty over sand and gravel and Ellington soils are in slightly lower areas and broad drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development, cultivated crops, or nursery stock.

This soil has few limitations for dwelling with basements, lawns and landscaping, and local roads and streets.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

34A—Merrimac sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on valleys, outwash plains on valleys, kames on valleys

Size of map unit: Areas commonly range from 5 to 75 acres.

Map Unit Composition

Merrimac and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; sandy loam

Bw1—9 to 16 inches; sandy loam

Bw2—16 to 24 inches; gravelly sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley and Windsor soils that are higher on the landscape. Hinckley soils are sandy and gravelly and Windsor soils are sandy throughout. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Ninigret and Sudbury soils are included in slightly lower areas of the landscape. Ninigret soils are loamy over sand and gravel and Sudbury soils are sandy and gravelly. Small areas of poorly drained Walpole soils and very poorly drained Scarboro soils are included in depressions and drainageways. A few areas include soils with a fine sandy loam surface texture. Reddish brown soils are included in the southern part of the town of Woodbury, in Litchfield County. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland, vegetable or nursery crops, or pasture.

This soil has few limitations for dwellings with basements, lawns and landscaping, and local roads and streets.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

34B—Merrimac sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 5 to 50 acres.

Map Unit Composition

Merrimac and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; sandy loam

Bw1—9 to 16 inches; sandy loam

Bw2—16 to 24 inches; gravelly sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley and Windsor soils that are higher on the landscape. Hinckley soils are sandy and gravelly and Windsor soils are sandy throughout. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Ninigret and Sudbury soils are included in slightly lower areas on the landscape. Ninigret soils are loamy

over sand and gravel and Sudbury soils are sandy and gravelly. Small areas of poorly drained Walpole soils and very poorly drained Scarboro soils are included in depressions and drainageways. A few areas include soils with a fine sandy loam surface texture. Reddish brown soils are included in the southern part of the town of Woodbury, in Litchfield County. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland, vegetable or nursery crops, or pasture.

This soil has few limitations for dwellings with basements, lawns and landscaping, and local roads and streets.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

34C—Merrimac sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: outwash plains on valleys, terraces on valleys, kames on valleys

Size of map unit: Areas commonly range from 5 to 40 acres.

Map Unit Composition

Merrimac and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; sandy loam

Bw1—9 to 16 inches; sandy loam

Bw2—16 to 24 inches; gravelly sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley and Windsor soils that are higher on the landscape. Hinckley soils are sandy and gravelly and Windsor soils are sandy throughout. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Ninigret and Sudbury soils are included in slightly lower areas of the landscape. Ninigret soils are loamy over sand and gravel and Sudbury soils are sandy and gravelly. Small areas of poorly drained Walpole soils and very poorly drained Scarboro soils are included in depressions and drainageways. A few areas include soils with a fine sandy loam surface texture. Reddish brown soils are included in the southern part of the town of Woodbury, in Litchfield County. Minoe components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland or pasture.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

35A—Penwood loamy sand, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Penwood and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; loamy sand

Bw1—8 to 18 inches; loamy sand

Bw2—18 to 30 inches; sand

C—30 to 60 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy glaciofluvial deposits derived from sandstone and shale

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Minor Components

Included with this soil in mapping are excessively drained Manchester soils and somewhat excessively drained Hartford soils in areas that are sandy and gravelly. Also included are areas of well drained Branford soils and moderately well drained Ellington soils in slightly lower areas of the landscape. Branford and Ellington soils are silty over sand and gravel. A few areas in New Haven County include soils with a gravelly substratum. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development. Some areas in Middlesex County are in cultivated cropland.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

35B—Penwood loamy sand, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Penwood and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; loamy sand

Bw1—8 to 18 inches; loamy sand

Bw2—18 to 30 inches; sand

C—30 to 60 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy glaciofluvial deposits derived from sandstone and shale

Permeability: rapid to very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Minor Components

Included with this soil in mapping are excessively drained Manchester soils and somewhat excessively drained Hartford soils in areas that are sandy and gravelly. Also included are areas of well drained Branford soils and moderately well drained Ellington soils in slightly lower areas of the landscape. Branford and Ellington soils are silty over sand and gravel. A few areas in New Haven County include soils with a gravelly substratum. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development. Some areas in Middlesex County are in cultivated cropland.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

36A—Windsor loamy sand, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Windsor and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loamy sand

Bw1—3 to 9 inches; loamy sand

Bw2—9 to 21 inches; loamy sand

Bw3—21 to 25 inches; sand

C—25 to 65 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: eolian sands over sandy glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are sandy and gravelly. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Deerfield, Ninigret, and Sudbury soils are included in slightly lower areas of the landscape. Ninigret soils are loamy over sand and gravel and Sudbury soils are sandy and gravelly. A few valleys in Litchfield County include areas with neutral or less acid subsoil. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cropland or community development. Some areas are in woodland, brushland, or pasture.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

36B—Windsor loamy sand, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: outwash plains on valleys, terraces on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Windsor and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loamy sand

Bw1—3 to 9 inches; loamy sand

Bw2—9 to 21 inches; loamy sand

Bw3—21 to 25 inches; sand

C—25 to 65 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: eolian sands over sandy glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are sandy and gravelly. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Deerfield, Ninigret, and Sudbury soils are included in slightly lower areas of the landscape. Ninigret soils are loamy over sand and gravel and Sudbury soils are sandy and gravelly. A few valleys in Litchfield County include areas with neutral or less acid subsoil. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cropland or community development. Some areas are in woodland, brushland, or pasture.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

36C—Windsor loamy sand, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: terraces on valleys, outwash plains on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Windsor and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loamy sand

Bw1—3 to 9 inches; loamy sand

Bw2—9 to 21 inches; loamy sand

Bw3—21 to 25 inches; sand

C—25 to 65 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: eolian sands over sandy glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are sandy and gravelly. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Deerfield, Ninigret, and Sudbury soils are included in slightly lower areas of the landscape. Ninigret soils are loamy over sand and gravel and Sudbury soils are sandy and gravelly. A few valleys in Litchfield County include areas with neutral or less acid subsoil. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cropland or community development. Some areas are in woodland, brushland, or pasture.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

37A—Manchester gravelly sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: eskers on valleys, kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Manchester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; gravelly sandy loam

Bw—9 to 18 inches; gravelly loamy sand

C—18 to 65 inches; stratified extremely gravelly coarse sand to very gravelly loamy sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Penwood soils that are sandy throughout. Also included are somewhat excessively drained Hartford soils, well drained Branford soils, and moderately well drained Ellington soils. Hartford soils are sandy loam over a sandy and gravelly substratum, Branford soils are silty over a sandy and gravelly substratum, and Ellington soils are in slightly lower areas and broad drainageways. In places, soils that lack a gravelly surface are included. A few areas in New Haven County have a gravelly loamy sand surface layer. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland or community development. Some areas are in woodland, nursery crops, gravel pits, or pasture.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

37C—Manchester gravelly sandy loam, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: terraces on valleys, eskers on valleys, kames on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Manchester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; gravelly sandy loam

Bw—9 to 18 inches; gravelly loamy sand

C—18 to 65 inches; stratified extremely gravelly coarse sand to very gravelly loamy sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Penwood soils that are sandy throughout. Also included are somewhat excessively drained Hartford soils, well drained Branford soils, and moderately well drained Ellington soils. Hartford soils are sandy loam over a sandy and gravelly substratum, Branford soils are silty over a sandy and gravelly substratum, and Ellington soils are in slightly lower areas and broad drainageways. In places, soils that lack a gravelly surface are included. A few areas in New Haven County have a gravelly loamy sand surface layer. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland or community development. Some areas are in woodland, nursery crops, gravel pits, or pasture.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Droughtiness is the main limitation for lawns and

landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

37E—Manchester gravelly sandy loam, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep to steep

Landscape: eskers on valleys, kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Manchester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; gravelly sandy loam

Bw—9 to 18 inches; gravelly loamy sand

C—18 to 65 inches; stratified extremely gravelly coarse sand to very gravelly loamy sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7e

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Penwood soils that are sandy throughout. Also included are somewhat excessively drained Hartford soils, well drained Branford soils, and moderately well drained Ellington soils. Hartford soils are sandy loam over a sandy and gravelly substratum, Branford soils are silty over a sandy and gravelly substratum, and Ellington soils are in slightly lower areas

and broad drainageways. In places, soils that lack a gravelly surface are included. A few areas in New Haven County have a gravelly loamy sand surface layer. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or gravel pits.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

38A—Hinckley gravelly sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on valleys, outwash plains on valleys, kames on valleys, eskers on valleys

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Hinckley and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 20 inches; very gravelly loamy sand

Bw2—20 to 27 inches; very gravelly sand

C1—27 to 42 inches; stratified cobbly coarse sand to extremely gravelly sand

C2—42 to 60 inches; stratified cobbly coarse sand to extremely gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: very low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Windsor soils which are sandy throughout. Also included are somewhat excessively drained Merrimac soils and well drained Agawam soils. Merrimac soils are sandy over sand and gravel and Agawam soils are loamy over sand and gravel. Small areas of moderately well drained Sudbury soils are included in slightly lower areas, poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. A few areas in Litchfield and Hartford counties include soils with a reddish brown color. Windham County includes some soils with a fine sandy loam surface. New London County includes some soils with less gravel or a gravelly silt loam surface and subsoil. New Haven County includes some soils with less gravel or a gravelly loamy sand surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland, pasture, vegetable or nursery crops, or are mined for sand and gravel.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

38C—Hinckley gravelly sandy loam, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: terraces on valleys, outwash plains on valleys, eskers on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Hinckley and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 20 inches; very gravelly loamy sand

Bw2—20 to 27 inches; very gravelly sand

C1—27 to 42 inches; stratified cobbly coarse sand to extremely gravelly sand

C2—42 to 60 inches; stratified cobbly coarse sand to extremely gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: very low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Windsor soils which are sandy throughout. Also included are somewhat excessively drained Merrimac soils and well drained Agawam soils. Merrimac soils are sandy over sand and gravel and Agawam soils are loamy over sand and gravel. Small areas of moderately well drained Sudbury soils are included in slightly lower areas, poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. A few areas in Litchfield and Hartford counties include soils with a reddish brown color. Windham County includes some soils with a fine sandy loam surface. New London County includes some soils with less gravel or a gravelly silt loam surface and subsoil. New Haven County includes some soils with less gravel or a gravelly loamy sand surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops or community development. Some areas are in woodland, pasture, vegetable or nursery crops, or are mined for sand and gravel.

Slope is the main limitation for dwellings with basements. Erosion is a hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

38E—Hinckley gravelly sandy loam, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep to steep

Landscape: eskers on valleys, terraces on valleys, outwash plains on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Hinckley and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 20 inches; very gravelly loamy sand

Bw2—20 to 27 inches; very gravelly sand

C1—27 to 42 inches; stratified cobbly coarse sand to extremely gravelly sand

C2—42 to 60 inches; stratified cobbly coarse sand to extremely gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: very low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6e

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Windsor soils which are sandy throughout. Also included are somewhat excessively drained Merrimac soils and well drained Agawam soils. Merrimac soils are sandy over sand and gravel and Agawam soils are loamy over sand and gravel. Small areas of moderately well drained Sudbury soils are included in slightly lower areas, poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. A few areas in Litchfield and Hartford counties include soils with a reddish brown color. Windham County includes some soils with a fine sandy loam surface. New London County includes some soils with less gravel or a gravelly silt loam surface and subsoil. New Haven County includes some soils with less gravel or a gravelly loamy sand surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland or community development. Some areas are in pasture, cropland or are mined for sand and gravel.

Slope is the main limitation for dwellings with basements. Erosion is a hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not

adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

39A—Groton gravelly sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: eskers on valleys, kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Groton and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 18 inches; very gravelly sandy loam

Bw2—18 to 24 inches; very gravelly loamy sand

Bw3—24 to 30 inches; very gravelly loamy sand

C1—30 to 52 inches; stratified extremely gravelly coarse sand to very gravelly loamy fine sand

C2—52 to 72 inches; stratified extremely gravelly coarse sand to gravelly loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderately rapid to very rapid

Available water capacity: low

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of well drained Copake soil that are loamy over sand and gravel. Also included are areas of moderately well drained Hero soils in slightly lower areas on the landscape. Poorly drained Fredon soils and very poorly drained Halsey soils are included in shallow depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in woodland, residential development, or are mined for sand and gravel.

This soil has few limitations for dwelling with basements. Small stones and droughtiness are limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

This soil has few limitations for local roads and streets.

39C—Groton gravelly sandy loam, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: eskers on valleys, kames on valleys, terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Groton and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 18 inches; very gravelly sandy loam

Bw2—18 to 24 inches; very gravelly loamy sand

Bw3—24 to 30 inches; very gravelly loamy sand

C1—30 to 52 inches; stratified extremely gravelly coarse sand to very gravelly loamy fine sand

C2—52 to 72 inches; stratified extremely gravelly coarse sand to gravelly loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderately rapid to very rapid

Available water capacity: low

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of well drained Copake soil that are loamy over sand and gravel. Also included are areas of moderately well drained Hero soils in slightly lower areas of the landscape. Poorly drained Fredon soils and very poorly drained Halsey soils are included in shallow depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in woodland, residential development, or are mined for sand and gravel.

Slope is the main limitation for dwelling with basements, lawns and landscaping. Small stones and droughtiness are also limitations for lawns and landscaping. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Erosion is a hazard during construction. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce these limitations.

39E—Groton gravelly sandy loam, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep to steep

Landscape: eskers on valleys, kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Groton and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 18 inches; very gravelly sandy loam

Bw2—18 to 24 inches; very gravelly loamy sand

Bw3—24 to 30 inches; very gravelly loamy sand

C1—30 to 52 inches; stratified extremely gravelly coarse sand to very gravelly loamy fine sand

C2—52 to 72 inches; stratified extremely gravelly coarse sand to gravelly loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderately rapid or very rapid

Available water capacity: low

Reaction: moderately acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6e
Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of well drained Copake soil that are loamy over sand and gravel. Also included are areas of moderately well drained Hero soils in slightly lower areas of the landscape. Poorly drained Fredon soils and very poorly drained Halsey soils are included in shallow depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland, residential development, or pasture. Some areas are mined for sand and gravel.

Slope is the main limitation for dwelling with basements, lawns and landscaping. Small stones and droughtiness are also limitations for lawns and landscaping. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Erosion is a hazard during construction. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is also a limitation for septic tank absorption. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce these limitations.

40A—Ludlow silt loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level
Landscape: hills on uplands, drumlins on uplands
Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ludlow and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam
 Bw1—8 to 20 inches; silt loam
 Bw2—20 to 26 inches; silt loam
 Cd—26 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Wethersfield soils that are higher on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are included in depressions and drainageways. Moderately well drained Watchaug soils and well drained Cheshire soils are in areas lacking a dense substratum. Moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches below the surface. Also included are soils with a stony surface and soils with a loam or fine sandy loam surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or farmland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil. Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

40B—Ludlow silt loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Ludlow and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 20 inches; silt loam

Bw2—20 to 26 inches; silt loam
 Cd—26 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale
Permeability: very slow to moderate
Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 40 inches to densic material
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e
Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Wethersfield soils that are higher on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are included in depressions and drainageways. Moderately well drained Watchaug soils and well drained Cheshire soils are in areas lacking a dense substratum. Moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches below the surface. Also included are soils with a stony surface and soils with a loam or fine sandy loam surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or farmland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil. Frost action is the main limitation for roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

41B—Ludlow silt loam, 2 to 8 percent slopes, very stony

Map Unit Setting

Slope: nearly level to gently sloping
Landscape: drumlins on uplands, hills on uplands
Surface cover: 0 to 3 percent stones
Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ludlow and similar soils: 80 percent

Minor components: 20 percent

Major Components

Ludlow and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; silt loam

Bw1—8 to 20 inches; silt loam

Bw2—20 to 26 inches; silt loam

Cd—26 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Wethersfield soils that are higher on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are included in depressions and drainageways. Moderately well drained Watchaug soils and well drained Cheshire soils are in areas lacking a dense substratum. Moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches below the surface. Also included are soils with a stony surface and soils with a loam or fine sandy loam surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or farmland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil. Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

42C—Ludlow silt loam, 2 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: nearly level to strongly sloping
Landscape: hills on uplands, drumlins on uplands
Surface cover: 3 to 15 percent stones
Size of map unit: 3 to 50 acres

Map Unit Composition

Ludlow and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; silt loam
Bw1—8 to 20 inches; silt loam
Bw2—20 to 26 inches; silt loam
Cd—26 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale
Permeability: very slow to moderate
Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 40 inches to densic material
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s
Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Wethersfield soils that are higher on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are included in depressions and drainageways. Moderately well drained Watchaug soils and well drained Cheshire soils are in areas lacking a dense substratum. Moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches below the surface. Also included are soils with a stony surface and soils with a loam or fine sandy loam surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or farmland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside

of basement walls, and diverting runoff from higher areas will reduce wetness. The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil. Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

43A—Rainbow silt loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas are commonly 3 to 30 acres.

Map Unit Composition

Rainbow and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 18 inches; silt loam

Bw2—18 to 26 inches; silt loam

2Cd—26 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook soils that are higher on the landscape. Also included are moderately well drained Sutton and Woodbridge soils. Sutton soils lack a dense substratum and Woodbridge soils are less silty. Poorly drained Ridgebury and Wilbraham soils are in depressions and drainageways. Ridgebury soils are less silty and Wilbraham soils are red. Well drained Narragansett soils are included in areas that lack a dense substratum. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland, cultivated crop, hay, or pasture. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil. Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

43B—Rainbow silt loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas are commonly 3 to 30 acres.

Map Unit Composition

Rainbow and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 18 inches; silt loam

Bw2—18 to 26 inches; silt loam

2Cd—26 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook soils that are higher on the landscape. Also included are moderately well drained Sutton and

Woodbridge soils. Sutton soils lack a dense substratum and Woodbridge soils are less silty. Poorly drained Ridgebury and Wilbraham soils are in depressions and drainageways. Ridgebury soils are less silty and Wilbraham soils are red. Well drained Narragansett soils are included in areas that lack a dense substratum. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland, cultivated crop, hay, or pasture. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil. Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

44B—Rainbow silt loam, 2 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas are commonly 3 to 30 acres.

Map Unit Composition

Rainbow and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 18 inches; silt loam

Bw2—18 to 26 inches; silt loam

2Cd—26 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook soils that are higher on the landscape. Also included are moderately well drained Sutton and Woodbridge soils. Sutton soils lack a dense substratum and Woodbridge soils are less silty. Poorly drained Ridgebury and Wilbraham soils are in depressions and drainageways. Ridgebury soils are less silty and Wilbraham soils are red. Well drained Narragansett soils are included in areas that lack a dense substratum. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland or pasture. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil. Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

45A—Woodbridge fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Woodbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; fine sandy loam

Bw1—7 to 18 inches; fine sandy loam

Bw2—18 to 26 inches; fine sandy loam

Bw3—26 to 30 inches; fine sandy loam

Cd1—30 to 43 inches; gravelly fine sandy loam

Cd2—43 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Paxton and Montauk soils that are higher on the landscape. Also included are areas of poorly drained Ridgebury soils and very poorly drained Whitman soils in depressions and along drainageways. Moderately well drained Sutton soils are included in areas lacking a dense substratum. Poorly drained Leicester soils are in depressions and lack a dense substratum. In Fairfield and Litchfield counties where the soil is less acid and lacks a dense substratum, some areas of well drained Stockbridge soils and moderately well drained Georgia soils are included. A few areas in New London County include a loamy sand substratum. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are used for cropland, hayland, pastureland, or woodland. Some areas are used for community development or woodland.

community development or woodland

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

45B—Woodbridge fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 80 acres.

Map Unit Composition

Woodbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; fine sandy loam
 Bw1—7 to 18 inches; fine sandy loam
 Bw2—18 to 26 inches; fine sandy loam
 Bw3—26 to 30 inches; fine sandy loam
 Cd1—30 to 43 inches; gravelly fine sandy loam
 Cd2—43 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss
Permeability: very slow to moderate
Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 40 inches to densic material
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w
Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Paxton and Montauk soils that are higher on the landscape. Also included are areas of poorly drained Ridgebury soils and very poorly drained Whitman soils in depressions and along drainageways. Moderately well drained Sutton soils are included in areas lacking a dense substratum. Poorly drained Leicester soils are in depressions and lack a dense substratum. In Fairfield and Litchfield Counties where the soil is less acid and lacks a dense substratum, some areas of well drained Stockbridge soils and moderately well drained Georgia soils are included. A few areas in New London County include a loamy sand substratum. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in cultivated crops, hay, pasture or woodland. Some areas are in community development.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

45C—Woodbridge fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Woodbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; fine sandy loam

Bw1—7 to 18 inches; fine sandy loam

Bw2—18 to 26 inches; fine sandy loam

Bw3—26 to 30 inches; fine sandy loam

Cd1—30 to 43 inches; gravelly fine sandy loam

Cd2—43 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Paxton and Montauk soils that are higher on the landscape. Also included are areas of poorly drained Ridgebury soils and very poorly drained Whitman soils in depressions and along drainageways. Moderately well drained Sutton soils are included in areas lacking a dense substratum. Poorly drained Leicester soils are in depressions and lack a dense substratum. In Fairfield and Litchfield Counties where the soil is less acid and lacks a dense substratum, some areas of well drained Stockbridge soils and moderately well drained Georgia soils are included. A few areas in New London County include a loamy sand substratum. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in cultivated crops, hay, pasture, or woodland. Some areas are in community development.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Slope is also a limitation for lawns and landscaping. Erosion is a hazard during construction. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

46B—Woodbridge fine sandy loam, 2 to 8 percent slopes, very stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Woodbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 7 inches; fine sandy loam

Bw1—7 to 18 inches; fine sandy loam

Bw2—18 to 26 inches; fine sandy loam

Bw3—26 to 30 inches; fine sandy loam

Cd1—30 to 43 inches; gravelly fine sandy loam

Cd2—43 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Paxton and Montauk soils that are higher on the landscape. Also included are areas of poorly drained Ridgebury soils and very poorly drained Whitman soils in depressions and along drainageways. Moderately well drained Sutton soils are included in areas lacking a dense substratum. Poorly drained Leicester soils are in depressions and lack a dense substratum. In Fairfield and Litchfield Counties where the soil is less acid and lacks a dense substratum, some areas of well drained Stockbridge soils and moderately well drained Georgia soils are included. A few areas in New London County include a loamy sand substratum. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or community development.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

46C—Woodbridge fine sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Woodbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 7 inches; fine sandy loam

Bw1—7 to 18 inches; fine sandy loam

Bw2—18 to 26 inches; fine sandy loam

Bw3—26 to 30 inches; fine sandy loam

Cd1—30 to 43 inches; gravelly fine sandy loam

Cd2—43 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Paxton and Montauk soils that are higher on the landscape. Also included are areas of poorly drained Ridgebury soils and very poorly drained Whitman soils in depressions and along drainageways. Moderately well drained Sutton soils are included in areas lacking a dense substratum. Poorly drained Leicester soils are in depressions and lack a dense substratum. In Fairfield and Litchfield Counties where the soil is less acid and lacks a dense substratum, some areas of well drained Stockbridge soils and moderately well drained Georgia soils are included. A few areas in New London County include a loamy sand substratum. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or community development.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Slope and large stones are also limitations for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

47C—Woodbridge fine sandy loam, 2 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: nearly level to strongly sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 80 acres.

Map Unit Composition

Woodbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 7 inches; fine sandy loam

Bw1—7 to 18 inches; fine sandy loam

Bw2—18 to 26 inches; fine sandy loam

Bw3—26 to 30 inches; fine sandy loam

Cd1—30 to 43 inches; gravelly fine sandy loam

Cd2—43 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Paxton and Montauk soils that are higher on the landscape. Also included are areas of poorly drained Ridgebury soils and very poorly drained Whitman soils in depressions and along drainageways. Moderately well drained Sutton soils are included in areas lacking a dense substratum. Poorly drained Leicester soils are in depressions and lack a dense substratum. In Fairfield and Litchfield counties where the soil is less acid and lacks a dense substratum, some areas of well drained Stockbridge soils and moderately well drained Georgia soils are included. A few areas in New London County include a loamy sand substratum. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland (fig. 10). Some areas are in pasture or community development.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Slope and large stones are also limitations for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.



Figure 10.—This extremely stony area of Woodbridge fine sandy loam is suited for woodland and wildlife habitat.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

48B—Georgia and Amenia silt loams, 2 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: hills, uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Georgia and similar soils: 50 percent

Amenia and similar soils: 35 percent

Minor components: 15 percent

Major Components

Georgia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam
 Bw1—8 to 14 inches; loam
 Bw2—14 to 24 inches; loam
 C—24 to 60 inches; gravelly fine sandy loam

Amenia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; silt loam
 Bw1—9 to 16 inches; silt loam
 Bw2—16 to 25 inches; silt loam
 C—25 to 60 inches; gravelly loam

Major Component Properties and Qualities

Georgia and similar soils

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy till derived from limestone and dolomite and/or schist
Permeability: moderately slow or moderate
Available water capacity: high
Reaction: strongly acid to neutral
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 36 inches
Flooding: none

Amenia and similar soils

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy till derived from limestone and dolomite and/or schist
Permeability: moderately slow to moderate
Available water capacity: high
Reaction: moderately acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 36 inches
Flooding: none

Interpretative Groups

Georgia and similar soils

Land capability classification (non-irrigated): 2e
Hydrologic group: B

Amenia and similar soils

Land capability classification (non-irrigated): 2e
Hydrologic group: B

Minor Components

Included with this unit in mapping are well drained Stockbridge and Nellis soils in higher areas of the landscape. Poorly drained Mudgepond soils and very poorly drained Alden soils are in depressions and along drainageways. Also included are well drained Paxton soils, moderately well drained Woodbridge soils, and poorly drained Ridgebury in areas with a dense substratum. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in cultivated cropland or woodland. Some areas, mostly in Fairfield County, are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

48C—Georgia and Amenia silt loams, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills, uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Georgia and similar soils: 50 percent

Amenia and similar soils: 35 percent

Minor components: 15 percent

Major Components

Georgia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; loam

Bw2—14 to 24 inches; loam

C—24 to 60 inches; gravelly fine sandy loam

Amenia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; silt loam

Bw1—9 to 16 inches; silt loam

Bw2—16 to 25 inches; silt loam

C—25 to 60 inches; gravelly loam

Major Component Properties and Qualities

Georgia and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow to moderate

Available water capacity: high

Reaction: strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Amenia and similar soils*Depth to bedrock:* very deep*Drainage class:* moderately well drained*Parent material:* coarse-loamy till derived from limestone and dolomite and/or schist*Permeability:* moderately slow to moderate*Available water capacity:* high*Reaction:* moderately acid to moderately alkaline*Depth to restrictive feature:* greater than 72 inches*Depth to seasonal water table:* 18 to 36 inches*Flooding:* none***Interpretative Groups*****Georgia and similar soils***Land capability classification (non-irrigated):* 3e*Hydrologic group:* B**Amenia and similar soils***Land capability classification (non-irrigated):* 3e*Hydrologic group:* B***Minor Components***

Included with this unit in mapping are well drained Stockbridge and Nellis soils in higher areas of the landscape. Poorly drained Mudgepond soils and very poorly drained Alden soils are in depressions and along drainageways. Also included are well drained Paxton soils, moderately well drained Woodbridge soils, and poorly drained Ridgebury in areas with a dense substratum. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in cultivated cropland or woodland. Some areas, mostly in Fairfield County, are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

49B—Georgia and Amenia silt loams, 3 to 8 percent slopes, very stony***Map Unit Setting****Slope:* gently sloping*Landscape:* uplands, hills*Surface cover:* 0 to 3 percent stones*Size of map unit:* Areas commonly range from 3 to 40 acres.

Map Unit Composition

Georgia and similar soils: 50 percent

Amenia and similar soils: 35 percent

Minor components: 15 percent

Major Components

Georgia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; loam

Bw2—14 to 24 inches; loam

C—24 to 60 inches; gravelly fine sandy loam

Amenia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 9 inches; silt loam

Bw1—9 to 16 inches; silt loam

Bw2—16 to 25 inches; silt loam

C—25 to 60 inches; gravelly loam

Major Component Properties and Qualities

Georgia and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow to moderate

Available water capacity: high

Reaction: strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Amenia and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow to moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Georgia and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Amenia and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this unit in mapping are well drained Stockbridge and Nellis soils in higher areas of the landscape. Poorly drained Mudgepond soils and very poorly

drained Alden soils are in depressions and along drainageways. Also included are well drained Paxton soils, moderately well drained Woodbridge soils, and poorly drained Ridgebury in areas with a dense substratum. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in cultivated cropland or woodland. Some areas, mostly in Fairfield County, are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large and small stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

49C—Georgia and Amenia silt loams, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: uplands, hills

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Georgia and similar soils: 50 percent

Amenia and similar soils: 35 percent

Minor components: 15 percent

Major Components

Georgia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; loam

Bw2—14 to 24 inches; loam

C—24 to 60 inches; gravelly fine sandy loam

Amenia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 9 inches; silt loam

Bw1—9 to 16 inches; silt loam

Bw2—16 to 25 inches; silt loam

C—25 to 60 inches; gravelly loam

Major Component Properties and Qualities

Georgia and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow to moderate

Available water capacity: high

Reaction: strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Amenia and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow to moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Georgia and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Amenia and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this unit in mapping are well drained Stockbridge and Nellis soils in higher areas of the landscape. Poorly drained Mudgepond soils and very poorly drained Alden soils are in depressions and along drainageways. Also included are well drained Paxton soils, moderately well drained Woodbridge soils, and poorly drained Ridgebury in areas with a dense substratum. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are in pasture. Other areas, mostly in Fairfield County, are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Slope and large and small stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

50A—Sutton fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: drainageways on uplands, depressions on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Sutton and similar soils: 80 percent

Minor components: 20 percent

Major Components

Sutton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

Bw1—6 to 12 inches; fine sandy loam

Bw2—12 to 24 inches; fine sandy loam

Bw3—24 to 28 inches; fine sandy loam

C1—28 to 36 inches; gravelly fine sandy loam

C2—36 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Paxton soils that are higher on the landscape. Canton soils are loamy over sandy, Charlton soils are sandy loam throughout, and Paxton soils have a dense substratum. Also included are small areas of poorly drained Leicester soils in depressions and drainageways. Small areas of moderately well drained Woodbridge soils are included in areas with a dense substratum. Some areas have a silt loam surface layer and subsoil. A few areas in New London County include well drained Narragansett soils and moderately well drained Rainbow soils. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, hay, pasture, or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

50B—Sutton fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: depressions on uplands, drainageways on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Sutton and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

Bw1—6 to 12 inches; fine sandy loam

Bw2—12 to 24 inches; fine sandy loam

Bw3—24 to 28 inches; fine sandy loam

C1—28 to 36 inches; gravelly fine sandy loam

C2—36 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Paxton soils that are higher on the landscape. Canton soils are loamy over sandy, Charlton soils are sandy loam throughout, and Paxton soils have a dense substratum. Also included are small areas of poorly drained Leicester soils in depressions and drainageways. Small areas of moderately well drained Woodbridge soils are included

in areas with a dense substratum. Some areas have a silt loam surface layer and subsoil. A few areas in New London County include well drained Narragansett soils and moderately well drained Rainbow soils. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, hay, pasture, or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

51B—Sutton fine sandy loam, 2 to 8 percent slopes, very stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: drainageways on uplands, depressions on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Sutton and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

Bw1—6 to 12 inches; fine sandy loam

Bw2—12 to 24 inches; fine sandy loam

Bw3—24 to 28 inches; fine sandy loam

C1—28 to 36 inches; gravelly fine sandy loam

C2—36 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Paxton soils that are higher on the landscape. Canton soils are loamy over sandy, Charlton soils are sandy loam throughout, and Paxton soils have a dense substratum. Also included are small areas of poorly drained Leicester soils in depressions and drainageways. Small areas of moderately well drained Woodbridge soils are included in areas with a dense substratum. Some areas have a silt loam surface layer and subsoil. A few areas in New London County include well drained Narragansett soils and moderately well drained Rainbow soils. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce this limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

52C—Sutton fine sandy loam, 2 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: nearly level to strongly sloping

Landscape: drainageways on uplands, depressions on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Sutton and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

Bw1—6 to 12 inches; fine sandy loam

Bw2—12 to 24 inches; fine sandy loam

Bw3—24 to 28 inches; fine sandy loam

C1—28 to 36 inches; gravelly fine sandy loam

C2—36 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Paxton soils that are higher on the landscape. Canton soils are loamy over sandy, Charlton soils are sandy loam throughout, and Paxton soils have a dense substratum. Also included are small areas of poorly drained Leicester soils in depressions and drainageways. Small areas of moderately well drained Woodbridge soils are included in areas with a dense substratum. Some areas have a silt loam surface layer and subsoil. A few areas in New London County include well drained Narragansett soils and moderately well drained Rainbow soils. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development and pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large stones and slope are also limitations for lawns and landscaping. Removing the stones and designing lawns to conform to the slope of the land will reduce these limitations. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

53A—Wapping very fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Wapping and similar soils: 80 percent

Minor components: 20 percent

Major Components

Wapping and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 11 inches; very fine sandy loam

Bw1—11 to 16 inches; very fine sandy loam

Bw2—16 to 20 inches; very fine sandy loam

2C1—20 to 28 inches; gravelly sandy loam

2C2—28 to 36 inches; gravelly loamy sand

2C3—36 to 80 inches; gravelly loamy sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Narragansett soils that are higher on the landscape. Poorly drained Leicester and Wilbraham soils, and very poorly drained Menlo soils are included in depressions and drainageways. Leicester soils do not have the dense substratum that Wilbraham and Menlo soils have. Also included are moderately well drained Watchaug and Ludlow soils, and well drained Cheshire soils in areas where the subsoil and substratum are red. Watchaug soils do not have the dense substratum that Ludlow soils have. Small areas of moderately deep, well drained Yalesville soils are included where the depth to bedrock is 20 to 40 inches below the surface. Small areas of soils with a stony surface are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in farmland or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

53B—Wapping very fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Wapping and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 11 inches; very fine sandy loam

Bw1—11 to 16 inches; very fine sandy loam

Bw2—16 to 20 inches; very fine sandy loam

2C1—20 to 28 inches; gravelly sandy loam

2C2—28 to 36 inches; gravelly loamy sand

2C3—36 to 80 inches; gravelly loamy sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Narragansett soils that are higher on the landscape. Poorly drained Leicester and Wilbraham soils, and very poorly drained Menlo soils are included in depressions and drainageways. Leicester soils do not have the dense substratum that Wilbraham and Menlo soils have. Also included are moderately well drained Watchaug and Ludlow soils, and well drained Cheshire soils in areas where the subsoil and substratum are red. Watchaug soils do not have the dense substratum that Ludlow soils have. Small areas of moderately deep, well drained Yalesville soils are included where the depth to bedrock is 20 to 40 inches below the surface. Small areas of soils with a stony surface are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in farmland or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

54B—Wapping very fine sandy loam, 2 to 8 percent slopes, very stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: till plains on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Wapping and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 11 inches; very fine sandy loam

Bw1—11 to 16 inches; very fine sandy loam

Bw2—16 to 20 inches; very fine sandy loam

2C1—20 to 28 inches; gravelly sandy loam

2C2—28 to 36 inches; gravelly loamy sand

2C3—36 to 80 inches; gravelly loamy sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Narragansett soils that are higher on the landscape. Poorly drained Leicester and Wilbraham soils, and very poorly drained Menlo soils are included in depressions and drainageways. Leicester soils do not have the dense substratum that Wilbraham and Menlo soils have. Also included are moderately well drained Watchaug and Ludlow soils, and well drained Cheshire soils in areas where the subsoil and substratum are red. Watchaug soils do not have the dense substratum that Ludlow soils have. Small areas of moderately deep, well drained Yalesville soils are included where the depth to bedrock is 20 to 40 inches below the surface. Small areas of soils with a non-stony surface are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

55A—Watchaug fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Watchaug and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 18 inches; fine sandy loam

Bw2—18 to 24 inches; fine sandy loam

C—24 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Cheshire soils that are higher on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Moderately well drained Ludlow soils are included in areas with a dense substratum. A few areas in New Haven County have a silt loam or stony surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cropland, pasture, or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with footing or foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal in places.

Potential frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

55B—Watchaug fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping
Landscape: till plains on uplands, hills on uplands
Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Watchaug and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam
 Bw1—8 to 18 inches; fine sandy loam
 Bw2—18 to 24 inches; fine sandy loam
 C—24 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Cheshire soils that are higher on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Moderately well drained Ludlow soils are included in areas with a dense substratum. A few areas in New Haven County have a silt loam or stony surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cropland, pasture, or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with footing or foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal in places.

Potential frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

56B—Watchaug fine sandy loam, 2 to 8 percent slopes, very stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: till plains on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Watchaug and similar soils: 80 percent

Minor components: 20 percent

Major Components

Watchaug and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 18 inches; fine sandy loam
 Bw2—18 to 24 inches; fine sandy loam
 C—24 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale
Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Cheshire soils that are higher on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Moderately well drained Ludlow soils are included in areas with a dense substratum. A few areas in New Haven County have a silt loam or stony surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Locating dwellings on the highest part of the unit with footing or foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal in places.

Potential frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

57B—Gloucester gravelly sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping
Landscape: hills on uplands
Size of map unit: Areas commonly range from 5 to 70 acres.

Map Unit Composition

Gloucester and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; gravelly sandy loam

Bw1—4 to 12 inches; gravelly sandy loam

Bw2—12 to 25 inches; very gravelly loamy sand

C1—25 to 35 inches; very gravelly loamy coarse sand

C2—35 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils formed in stratified glacial outwash. Well drained Canton, Charlton, and Paxton soils are included in areas of finer textured soils. Paxton soils have a dense substratum. Areas of moderately well drained Sutton soils are in slightly lower areas of the landscape, and poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, pasture, or woodland. Some areas are in community development.

Large stones are the main limitation for dwellings with basements. Removing the stones will reduce the limitation. Droughtiness and small stones are the main limitation for lawns and landscaping. Removing the stones will reduce this limitation. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas of Gloucester soils.

Large stones are the main limitation for local roads and streets. Removing the stones will reduce the limitation.

57C—Gloucester gravelly sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands

Size of map unit: Areas commonly range from 5 to 70 acres.

Map Unit Composition

Gloucester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; gravelly sandy loam

Bw1—4 to 12 inches; gravelly sandy loam

Bw2—12 to 25 inches; very gravelly loamy sand

C1—25 to 35 inches; very gravelly loamy coarse sand

C2—35 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils formed in stratified glacial outwash. Well drained Canton, Charlton, and Paxton soils are included in areas of finer textured soils. Paxton soils have a dense substratum. Areas of moderately well drained Sutton soils are in slightly lower areas, and poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, pasture, or woodland. Some areas are in community development.

Large stones are the main limitation for dwellings with basements. Removing the stones and designing dwellings to conform to the slope of the land will reduce these limitations. Erosion is a hazard during construction.

Slope, droughtiness, and small stones are the main limitation for lawns and landscaping. Removing the stones will reduce this limitation. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas of Gloucester soils.

Slope and large stones are the main limitation for local roads and streets. Constructing roads on the contour and removing the stones will reduce the limitations.

57D—Gloucester gravelly sandy loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands

Size of map unit: Areas commonly range from 5 to 70 acres.

Map Unit Composition

Gloucester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; gravelly sandy loam

Bw1—4 to 12 inches; gravelly sandy loam

Bw2—12 to 25 inches; very gravelly loamy sand

C1—25 to 35 inches; very gravelly loamy coarse sand

C2—35 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils formed in stratified glacial outwash. Well drained Canton, Charlton, and Paxton soils are included in areas of finer textured soils. Paxton soils have a dense substratum. Areas of moderately well drained Sutton soils are in slightly lower areas, and poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in hay and pasture or woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Designing dwellings to conform to the slope of the land will reduce this limitation. Erosion is a severe hazard during construction.

Slope and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable

substratum does not adequately filter effluent. Specially designed systems are necessary in some areas of Gloucester soils.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the limitation.

58B—Gloucester gravelly sandy loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 5 to 70 acres.

Map Unit Composition

Gloucester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; gravelly sandy loam

Bw1—4 to 12 inches; gravelly sandy loam

Bw2—12 to 25 inches; very gravelly loamy sand

C1—25 to 35 inches; very gravelly loamy coarse sand

C2—35 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils formed in stratified glacial outwash. Well drained Canton, Charlton, and Paxton soils are included in areas of finer textured soils. Paxton soils have a dense substratum. Areas of moderately well drained Sutton soils are in slightly lower areas, and poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture or woodland. Some areas are in community development.

Large stones are the main limitation for dwellings with basements. Removing the stones will reduce the limitation. Droughtiness and small stones are the main limitation for lawns and landscaping. Removing the stones will reduce this limitation. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas of Gloucester soils.

Large stones are the main limitation for local roads and streets. Removing the stones will reduce the limitation.

58C—Gloucester gravelly sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 5 to 70 acres.

Map Unit Composition

Gloucester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; gravelly sandy loam

Bw1—4 to 12 inches; gravelly sandy loam

Bw2—12 to 25 inches; very gravelly loamy sand

C1—25 to 35 inches; very gravelly loamy coarse sand

C2—35 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils formed in stratified glacial outwash. Well drained Canton, Charlton, and Paxton soils are included in areas of finer textured soils. Paxton soils have a dense substratum. Areas of moderately well drained Sutton soils are in slightly lower areas, and poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or community development.

Large stones are the main limitation for dwellings with basements. Removing the stones and designing dwellings to conform to the slope of the land will reduce these limitations. Erosion is a hazard during construction.

Slope, droughtiness, and small stones are the main limitation for lawns and landscaping. Removing the stones will reduce this limitation. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas of Gloucester soils.

Slope and large stones are the main limitation for local roads and streets. Constructing roads on the contour and removing the stones will reduce the limitations.

59C—Gloucester gravelly sandy loam, 3 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 5 to 70 acres.

Map Unit Composition

Gloucester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; gravelly sandy loam

Bw1—4 to 12 inches; gravelly sandy loam

Bw2—12 to 25 inches; very gravelly loamy sand

C1—25 to 35 inches; very gravelly loamy coarse sand

C2—35 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils formed in stratified glacial outwash. Well drained Canton, Charlton, and Paxton soils are included in areas of finer textured soils. Paxton soils have a dense substratum. Areas of moderately well drained Sutton soils are in slightly lower areas, and poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or community development.

Large stones are the main limitation for dwellings with basements. Removing the stones and designing dwellings to conform to the slope of the land will reduce these limitations. Erosion is a moderate hazard during construction.

Slope, droughtiness, and small stones are the main limitation for lawns and landscaping. Removing the stones will reduce this limitation. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas of Gloucester soils.

Slope and large stones are the main limitation for local roads and streets. Constructing roads on the contour and removing the stones will reduce the limitations.

59D—Gloucester gravelly sandy loam, 15 to 35 percent slopes, extremely stony

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 5 to 70 acres.

Map Unit Composition

Gloucester and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; gravelly sandy loam

Bw1—4 to 12 inches; gravelly sandy loam

Bw2—12 to 25 inches; very gravelly loamy sand

C1—25 to 35 inches; very gravelly loamy coarse sand

C2—35 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: A

Minor Components

Included with this soil in mapping are areas of excessively drained Hinckley soils formed in stratified glacial outwash. Well drained Canton, Charlton, and Paxton soils are included in areas of finer textured soils. Paxton soils have a dense substratum. Areas of moderately well drained Sutton soils are in slightly lower areas, and poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Designing dwellings to conform to the slope of the land will reduce this limitation. Erosion is a severe hazard during construction.

Slope and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed systems are necessary in some areas of Gloucester soils.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the limitation.

60B—Canton and Charlton soils, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Canton and similar soils: 45 percent

Charlton and similar soils: 35 percent

Minor components: 20 percent

Major Components

Canton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; gravelly fine sandy loam

Bw1—3 to 15 inches; gravelly loam

Bw2—15 to 24 inches; gravelly loam

Bw3—24 to 30 inches; gravelly loam

2C—30 to 60 inches; very gravelly loamy sand

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Canton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: high

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Canton and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Charlton and similar soils*Land capability classification (non-irrigated): 2e**Hydrologic group: B***Minor Components**

Included with these soils in mapping are areas of moderately well drained Sutton soils in slight depressions on the landscape, and poorly drained Leicester soils in depressions and drainageways. Also included are areas of moderately deep, somewhat excessively drained and well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Shallow, somewhat excessively drained and well drained Hollis soils are in small areas where bedrock is 10 to 20 inches below the surface. A few areas in Litchfield County include soils with a silt loam surface and subsoil. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in cultivated crops, pasture, residential development, or woodland.

This unit has few limitations for dwellings with basements. Large stones are a limitation for lawns and landscaping in areas of Canton soils. Removing the stones will reduce the limitation.

Charlton soils have few limitations for septic tank absorption fields. Poor filtering is the main limitation for septic tank absorption fields in areas of Canton soils. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places. This unit has few limitations for local roads and streets.

60C—Canton and Charlton soils, 8 to 15 percent slopes**Map Unit Setting***Slope:* strongly sloping*Landscape:* hills on uplands*Size of map unit:* Areas commonly range from 3 to 100 acres.**Map Unit Composition***Canton and similar soils:* 45 percent*Charlton and similar soils:* 35 percent*Minor components:* 20 percent**Major Components****Canton and similar soils**

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; gravelly fine sandy loam

Bw1—3 to 15 inches; gravelly loam

Bw2—15 to 24 inches; gravelly loam

Bw3—24 to 30 inches; gravelly loam

2C—30 to 60 inches; very gravelly loamy sand

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Canton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid or very rapid

Available water capacity: high

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Canton and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Charlton and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils in slight depressions on the landscape, and poorly drained Leicester soils in depressions and drainageways. Also included are areas of moderately deep, somewhat excessively drained and well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Shallow, somewhat excessively drained and well drained Hollis soils are in small areas where bedrock is 10 to 20 inches below the surface. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in cultivated crops, pasture, residential development, or woodland.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Large stones are a limitation for lawns and landscaping in areas of Canton soils. Removing the stones will reduce the limitation.

Slope is the main limitation for septic tank absorption fields in areas of Charlton soils. Poor filtering is the main limitation for septic tank absorption fields in areas of Canton soils. There is a hazard of groundwater pollution because the rapidly

permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas of Canton soils.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

60D—Canton and Charlton soils, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Canton and similar soils: 45 percent

Charlton and similar soils: 35 percent

Minor components: 20 percent

Major Components

Canton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; gravelly fine sandy loam

Bw1—3 to 15 inches; gravelly loam

Bw2—15 to 24 inches; gravelly loam

Bw3—24 to 30 inches; gravelly loam

2C—30 to 60 inches; very gravelly loamy sand

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Canton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: high

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate to moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Canton and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Charlton and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils in slight depressions on the landscape, and poorly drained Leicester soils in depressions and drainageways. Also included are areas of moderately deep, somewhat excessively drained and well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Shallow, somewhat excessively drained and well drained Hollis soils are in small areas where bedrock is 10 to 20 inches below the surface. A few areas in Litchfield County include soils with a silt loam surface and subsoil. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or residential development.

Slope is the main limitations for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas of Canton soils.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

61B—Canton and Charlton soils, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Canton and similar soils: 45 percent

Charlton and similar soils: 35 percent

Minor components: 20 percent

Major Components

Canton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; gravelly fine sandy loam

Bw1—3 to 15 inches; gravelly loam

Bw2—15 to 24 inches; gravelly loam

Bw3—24 to 30 inches; gravelly loam

2C—30 to 60 inches; very gravelly loamy sand

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap— 0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Canton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: high

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Canton and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Charlton and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils in slight depressions on the landscape, and poorly drained Leicester soils in depressions and drainageways. Also included are areas of moderately deep, somewhat excessively drained and well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Shallow, somewhat excessively drained and well drained Hollis soils are in small areas where bedrock is 10 to 20 inches below the surface. A few areas in Litchfield County include soils with a silt loam surface and subsoil. Minor inclusions make up about 20 percent of the map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture.

This unit has few limitations for dwellings with basements. Large stones are a limitation for lawns and landscaping in areas of Canton soils. Removing the stones will reduce the limitation.

Charlton soils have few limitations for septic tank absorption fields. Poor filtering is the main limitation for septic tank absorption fields in areas of Canton soils. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

This unit has few limitations for local roads and streets.

61C—Canton and Charlton soils, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Canton and similar soils: 45 percent

Charlton and similar soils: 35 percent

Minor components: 20 percent

Major Components

Canton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; gravelly fine sandy loam

Bw1—3 to 15 inches; gravelly loam

Bw2—15 to 24 inches; gravelly loam

Bw3—24 to 30 inches; gravelly loam

2C—30 to 60 inches; very gravelly loamy sand

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam
 Bw3—19 to 27 inches; gravelly fine sandy loam
 C—27 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Canton and similar soils

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy over sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss
Permeability: moderately rapid to very rapid
Available water capacity: high
Reaction: extremely acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Charlton and similar soils

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss
Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Canton and similar soils

Land capability classification (non-irrigated): 6s
Hydrologic group: B

Charlton and similar soils

Land capability classification (non-irrigated): 6s
Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils in slight depressions on the landscape, and poorly drained Leicester soils in depressions and drainageways. Also included are areas of moderately deep, somewhat excessively drained and well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Shallow, somewhat excessively drained and well drained Hollis soils are in small areas where bedrock is 10 to 20 inches below the surface. Minor components make up about 20 percent of the mapping unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or residential development.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Large stones are a limitation for lawns and landscaping in areas of Canton soils. Removing the stones will reduce the limitation.

Slope is the main limitation for septic tank absorption fields in areas of Charlton soils. Poor filtering is the main limitation for septic tank absorption fields in areas of

Canton soils. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas of Canton soils.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

62C—Canton and Charlton soils, 3 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Canton and similar soils: 45 percent

Charlton and similar soils: 35 percent

Minor components: 20 percent

Major Components

Canton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; gravelly fine sandy loam

Bw1—3 to 15 inches; gravelly loam

Bw2—15 to 24 inches; gravelly loam

Bw3—24 to 30 inches; gravelly loam

2C—30 to 60 inches; very gravelly loamy sand

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap— 0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Canton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: high

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Charlton and similar soils*Depth to bedrock:* very deep*Drainage class:* well drained*Parent material:* coarse-loamy melt-out till derived from granite and/or schist and/or gneiss*Permeability:* moderate or moderately rapid*Available water capacity:* high*Reaction:* very strongly acid to moderately acid*Depth to restrictive feature:* greater than 72 inches*Depth to seasonal water table:* greater than 6 feet*Flooding:* none***Interpretative Groups*****Canton and similar soils***Land capability classification (non-irrigated):* 7s*Hydrologic group:* B**Charlton and similar soils***Land capability classification (non-irrigated):* 7s*Hydrologic group:* B***Minor Components***

Included with these soils in mapping are areas of moderately well drained Sutton soils in slight depressions on the landscape, and poorly drained Leicester soils in depressions and drainageways. Also included are areas of moderately deep, somewhat excessively drained and well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Shallow, somewhat excessively drained and well drained Hollis soils are in small areas where bedrock is 10 to 20 inches below the surface. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Large stones are a limitation for lawns and landscaping in areas of Canton soils. Removing the stones will reduce the limitation.

Slope is the main limitation for septic tank absorption fields in areas of Charlton soils. Poor filtering is the main limitation for septic tank absorption fields in areas of Canton soils. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas of Canton soils.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

62D—Canton and Charlton soils, 15 to 35 percent slopes, extremely stony***Map Unit Setting****Slope:* moderately steep or steep*Landscape:* hills on uplands*Surface cover:* 3 to 15 percent stones*Size of map unit:* Areas commonly range from 3 to 100 acres.

Map Unit Composition

Canton and similar soils: 45 percent

Charlton and similar soils: 35 percent

Minor components: 20 percent

Major Components

Canton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; gravelly fine sandy loam

Bw1—3 to 15 inches; gravelly loam

Bw2—15 to 24 inches; gravelly loam

Bw3—24 to 30 inches; gravelly loam

2C—30 to 60 inches; very gravelly loamy sand

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Canton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy over sandy and gravelly melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: high

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Canton and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Charlton and similar soils*Land capability classification (non-irrigated): 7s**Hydrologic group: B****Minor Components***

Included with these soils in mapping are areas of moderately well drained Sutton soils in slight depressions on the landscape, and poorly drained Leicester soils in depressions and drainageways. Also included are areas of moderately deep, somewhat excessively drained and well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Shallow, somewhat excessively drained and well drained Hollis soils are in small areas where bedrock is 10 to 20 inches below the surface. A few areas in Litchfield County include soils with a silt loam surface and subsoil. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Slope is the main limitations for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas of Canton soils.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

63B—Cheshire fine sandy loam, 3 to 8 percent slopes***Map Unit Setting****Slope: gently sloping**Landscape: till plains on uplands, hills on uplands**Size of map unit: Areas commonly range from 3 to 100 acres.****Map Unit Composition****Cheshire and similar soils: 80 percent**Minor components: 20 percent****Major Components***

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

C—26 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities*Depth to bedrock: very deep**Drainage class: well drained**Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale**Permeability: moderate or moderately rapid**Available water capacity: high*

Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Watchaug soils in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Also included are areas of well drained Wethersfield and Yalesville soils. Wethersfield soils have a dense substratum; Yalesville soils have bedrock between 20 and 40 inches below the surface. Some soils with slopes less than 3 percent and soils with strong brown subsoil are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, residential development, or woodland. Some areas are in pasture.

This soil has few limitations for dwellings with basements, lawns and landscaping, septic tank absorption fields, or local roads and streets.

63C—Cheshire fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping
Landscape: till plains on uplands, hills on uplands
Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Cheshire and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam
 Bw1—8 to 16 inches; fine sandy loam
 Bw2—16 to 26 inches; fine sandy loam
 C—26 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale
Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Watchaug soils in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Also included are areas of well drained Wethersfield and Yalesville soils. Wethersfield soils have a dense substratum; Yalesville soils have bedrock between 20 and 40 inches below the surface. Some soils with slopes less than 3 percent and soils with strong brown subsoil are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture or cultivated cropland.

Slope is the main limitation for dwellings with basements, septic tank absorption fields, local roads, streets, and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Placing the septic tank absorption field distribution lines on the contour increases the efficiency of the system. Construction roads on the contour will reduce the slope limitation.

63D—Cheshire fine sandy loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Cheshire and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

C—26 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Watchaug soils in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Also included are areas of well drained Wethersfield and Yalesville soils. Wethersfield soils have a dense substratum; Yalesville soils have bedrock between 20 and 40 inches below the surface. Some soils with slopes less than 3 percent and soils with strong brown subsoil are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture or cultivated cropland.

Slope is the main limitation for dwellings with basements, septic tank absorption fields, local roads, streets, and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Placing the septic tank absorption field distribution lines on the contour increases the efficiency of the system. Construction roads on the contour will reduce the slope limitation.

64B—Cheshire fine sandy loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping
Landscape: till plains on uplands, hills on uplands
Surface cover: 0 to 3 percent stones
Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Cheshire and similar soils: 80 percent
Minor components: 20 percent

Major Components

Cheshire and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam
 Bw1—8 to 16 inches; fine sandy loam
 Bw2—16 to 26 inches; fine sandy loam
 C—26 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale
Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Watchaug soils in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Also included are areas of well drained Wethersfield and Yalesville soils. Wethersfield soils have a dense substratum; Yalesville soils have bedrock between 20 and 40 inches below the surface. Some soils with slopes less than 3 percent and soils with strong brown subsoil are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture or woodland. Some areas are in residential development.

This soil has few limitations for dwellings with basements, septic tank absorption fields, or local roads and streets.

Large stones are the main limitation for lawns and landscaping. Removing the stones will reduce the limitation.

64C—Cheshire fine sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping
Landscape: till plains on uplands, hills on uplands
Surface cover: 0 to 3 percent stones
Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Cheshire and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam
 Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam
 C—26 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Watchaug soils in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Also included are areas of well drained Wethersfield and Yalesville soils. Wethersfield soils have a dense substratum; Yalesville soils have bedrock between 20 and 40 inches below the surface. Some soils with slopes less than 3 percent and soils with strong brown subsoil are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture.

Slope is the main limitation for dwellings with basements, septic tank absorption fields, local roads, streets, and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Placing the septic tank absorption field distribution lines on the contour increases the efficiency of the system. Construction roads on the contour will reduce the slope limitation.

65C—Cheshire fine sandy loam, 3 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: till plains on uplands, hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Cheshire and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap— 0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

C—26 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Watchaug soils in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Also included are areas of well drained Wethersfield and Yalesville soils. Wethersfield soils have a dense substratum; Yalesville soils have bedrock between 20 and 40 inches below the surface. Some soils with slopes less than 3 percent and soils with strong brown subsoil are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture.

Slope is the main limitation for dwellings with basements, septic tank absorption fields, local roads, streets, and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Placing the septic tank absorption field distribution lines on the contour increases the efficiency of the system. Construction roads on the contour will reduce the slope limitation.

65D—Cheshire fine sandy loam, 15 to 35 percent slopes, extremely stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: till plains on uplands, hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Cheshire and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

C—26 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Watchaug soils in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are in depressions and drainageways. Also included are areas of well drained Wethersfield and Yalesville soils. Wethersfield soils have a dense substratum; Yalesville soils have bedrock between 20 and 40 inches below the surface. Some soils with slopes less than 3 percent and soils with strong brown subsoil are also included. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture.

Slope is the main limitation for dwellings with basements, septic tank absorption fields, local roads, streets, and lawns and landscaping. Erosion is a severe hazard during construction. A site should be selected on a less sloping portion of the unit or nearby soil. Construction roads on the contour will reduce the slope limitation.

66B—Narragansett silt loam, 2 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Narragansett and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook, Canton, and Charlton soils. Broadbrook soils have a dense substratum. Canton soils are coarser textured and Charlton soils are loamy throughout. Also included are small areas of moderately well drained Wapping and Sutton soils in slight depressions lower on the landscape. Wapping soils are silty over a sandy substratum; Sutton soils are loamy throughout. Poorly drained Leicester soils are in depressions and drainageways. A few areas in northeast Hartford County include red substrata. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are cropland or pasture. Some areas are in community development or woodland.

This soil has few limitations for dwellings with basements or lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas of Narragansett soils.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

66C—Narragansett silt loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Narragansett and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook, Canton, and Charlton soils. Broadbrook soils have a dense substratum. Canton soils are coarser textured and Charlton soils are loamy throughout. Also included are small areas of moderately well drained Wapping and Sutton soils in slight depressions lower on the landscape. Wapping soils are silty over a sandy substratum; Sutton soils are loamy throughout. Poorly drained Leicester soils are in depressions and drainageways. A few areas in northeast Hartford County include red substrata. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are cropland or pasture. Some areas are in community development or woodland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas of Narragansett soils.

Slope and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth and constructing roads on the contour will reduce these limitations.

67B—Narragansett silt loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: till plains on uplands, hills on uplands

Surface cover: 0 to 3 percent boulders

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Narragansett and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate or very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook, Canton, and Charlton soils. Broadbrook soils have a dense substratum. Canton soils are coarser textured and Charlton soils are loamy throughout. Also included are small areas of moderately well drained Wapping and Sutton soils in slight depressions

lower on the landscape. Wapping soils are silty over a sandy substratum; Sutton soils are loamy throughout. Poorly drained Leicester soils are in depressions and drainageways. A few areas in northeast Hartford County include red substrata. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

This soil has few limitations for dwellings with basements or lawns and landscaping. Large stones are the main limitation for lawns and landscaping. Removing the large stones will reduce the limitation.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas of Narragansett soils.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

67C—Narragansett silt loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: till plains on uplands, hills on uplands

Surface cover: 0 to 3 percent boulders

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Narragansett and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate or very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook, Canton, and Charlton soils. Broadbrook soils have a dense substratum. Canton soils are coarser textured and Charlton soils are loamy throughout. Also included are small areas of moderately well drained Wapping and Sutton soils in slight depressions lower on the landscape. Wapping soils are silty over a sandy substratum; Sutton soils are loamy throughout. Poorly drained Leicester soils are in depressions and drainageways. A few areas in northeast Hartford County include red substrata. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas of Narragansett soils.

Slope and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth and constructing roads on the contour will reduce these limitations.

68C—Narragansett silt loam, 3 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: hills on uplands, till plains on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Narragansett and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook, Canton, and Charlton soils. Broadbrook soils have a dense substratum. Canton soils are coarser textured and Charlton soils are loamy throughout. Also included are small areas of moderately well drained Wapping and Sutton soils in slight depressions lower on the landscape. Wapping soils are silty over a sandy substratum; Sutton soils are loamy throughout. Poorly drained Leicester soils are in depressions and drainageways. A few areas in northeast Hartford County include red substrata. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the large stones will reduce the limitation. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas of Narragansett soils.

Slope and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth and constructing roads on the contour will reduce these limitations.

68D—Narragansett silt loam, 15 to 25 percent slopes, extremely stony

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands, till plains on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Narragansett and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook, Canton, and Charlton soils. Broadbrook soils have a dense substratum. Canton soils are coarser textured and Charlton soils are loamy throughout. Also included are small areas of moderately well drained Wapping and Sutton soils in slight depressions lower on the landscape. Wapping soils are silty over a sandy substratum; Sutton soils are loamy throughout. Poorly drained Leicester soils are in depressions and drainageways. A few areas in northeast Hartford County include red substrata. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas of Narragansett soils.

Slope and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth and constructing roads on the contour will reduce these limitations.

69B—Yalesville fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: bedrock controlled hills on uplands, bedrock controlled ridges on uplands

Size of map unit: Areas commonly range from 3 to 80 acres.

Map Unit Composition

Yalesville and similar soils: 75 percent

Minor components: 25 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; loam

C—25 to 36 inches; gravelly sandy loam

2R—36 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of shallow well drained Holyoke soils in areas where the bedrock is between 10 to 20 inches deep. Very deep, well drained Cheshire soils are included where the bedrock is deeper than 60 inches. Also included are very deep, well drained Wethersfield soils in areas where the substratum is very firm. Moderately well drained Watchaug and Ludlow soils are included in slightly lower areas. Poorly drained Wilbraham soils are included in depressions and along drainageways. Also included are areas with a silt loam surface texture in Middlesex and New Haven counties. Areas with slopes less than 3 percent are included in New Haven County. Minor components make up about 25 percent of the unit.

Use and Management

Most areas are in cultivated cropland, nursery crops, orchards, or woodland. Some areas are in community development.

Depth to bedrock is the main limitation for dwellings with basements and septic tank absorption fields. The short uneven slopes and variable depth to bedrock reduce

site selection options. Where possible, dwellings with basements and septic tank absorption fields should be constructed in very deep inclusions. This soil has few limitations for lawns and landscaping.

Depth to bedrock is the main limitation for local roads and streets. Careful planning of road locations will avoid some removal of rock.

69C—Yalesville fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: bedrock-controlled ridges on uplands, bedrock controlled hills on uplands

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Yalesville and similar soils: 75 percent

Minor components: 25 percent

Major Components

Yalesville and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; loam

C—25 to 36 inches; gravelly sandy loam

2R—36 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Depth to bedrock: moderately deep or deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land Capability Classification (non-irrigated): 3e

Hydrologic Group: C

Minor Components

Included with this soil in mapping are areas of shallow well drained Holyoke soils in areas where the bedrock is between 10 to 20 inches deep. Very deep, well drained Cheshire soils are included where the bedrock is deeper than 60 inches. Also included are very deep, well drained Wethersfield soils in areas where the substratum is very firm. Moderately well drained Watchaug and Ludlow soils are included in slightly lower areas. Poorly drained Wilbraham soils are included in depressions and along drainageways. Also included are areas with a silt loam surface texture in Middlesex and New Haven counties. Minor components make up about 25 percent of the unit.

Use and Management

Most areas are in cultivated cropland, nursery crops, orchards, or woodland. Some areas are in community development.

Depth to bedrock is the main limitation for dwellings with basements and septic tank absorption fields. The short uneven slopes and variable depth to bedrock reduce site selection options. Where possible, dwellings with basements and septic tank absorption fields should be constructed in very deep inclusions. This soil has few limitations for lawns and landscaping.

Depth to bedrock is the main limitation for local roads and streets. Careful planning of road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

70C—Branford—Holyoke complex, 3 to 15 percent slopes, very rocky

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Branford and similar soils: 50 percent

Holyoke and similar soils: 30 percent

Minor components: 20 percent

Major Components

Branford and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; loam

Bw2—18 to 24 inches; gravelly loam

2C—24 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Holyoke and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; silt loam

Bw1—3 to 8 inches; silt loam

Bw2—8 to 18 inches; gravelly silt loam

2R—18 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Branford and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Holyoke and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy eolian deposits over melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate

Available water capacity: low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Branford and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Holyoke and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with this unit in mapping are areas of well drained Haven and Enfield soils, moderately well drained Ellington soils and poorly drained Raypol soils. Enfield soils are silty over sand and gravel and Haven soils are loamy over sand and gravel. Ellington soils are in slight depressions and broad drainageways and Raypol soils are in low depressions on the landscape. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Branford soils have few limitations for dwellings with basements. Shallow depth to bedrock in areas of Holyoke soils and frequent rock outcroppings are the main limitations for dwellings with basements and lawns and landscaping. Slope is also a limitation for lawns and landscaping in areas of Branford soil where the slope exceeds 8 percent. Erosion is a moderate hazard during construction. Dwellings can be built above the rock and landscaped with additional fill. Where possible, dwellings with basements should be constructed in inclusions of very deep Branford soils or a near soil.

Poor filtering is the main limitation for septic tank absorption fields in areas of Branford soils. Shallow depth to bedrock and frequent rock outcroppings are the main limitations in areas of Holyoke soils for septic tank absorption fields. There is a hazard of groundwater pollution in Holyoke and Branford soils because the substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas of Branford soils. A more suitable site should be selected in an inclusion of deeper, loamier soil or in a nearby soil.

Frost action is the main limitation for local roads and streets in areas of Branford soils. Providing a coarse textured subgrade will reduce this limitation. Depth to bedrock is the main limitation for local roads and streets in areas of Holyoke soils. Planning of grades and road locations will avoid some removal of rock in areas of Holyoke soils.

71C—Brookfield–Brimfield–Rock outcrop complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock-controlled hills on uplands, bedrock-controlled ridges on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Brookfield and similar soils: 45 percent

Brimfield and similar soils: 30 percent

Rock outcrop and similar soils: 15 percent

Minor components: 10 percent

Major Components

Brookfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; fine sandy loam

Bw1—3 to 13 inches; gravelly fine sandy loam

Bw2—13 to 27 inches; gravelly fine sandy loam

C—27 to 60 inches; gravelly sandy loam

Brimfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; fine sandy loam

Bw1—3 to 6 inches; fine sandy loam

Bw2—6 to 17 inches; gravelly fine sandy loam

2R—17 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Brookfield and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from mica schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Brimfield and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from mica schist

Permeability: moderate to rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Brookfield and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Brimfield and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton and Woodbridge soils and poorly drained Leicester soils. Sutton and Woodbridge soils are in slight depressions on the landscape, and Leicester soils are in depressions and drainageways. Woodbridge soils are in areas with a dense substratum. Also included are shallow, somewhat excessively drained Hollis soils where bedrock is 10 to 20 inches below the surface; Hollis soils are more yellow in the subsoil than Brimfield soils. Some areas of moderately deep, well drained Chatfield soils are included where bedrock is 20 to 40 inches below the surface. Areas of well drained Canton, Charlton, and Paxton soils are also included. These soils are more yellow in the subsoil than Brookfield soils, and Paxton soils have a dense substratum. Minor components make up 10 percent of the map unit.

Use and Management

This unit is mostly in woodland. Some areas are in pasture or community development.

The depth to bedrock in Brimfield soils and areas of Rock outcrop are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slope is also a limitation for dwellings with basements and septic tank absorption fields on the steeper portions of the unit. A pollution hazard exists for septic tank absorption fields in Brimfield soils because the soil is not thick enough to filter effluent. Large stones are a limitation for lawns and landscaping. Erosion is a moderate hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements and septic tank absorption fields should be constructed in very deep Brookfield soils.

Slope, frost action, and variable depth to bedrock are the main limitations for local roads and streets. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

71E—Brookfield–Brimfield–Rock outcrop complex, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep or steep

Landscape: bedrock-controlled hills on uplands, bedrock-controlled ridges on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Brookfield and similar soils: 45 percent

Brimfield and similar soils: 30 percent

Rock outcrop and similar soils: 15 percent

Minor components: 10 percent

Major Components

Brookfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; fine sandy loam

Bw1—3 to 13 inches; gravelly fine sandy loam

Bw2—13 to 27 inches; gravelly fine sandy loam

C—27 to 60 inches; gravelly sandy loam

Brimfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; fine sandy loam

Bw1—3 to 6 inches; fine sandy loam

Bw2—6 to 17 inches; gravelly fine sandy loam

2R—17 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Brookfield and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from mica schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Brimfield and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from mica schist

Permeability: moderate to rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Brookfield and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Brimfield and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton and Woodbridge soils and poorly drained Leicester soils. Sutton and Woodbridge soils are in slight depressions on the landscape, and Leicester soils are in depressions and drainageways. Woodbridge soils are in areas with a dense substratum. Also included are shallow, somewhat excessively drained Hollis soils where bedrock is 10 to 20 inches below the surface; Hollis soils are more yellow in the subsoil than Brimfield soils. Some areas of moderately deep, well drained Chatfield soils are included where bedrock is 20 to 40 inches below the surface. Areas of well drained Canton, Charlton, and Paxton soils are also included. These soils are more yellow in the subsoil than Brookfield soils, and Paxton soils have a dense substratum. Minor components make up about 10 percent of the map unit.

Use and Management

This unit is mostly in woodland.

Steep slope and the depth to bedrock in Brimfield soils and areas of Rock outcrop are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. A pollution hazard exists for septic tank absorption fields in Brimfield soils because the soil is not thick enough to filter effluent. Erosion is a severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection.

Steep slope and variable depth to bedrock are the main limitations for local roads and streets. Constructing roads on the contour or locating them on less sloping inclinations will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

73C—Charlton–Chatfield complex, 3 to 15 percent slopes, very rocky

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock-controlled hills, bedrock-controlled uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Charlton and similar soils: 45 percent

Chatfield and similar soils: 30 percent

Minor components: 25 percent

Major Components

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities**Charlton and similar soils**

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Chatfield and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups**Charlton and similar soils**

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Chatfield and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils and poorly drained Leicester soils. Sutton soils are in slight depressions in the landscape; Leicester soils are in depressions and drainageways. Also included are small areas of shallow, somewhat excessively drained Hollis soils where bedrock is 10 to 20 inches below the surface. A few areas in Litchfield County have a yellowish red surface layer and subsoil. Other areas in Litchfield County include sandier soils over bedrock. Minor components make up about 25 percent of the map unit.

Use and Management

Most areas are in woodland or residential development. Some areas are in pasture.

Depth to bedrock in areas of Chatfield soils and rock outcrops over portions of the unit are the main limitations for dwellings with basements. Slope is also a limitation. Erosion is a moderate hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in areas of very deep Charlton soils.

Slope is the main limitation for lawns and landscaping. Large stones are a limitation in areas of Charlton soils, and the thin soil layer is a limitation in areas of Chatfield soils. Droughtiness can make establishment and maintenance of lawns difficult. Addition of fill material and removing the stones will reduce these limitations.

Slope is the main limitation for septic tank absorption fields in areas of Charlton soils. Depth to bedrock in areas of Chatfield soils, and rock outcrops over portions of the landscape are also limitations. Where possible, septic tank absorption fields should be constructed in areas of very deep Charlton soils. Placing distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Depths to bedrock and frost action are also limitations in areas of Chatfield soils. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

73E—Charlton–Chatfield complex, 15 to 45 percent slopes, very rocky

Map Unit Setting

Slope: moderately steep or steep

Landscape: bedrock-controlled hills, uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Charlton and similar soils: 45 percent

Chatfield and similar soils: 30 percent

Minor components: 25 percent

Major Components

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Chatfield and similar soils

Depth to bedrock: moderately deep or deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Charlton and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Chatfield and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils and poorly drained Leicester soils. Sutton soils are in slight depressions on the landscape; Leicester soils are in depressions and drainageways. Also included are small areas of shallow, somewhat excessively drained Hollis soils where bedrock is 10 to 20 inches below the surface. A few areas in Litchfield County have a yellowish red surface layer and subsoil. Other areas in Litchfield County include sandier soils over bedrock. Minor components make up about 25 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Slope and depth to bedrock in areas of Chatfield soils and Rock outcrops over portions of the unit are the main limitations for dwellings with basements. Erosion is a severe or very severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in areas of very deep Charlton soils, a less sloping inclusion, or nearby soil.

Slope is the main limitation for lawns and landscaping. A site should be selected on a less sloping portion of the unit or nearby soil.

Slope is the main limitation for septic tank absorption fields in areas of Charlton soils. Depth to bedrock in areas of Chatfield soils and Rock outcrops over portions of the landscape are also limitations. Where possible, septic tank absorption fields should be constructed in areas of very deep Charlton soils on a less sloping portion of the unit. Placing distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

74C—Narragansett-Hollis complex, 3 to 15 percent slopes, very rocky

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: hills on uplands, till plains on uplands

Surface cover: 0 to 3 percent boulders

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Narragansett and similar soils: 55 percent

Hollis and similar soils: 20 percent

Minor components: 25 percent

Major Components

Narragansett and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Hollis and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 9 inches; channery fine sandy loam

Bw2—9 to 15 inches; gravelly fine sandy loam

2R—15 to 80 inches;

Major Component Properties and Qualities

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Chatfield soils in areas where the depth to bedrock is 20 to 40 inches below the surface. Also included are some areas of well drained Canton and Charlton soils. Canton soils are coarser textured; Charlton soils are loamy throughout. Areas of moderately well drained Wapping and Sutton soils are included in slight depressions on the landscape. Wapping soils are silty over a sandy substratum; and Sutton soils are loamy throughout. Poorly drained Leicester soils are in depressions and drainageways. Minor components make up about 25 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Shallow depth to bedrock and many Rock outcrops in areas of Hollis soils are the main limitations for dwellings with basements and lawns and landscaping. The short, uneven slopes are also a limitation. Erosion is a moderate hazard during construction. Dwellings can be built above the rock and landscaped with additional fill. Droughtiness can make establishment and maintenance of lawns difficult. Where possible, dwellings with basements should be constructed in areas of very deep Narragansett soils.

Poor filtering, shallow depth to bedrock, and Rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent, or the soil is not thick enough to filter effluent. Specially designed septic systems are necessary in some areas of Narragansett soils. A more suitable site should be selected in a deeper, loamier inclusion or nearby soil.

Shallow depth to bedrock and many Rock outcrops are the main limitations for local roads and streets. Slope and frost action are also limitations. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the

contour will reduce the slope limitation. Providing a coarse grained subgrade will reduce frost action.

75C—Hollis-Chatfield-Rock outcrop complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: uplands, bedrock controlled hills, bedrock controlled ridges

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Hollis and similar soils: 35 percent

Chatfield and similar soils: 30 percent

Rock outcrop and similar soils: 15 percent (fig. 11)

Minor components: 20 percent

Major Components

Hollis and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 9 inches; channery fine sandy loam

Bw2—9 to 15 inches; gravelly fine sandy loam

2R—15 to 80 inches



Figure 11. —Rock outcrop in an area of Hollis-Chatfield-Rock outcrop complex.

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities**Hollis and similar soils**

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Chatfield and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups**Hollis and similar soils**

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Chatfield and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils and poorly drained Leicester soils. Sutton soils are in slight depressions and Leicester soils are in depressions and drainageways. Also included are very deep, well drained Charlton soils adjacent to Hollis and Chatfield soils. Areas of shallow, somewhat excessively drained Brimfield soils are included where the underlying bedrock is micaceous schist at a depth of 10 to 20 inches below the surface. A few areas in Litchfield County have a yellowish red surface layer and subsoil. Other areas in Litchfield County include sandier soils above bedrock. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Shallow depth to bedrock and many Rock outcrops in areas of Hollis soils are the main limitations for dwellings with basements and lawns and landscaping. The short, uneven slopes are also a limitation. Erosion is a moderate to severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper, less sloping inclusion or nearby soil. Droughtiness can make establishment and maintenance of lawns difficult in areas of Chatfield soils. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Shallow depth to bedrock and many Rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent, or the soil is not thick enough to filter effluent. A more suitable site should be selected in a deeper inclusion or nearby soil.

Shallow depth to bedrock and many Rock outcrops are the main limitations for local roads and streets. Frost action is also a limitation for Chatfield soils. Slope is a limitation in steeper areas at Chatfield. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation. Providing a coarse grained subgrade will reduce frost action.

75E—Hollis-Chatfield-Rock outcrop complex, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep or steep

Landscape: bedrock controlled hills, bedrock controlled ridges, uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Hollis and similar soils: 35 percent

Chatfield and similar soils: 30 percent

Rock outcrop and similar soils: 15 percent

Minor components: 20 percent

Major Components

Hollis and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 9 inches; channery fine sandy loam

Bw2—9 to 15 inches; gravelly fine sandy loam

2R—15 to 80 inches

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Hollis and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Chatfield and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Hollis and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Chatfield and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of moderately well drained Sutton soils and poorly drained Leicester soils. Sutton soils are in slight depressions and Leicester soils are in depressions and drainageways. Also included are very deep, well drained Charlton soils adjacent to Hollis and Chatfield soils. Areas of shallow, somewhat excessively drained Brimfield soils are included where the underlying bedrock is micaceous schist at a depth of 10 to 20 inches below the surface. A few areas in Litchfield County have a yellowish red surface layer and subsoil. Other areas in Litchfield County include sandier soils above bedrock. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Shallow depth to bedrock in areas of Hollis soils, and many Rock outcrops are also limitations. Erosion is a very severe hazard during construction. Additional fill will reduce the depth limitation. A more suitable site should be selected on a less sloping, deeper portion of the unit or nearby soil.

Slope, shallow depth to bedrock, and many Rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because

the soil is not thick enough to filter effluent. A more suitable site should be selected in a less sloping, deeper inclusion or nearby soil.

Slope and many rock outcrops are the main limitations for local roads and streets. Shallow depth to bedrock is also a limitation in areas of Hollis soils. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

76E—Rock outcrop-Hollis complex, 3 to 45 percent slopes

Map Unit Setting

Slope: gently sloping to steep

Landscape: bedrock-controlled hills, bedrock-controlled ridges, bedrock-controlled uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Rock outcrop and similar soils: 55 percent

Hollis and similar soils: 25 percent

Minor components: 20 percent

Major Components

Hollis and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 9 inches; channery fine sandy loam

Bw2—9 to 15 inches; gravelly fine sandy loam

2R—15 to 80 inches

Major Component Properties and Qualities

Hollis and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Hollis and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Also included are very deep, well drained Charlton soils adjacent to Hollis soils and moderately well drained Sutton soils in slight depressions. Poorly drained Leicester soils are included in depressions and drainageways. Areas of shallow, somewhat excessively drained

Brimfield soils are included where the underlying bedrock is micaceous schist at a depth of 10 to 20 inches below the surface. Minor components make up 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Many Rock outcrops and shallow depth to bedrock are the main limitations for dwellings with basements and lawns and landscaping. Slope is also a main limitation in steeper areas of the unit. Uneven slopes and variable depth to bedrock reduce site selection. Erosion is a severe to very severe hazard during construction. Dwellings can be built above the rock and landscaped in with additional fill to reduce the depth limitation. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. A more suitable site should be selected in a deeper, less sloping inclusion or nearby soil.

Shallow depth to bedrock and many Rock outcrops are the main limitations for septic tank absorption fields. Slope is also a main limitation in steeper areas. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. Placing septic tank absorption field distribution lines on the contour increases the efficiency of the system. A more suitable site should be selected in a less sloping, deeper inclusion or nearby soil.

Shallow depth to bedrock and many Rock outcrops are the main limitations for local roads and streets. Slope is also a limitation in steeper areas. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

76F—Rock outcrop-Hollis complex, 45 to 60 percent slopes

Map Unit Setting

Slope: very steep

Landscape: bedrock-controlled uplands, bedrock-controlled ridges, bedrock-controlled hills

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Rock outcrop and similar soils: 55 percent

Hollis and similar soils: 25 percent

Minor components: 20 percent

Major Components

Hollis and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 9 inches; channery fine sandy loam

Bw2—9 to 15 inches; gravelly fine sandy loam

2R—15 to 80 inches

Major Component Properties and Qualities

Hollis and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Hollis and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Chatfield soils where bedrock is 20 to 40 inches below the surface. Also included are very deep, well drained Charlton soils adjacent to Hollis soils. Poorly drained Leicester soils are included in depressions and drainageways. Areas of shallow, somewhat excessively drained Brimfield soils are included where the underlying bedrock is micaceous schist at a depth of 10 to 20 inches below the surface. Minor components make up 20 percent of the map unit.

Use and Management

Most areas are in woodland.

Many rock outcrops, slope, and shallow depth to bedrock are the main limitations for dwellings with basements and lawns and landscaping. Erosion is a very severe hazard during construction. Addition of fill will reduce the depth limitation. A more suitable site should be selected in a deeper, less sloping inclusion or nearby soil.

Many Rock outcrops, slope, and shallow depth to bedrock are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a less sloping, deeper inclusion or nearby soil.

Shallow depth to bedrock, slope, and many Rock outcrops are the main limitations for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

77C—Cheshire-Holyoke complex, 3 to 15 percent slopes, very rocky

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: hills on uplands, till plains on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 125 acres.

Map Unit Composition

Cheshire and similar soils: 45 percent

Holyoke and similar soils: 35 percent

Minor components: 20 percent

Major Components

Cheshire and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

- Ap—0 to 8 inches; fine sandy loam
- Bw1—8 to 16 inches; fine sandy loam
- Bw2—16 to 26 inches; fine sandy loam
- C—26 to 65 inches; gravelly sandy loam

Holyoke and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

- Oe—0 to 1 inch; moderately decomposed plant material
- A—1 to 3 inches; silt loam
- Bw1—3 to 8 inches; silt loam
- Bw2—8 to 18 inches; gravelly silt loam
- 2R—18 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Cheshire and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Holyoke and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy eolian deposits over melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate

Available water capacity: low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Cheshire and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Holyoke and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Yalesville soils where bedrock is 20 to 40 inches below the surface. In places, the Yalesville soils may make up as much as 20 percent of the unit. Also included are well

drained Wethersfield soils in areas with a dense substratum. Moderately well drained Watchaug soils are in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are included in drainageways and depressions.

Use and Management

Most areas are in woodland. Some areas are in residential development, pasture, or orchards.

Shallow depth to bedrock in areas of Holyoke soils, and Rock outcrops over portions of the unit are the main limitations for dwellings with basements. Slope is a limitation in steeper areas of Cheshire soil. Erosion is a moderate to severe hazard during construction. Dwellings can be built above the rock and landscaped in with additional fill to reduce the depth limitation. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in inclusions of very deep Cheshire soils.

Shallow depth to bedrock in areas of Holyoke soils, and Rock outcrops over portions of the unit are the main limitation for lawns and landscaping. Large stones are the main limitation in areas of Cheshire soils. Addition of fill material will reduce the depth limitation. Removing the stones will reduce the stone limitation. Slope is also a limitation in the steeper Cheshire soils.

Shallow depth to bedrock in areas of Holyoke soils, and Rock outcrops over portions of the unit are the main limitations for septic tank absorption fields. Slope is also a limitation in the steeper Cheshire soils. A pollution hazard exists because the Holyoke soil is not thick enough to filter effluent. Placing the distribution lines on the contour increases the efficiency of the system. Where possible, septic tank absorption fields should be constructed in areas of very deep Cheshire soils.

Shallow depth to bedrock in areas of Holyoke soils, and Rock outcrops over portions of the unit are the main limitation for local roads and streets. Slope is also a limitation in the steeper Cheshire soils. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

77D—Cheshire-Holyoke complex, 15 to 35 percent slopes, very rocky

Map Unit Setting

Slope: moderately steep to steep

Landscape: till plains on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 125 acres.

Map Unit Composition

Cheshire and similar soils: 45 percent

Holyoke and similar soils: 35 percent

Minor components: 20 percent

Major Components

Cheshire and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam
 C—26 to 65 inches; gravelly sandy loam

Holyoke and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material
 A—1 to 3 inches; silt loam
 Bw1—3 to 8 inches; silt loam
 Bw2—8 to 18 inches; gravelly silt loam
 2R—18 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Cheshire and similar soils

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale
Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Holyoke and similar soils

Depth to bedrock: shallow to moderately deep
Drainage class: well drained
Parent material: loamy eolian deposits over melt-out till derived from basalt and/or sandstone and shale
Permeability: moderate
Available water capacity: low
Reaction: extremely acid to moderately acid
Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Cheshire and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: B

Holyoke and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Yalesville soils where bedrock is 20 to 40 inches below the surface. In places, the Yalesville soils may make up as much as 20 percent of the unit. Also included are well drained Wethersfield soils in areas with a dense substratum. Moderately well drained Watchaug soils are in slight depressions on the landscape. Poorly drained Wilbraham soils and very poorly drained Menlo soils are included in drainageways and depressions. Minor components make up about 20 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in residential development, pasture, or orchards.

Slope is the main limitations for dwellings with basements and lawns and landscaping. Shallow depth to bedrock in areas of Holyoke soils, and Rock outcrops over portions of the unit are also limitations. Erosion is a very severe hazard during construction. Addition of fill will reduce the depth limitation. A site should be selected in less sloping, very deep Cheshire soils or nearby soil.

Slope is the main limitation for septic tank absorption fields. Shallow depth to bedrock in areas of Holyoke soils, and Rock outcrops over portions of the unit are also limitations. There is the hazard of groundwater pollution because the Holyoke soil is not thick enough to filter effluent. Placing distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, very deep Cheshire soil or nearby soil.

Slope is the main limitation for local roads and streets. Shallow depth to bedrock in areas of Holyoke soils, and Rock outcrops over portions of the unit are also limitations. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

78C—Holyoke-Rock outcrop complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock-controlled hills on uplands, bedrock-controlled ridges on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Holyoke and similar soils: 50 percent

Rock outcrop and similar soils: 25 percent

Minor components: 25 percent

Major Components

Holyoke and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; silt loam

Bw1—3 to 8 inches; silt loam

Bw2—8 to 18 inches; gravelly silt loam

2R—18 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Holyoke and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy eolian deposits over melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate

Available water capacity: low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Holyoke and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Yalesville soils where bedrock is 20 to 40 inches below the surface. Also included are very deep Wethersfield and Cheshire soils. Moderately well drained Ludlow and Watchaug soils are in slight depressions and along drainageways. In New Haven County, soils with a fine sandy loam surface and subsoil are included. Minor components make up about 25 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Shallow depth to bedrock and frequent rock outcroppings are the main limitations for dwellings with basements and lawns and landscaping. Erosion is a moderate to severe hazard during construction. Dwellings can be built above the rock and landscaped in with additional fill to reduce the depth limitation. Addition of fill materials will reduce the depth limitation for lawns and landscaping. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper inclusions or a nearby soil.

Shallow depth to bedrock and frequent rock outcroppings are the main limitations for septic tank absorption fields. A pollution hazard exists because the soil is not thick enough to filter effluent. Where possible, septic tank absorption fields should be constructed in deeper inclusion or nearby soil.

Shallow depth to bedrock and frequent rock outcroppings are the main limitations for local roads and streets. Careful planning of grades and road locations will avoid some removal of rock.

78E—Holyoke-Rock outcrop complex, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep or steep

Landscape: bedrock-controlled hills on uplands, bedrock-controlled ridges on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Holyoke and similar soils: 50 percent

Rock outcrop and similar soils: 25 percent

Minor components: 25 percent

Major Components

Holyoke and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material
 A—1 to 3 inches; silt loam
 Bw1—3 to 8 inches; silt loam
 Bw2—8 to 18 inches; gravelly silt loam
 2R—18 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Holyoke and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy eolian deposits over melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate

Available water capacity: low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Holyoke and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Yalesville soils where bedrock is 20 to 40 inches below the surface. Also included are very deep Wethersfield and Cheshire soils. Very poorly drained Menlo soils are included along drainageways. Some areas have slopes up to 90 percent or less than 15 percent. In New Haven County, soils with a fine sandy loam surface and subsoil are included. Minor components make up about 25 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Slope, shallow to bedrock, and rock outcroppings are the main limitations for dwellings with basements and lawns and landscaping. Erosion is a very severe hazard during construction. Addition of fill will reduce the depth limitation. A site should be selected in deeper, less sloping inclusion or nearby soil.

Slope, shallow to bedrock, and rock outcropping are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be considered in a deeper, less sloping inclusion or nearby soil.

Slope, shallow depth to bedrock, and rock outcroppings are the main limitations for local roads and streets. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

79E—Rock outcrop-Holyoke complex, 3 to 45 percent slopes

Map Unit Setting

Slope: gently sloping to steep

Landscape: bedrock-controlled uplands, bedrock-controlled ridges, bedrock-controlled hills

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Rock outcrop and similar soils: 55 percent

Holyoke and similar soils: 25 percent

Minor components: 20 percent

Major Components

Holyoke and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; silt loam

Bw1—3 to 8 inches; silt loam

Bw2—8 to 18 inches; gravelly silt loam

2R—18 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Holyoke and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy eolian deposits over melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate

Available water capacity: low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Holyoke and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately deep, well drained Yalesville soils where bedrock is 20 to 40 inches below the surface. Also included are very deep Wethersfield and Cheshire soils. Very poorly drained Menlo soils are included along drainageways. Some areas have slopes up to 90 percent. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland.

Shallow to bedrock and frequent rock outcroppings are the main limitations for dwellings with basements and lawns and landscaping. Slope is also a main limitation in steeper areas. Erosion is a severe to very severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Addition of fill will reduce the depth limitation. A site should be selected in deeper, less sloping inclusion or nearby soil.

Shallow to bedrock and frequent rock outcropping are the main limitations for septic tank absorption fields. Slope is also a limitation in steeper areas. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be considered in a deeper, less sloping inclusion or nearby soil.

Shallow depth to bedrock and frequent rock outcroppings are the main limitations for local roads and streets. Slope is also a limitation in steeper areas. Careful

planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour or locating them in less sloping inclinations will reduce the slope limitation.

80B—Bernardston silt loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Bernardston and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; channery silt loam

Bw2—14 to 24 inches; channery silt loam

BC—24 to 26 inches; channery silt loam

Cd—26 to 60 inches; channery silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from phyllite and/or schist

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 24 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with this soil are areas of well drained Paxton and Lanesboro soils, moderately well drained Woodbridge and Fullam soils, and poorly drained Brayton soils. Paxton and Woodbridge soils are included where the soil is less silty. Lanesboro, Fullam, and Brayton soils are included where the mean soil temperature is less than 48 degrees F. Also included are areas with slopes outside of the range. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, residential development, or woodland.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls and diverting

runoff from higher areas will reduce wetness. This soil has few limitations for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

80C—Bernardston silt loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Bernardston and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; channery silt loam

Bw2—14 to 24 inches; channery silt loam

BC—24 to 26 inches; channery silt loam

Cd—26 to 60 inches; channery silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from phyllite and/or schist

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 24 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Minor Components

Included with this soil are areas of well drained Paxton and Lanesboro soils, moderately well drained Woodbridge and Fullam soils, and poorly drained Brayton soils. Paxton and Woodbridge soils are included where the soil is less silty. Lanesboro, Fullam, and Brayton soils are included where the mean soil temperature is less than 48 degrees F. Also included are areas with slopes outside of the range. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, residential development, or woodland.

The seasonal high water table and slope are the main limitations for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls and diverting runoff from higher areas will reduce wetness. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Slope is the main limitation for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

81C—Bernardston silt loam, 3 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: uplands, hills

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Bernardston and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; silt loam

Bw1—8 to 14 inches; channery silt loam

Bw2—14 to 24 inches; channery silt loam

BC—24 to 26 inches; channery silt loam

Cd—26 to 60 inches; channery silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from phyllite and/or schist

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 24 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: C

Minor Components

Included with this soil are areas of well drained Paxton and Lanesboro soils, moderately well drained Woodbridge and Fullam soils, and poorly drained Brayton soils. Paxton and Woodbridge soils are included where the soil is less silty. Lanesboro, Fullam, and Brayton soils are included where the mean soil temperature is less than 48 degrees F. Also included are areas with slopes outside of the range. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland.

The seasonal high water table and slope are the main limitations for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls and diverting runoff from higher areas will reduce wetness. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Slope and large stones are the main limitations for lawns and landscaping. Removing the stones will reduce this limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

81D—Bernardston silt loam, 15 to 25 percent slopes, extremely stony

Map Unit Setting

Slope: moderately steep

Landscape: uplands, hills

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Bernardston and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; silt loam

Bw1—8 to 14 inches; channery silt loam

Bw2—14 to 24 inches; channery silt loam

BC—24 to 26 inches; channery silt loam

Cd—26 to 60 inches; channery silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from phyllite and/or schist

Permeability: very slow to moderate

Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 30 inches to densic material
Depth to seasonal water table: 18 to 24 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s
Hydrologic group: C

Minor Components

Included with this soil are areas of well drained Paxton and Lanesboro soils, moderately well drained Woodbridge and Fullam soils, and poorly drained Brayton soils. Paxton and Woodbridge soils are included where the soil is less silty. Lanesboro, Fullam, and Brayton soils are included where the mean soil temperature is less than 48 degrees F. Also included are areas with slopes outside of the range. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland.

Slope is the main limitation for dwellings with basements. Erosion is a severe hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation. A site should be selected on a less sloping portion of the unit or nearby soil.

Slope and large stones are the main limitations for lawns and landscaping.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less dense inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

82B—Broadbrook silt loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: till plains on uplands, hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Broadbrook and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; silt loam

Bw2—14 to 25 inches; silt loam

2Cd—25 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Rainbow soils in slight depressions in the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Areas of well drained Narragansett soils and Wethersfield soils are included. Narragansett soils lack a dense substratum; Wethersfield soils have a redder color in the subsoil. Small areas of shallow, well drained Holyoke soils are included where bedrock is 10 to 20 inches below the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, residential development, or woodland.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls and diverting runoff from higher areas will reduce wetness. This soil has few limitations for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

82C—Broadbrook silt loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: till plains on uplands, hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Broadbrook and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; silt loam

Bw2—14 to 25 inches; silt loam

2Cd—25 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Rainbow soils, poorly drained Wilbraham soils, and very poorly drained Menlo soils. Rainbow soils are in slightly lower areas. Wilbraham and Menlo soils in depressions and drainageways. Also included are small areas of Holyoke, Narragansett, and Wethersfield soils. Narragansett soils lack a dense substratum; Wethersfield soils have a redder color in the subsoil; Holyoke soils are shallow to bedrock. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, residential development, or woodland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. The seasonal high water table is also a limitation for dwellings with basements. Erosion is a moderate hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls and diverting runoff from higher areas will reduce wetness.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

82D—Broadbrook silt loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: drumlins on uplands, hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Broadbrook and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; silt loam

Bw2—14 to 25 inches; silt loam

2Cd—25 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Rainbow soils, poorly drained Wilbraham soils, and very poorly drained Menlo soils. Rainbow soils are in slightly lower areas. Wilbraham and Menlo soils in depressions and drainageways. Also included are small areas of Holyoke, Narragansett, and Wethersfield soils. Narragansett soils lack a dense substratum; Wethersfield soils have a redder color in the subsoil; Holyoke soils are shallow to bedrock. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. A more suitable site should be considered on a less sloping inclusion or nearby soil.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places. Placing distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation.

83B—Broadbrook silt loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands, till plains on uplands, drumlins on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Broadbrook and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; silt loam

Bw2—14 to 25 inches; silt loam

2Cd—25 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Rainbow soils in slight depressions on the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Areas of well drained Narragansett soils and Wethersfield soils are included. Narragansett soils lack a dense substratum; Wethersfield soils have a redder color in the subsoil. Small areas of shallow, well drained Holyoke soils are included where bedrock is 10 to 20 inches below the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls and diverting runoff from higher areas will reduce wetness.

Large stones are the main limitation for lawns and landscaping. Removing the large stones will reduce this limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

83C—Broadbrook silt loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: till plains on uplands, hills on uplands, drumlins on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Broadbrook and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; silt loam

Bw2—14 to 25 inches; silt loam

2Cd—25 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Rainbow soils, poorly drained Wilbraham soils, and very poorly drained Menlo soils. Rainbow soils are in slightly lower areas. Wilbraham and Menlo soils in depressions and drainageways. Also included are small areas of Holyoke, Narragansett, and Wethersfield soils. Narragansett soils lack a dense substratum; Wethersfield soils

have a redder color in the subsoil; Holyoke soils are shallow to bedrock. Minor Inclusions make up about 20 percent of this map unit.

Use and Management

Most areas are in residential development or woodland. Some areas are in pasture.

Slope is the main limitation for dwellings with basements and lawns and landscaping. The seasonal high water table is also a limitation for dwellings with basements. Erosion is a moderate hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls and diverting runoff from higher areas will reduce wetness.

Large stones are the main limitation for lawns and landscaping. Removing the large stones will reduce this limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

84B—Paxton and Montauk fine sandy loams, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands, till plains on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 85 acres.

Map Unit Composition

Paxton and similar soils: 55 percent

Montauk and similar soils: 30 percent

Minor components: 15 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Montauk and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; fine sandy loam

Bw1—4 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; sandy loam

2Cd1—25 to 39 inches; gravelly loamy coarse sand

2Cd2—39 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Paxton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Montauk and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from granite

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 38 inches to densic material

Depth to seasonal water table: 24 to 30 inches

Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Montauk and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of moderately well drained Woodbridge soils in slight depressions on the landscape. Also included are poorly drained Ridgebury soils in depressions and along drainageways. Well drained Canton and Charlton soils are included in areas lacking a dense substratum. Well drained Stockbridge soils are included in areas of Litchfield and Fairfield counties with carbonates below 40 inches. Also included are areas of nearly level soils and soils with a stony surface. A few areas in Hartford, Middlesex, and New Haven counties include soils with a red substratum. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in cultivated crops, pasture, or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Droughtiness is the main limitation for lawns and landscaping in areas of Montauk soils. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Slow percolation is the main limitation for septic tank absorption fields. The seasonal high water table is also a limitation in areas of Montauk soil. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

84C—Paxton and Montauk fine sandy loams, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: drumlins on uplands, hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Paxton and similar soils: 55 percent

Montauk and similar soils: 30 percent

Minor components: 15 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Montauk and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; fine sandy loam

Bw1—4 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; sandy loam

2Cd1—25 to 39 inches; gravelly loamy coarse sand

2Cd2—39 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Paxton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Montauk and similar soils*Depth to bedrock:* very deep*Drainage class:* well drained*Parent material:* coarse-loamy lodgment till derived from granite and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from granite*Permeability:* very slow to moderately rapid*Available water capacity:* moderate*Reaction:* very strongly acid to moderately acid*Depth to restrictive feature:* 20 to 38 inches to densic material*Depth to seasonal water table:* 24 to 30 inches*Flooding:* none***Interpretative Groups*****Paxton and similar soils***Land capability classification (non-irrigated):* 3e*Hydrologic group:* C**Montauk and similar soils***Land capability classification (non-irrigated):* 3e*Hydrologic group:* C***Minor Components***

Included with these soils in mapping are areas of moderately well drained Woodbridge soils in slight depressions on the landscape. Also included are poorly drained Ridgebury soils in depressions and along drainageways. Well drained Canton and Charlton soils are included in areas lacking a dense substratum. Well drained Stockbridge soils are included in some areas of Litchfield and Fairfield counties with free carbonates below 40 inches. Also included are soils with a stony surface. A few areas in Hartford, Middlesex, and New Haven counties include soils with a red substratum. Minor Components make up about 15 percent of the map unit.

Use and Management

Most areas are in cultivated crops, pasture, or woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements. Slope is also a limitation in areas of Paxton soil. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Slope is the main limitation for lawns and landscaping in areas of Montauk soils. Droughtiness is the main limitation in areas dominated by Montauk soils. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Slow percolation is the main limitation for septic tank absorption fields. The seasonal high water table is also a limitation in areas of Montauk soil. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation. Constructing roads on the contour will reduce the slope limitation.

84D—Paxton and Montauk fine sandy loams, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands, till plains on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Paxton and similar soils: 55 percent

Montauk and similar soils: 30 percent

Minor components: 15 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Montauk and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; fine sandy loam

Bw1—4 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; sandy loam

2Cd1—25 to 39 inches; gravelly loamy coarse sand

2Cd2—39 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Paxton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Montauk and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from granite

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 38 inches to densic material

Depth to seasonal water table: 24 to 30 inches

Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: C

Montauk and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of moderately well drained Woodbridge soils in slight depressions on the landscape. Also included are poorly drained Ridgebury soils in depressions and along drainageways. Well drained Canton and Charlton soils are included in areas lacking a dense substratum. Well drained Stockbridge soils are found in areas of Litchfield and Fairfield Counties with free carbonates below 40 inches. Also included are small areas with soils steeper than 25 percent slopes and soils with a stony surface. A few areas in Hartford, Middlesex, and New Haven counties include soils with a red substratum. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development, cultivated crops, or pasture.

Slope is the main limitation for dwellings with basements. The seasonal high water table is also a limitation in areas of Montauk soils. Erosion is a severe hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the slope of the land will reduce the slope limitation. A site should be selected on a less sloping portion of the unit or nearby soil.

Slope is the main limitation for lawns and landscaping. Droughtiness is also a limitation in areas dominated by Montauk soils. Planting early in the spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Slow percolation and slope are the main limitations for septic tank absorption fields. The seasonal high water table is also a limitation in areas of Montauk soil. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less dense inclusion or nearby soil.

Slope is the main limitations for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

85B—Paxton and Montauk fine sandy loams, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, till plains on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Paxton and similar soils: 55 percent

Montauk and similar soils: 30 percent

Minor components: 15 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Montauk and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; fine sandy loam

Bw1—4 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; sandy loam

2Cd1—25 to 39 inches; gravelly loamy coarse sand

2Cd2—39 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Paxton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Montauk and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from granite

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 38 inches to densic material

Depth to seasonal water table: 24 to 30 inches

Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Montauk and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of moderately well drained Woodbridge soils in slight depressions on the landscape. Also included are poorly drained Ridgebury soils in depressions and along drainageways. Well drained Canton and Charlton soils are included in areas lacking a dense substratum. Well drained Stockbridge soils are included in areas of Litchfield and Fairfield counties with free carbonates below 40 inches. Also included are areas of nearly level soils and soils with a stony surface. A few areas in Hartford, Middlesex, and New Haven counties include soils with a red substratum. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Small and large stones are the main limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. The seasonal high water table is also a limitation in areas of Montauk soil. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

85C—Paxton and Montauk fine sandy loams, 8 to 15 percent slopes, very stony**Map Unit Setting**

Slope: strongly sloping

Landscape: till plains on uplands, drumlins on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Paxton and similar soils: 55 percent

Montauk and similar soils: 30 percent

Minor components: 15 percent

Major Components**Paxton and similar soils**

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam
 Bw2—15 to 26 inches; fine sandy loam
 Cd—26 to 65 inches; gravelly fine sandy loam

Montauk and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; fine sandy loam
 Bw1—4 to 14 inches; fine sandy loam
 Bw2—14 to 25 inches; sandy loam
 2Cd1—25 to 39 inches; gravelly loamy coarse sand
 2Cd2—39 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Paxton and similar soils

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss
Permeability: very slow to moderate
Available water capacity: moderate
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: 20 to 40 inches to densic material
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Montauk and similar soils

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy lodgment till derived from granite and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from granite
Permeability: very slow to moderately rapid
Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 38 inches to densic material
Depth to seasonal water table: 24 to 30 inches
Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 6s
Hydrologic group: C

Montauk and similar soils

Land capability classification (non-irrigated): 6s
Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of moderately well drained Woodbridge soils in slight depressions on the landscape. Also included are poorly drained Ridgebury soils in depressions and along drainageways. Well drained Canton and Charlton soils are included in areas lacking a dense substratum. Well drained Stockbridge soils are included in areas of Litchfield and Fairfield counties in soils that have free carbonate below 40 inches. Also included are soils with a stony surface. A

few areas in Hartford, Middlesex, and New Haven counties include soils with a red substratum. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements. Slope is also a limitation in areas of Paxton soil. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Large stones and slope are the main limitations for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. The seasonal high water table is also a limitation in areas of Montauk soil. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation. Constructing roads on the contour will reduce the slope limitation.

86C—Paxton and Montauk fine sandy loams, 3 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: drumlins on uplands, hills on uplands, till plains on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Paxton and similar soils: 55 percent

Montauk and similar soils: 30 percent

Minor components: 15 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Montauk and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; fine sandy loam

Bw1—4 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; sandy loam

2Cd1—25 to 39 inches; gravelly loamy coarse sand

2Cd2—39 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Paxton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Montauk and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from granite

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 38 inches to densic material

Depth to seasonal water table: 24 to 30 inches

Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: C

Montauk and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of moderately well drained Woodbridge soils in slight depressions on the landscape. Also included are poorly drained Ridgebury soils in depressions and along drainageways. Well drained Canton and Charlton soils are included in areas lacking a dense substratum. Well drained Stockbridge soils are included in litchfield and Fairfield counties in areas with free carbonates below 40 inches. Also included are soils with a stony surface. A few areas in Hartford, Middlesex, and New Haven counties include soils with a red substratum. Minor inclusions make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements. Slope is also a limitation in areas of Paxton soil. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas

will reduce wetness. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Slope and large stones are the main limitations for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. The seasonal high water table is also a limitation in areas of Montauk soil. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation. Constructing roads on the contour will reduce the slope limitation.

86D—Paxton and Montauk fine sandy loams, 15 to 35 percent slopes, extremely stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: till plains on uplands, drumlins on uplands, hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Paxton and similar soils: 55 percent

Montauk and similar soils: 30 percent

Minor components: 15 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Montauk and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; fine sandy loam

Bw1—4 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; sandy loam

2Cd1—25 to 39 inches; gravelly loamy coarse sand

2Cd2—39 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Paxton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: 20 to 40 inches to densic material
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Montauk and similar soils

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy lodgment till derived from granite and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from gneiss and/or coarse-loamy lodgment till derived from granite
Permeability: very slow to moderately rapid
Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 38 inches to densic material
Depth to seasonal water table: 24 to 30 inches
Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: C

Montauk and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of moderately well drained Woodbridge soils in slight depressions on the landscape. Also included are poorly drained Ridgebury soils in depressions and along drainageways. Well drained Canton and Charlton soils are included in areas lacking a dense substratum. Well drained Stockbridge Soils are included in litchfield and Fairfield counties in areas with free carbonates below 40 inches. Also included are soils with less stones on the surface. A few areas in Hartford, Middlesex, and New Haven counties include soils with a red substratum. Minor componets make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

Slope is the main limitation for dwellings with basements. Erosion is a severe hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation. A site should be selected on a less sloping portion of the unit or nearby soil.

Slope is the main limitation for lawns and landscaping.

Slow percolation and slope are the main limitations for septic tank absorption fields. The seasonal high water table is also a limitation in areas of Montauk soil. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less dense inclusion or nearby soil.

Slope is the main limitations for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

87B—Wethersfield loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Wethersfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Ludlow soils in slight depressions on the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Well drained Cheshire soils are included in areas that lack a dense substratum. Small areas of moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches from the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are in vegetables, orchards (fig. 12), nursery stock, or woodland.

The seasonal high water table is the main limitation for dwellings with basements. This soil has few limitations for lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.



Figure 12.—Wethersfield loam (foreground) is well suited for orchards.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce this limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

87C—Wethersfield loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Wethersfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Ludlow soils in slight depressions on the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Well drained Cheshire soils are included in areas that lack a dense substratum. Small areas of moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches from the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are in vegetables, orchards, nursery stock, or woodland.

The seasonal high water table is the main limitation for dwellings with basements. Slope is also a main limitation for dwellings with basements and for lawns and landscaping. Erosion is a moderate hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce this limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation and constructing roads on the contour will reduce the slope limitation.

87D—Wethersfield loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Wethersfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Ludlow soils in slight depressions on the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Well drained Cheshire soils are included in areas that lack a dense substratum. Small areas of moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches from the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are in vegetables, orchards, nursery stock, or woodland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal. Placing distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

88B—Wethersfield loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Wethersfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Ludlow soils in slight depressions on the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Well drained Cheshire soils are included in areas that lack a dense substratum. Small areas of moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches from the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are in vegetables, orchards, nursery stock, or woodland.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Large stones are a limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce this limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

88C—Wethersfield loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands, drumlins on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Wethersfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Ludlow soils in slight depressions on the landscape. Also included are poorly drained

Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Well drained Cheshire soils are included in areas that lack a dense substratum. Small areas of moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches from the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in community development or cultivated crops, hay, or pasture. Some areas are in vegetables, orchards, nursery stock, or woodland.

The seasonal high water table and slope are the main limitations for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope and large stones are limitations for lawns and landscaping. Removing the large stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce this limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation and constructing roads on the contour will reduce the slope limitation.

89C—Wethersfield loam, 3 to 15 percent slopes, extremely stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 90 acres.

Map Unit Composition

Wethersfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Ludlow soils in slight depressions on the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Well drained Cheshire soils are included in areas that lack a dense substratum. Small areas of moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches from the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are cleared and used for community development, hay, or pasture. Some areas are in orchards, nursery stock, or woodland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Slope is a limitation in the steeper areas. Erosion is a moderate hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Large stones are limitations for lawns and landscaping. Removing the large stones will reduce the limitation. Slope is a limitation in the steeper areas.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal.

The seasonal high water table and frost action are the main limitations for local roads and streets. Slope is a limitation in the steeper areas. Construction on raised fill materials and installing a drainage system will reduce this limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation and constructing roads on the contour will reduce the slope limitation.

89D—Wethersfield loam, 15 to 35 percent slopes, extremely stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: drumlins on uplands, hills on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 90 acres.

Map Unit Composition

Wethersfield and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of moderately well drained Ludlow soils in slight depressions on the landscape. Also included are poorly drained Wilbraham soils and very poorly drained Menlo soils in depressions and drainageways. Well drained Cheshire soils are included in areas that lack a dense substratum. Small areas of moderately deep, well drained Yalesville soils are included where bedrock is 20 to 40 inches from the surface. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are cleared and used for community development or pasture. Some areas are in orchards, nursery stock, or woodland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe to very severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. A site should be selected on a less sloping portion of the unit or nearby soil.

Slope and large stones are also limitations for lawns and landscaping. Removing the large stones will reduce the limitation.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal. Placing the distribution lines on the contour will increase the efficiency of the system and reduces the slope limitation. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

The slope is the main limitation for local roads and streets. Constructing roads on the contour or locating them on less sloping inclinations will reduce the slope limitation.

90B—Stockbridge loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Stockbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow to moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Georgia soils are in slightly lower areas, Mudgepond and Alden soils are in depressions and along drainageways. Also included are Nellis soils in areas where the soil is calcareous within 40 inches. Areas of shallow well drained Farmington soils are included where the underlying limestone bedrock is between 10 and 20 inches below the surface. Paxton soils are included where the substratum is denser and more acid. Also included in Litchfield County are some areas with slopes less than 3 percent. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in cropland or pasture. Other areas are in woodland or community development.

This soil has few limitations for dwellings with basements and lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

90C—Stockbridge loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Stockbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Georgia soils are in slightly lower areas, Mudgepond and Alden soils are in depressions and along drainageways. Also included are Nellis soils in areas where the soil is calcareous within 40 inches. Areas of shallow well drained Farmington soils are included where the underlying limestone bedrock is between 10 and 20 inches below

the surface. Paxton soils are included where the substratum is denser and more acid. Also included in Litchfield County are some areas with slopes less than 3 percent. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in cropland or pasture. Other areas are in woodland or community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation (fig. 13).

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action and slope are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation. Constructing roads on the contour will reduce the slope limitation.

90D—Stockbridge loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Stockbridge and similar soils: 80 percent

Minor components: 20 percent



Figure 13.—Stripcropping reduces soil erosion on steeper Stockbridge soils (background).

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

- Ap—0 to 10 inches; loam
- Bw1—10 to 20 inches; loam
- Bw2—20 to 28 inches; loam
- C1—28 to 42 inches; gravelly loam
- C2—42 to 48 inches; gravelly loam
- C3—48 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Georgia soils are in slightly lower areas, Mudgepond and Alden soils are in depressions and along drainageways. Also included are Nellis soils in areas where the soil is calcareous within 40 inches. Areas of shallow well drained Farmington soils are included where the underlying limestone bedrock is between 10 and 20 inches below the surface. Paxton soils are included where the substratum is denser and more acid. Also included in Litchfield County are some areas with slopes less than 3 percent. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in hay, woodland, or pasture. Other areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Frost action and slope are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation.

91B—Stockbridge loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Stockbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Georgia soils are in slightly lower areas, Mudgepond and Alden soils are in depressions and along drainageways. Also included are Nellis soils in areas where the soil is calcareous within 40 inches. Areas of shallow well drained Farmington soils are included where the underlying limestone bedrock is between 10 and 20 inches below the surface. Paxton soils are included where the substratum is denser and more acid. Also included in Litchfield County are some areas with slopes less than 3 percent. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland or pasture. Other areas are in community development.

This soil has few limitations for dwellings with basements and lawns and landscaping. Large stones are the main limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

91C—Stockbridge loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Stockbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Georgia soils are in slightly lower areas, Mudgepond and Alden soils are in depressions and along drainageways. Also included are Nellis soils in areas where the soil is calcareous within 40 inches. Areas of shallow well drained Farmington soils are included where the underlying limestone bedrock is between 10 and 20 inches below

the surface. Paxton soils are included where the substratum is denser and more acid. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland or pasture. Other areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce this limitation. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action and slope are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation. Constructing roads on the contour will reduce the slope limitation.

91D—Stockbridge loam, 15 to 35 percent slopes, very stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Stockbridge and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Georgia soils are in slightly lower areas, Mudgepond and Alden soils are in depressions and along drainageways. Also included are Nellis soils in areas where the soil is calcareous within 40 inches. Areas of shallow well drained Farmington soils are included where the underlying limestone bedrock is between 10 and 20 inches below the surface. Paxton soils are included where the substratum is denser and more acid. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland. Other areas are in pasture or community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation.

92B—Nellis fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Nellis and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; fine sandy loam

BC—25 to 27 inches; loam

C—27 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of well drained Farmington soils, well drained Stockbridge soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Farmington soils are in areas underlain by limestone bedrock at depths of 10 to 20 inches and Stockbridge soils are in areas where the soil lack carbonates within 40 inches of the surface. Amenia and Georgia soils are in slightly lower areas. Mudgepond and Alden soils are in depressions and along drainageways. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are cleared and are in crops, pasture, or hay. Some areas are in residential development or woodland.

This unit has few limitations for dwellings with basements. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduced seedling mortality. Lawns need watering in the summer.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

92C—Nellis fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills, uplands

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Nellis and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; fine sandy loam

BC—25 to 27 inches; loam
 C—27 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy melt-out till derived from limestone and dolomite and/or schist
Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: moderately acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e
Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of well drained Farmington soils, well drained Stockbridge soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Farmington soils are in areas underlain by limestone bedrock at depths of 10 to 20 inches and Stockbridge soils are in areas where the soil lack carbonates within 40 inches of the surface. Amenia and Georgia soils are in slightly lower areas. Mudgepond and Alden soils are in depressions and along drainageways. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are cleared and are in crops, pasture, or hay. Some areas are in residential development or woodland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Droughtiness is also a limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduced seedling mortality. Lawns need watering in the summer.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action and slope are the main limitations for local roads and streets. Constructing roads on the contour will reduce the slope limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

92D—Nellis fine sandy loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep
Landscape: uplands, hills
Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Nellis and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; fine sandy loam

BC—25 to 27 inches; loam

C—27 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of well drained Farmington soils, well drained Stockbridge soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Farmington soils are in areas underlain by limestone bedrock at depths of 10 to 20 inches and Stockbridge soils are in areas where the soil lack carbonates within 40 inches of the surface. Amenia and Georgia soils are in slightly lower areas. Mudgepond and Alden soils are in depressions and along drainageways. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are cleared and are in crops, pasture, or hay. Some areas are in residential development or woodland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Droughtiness is also a limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduced seedling mortality. Lawns need watering in the summer.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action and slope are the main limitations for local roads and streets. Constructing roads on the contour will reduce the slope limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

93C—Nellis fine sandy loam, 3 to 15 percent slopes, very stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: uplands, hills

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Nellis and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; fine sandy loam

BC—25 to 27 inches; loam

C—27 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of well drained Farmington soils, well drained Stockbridge soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Farmington soils are in areas underlain by limestone bedrock at depths of 10 to 20 inches and Stockbridge soils are in areas where the soil lack carbonates within 40 inches of the surface. Amenia and Georgia soils are in slightly lower areas. Mudgepond and Alden soils are in depressions and along drainageways. Minor components make up about 15 percent of the unit.

Use and Management

Most areas in residential development or woodland. Some areas are in pasture.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Large

stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Slope is a limitation on steeper areas. Constructing roads on the contour will reduce the slope limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

94C—Farmington-Nellis complex, 3 to 15 percent slopes, very rocky

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock-controlled ridges on glaciated uplands, bedrock-controlled hills on glaciated uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Farmington and similar soils: 40 percent

Nellis and similar soils: 35 percent

Minor components: 25 percent

Major Components

Farmington and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 3 inches; fine sandy loam

Bw1—3 to 8 inches; fine sandy loam

Bw2—8 to 17 inches; fine sandy loam

2R—17 to 80 inches; bedrock

Nellis and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; fine sandy loam

BC—25 to 27 inches; loam

C—27 to 60 inches; sandy loam

Major Component Properties and Qualities

Farmington and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Nellis and similar soils*Depth to bedrock:* very deep*Drainage class:* well drained*Parent material:* coarse-loamy melt-out till derived from limestone and dolomite and/or schist*Permeability:* moderate or moderately rapid*Available water capacity:* high*Reaction:* moderately acid to moderately alkaline*Depth to restrictive feature:* greater than 72 inches*Depth to seasonal water table:* greater than 6 feet*Flooding:* none***Interpretative Groups*****Farmington and similar soils***Land capability classification (non-irrigated):* 6s*Hydrologic group:* D**Nellis and similar soils***Land capability classification (non-irrigated):* 6s*Hydrologic group:* B***Minor Components***

Included with this unit in mapping are areas of somewhat excessively drained, well drained Stockbridge soils, well drained Charlton soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Hollis soils are in areas where the underlying bedrock is less than 20 inches from the surface and is schist, granite, or gneiss. Charlton and Stockbridge soils are in areas where the soil is more acid. Amenia and Georgia soils are in slightly lower areas on the landscape. Mudgepond and Alden soils are in depressions and along drainageways. Soils with a silt loam surface are included in areas of Litchfield County and Farmington soils with slightly redder subsoils are included in areas of Fairfield County. Minor components make up about 25 percent of the unit.

Use and Management

Most areas in pasture or woodland. Some areas are in community development.

Shallow depths to bedrock and rock outcroppings in areas of Farmington soils are the main limitations for dwellings with basements. Erosion is a moderate hazard during construction. Slope is a limitation in steeper areas of Nellis soils. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Where possible, dwellings with basements should be constructed in areas of very deep, less sloping Nellis soils or nearby soils.

Droughtiness is the main limitation for lawns and landscaping in areas of Nellis soils. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer. Slope is also a limitation in the steeper areas of Nellis soils. Large rocks are also a limitation. Removing the rocks will reduce the limitation.

Slow percolation in areas of Nellis soils and shallow depth to bedrock and rock outcroppings in areas of Farmington soils are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the Farmington soils are not thick enough to filter effluent. A more suitable site should be considered in areas of very deep Nellis soils. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in areas of Nellis soils.

Shallow depths to bedrock and rock outcroppings are the main limitations for local roads and streets in areas of Farmington soils. Frost action is a main limitation for

Nellis soils. Slope is a limitation on steeper areas. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

94E—Farmington-Nellis complex, 15 to 35 percent slopes, very rocky

Map Unit Setting

Slope: moderately steep to steep

Landscape: bedrock-controlled ridges on glaciated uplands, bedrock-controlled hills on glaciated uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Farmington and similar soils: 40 percent

Nellis and similar soils: 35 percent

Minor components: 25 percent

Major Components

Farmington and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 3 inches; fine sandy loam

Bw1—3 to 8 inches; fine sandy loam

Bw2—8 to 17 inches; fine sandy loam

2R—17 to 80 inches; bedrock

Nellis and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; fine sandy loam

BC—25 to 27 inches; loam

C—27 to 60 inches; sandy loam

Major Component Properties and Qualities

Farmington and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Nellis and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Farmington and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Nellis and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of somewhat excessively drained Hollis, well drained Stockbridge soils, well drained Charlton soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Hollis soils are in areas where the underlying bedrock is less than 20 inches from the surface and is schist, granite, or gneiss. Charlton and Stockbridge soils are in areas where the soil is more acid. Amenia and Georgia soils are in slightly lower areas on the landscape. Mudgepond and Alden soils are in depressions and along drainageways. Soils with a silt loam surface are included in areas of Litchfield County and Farmington soils with slightly redder subsoils are included in areas of Fairfield County. Minor components make up about 25 percent of the unit.

Use and Management

Most areas in woodland. Some areas are in pasture or community development.

Slope and rock outcroppings are the main limitations for dwellings with basements and lawns and landscaping. Shallow to depth bedrock is also a limitation in areas of Farmington soils for dwellings with basements. Erosion is a severe to very severe hazard during construction. Where possible, dwellings with basements should be constructed in areas of very deep, less sloping Nellis soils or nearby soils. Large rocks are also a limitation for lawns and landscaping. Removing the rocks will reduce the limitation.

Slope and rock outcroppings are the main limitations for septic tank absorption fields. Shallow depth to bedrock in areas of Farmington soils and slow percolation in areas of Nellis soils are also limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the Farmington soils are not thick enough to filter effluent. A more suitable site should be considered in areas of very deep Nellis soils. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in areas of Nellis soils.

Slope is the main limitation for local roads and streets. Shallow depths to bedrock and rock outcroppings are also limitations for local roads and streets in areas of Farmington soils. Frost action is a main limitation for Nellis soils. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

95C—Farmington-Rock outcrop complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock controlled ridges on glaciated uplands, bedrock controlled hills on glaciated uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Farmington and similar soils: 60 percent

Rock outcrop and similar soils: 20 percent

Minor components: 20 percent

Major Components

Farmington and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 3 inches; fine sandy loam

Bw1—3 to 8 inches; fine sandy loam

Bw2—8 to 17 inches; fine sandy loam

2R—17 to 80 inches; bedrock

Major Component Properties and Qualities

Farmington and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Farmington and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with this unit in mapping are areas of somewhat excessively drained Hollis, well drained Stockbridge soils, well drained Nellis soils, well drained Charlton soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Hollis soils are in areas where the underlying bedrock is less than 20 inches from the surface and is schist, granite, or gneiss. Charlton, Nellis, and Stockbridge soils are in areas where the bedrock is deeper than 60 inches. Amenia and Georgia soils are in slightly lower areas on the landscape. Mudgepond and Alden soils are in depressions and along drainageways. Soils with slightly redder subsoils are included in areas of Fairfield County. Minor components make up about 20 percent of the unit.

Use and Management

Most areas in woodland. Some areas are in pasture or community development.

Shallow depths to bedrock and rock outcroppings are the main limitations for dwellings with basements and lawns and landscaping. Slope is also a limitation. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Where possible, dwellings with basements should be constructed in areas of very deeper inclusions or nearby soils.

Shallow depths to bedrock and rock outcroppings are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soils are not thick enough to filter effluent. A more suitable site should be considered in areas of deeper soils.

Shallow depths to bedrock and rock outcroppings are the main limitations for local roads and streets. Slope is a limitation on steeper areas. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

95E—Farmington-Rock outcrop complex, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep to steep

Landscape: bedrock-controlled ridges on glaciated uplands, bedrock-controlled hills on glaciated uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Farmington and similar soils: 60 percent

Rock outcrop and similar soils: 20 percent

Minor components: 20 percent

Major Components

Farmington and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 3 inches; fine sandy loam

Bw1—3 to 8 inches; fine sandy loam

Bw2—8 to 17 inches; fine sandy loam

2R—17 to 80 inches; bedrock

Major Component Properties and Qualities

Farmington and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: well drained

Parent material: loamy melt-out till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Farmington and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with this unit in mapping are areas of somewhat excessively drained Hollis, well drained Stockbridge soils, well drained Nellis soils, well drained Charlton soils, moderately well drained Amenia and Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Hollis soils are in areas where the underlying bedrock is less than 20 inches from the surface and is schist, granite, or gneiss. Charlton, Nellis, and Stockbridge soils are in areas where the bedrock is deeper than 60 inches. Amenia and Georgia soils are in slightly lower areas on the landscape. Mudgepond and Alden soils are in depressions and along drainageways. Soils with slightly redder subsoils are included in areas of Fairfield County. Minor components make up about 20 percent of the unit.

Use and Management

Most areas in woodland. Some areas are in community development.

Slope, shallow depths to bedrock, and rock outcroppings are the main limitations for dwellings with basements and lawns and landscaping. Erosion is a very severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Where possible, dwellings with basements should be constructed in areas of very deeper inclusions or nearby soils.

Slope, shallow depths to bedrock, and rock outcroppings are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soils are not thick enough to filter effluent. A more suitable site should be considered in areas of deeper soils.

Slope, shallow depths to bedrock, and rock outcroppings are the main limitations for local roads and streets. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

96—Ipswich mucky peat

Map Unit Setting

Slope: nearly level

Landscape: salt marshes on coastal plains, tidal marshes on coastal plains (fig. 14)

Size of map unit: Areas commonly range from 3 to 25 acres.

Map Unit Composition

Ipswich and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe1—0 to 16 inches; mucky peat

Oe2—16 to 23 inches; mucky peat

Oe3—23 to 64 inches; mucky peat

Oa—64 to 80 inches; muck




Figure 14.—Typical pattern of soils and parent material in coastal tidal marshes and glaciofluvial areas adjacent to Long Island Sound.

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: very poorly drained
Parent material: herbaceous organic material
Permeability: moderate to very rapid
Available water capacity: very high
Reaction: strongly acid to slightly alkaline
Depth to restrictive feature: 20 to 40 inches to salic; 20 to 40 inches to sulfuric
Ponding depth: 0 to 12 inches above surface
Depth to seasonal water table: 0 to 12 inches
Flooding: very frequent

Interpretative Groups

Land capability classification (non-irrigated): 8
Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of very poorly drained Pawcatuck soils where the mucky peat is between 16 to 51 inches thick over sandy substratum. Also included are very poorly drained Westbrook soils where the mucky peat is between 16 to 51 inches thick over loamy substratum. Some areas include Udorthents formed in dredge spoils. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas of this soil are undisturbed and provide habitat for wildlife and are suited to the reproduction of shellfish.

Flooding and ponding are the main limitations for dwellings with basements, lawns and landscaping, septic tank absorption fields, local roads, and streets. Low strength

is also a limitation for dwellings with basements, local roads, and streets. Excess salt and sulfur are also limitations for lawns and landscaping. A more suitable site for all of these uses should be selected on a drier soil not subject to tidal inundation.

97—Pawcatuck mucky peat

Map Unit Setting

Slope: nearly level

Landscape: salt marshes on coastal plains, tidal marshes on coastal plains

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Pawcatuck and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe1—0 to 12 inches; mucky peat

Oe2—12 to 40 inches; mucky peat

Oe3—40 to 46 inches; mucky peat

2Cg1—46 to 50 inches; very fine sandy loam

2Cg2—50 to 60 inches; loamy sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: herbaceous organic material over sandy and gravelly glaciofluvial deposits

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: 0 to 60 inches to salic; 0 to 60 inches to sulfuric

Ponding depth: 0 to 12 inches above surface

Depth to seasonal water table: 0 to 12 inches

Flooding: very frequent

Interpretative Groups

Land capability classification (non-irrigated): 8

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of very poorly drained Ipswich soils where the mucky peat is greater than 51 inches thick. Also included are very poorly drained Westbrook soils where the mucky peat is between 16 to 51 inches thick over loamy substratum. Some areas include Udorthents formed in dredge spoils. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas of this soil are undisturbed and provide habitat for wildlife and are suited to the reproduction of shellfish.

Flooding and ponding are the main limitations for dwellings with basements, lawns and landscaping, septic tank absorption fields, local roads, and streets. Poor filtering

is also a limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Excess salt and sulfur are also limitations for lawns and landscaping. A more suitable site for all of these uses should be selected on a drier soil not subject to tidal inundation.

98—Westbrook mucky peat

Map Unit Setting

Slope: nearly level

Landscape: salt marshes on coastal plains, tidal marshes on coastal plains

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Westbrook and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe1—0 to 10 inches; mucky peat

Oe2—10 to 40 inches; mucky peat

Oe3—40 to 48 inches; mucky peat

Cg1—48 to 64 inches; silt loam

Cg2—64 to 99 inches; silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: herbaceous organic material over loamy drift and/or marine deposits

Permeability: very slow to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: 0 to 51 inches to salic; 0 to 51 inches to sulfuric

Ponding depth: 0 to 12 inches above surface

Depth to seasonal water table: 0 to 12 inches

Flooding: very frequent

Interpretative Groups

Land capability classification (non-irrigated): 8

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of very poorly drained Pawcatuck and Timakwa soils. Pawcatuck soils have mucky peat between 16 to 51 inches thick over sandy substratum; Timakwa soils have muck between 16 and 51 inches thick over a sandy substratum and have a lower salt content. Also included are very poorly drained Ipswich and Natchaug soils. Ipswich soils have mucky peat greater than 51 inches thick; Natchaug soils have muck between 16 to 51 inches thick over loamy substratum and have a lower salt content. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas of this soil are undisturbed and provide wildlife habitat. Some areas have been filled and are in community development.

Flooding and ponding are the main limitations for dwellings with basements, septic tank absorption fields, lawns and landscaping, local roads, and streets. Excess salt and sulfur are also limitations for lawns and landscaping. A more suitable site for all of these uses should be selected on a drier soil not subject to tidal inundation.

99—Westbrook mucky peat, low salt

Map Unit Setting

Slope: nearly level

Landscape: salt marshes on coastal plains, tidal marshes on coastal plains

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Westbrook and similar soils: 80 percent ([fig. 15](#))

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe1—0 to 10 inches; mucky peat

Oe2—10 to 40 inches; mucky peat

Oe3—40 to 48 inches; mucky peat

Cg1—48 to 64 inches; silt loam

Cg2—64 to 99 inches; silt loam



Figure 15.—Areas of Westbrook mucky peat are subject to daily tidal inundation.

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: herbaceous organic material over loamy drift and/or marine deposits

Permeability: very slow to very rapid

Available water capacity: very high

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 12 inches above surface

Depth to seasonal water table: 0 to 12 inches

Flooding: very frequent

Interpretative Groups

Land capability classification (non-irrigated): 8

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of very poorly drained Pawcatuck and Timakwa soils. Pawcatuck soils have mucky peat between 16 to 51 inches thick over sandy substratum; Timakwa soils have muck between 16 to 51 inches thick over a sandy substratum and have a lower salt content. Also included are very poorly drained Ipswich and Natchaug soils. Ipswich soils have mucky peat greater than 51 inches thick; Natchaug soils have muck between 16 to 51 inches thick over loamy substratum and have a lower salt content. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas of this soil are undisturbed and provide wildlife habitat. Some areas have been filled and are in community development.

Flooding and ponding are the main limitations for dwellings with basements, septic tank absorption fields, lawns and landscaping, local roads, and streets. Excess sulfur is a limitation for lawns and landscaping. A more suitable site for all of these uses should be selected on a drier soil not subject to tidal inundation.

100—Suncook loamy fine sand

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Suncook and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; loamy fine sand

C1—7 to 15 inches; stratified coarse sand to loamy fine sand

C2—15 to 22 inches; stratified coarse sand to loamy fine sand

C3—22 to 32 inches; stratified coarse sand to loamy fine sand

C4—32 to 42 inches; stratified coarse sand to loamy fine sand

C5—42 to 65 inches; stratified gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy alluvium

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 60 to 72 inches

Flooding: occasional

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Minor Components

Included with this soil are areas of well drained Occum soils adjacent to natural levees and moderately well drained Pootatuck soils in slightly lower portions of the flood plain. Areas of poorly drained Rippowam soils and very poorly drained Saco soils are included in depressions and channel scars on the flood plain. Also included are Fluvaquents-Udfluvents complex in areas subject to frequent flooding and some soils that are not subject to flooding. A few areas in Litchfield County include soils with a fine sandy loam or very fine sandy loam surface layer. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture or woodland. Some areas are in cultivated cropland.

Flooding is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Poor filtering is also a limitation for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site for these uses should be selected on a soil that does not flood.

Flooding is the main limitation for local roads and streets. Providing drainage and building on raised fill will reduce this limitation. A more suitable site should be considered on a soil that does not flood.

101—Occum fine sandy loam

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Occum and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 10 inches; fine sandy loam

Bw1—10 to 17 inches; fine sandy loam

Bw2—17 to 28 inches; sandy loam

C1—28 to 32 inches; stratified very gravelly coarse sand to loamy fine sand

C2—32 to 42 inches; stratified very gravelly coarse sand to loamy fine sand

C3—42 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy alluvium

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 60 to 72 inches

Flooding: occasional

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil are excessively drained Suncook soils on natural flood plain levees, well drained Agawam soils on nearby outwash plains and terraces, and moderately well drained Pootatuck soils in slightly lower portions of the flood plain. Areas of poorly drained Rippowam soils are included in depressions and channel scars on the flood plain. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. Some areas are in woodland or urban development.

Flooding is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Poor filtering is also a limitation for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site for these uses should be selected on a soil that does not flood.

Flooding is the main limitation for local roads and streets. Providing drainage and building on raised fill will reduce this limitation. A more suitable site should be considered on a soil that does not flood.

102—Pootatuck fine sandy loam

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Pootatuck and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 16 inches; fine sandy loam

Bw2—16 to 21 inches; fine sandy loam

Bw3—21 to 29 inches; sandy loam

C1—29 to 35 inches; stratified very gravelly coarse sand to loamy fine sand

C2—35 to 40 inches; stratified very gravelly coarse sand to loamy fine sand

C3—40 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy alluvium

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: frequent

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil are areas of excessively drained Suncook soils on natural flood plain levees and well drained Occum soils on slightly higher portions of the flood plain. Areas of poorly drained Rippowam, Lim, and Limerick soils and very poorly drained Saco soils are included in depressions and channel scars on the flood plain. Lim and Limerick soils do not have the fine sandy loam or coarser textures throughout the substratum that Rippowam soils have. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are cleared and in cultivated crops, hay, or pasture.

Flooding is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Poor filtering is also a limitation for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site for these uses should be selected on a soil that does not flood.

Flooding is the main limitation for local roads and streets. Providing drainage and building on raised fill will reduce this limitation. A more suitable site should be considered on a soil that does not flood.

103—Rippowam fine sandy loam

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Rippowam and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; fine sandy loam

Bg1—5 to 12 inches; fine sandy loam

Bg2—12 to 19 inches; fine sandy loam

BCg1—19 to 24 inches; sandy loam

BCg2—24 to 27 inches; sandy loam

Cg1—27 to 31 inches; loamy sand

Cg2—31 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy alluvium

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 18 inches

Flooding: frequent

Interpretative Groups

Land capability classification (non-irrigated): not specified

Hydrologic group: D

Minor Components

Included with this soil are areas of excessively drained, sandy Suncook soils on natural flood plain levees, well drained, loamy Occum soils and moderately well drained, loamy Pootatuck soils on slightly lower portions of the flood plain. Areas of poorly drained, silty Lim and Limerick soils are included. Lim and Limerick soils do not have the fine sandy loam or coarser textures throughout the substratum that Rippowam soils have. Very poorly drained, silty Saco soils are included in the lowest lying areas of the flood plain. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in cultivated crops, hay, or pasture.

Flooding and the seasonal high water table are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Poor filtering is also a limitation for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not

adequately filter effluent. A more suitable site for these uses should be selected on a soil that does not flood.

Flooding, wetness, and potential frost action are the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

104—Bash silt loam

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 80 acres.

Map Unit Composition

Bash and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 11 inches; silt loam

Bw1—11 to 21 inches; silt loam

Bw2—21 to 28 inches; silt loam

C—28 to 60 inches; silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat poorly drained

Parent material: coarse-loamy alluvium derived from sandstone and shale

Permeability: moderately slow to moderate

Available water capacity: very high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 6 to 18 inches

Flooding: frequent

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of well drained Hadley soils and moderately well drained Winooski soils on slightly higher portions of the flood plain above Bash soils. Also included are poorly drained Limerick and Lim soils and very poorly drained Saco soils in depressions and channel scars on the flood plain. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are cleared and in cultivated crops and pasture. A few scattered areas are filled and in community development.

Flooding and the seasonal high water table are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. A more suitable site should be selected on a drier soil that does not flood.

Flooding, wetness, and potential frost action are the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

105—Hadley silt loam

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Hadley and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 12 inches; silt loam

C1—12 to 29 inches; stratified very fine sand to silt loam

C2—29 to 40 inches; stratified very fine sand to silt loam

C3—40 to 45 inches; stratified sand to silt loam

C4—45 to 60 inches; stratified sand to silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-silty alluvium

Permeability: moderate to very rapid

Available water capacity: very high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 60 to 72 inches

Flooding: occasional

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of moderately well drained Winooski soils on slightly lower portions of the flood plain. Also included are somewhat poorly drained Bash soils that are redder in color. Small areas of poorly drained Limerick and Lim soils and very poorly drained Saco soils are included in depressions and channel scars on the flood plain. A few areas in Litchfield County include soils with sand and gravel at 24 to 36 inches. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated crops, hay, or pasture. A few areas are wooded or in community development.

Flooding is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. A more suitable site should be selected on a drier soil that does not flood.

Flooding and potential frost action is the main limitation for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

106—Winooski silt loam

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 25 acres.

Map Unit Composition

Winooski and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 12 inches; silt loam

Bw1—12 to 18 inches; silt loam

Bw2—18 to 36 inches; silt loam

C1—36 to 52 inches; very fine sandy loam

C2—52 to 65 inches; silt loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-silty alluvium

Permeability: moderate or moderately rapid

Available water capacity: very high

Reaction: moderately acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: frequent

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of well drained Hadley soils on slightly higher portions of the flood plain. Also included are somewhat poorly drained Bash soils that have a red color. Small areas of poorly drained Limerick and Lim soils and very poorly drained Saco soils are included in depressions and channel scars on the flood plain. A few areas in Litchfield County include some strongly acid soils and some soils with coarse textured materials at 24 to 30 inches. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in cultivated cropland. Some areas are in woodland or pasture.

Flooding and the seasonal high water table are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. A more suitable site should be selected on a drier soil that does not flood.

Flooding and potential frost action is the main limitation for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

107—Limerick and Lim soils

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Limerick and similar soils: 50 percent

Lim and similar soils: 30 percent

Minor components: 20 percent

Major Components

Limerick and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

BCg1—8 to 20 inches; silt loam

BCg2—20 to 36 inches; silt loam

BCg3—36 to 54 inches; silt loam

Cg—54 to 65 inches; silt loam

Lim and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 6 inches; very fine sandy loam

Bg1—6 to 11 inches; very fine sandy loam

Bg2—11 to 15 inches; very fine sandy loam

Bg3—15 to 22 inches; silt loam

Bg4—22 to 29 inches; fine sandy loam

CB—29 to 42 inches; stratified very gravelly coarse sand to loamy fine sand

Cg1—42 to 50 inches; stratified very gravelly coarse sand to loamy fine sand

Cg2—50 to 57 inches; stratified very gravelly coarse sand to loamy fine sand

Cg3—57 to 65 inches; stratified very gravelly coarse sand to loamy sand

Major Component Properties and Qualities

Limerick and similar soils

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-silty alluvium

Permeability: moderate

Available water capacity: very high

Reaction: strongly acid to neutral
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 0 to 18 inches
Flooding: frequent

Lim and similar soils

Depth to bedrock: very deep
Drainage class: poorly drained
Parent material: coarse-loamy alluvium
Permeability: moderate to very rapid
Available water capacity: high
Reaction: strongly acid to neutral
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 0 to 18 inches
Flooding: frequent

Interpretative Groups

Limerick and similar soils

Land capability classification (non-irrigated): not specified
Hydrologic group: D

Lim and similar soils

Land capability classification (non-irrigated): 4w
Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of very poorly drained Saco soils in the lowest lying areas of the flood plain. Saco soils are greater than 40 inches to coarse textured substratum. Also included are areas of poorly drained Rippowam soils and somewhat poorly drained Bash soils. Rippowam soils are loamier throughout and Bash soils have red color. Areas of moderately well drained Winooski soils and well drained Hadley soils are on higher portions of the flood plain. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland or in marsh grasses and sedges. Some areas are drained or cleared. Cleared areas are in cultivated cropland or pasture and other areas are in residential development.

Flooding and the seasonal high water table are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Poor filtering is also a limitation for septic tank absorption fields in areas of Lim soils. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be selected on a drier soil that does not flood.

Flooding, potential frost action, and the seasonal high water table are the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

108—Saco silt loam

Map Unit Setting

Slope: nearly level
Landscape: flood plains
Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Saco and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 12 inches; silt loam

Cg1—12 to 32 inches; silt loam

Cg2—32 to 48 inches; silt loam

2Cg3—48 to 60 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: coarse-silty alluvium

Permeability: moderate to very rapid

Available water capacity: very high

Reaction: strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 18 inches above surface

Depth to seasonal water table: 0 to 6 inches

Flooding: frequent

Interpretative Groups

Land capability classification (non-irrigated): 6w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of poorly drained Limerick, Lim, and Rippowam soils. Limerick soils are siltier throughout, Lim soils are less than 40 inches to coarse textured substratum, and Rippowam soils are loamier throughout. Somewhat poorly drained Bash soils are included in areas where the soil color is red due to parent material. Also included are moderately well drained Winooski soils and well drained Hadley soils on higher portions of the flood plain. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland or marsh, and are wetland wildlife habitat.

Flooding and the seasonal high water table are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. A more suitable site should be selected on a drier soil that does not flood.

Flooding, potential frost action, and the seasonal high water table are the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

109—Fluvaquents-Udifluents complex, frequently flooded

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Fluvaquents and similar soils: 50 percent

Udifuvents and similar soils: 35 percent

Minor components: 15 percent

Major Components

Fluvaquents and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 4 inches; silt loam

Cg1—4 to 14 inches; fine sand

Cg2—14 to 21 inches; very fine sand

Ab1—21 to 38 inches; silt loam

Ab2—38 to 45 inches; fine sandy loam

C'g3—45 to 55 inches; sand

A'b3—55 to 60 inches; fine sandy loam

Udifuvents and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 2 inches; fine sandy loam

C—2 to 4 inches; loamy fine sand

Ap—4 to 12 inches; fine sandy loam

AC—12 to 18 inches; fine sandy loam

C1—18 to 35 inches; loamy sand

C2—35 to 38 inches; very gravelly loamy sand

C3—38 to 60 inches; very gravelly coarse sand

Major Component Properties and Qualities

Fluvaquents and similar soils

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: alluvium

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: frequent

Udifuvents and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: alluvium

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 72 inches

Flooding: frequent

Interpretative Groups

Fluvaquents and similar soils

Land capability classification (non-irrigated): 6w

Hydrologic group: D

Udfluvents and similar soils

Land capability classification (non-irrigated): 6w

Hydrologic group: B

Minor Components

Included with this complex in mapping are areas of Riverwash and other gravelly deposits. This map unit includes areas very poorly drained Saco soils, poorly drained Rippowam soils, moderately well drained Pootatuck soils, and well drained Occum soils. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. A few areas are in pasture.

Flooding is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. The seasonal high water table is also a limitation for septic tank adsorption fields. A more suitable site should be selected on a drier soil that does not flood. Flooding and frost action are the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

221A—Ninigret-Urban land complex, 0 to 5 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ninigret and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Ninigret and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

2C—26 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: high
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Ninigret and similar soils

Land capability classification (non-irrigated): 2w
Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of somewhat excessively drained Merrimac soils and well drained Agawam and Haven soils that are higher on the landscape. Agawam soils are loamy over sand and gravel and Haven soils are silty over sand and gravel. Also included are moderately well drained Sudbury soils that are sandy and gravelly throughout. Small areas poorly drained Raypol soils are included in shallow depressions and drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce the wetness.

Poor filtering and the seasonal high water table are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal. There is also a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

224A—Deerfield-Urban land complex, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level
Landscape: terraces on valleys, outwash plains on valleys
Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Deerfield and similar soils: 40 percent
Urban land and similar soils: 35 percent
Minor components: 25 percent

Major Components

Deerfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; loamy fine sand

Bw1—8 to 16 inches; loamy sand

Bw2—16 to 28 inches; loamy sand

C1—28 to 34 inches; fine sand

C2—34 to 60 inches; fine sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H— 0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: sandy glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Deerfield and similar soils

Land capability classification (non-irrigated): 2w

Hydrologic group: A

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Windsor and Penwood soils that are higher on the landscape. Windsor soils are sandy throughout and Penwood soils have a red subsoil. Small areas of poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Droughtiness can make establishment and maintenance of lawns difficult. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Modifying a conventional

system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

225B—Brancroft-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Brancroft and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Bancroft and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 17 inches; silt loam

Bw2—17 to 22 inches; silty clay loam

Bw3—22 to 32 inches; silt loam

C1—32 to 43 inches; silty clay loam

C2—43 to 66 inches; silt loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: fine-silty glaciolacustrine deposits

Permeability: very slow to moderate

Available water capacity: very high

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Bancroft and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are some areas of moderately well drained Elmridge and Berlin soils. Elmridge soils have a loamy over clayey substratum and Berlin soils are reddish brown in color due to parent material. Poorly drained Scitico soils and very poorly drained Maybid soils are included in deep depressions and drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Low strength and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

226B—Berlin-Urban land complex, 0 to 8 percent slopes**Map Unit Setting**

Slope: nearly level to gently sloping

Landscape: terraces on lake plains

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Berlin and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components**Berlin and similar soils**

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 12 inches; silt loam

Bw2—12 to 20 inches; silty clay loam

Bw3—20 to 34 inches; silty clay loam

C1—34 to 48 inches; silty clay loam

C2—48 to 65 inches; silty clay loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: fine-silty glaciolacustrine deposits
Permeability: very slow to moderate
Available water capacity: very high
Reaction: very strongly acid to neutral
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 12 to 30 inches
Flooding: none

Interpretative Groups

Berlin and similar soils

Land capability classification (non-irrigated): 2e
Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of moderately well drained Brancroft, Elmridge, and Belgrade soils. These soils are yellower in the subsoil and substratum. Also included are areas of moderately well drained Ludlow soils, which have a dense substratum. Well drained Wethersfield soils are included in areas that are higher on the landscape and have a dense substratum. Small areas of poorly drained Scitico soils and very poorly drained Maybid soils are included in depressions and along drainageways. Areas of Udorthents are included adjacent to building and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Low strength and frost action are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

228B—Elmridge-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping
Landscape: terraces on lake plains
Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Elmridge and similar soils: 40 percent
Urban land and similar soils: 35 percent
Minor components: 25 percent

Major Components

Elmridge and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam
 Bw1—6 to 10 inches; fine sandy loam
 Bw2—10 to 18 inches; fine sandy loam
 Bw3—18 to 25 inches; sandy loam
 2C—25 to 65 inches; silty clay

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over clayey glaciolacustrine deposits

Permeability: very slow to moderately rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Elmridge and similar soils

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of moderately well drained Brancroft, Sudbury, Ninigret, Berlin, and Belgrade soils. Brancroft soils are silty and clayey; Sudbury soils are sandy and gravelly; Ninigret soils are loamy over sand and gravel; Berlin soils are redder; and Belgrade soils are silty throughout. Also included are small areas of poorly drained Shaker and Scitico soils in broad, flat low-lying or slightly concave areas. Small areas of very poorly drained Maybid soils are also included in depressions and along drainageways. A few areas include soils with reddish brown to yellowish red subsoil. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. Open areas are in lawns, gardens, and woodland or brushland structures.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in places.

Frost action is the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

229B—Agawam-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Agawam and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Agawam and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to fine sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid or very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Agawam and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are higher on the landscape; Hinckley and Merrimac soils are sandier in the subsoil. Also included are some moderately well drained Ninigret soils in slightly lower areas below Agawam soils. Poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. Scarboro soils have a mucky surface layer. A few areas in Hartford County include soils with red subsoil and substratum. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This soil has few limitations for dwellings with basements and lawns and landscaping. Droughtiness can make establishment and maintenance of lawns difficult.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

This soil has few limitations for local roads and streets.

229C—Agawam-Urban land complex, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Agawam and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Agawam and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to fine sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Agawam and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are higher on the landscape; Hinckley and Merrimac soils are sandier in the subsoil. Also included are some moderately well drained Ninigret soils in slightly lower areas below Agawam soils. Poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. Scarboro soils have a mucky surface layer. A few areas in Hartford County include soils with red subsoil and substratum. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

Slope is the main limitation for dwellings with basements and for lawns and landscaping. Erosion is a moderate hazard during construction.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

230B—Branford-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Branford and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Branford and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; loam

Bw2—18 to 24 inches; gravelly loam

2C—24 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Branford and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of well drained Haven and Enfield soils. Enfield soils are coarse-silty over sand and gravel and Haven soils are coarse-loamy over sand and gravel. Also included are moderately well drained Ellington soils in slightly lower areas on the landscape. Small areas of poorly drained Raypol soils are included in shallow depressions and along drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This soil has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

230C—Branford-Urban land complex, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Branford and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Branford and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 18 inches; loam

Bw2—18 to 24 inches; gravelly loam

2C—24 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Branford and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of well drained Haven and Enfield soils. Enfield soils are coarse-silty over sand and gravel and Haven soils are coarse-loamy over sand and gravel. Also included are moderately well drained Ellington soils in slightly lower areas below Branford soils. Small areas of very poorly drained Raypol soils are included in shallow depressions and along drainageways. Areas of

Udorthents are included adjacent to buildings and other structures. Minor componets make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

232B—Haven-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Haven and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Haven and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; silt loam

Bw1—7 to 14 inches; silt loam

Bw2—14 to 20 inches; silt loam

BC—20 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly sand to gravelly fine sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Haven and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of well drained Branford and Agawam soils. Branford soils are silty over sand and gravel, and are red in color. Agawam soils are sandier in the surface layer and subsoil. Also included are moderately well drained Ninigret and Tisbury soils in slightly lower areas below Haven and Enfield soils. Poorly drained Raypol soils are included in shallow depressions and drainageways. A few areas in New London County include soils with a gravelly surface layer. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This unit has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

234B—Merrimac-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: terraces on valleys, kames on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 5 to 75 acres.

Map Unit Composition

Merrimac and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Merrimac and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; sandy loam

Bw1—9 to 16 inches; sandy loam

Bw2—16 to 24 inches; gravelly sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to gravelly sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups**Merrimac and similar soils**

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Hinckley and Windsor soils that are higher on the landscape. Hinckley soils are sandy and gravelly and Windsor soils are sandy throughout. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Ninigret and Sudbury soils are included in slightly lower areas. Ninigret soils are loamy over sand and gravel and Sudbury soils are sandy and gravelly. Small areas of poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This soil has few limitations for dwellings with basements, lawns and landscaping, and local roads and streets.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

235B—Penwood-Urban land complex, 0 to 8 percent slopes***Map Unit Setting***

Slope: nearly level to gently sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Penwood and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Penwood and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; loamy sand

Bw1—8 to 18 inches; loamy sand

Bw2—18 to 30 inches; sand

C—30 to 60 inches; sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy glaciofluvial deposits derived from sandstone and shale

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Penwood and similar soils

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are excessively drained Manchester soils and somewhat excessively drained Hartford soils in areas that are sandy and gravelly. Also included are areas of well drained Branford soils and moderately well drained Ellington soils. Branford and Ellington soils are silty over sand and gravel. A few areas in New Haven County include soils with a gravelly substratum. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

236B—Windsor-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: terraces on valleys, outwash plains on valleys, kames on valleys

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Windsor and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Windsor and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 3 inches; loamy sand

Bw1—3 to 9 inches; loamy sand

Bw2—9 to 21 inches; loamy sand

Bw3—21 to 25 inches; sand

C—25 to 65 inches; sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: eolian sands over sandy glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Windsor and similar soils

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Hinckley soils and somewhat excessively drained Merrimac soils that are sandy and gravelly. Also included are well drained Agawam soils that are loamy over sand and gravel. Moderately well drained Deerfield, Ninigret, and Sudbury soils are included in slightly lower areas. Ninigret soils are loamy over sand and gravel and Sudbury soils are sandy and gravelly. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

237A—Manchester-Urban land complex, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: outwash plains on valleys, kames on valleys, terraces on valleys, eskers on valleys

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Manchester and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Manchester and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; gravelly sandy loam

Bw—9 to 18 inches; gravelly loamy sand

C—18 to 65 inches; stratified extremely gravelly coarse sand to very gravelly loamy sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Manchester and similar soils

Land capability classification (non-irrigated): 2s

Hydrologic group: A

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Penwood soils that are sandy throughout. Also included are somewhat excessively drained Hartford soils, well drained Branford soils, and moderately well drained Ellington soils. Hartford soils are sandy loam over a sandy and gravelly substratum, Branford soils are silty over a sandy and gravelly substratum, and Ellington soils are in slightly lower areas and broad drainageways. In places, soils that lack a gravelly surface are included. A few areas in New Haven County have a gravelly loamy sand surface layer. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

237C—Manchester-Urban land complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: kames on valleys, eskers on valleys, terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Manchester and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Manchester and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; gravelly sandy loam

Bw—9 to 18 inches; gravelly loamy sand

C—18 to 65 inches; stratified extremely gravelly coarse sand to very gravelly loamy sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from sandstone and shale and/or basalt

Permeability: rapid or very rapid

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Manchester and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: A

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Penwood soils that are sandy throughout. Also included are somewhat excessively drained Hartford soils, well drained Branford soils, and moderately well drained Ellington soils. Hartford soils are sandy loam over a sandy and gravelly substratum, Branford soils are silty over a sandy and gravelly substratum, and Ellington soils are in slightly lower areas and broad drainageways. A few areas in New Haven County have a gravelly loamy sand surface layer. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Droughtiness is the main limitation for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not

adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

238A—Hinckley-Urban land complex, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: kames on valleys, eskers on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Hinckley and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Hinckley and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 20 inches; very gravelly loamy sand

Bw2—20 to 27 inches; very gravelly sand

C1—27 to 42 inches; stratified cobbly coarse sand to extremely gravelly sand

C2—42 to 60 inches; stratified cobbly coarse sand to extremely gravelly sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: very low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Hinckley and similar soils

Land capability classification (non-irrigated): 3s

Hydrologic group: A

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Windsor soils which are sandy throughout. Also included are somewhat excessively drained Merrimac soils and well drained Agawam soils. Merrimac soils are sandy over sand and gravel and Agawam soils are loamy over sand and gravel. Small areas of moderately well drained Sudbury soils are included in slightly lower areas, poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

238C—Hinckley-Urban land complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: kames on valleys, outwash plains on valleys, terraces on valleys, eskers on valleys

Size of map unit: Areas commonly range from 5 to 200 acres.

Map Unit Composition

Hinckley and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Hinckley and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; gravelly sandy loam

Bw1—8 to 20 inches; very gravelly loamy sand

Bw2—20 to 27 inches; very gravelly sand

C1—27 to 42 inches; stratified cobbly coarse sand to extremely gravelly sand

C2—42 to 60 inches; stratified cobbly coarse sand to extremely gravelly sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: rapid or very rapid

Available water capacity: very low

Reaction: extremely acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Hinckley and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: A

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this complex in mapping are areas of excessively drained Windsor soils which are sandy throughout. Also included are somewhat excessively drained Merrimac soils and well drained Agawam soils. Merrimac soils are sandy over sand and gravel and Agawam soils are loamy over sand and gravel. Small areas of moderately well drained Sudbury soils are included in slightly lower areas, poorly drained Walpole soils and very poorly drained Scarboro soils are included in shallow depressions and drainageways. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of this map unit.

Use and Management

This unit is in urban and suburban development. The open areas are in lawns, gardens, and woodland or brushland between structures.

Slope is the main limitation for dwellings with basements. Erosion is a hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

240B—Ludlow-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ludlow and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Ludlow and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 20 inches; silt loam

Bw2—20 to 26 inches; silt loam

Cd—26 to 65 inches; gravelly loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Ludlow and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are well drained Wethersfield soils, well drained Cheshire soils, well drained Yalesville soils, moderately well drained Watchaug soils, poorly drained Wilbraham, and very poorly drained Menlo. Wethersfield soils are on higher areas. Yalesville soils are moderately deep to bedrock. Watchaug and Cheshire soils are in areas where the substratum is friable. Wilbraham and Menlo soils are in depressions and along drainageways. Soils with a loam or fine sandy loam surface are included in New Haven County. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with

foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

243B—Rainbow-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Rainbow and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 20 percent

Major Components

Rainbow and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 18 inches; silt loam

Bw2—18 to 26 inches; silt loam

2Cd—26 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Rainbow and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Narragansett soils, well drained Broadbrook soils, and poorly drained Wilbraham soils. Narragansett soils are in areas of very friable to firm sandy glacial till. Broadbrook soils are on the highest parts of the landscape. Wilbraham soils are in depressions and along drainageways. Soils with a stony surface are included in some areas. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

245B—Woodbridge-Urban land complex, 0 to 8 percent slopes**Map Unit Setting**

Slope: nearly level to gently sloping

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Woodbridge and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components**Woodbridge and similar soils**

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; fine sandy loam

Bw1—7 to 18 inches; fine sandy loam

Bw2—18 to 26 inches; fine sandy loam

Bw3—26 to 30 inches; fine sandy loam

Cd1—30 to 43 inches; gravelly fine sandy loam

Cd2—43 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Woodbridge and similar soils

Land capability classification (non-irrigated): 2w

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of well drained Paxton soils, well drained Montauk soils, moderately well drained Sutton and Rainbow soils, poorly drained Ridgebury and Leicester soils, and very poorly drained Whitman soils. Paxton and Montauk soils are on higher areas. Sutton soils have a friable substratum and Rainbow soils are finer in texture. Ridgebury, Leicester, and Whitman soils are in depressions and along drainageways. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

Potential frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

245C—Woodbridge-Urban land complex, 8 To 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Woodbridge and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Woodbridge and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 7 inches; fine sandy loam

Bw1—7 to 18 inches; fine sandy loam

Bw2—18 to 26 inches; fine sandy loam

Bw3—26 to 30 inches; fine sandy loam

Cd1—30 to 43 inches; gravelly fine sandy loam

Cd2—43 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Woodbridge and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of well drained Paxton soils, well drained Montauk soils, moderately well drained Sutton and Rainbow soils, poorly drained Ridgebury and Leicester soils, and very poorly drained Whitman soils. Paxton and Montauk soils are on higher areas. Sutton soils have a friable substratum and Rainbow soils are finer in texture. Ridgebury, Leicester, and Whitman soils are in depressions and along drainageways. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

Potential frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

248B—Georgia-Urban land complex, 2 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: hills, uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Georgia and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Georgia and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; loam

Bw2—14 to 24 inches; loam

C—24 to 60 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow to moderate

Available water capacity: high

Reaction: strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Georgia and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of well drained Stockbridge, Nellis, and Paxton soils, moderately well drained Amenia and Woodbridge soils, poorly drained Mudgepond and Ridgebury soils, and very poorly drained Alden soils. Amenia soils are less acid than Georgia soils and are calcareous within 40 inches. Stockbridge and Nellis soils are in higher areas. Mudgepond and Alden soils are in depressions and along drainageways. Paxton, Woodbridge, and Ridgebury soils are in areas with dense substratum. Udorthents are included in areas adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements. Large stones are a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, water proofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal in most places.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

250B—Sutton-Urban land complex, 0 to 8 percent slopes**Map Unit Setting**

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Sutton and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components**Sutton and similar soils**

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; fine sandy loam

Bw1—6 to 12 inches; fine sandy loam

Bw2—12 to 24 inches; fine sandy loam

Bw3—24 to 28 inches; fine sandy loam

C1—28 to 36 inches; gravelly fine sandy loam

C2—36 to 65 inches; gravelly sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups**Sutton and similar soils**

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Paxton soils and moderately well drained Woodbridge soils. Canton soils are loamy over sandy; Charlton soils are sandy loam throughout; Paxton and Woodbridge soils have a dense substratum. Some areas have a silt loam surface layer and subsoil. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the map unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

253B—Wapping-Urban land complex, 0 to 8 percent slopes***Map Unit Setting***

Slope: nearly level to gently sloping

Landscape: till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Wapping and similar soils: 40 percent
Urban land and similar soils: 35 percent
Minor components: 25 percent

Major Components

Wapping and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 11 inches; very fine sandy loam
 Bw1—11 to 16 inches; very fine sandy loam
 Bw2—16 to 20 inches; very fine sandy loam
 2C1—20 to 28 inches; gravelly sandy loam
 2C2—28 to 36 inches; gravelly loamy sand
 2C3—36 to 80 inches; gravelly loamy sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Wapping and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of well drained Narragansett, Cheshire, and Yalesville soils, moderately well drained Watchaug and Ludlow soils, poorly drained Leicester and Wilbraham soils, and very poorly drained Menlo soils. Narragansett soils are in higher areas and Yalesville soils are in areas moderately deep to bedrock. Cheshire, Watchaug, and Ludlow soils are in areas where the subsoil and substratum are red due to the parent material. Wilbraham, Leicester, and Menlo soils are in depressions and along drainageways. Also included are soils with a stony surface. Minor components make up about 25 percent of the unit.

Use and Management

Most areas are in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with footing or foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of the basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action and low strength are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

255B—Watchaug-Urban land complex, 0 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Watchaug and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Watchaug and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 18 inches; fine sandy loam

Bw2—18 to 24 inches; fine sandy loam

C—24 to 65 inches; gravelly sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate to moderately rapid

Available water capacity: high

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Watchaug and similar soils

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of well drained Cheshire soils, moderately well drained Ludlow soils, poorly drained Wilbraham and Watchaug soils, and very poorly drained Menlo soils. Cheshire soils are on higher areas and Ludlow soils are included in areas with dense substratum. Wilbraham and Menlo soils are in depressions and along drainageways. Areas of Watchaug soils with a silt loam surface or stony surface are included in New Haven County. Udorthents are included adjacent to buildings and other structures. Minor componets make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings on the highest part of the unit with footing or foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff form higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Potential frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

260B—Charlton-Urban land complex, 3 to 8 percent slopes**Map Unit Setting**

Slope: gently sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Charlton and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components**Charlton and similar soils**

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate to moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups**Charlton and similar soils**

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of somewhat excessively drained and well drained Chatfield and Hollis soils, moderately well drained Sutton soils, and poorly drained Leicester soils. Moderately deep Chatfield soils and shallow Hollis soils are in areas underlain by bedrock. Sutton soils are on slightly lower areas and Leicester soils are in depressions and along drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the map unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

This unit has few limitations for dwellings with basements, lawns and landscaping, septic tank absorption fields, and local roads and streets.

260C—Charlton-Urban land complex, 8 to 15 percent slopes***Map Unit Setting***

Slope: strongly sloping

Landscape: hills, uplands

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Charlton and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

- Ap—0 to 4 inches; fine sandy loam
- Bw1—4 to 7 inches; fine sandy loam
- Bw2—7 to 19 inches; fine sandy loam
- Bw3—19 to 27 inches; gravelly fine sandy loam
- C—27 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

- H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Charlton and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of somewhat excessively drained and well drained Chatfield and Hollis soils, moderately well drained Sutton soils, and poorly drained Leicester soils. Moderately deep Chatfield soils and shallow Hollis soils are in areas underlain by bedrock. Sutton soils are on slightly lower areas and Leicester soils are in depressions and along drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the map unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements, lawns and landscaping, septic tank absorption fields, and local roads and streets. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Placing the septic tank absorption field distribution lines on the contour increases the efficiency of the system. Constructing roads on the contour will reduce the slope limitation.

260D—Charlton-Urban land complex, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Charlton and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Charlton and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of somewhat excessively drained and well drained Chatfield and Hollis soils, moderately well drained Sutton soils, and poorly drained Leicester soils. Moderately deep Chatfield soils and shallow Hollis soils are in areas underlain by bedrock. Sutton soils are on slightly lower areas and

Leicester soils are in depressions and along drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor componets make up about 25 percent of the map unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements, lawns and landscaping, septic tank absorption fields, and local roads and streets. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Placing the septic tank absorption field distribution lines on the contour increases the efficiency of the system. Constructing roads on the contour will reduce the slope limitation. A more suitable site should be considered on a less sloping portion of the unit or nearby soil.

263B—Cheshire-Urban land complex, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Cheshire and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Cheshire and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

C—26 to 65 inches; gravelly sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Cheshire and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Wethersfield and Yalesville soils, moderately well drained Watchaug soils, poorly drained Wilbraham soils, and very poorly drained Menlo soils. Wethersfield soils have a firm substratum and Yalesville soils are moderately deep to bedrock. Watchaug soils are on slightly lower areas. Wilbraham and Menlo soils are along drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

This unit has few limitations for dwellings with basements, lawns and landscaping, septic tank absorption fields, or local roads and streets.

263C—Cheshire-Urban land complex, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Cheshire and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Cheshire and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

C—26 to 65 inches; gravelly sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Cheshire and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Wethersfield and Yalesville soils, moderately well drained Watchaug soils, poorly drained Wilbraham soils, and very poorly drained Menlo soils. Wethersfield soils have a firm substratum and Yalesville soils are moderately deep to bedrock. Watchaug soils are on slightly lower areas. Wilbraham and Menlo soils are along drainageways. Areas of Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements, lawns and landscaping, septic tank absorption fields, or local roads and streets. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Placing the septic tank absorption field distribution lines on the contour increases the efficiency of the system. Constructing roads on the contour will reduce the slope limitation.

266B—Narragansett-Urban land complex, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Narragansett and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Narragansett and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 6 inches; silt loam

Bw1—6 to 15 inches; silt loam

Bw2—15 to 24 inches; silt loam

Bw3—24 to 28 inches; gravelly silt loam

2C—28 to 60 inches; very gravelly loamy coarse sand

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly melt-out till derived from gneiss and/or schist and/or sandstone and shale

Permeability: moderate to very rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Narragansett and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Broadbrook, Canton, and Charlton soils, moderately well drained Wapping and Sutton soils, and poorly drained Leicester soils. Broadbrook soils have a dense substratum, Canton soils are coarser textured, and Charlton soils are loamy throughout. Wapping and Sutton soils are on slightly lower areas. Leicester soils are in depressions and along drainageways. Soils with reddish color are included in northeast Hartford County. Minor components make up about 20 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

This unit has few limitations for dwellings with basements or lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce these limitations.

269B—Yalesville-Urban land complex, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: bedrock-controlled hills on uplands, bedrock-controlled ridges on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Yalesville and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Yalesville and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; loam

C—25 to 36 inches; gravelly sandy loam

2R—36 to 80 inches; unweathered bedrock

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Yalesville and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Holyoke soils, well drained Cheshire and Wethersfield soils, and moderately well drained Watchaug and Ludlow soils. Holyoke soils are in areas where the bedrock is between 10 and 20 inches deep and Wethersfield soils are in areas where the substratum is very firm.

Watchaug and Ludlow soils are on slightly lower areas. Also included are areas with a silt loam surface texture in Middlesex and New Haven counties. Areas with slopes less than 3 percent are included in New Haven County. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Shallow depth to bedrock is the main limitation for dwellings with basements. Where possible, dwellings with basements should be constructed in a deeper inclusion or nearby soil. This unit has few limitations for lawns and landscaping.

Shallow depth to bedrock is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the soil is not thick enough to adequately filter effluent. A more suitable site should be considered in a deeper inclusion or nearby soil.

Shallow depth to bedrock is the main limitation for local roads and streets. Careful planning of grades and road locations will avoid some removal of rock.

269C—Yalesville-Urban land complex, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: bedrock-controlled ridges on uplands, bedrock-controlled hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Yalesville and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Yalesville and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 25 inches; loam

C—25 to 36 inches; gravelly sandy loam

2R—36 to 80 inches; unweathered bedrock

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: moderately deep or deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from basalt and/or sandstone and shale

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Yalesville and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Holyoke soils, well drained Cheshire and Wethersfield soils, and moderately well drained Watchaug and Ludlow soils. Holyoke soils are in areas where the bedrock is between 10 and 20 inches deep and Wethersfield soils are in areas where the substratum is very firm. Watchaug and Ludlow soils are on slightly lower areas. Also included are areas with a silt loam surface texture in Middlesex and New Haven counties. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Shallow depth to bedrock is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Where possible, dwellings with basements should be constructed in a deeper inclusion or nearby soil. Slope is the main limitation for lawns and landscaping.

Shallow depth to bedrock is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the soil is not thick enough to adequately filter effluent. A more suitable site should be considered in a deeper inclusion or nearby soil.

Shallow depth to bedrock and slope are the main limitations for local roads and streets. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation.

273C—Urban land-Charlton-Chatfield complex, rocky, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: ridges, hills, uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Urban land and similar soils: 35 percent

Charlton and similar soils: 25 percent

Chatfield and similar soils: 15 percent

Minor components: 25 percent

Major Components

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa— 0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Chatfield and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Urban land and similar soils

Land capability classification (non-irrigated): 8

Charlton and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Chatfield and similar soils*Land capability classification (non-irrigated): 3e**Hydrologic group: B****Minor Components***

Included with this soil in mapping are areas of somewhat excessively drained Hollis soils, moderately well drained Sutton soils and poorly drained Leicester soils. Hollis soils are in areas where the bedrock is between 10 and 20 inches deep. Sutton soils are on slightly lower areas and Leicester soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components and rock outcrops make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Shallow depths to bedrock in areas of Chatfield soils and rock outcroppings are the main limitations for dwellings with basements. Slope is also a limitation in areas of Charlton soils. Erosion is a slight to moderate hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper inclusion or nearby soil.

Slope is the main limitation for lawns and landscaping. Large stones are also a limitation in areas of Charlton soils and the thin soil layer are also limitations in areas of Chatfield soils. Droughtiness can make establishment and maintenance of lawns difficult. Addition of fill material and removing the stones will reduce these limitations.

Slope in areas of Charlton soils is the main limitation for septic tank absorption fields. Depths to bedrock in areas of Chatfield soils and rock outcroppings over portions of the landscape are also limitations. There is a hazard of groundwater pollution in areas of Chatfield soils because the soil layer is not thick enough to adequately filter effluent. A more suitable site should be considered in a deeper inclusion or nearby soil. Placing distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Depths to bedrock and frost action are also limitations in areas of Chatfield soils. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

273E—Urban land-Charlton-Chatfield complex, rocky, 15 to 45 percent slopes***Map Unit Setting****Slope:* moderately steep to steep*Landscape:* ridges, hills, uplands*Size of map unit:* Areas commonly range from 3 to 500 acres.***Map Unit Composition****Urban land and similar soils:* 35 percent*Charlton and similar soils:* 25 percent*Chatfield and similar soils:* 15 percent*Minor components:* 25 percent

Major Components

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Charlton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 4 inches; fine sandy loam

Bw1—4 to 7 inches; fine sandy loam

Bw2—7 to 19 inches; fine sandy loam

Bw3—19 to 27 inches; gravelly fine sandy loam

C—27 to 65 inches; gravelly fine sandy loam

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Charlton and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Chatfield and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Urban land and similar soils

Land capability classification (non-irrigated): 8

Charlton and similar soils

Land capability classification (non-irrigated): 6e

Hydrologic group: B

Chatfield and similar soils*Land capability classification (non-irrigated): 6e**Hydrologic group: B****Minor Components***

Included with this soil in mapping are areas of somewhat excessively drained Hollis soils, moderately well drained Sutton soils and poorly drained Leicester soils. Hollis soils are in areas where the bedrock is between 10 and 20 inches deep. Sutton soils are on slightly lower areas and Leicester soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components and rock outcrop make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Shallow depth to bedrock in areas of Chatfield soils and rock outcroppings is also a limitation for dwellings with basements. Erosion is a severe or very severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper inclusion, in a less sloping inclusion, or nearby soil.

Slope is the main limitation for lawns and landscaping. Large stones are also a limitation in areas of Charlton soils and the thin soil layer are also limitations in areas of Chatfield soils. Droughtiness can make establishment and maintenance of lawns difficult. Addition of fill material and removing the stones will reduce these limitations.

Slope in areas of Charlton soils is the main limitation for septic tank absorption fields. Depth to bedrock in areas of Chatfield soils and rock outcroppings over portions of the landscape are also limitations. There is a hazard of groundwater pollution in areas of Chatfield soils because the soil layer is not thick enough to adequately filter effluent. A more suitable site should be considered in a deeper inclusion or nearby soil. Placing distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Depths to bedrock and frost action are also limitations in areas of Chatfield soils. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

275C—Urban land-Chatfield complex, rocky, 3 to 15 percent slopes***Map Unit Setting****Slope:* gently sloping to strongly sloping*Landscape:* ridges, uplands, hills*Size of map unit:* Areas commonly range from 3 to 500 acres.***Map Unit Composition****Urban land and similar soils:* 45 percent*Chatfield and similar soils:* 30 percent*Minor components:* 25 percent

Major Components

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Urban land and similar soils

Land capability classification (non-irrigated): 8

Chatfield and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of somewhat excessively drained Hollis soils, well drained Charlton soils, moderately well drained Sutton soils and poorly drained Leicester soils. Hollis soils are in areas where the bedrock is between 10 and 20 inches deep and Charlton soils are very deep. Sutton soils are on slightly lower areas and Leicester soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components and rock outcrops make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Shallow depth to bedrock and rock outcroppings are the main limitations for dwellings with basements. Slope is also a limitation in areas of Charlton soils. Erosion is a slight to moderate hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper inclusion or nearby soil.

Slope and a thin soil layer are the main limitations for lawns and landscaping. Droughtiness can make establishment and maintenance of lawns difficult. Addition of fill material will reduce these limitations.

Depth to bedrock and rock outcroppings are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the soil layer is not thick enough to adequately filter effluent. A more suitable site should be considered in a deeper inclusion or nearby soil.

Depth to bedrock and frost action are the main limitations for local roads and streets. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

275E—Urban land-Chatfield-Rock outcrop complex, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep or steep

Landscape: hills, ridges, uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Urban land and similar soils: 35 percent

Chatfield and similar soils: 25 percent

Rock outcrop and similar soils: 15 percent

Minor components: 25 percent

Major Components

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Chatfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 6 inches; gravelly fine sandy loam

Bw1—6 to 15 inches; gravelly fine sandy loam

Bw2—15 to 29 inches; gravelly fine sandy loam

2R—29 to 80 inches; unweathered bedrock

Major Component Properties and Qualities

Depth to bedrock: moderately deep or deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Urban land and similar soils

Land capability classification (non-irrigated): 8

Chatfield and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of somewhat excessively drained Hollis soils, moderately well drained Sutton soils and poorly drained Leicester soils. Hollis soils are in areas where the bedrock is between 10 and 20 inches deep. Sutton soils are on slightly lower areas and Leicester soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components and rock outcrop make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Shallow depth to bedrock and rock outcroppings is also a limitation for dwellings with basements. Erosion is a severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper inclusion, in a less sloping inclusion, or nearby soil.

Slope, depth to bedrock, and rock outcroppings are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the soil layer is not thick enough to adequately filter effluent. A more suitable site should be considered in a deeper inclusion or nearby soil. Placing distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

282B—Broadbrook-Urban land complex, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Broadbrook and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Broadbrook and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bw1—8 to 14 inches; silt loam

Bw2—14 to 25 inches; silt loam
 2Cd—25 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: eolian deposits over coarse-loamy lodgment till derived from gneiss and/or schist and/or sandstone and/or basalt

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Broadbrook and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Holyoke soils, well drained Narragansett and Wethersfield soils, moderately well drained Rainbow soils, poorly drained Wilbraham soils, and very poorly drained Menlo soils. Holyoke soils are in areas where the bedrock is between 10 and 20 inches deep. Narragansett soils developed on very friable to firm sandy till and Wethersfield soils are 7.5YR or redder in the B horizon. Rainbow soils are on slightly lower areas. Wilbraham and Menlo soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components and rock outcrop make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. This unit has few limitations for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines will allow on site sewage disposal in many places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade will reduce these limitations.

284B—Paxton-Urban land complex, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, till plains on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 85 acres.

Map Unit Composition

Paxton and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Stockbridge soils, moderately well drained Woodbridge soils and poorly drained Ridgebury soils. Canton, Charlton, and Stockbridge soils are in areas lacking a firm substratum. Woodbridge soils are on slightly lower areas and Ridgebury soils are in depressions and along drainageways. Soils with a redder color in the substratum are

included in Hartford, Middlesex, and New Haven counties. Udorthents are included adjacent to buildings and other structures. Minor componets make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. This unit has few limitations for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines will allow on site sewage disposal in many places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade will reduce these limitations.

284C—Paxton-Urban land complex, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: drumlins on uplands, hills on uplands, till plains on uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Paxton and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid
Depth to restrictive feature: 20 to 40 inches to densic material
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 3e
Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Stockbridge soils, moderately well drained Woodbridge soils and poorly drained Ridgebury soils. Canton, Charlton, and Stockbridge soils are in areas lacking a firm substratum. Woodbridge soils are on slightly lower areas and Ridgebury soils are in depressions and along drainageways. Soils with a redder color in the substratum are included in Hartford, Middlesex, and New Haven counties. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table and slope are the main limitations for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Erosion is a moderate hazard during construction. Slope is the main limitation for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines will allow on site sewage disposal in many places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, constructing roads on the contour, and providing a coarse grained subgrade will reduce these limitations.

284D—Paxton-Urban land complex, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep
Landscape: drumlins on uplands, hills on uplands, till plains on uplands
Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Paxton and similar soils: 40 percent
Urban land and similar soils: 35 percent
Minor components: 25 percent

Major Components

Paxton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 15 inches; fine sandy loam

Bw2—15 to 26 inches; fine sandy loam

Cd—26 to 65 inches; gravelly fine sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Paxton and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Canton, Charlton, and Stockbridge soils, moderately well drained Woodbridge soils and poorly drained Ridgebury soils. Canton, Charlton, and Stockbridge soils are in areas lacking a firm substratum. Woodbridge soils are on slightly lower areas and Ridgebury soils are in depressions and along drainageways. Soils with a redder color in the substratum are included in Hartford, Middlesex, and New Haven counties. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements. Designing dwellings to conform to the slope of the land will reduce the slope limitation. Erosion is a severe hazard during construction. Slope is the main limitation for lawns and landscaping. A site should be selected on a less sloping portion of the unit or nearby soil.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum may allow on site sewage

disposal. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

287B—Wethersfield-Urban land complex, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Wethersfield and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Wethersfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Wethersfield and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Cheshire and Yalesville soils, moderately well drained Ludlow soils, poorly drained Wilbraham soils and very poorly drained Menlo soils. Cheshire soils have a friable substratum and Yalesville soils are moderately deep to bedrock. Ludlow soils are on slightly lower areas. Wilbraham and Menlo soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. This unit has few limitations for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines will allow on site sewage disposal in many places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade will reduce these limitations.

287C—Wethersfield-Urban land complex, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Wethersfield and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Wethersfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Wethersfield and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Cheshire and Yalesville soils, moderately well drained Ludlow soils, poorly drained Wilbraham soils and very poorly drained Menlo soils. Cheshire soils have a friable substratum and Yalesville soils are moderately deep to bedrock. Ludlow soils are on slightly lower areas. Wilbraham and Menlo soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

The seasonal high water table and slope are the main limitations for dwellings with basements. Erosion is a moderate hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Slope is the main limitation for lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines will allow on site sewage disposal in many places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, constructing roads on the contour, and providing a coarse grained subgrade will reduce these limitations.

287D—Wethersfield-Urban land complex, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Wethersfield and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Wethersfield and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 3 inches; loam

Bw1—3 to 13 inches; loam

Bw2—13 to 27 inches; gravelly loam

Cd—27 to 65 inches; gravelly loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from basalt and/or sandstone and shale

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: 20 to 40 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Wethersfield and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: C

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Cheshire and Yalesville soils, moderately well drained Ludlow soils, poorly drained Wilbraham soils and very poorly drained Menlo soils. Cheshire soils have a friable substratum and Yalesville soils are moderately deep to bedrock. Ludlow soils are on slightly lower areas. Wilbraham and Menlo soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements. Erosion is a severe hazard during construction. Designing dwellings to conform to the slope of the land will reduce this limitation. This unit has few limitations for lawns and landscaping.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce this limitation.

290B—Stockbridge-Urban land complex, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Stockbridge and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Stockbridge and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Stockbridge and similar soils

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Farmington, well drained Nellis and Paxton soils, moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Farmington soils are in areas where the underlying limestone bedrock is between 10 and 20 inches below the surface. Paxton soils have a dense substratum and are more acid. Nellis soils are in areas where the soil is calcareous within 40 inches. Georgia soils are on slightly lower areas. Mudgepond and Alden soils are in depressions and along drainageways. Also included in Litchfield County are some areas with slopes less than 3 percent. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

This unit has few limitations for dwellings with basements and lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

290C—Stockbridge-Urban land complex, 8 to 15 percent slopes**Map Unit Setting**

Slope: strongly sloping

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Stockbridge and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components**Stockbridge and similar soils**

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Stockbridge and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Farmington, well drained Nellis and Paxton soils, moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Farmington soils are in areas where the underlying limestone bedrock is between 10 and 20 inches below the surface. Paxton soils have a dense substratum and are more acid. Nellis soils are in areas where the soil is calcareous within 40 inches. Georgia soils are on slightly lower areas. Mudgepond and Alden soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines will allow on site sewage disposal.

Frost action and slope are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth and constructing roads on the contour will reduce these limitations.

290D—Stockbridge-Urban land complex, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Stockbridge and similar soils: 40 percent

Urban land and similar soils: 35 percent

Minor components: 25 percent

Major Components

Stockbridge and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 10 inches; loam

Bw1—10 to 20 inches; loam

Bw2—20 to 28 inches; loam

C1—28 to 42 inches; gravelly loam

C2—42 to 48 inches; gravelly loam

C3—48 to 65 inches; gravelly loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Stockbridge and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this soil in mapping are areas of well drained Farmington, well drained Nellis and Paxton soils, moderately well drained Georgia soils, poorly drained Mudgepond soils, and very poorly drained Alden soils. Farmington soils are in areas where the underlying limestone bedrock is between 10 and 20 inches below the surface. Paxton soils have a dense substratum and are more acid. Nellis soils are in areas where the soil is calcareous within 40 inches. Georgia soils are on slightly lower areas. Mudgepond and Alden soils are in depressions and along drainageways. Udorthents are included adjacent to buildings and other structures. Minor components make up about 25 percent of the unit.

Use and Management

This unit is in urban and suburban development. The open areas between structures are in lawns, gardens, woodland, or brushland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. A site should be selected on a less sloping portion of the unit or nearby site.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional septic system by extending the length of distribution lines and adding fill may allow on site sewage disposal. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce these limitations.

301—Beaches-Udipsamments complex, coastal

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: beaches on coastal plains

Size of map unit: Areas are long and narrow and range from 3 to 100 acres.

Map Unit Composition

Beaches and similar soils: 50 percent

Udipsamments and similar soils: 35 percent

Minor components: 15 percent

Major Components

Beaches and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

C—0 to 65 inches; gravelly sand

Udipsamments and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

C1—0 to 38 inches; sand

C2—38 to 50 inches; coarse sand

C3—50 to 65 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: beach sand

Permeability: rapid or very rapid

Available water capacity: low

Reaction: moderately acid to neutral

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 48 to 72 inches

Flooding: occasional

Interpretative Groups

Beaches and similar soils

Land capability classification (non-irrigated): 8

Udipsamments and similar soils*Land capability classification (non-irrigated): 3s**Hydrologic group: A****Minor Components***

Included with this complex in mapping are small areas of very poorly drained Westbrook, Pawcatuck, and Ipswich soils in tidal marshes. Also included are areas of Udorthents, urbanized areas, and a few areas of rock outcrop. Udorthents are in areas that have been altered by cutting and filling. Minor components make up about 15 percent of this map unit.

Use and Management

This complex is poorly suited for most uses other than for recreation. On site investigation is required to determine the suitability for proposed uses.

Coastal flooding is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. The high water table is also a limitation for septic tank adsorption fields. A more suitable site should be selected on a drier soil that does not flood. Coastal flooding is the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade will reduce this limitation. A more suitable site should be considered on a soil that does not flood during coastal storms.

302—Dumps***Map Unit Setting****Slope: nearly level to strongly sloping**Landscape: None assigned**Size of map unit: Areas commonly range from 3 to 100 acres.****Map Unit Composition****Dumps and similar soils: 95 percent**Minor components: 5 percent****Major Components***

The typical sequence, depth, and composition of the layers of the soil are as follows—

C—0 to 65 inches; variable

Interpretative Groups*Land capability classification (non-irrigated): 8****Minor Components***

Included in mapping are small areas of Westbrook soils and small areas of Udorthents. Westbrook soils are in very poorly drained tidal marshes. Udorthents are soils that have been altered by cutting and filling. In a few small dumps there are rock outcrops. A few dumps along the larger streams are subject to flooding. Minor components make up about 5 percent of this map unit.

Use and Management

This unit is in both active and inactive landfills.

Dumps require onsite investigation and evaluation if considered for other uses. An important item to consider is the leachate and from the land fill and its environmental impacts.

303—Pits, Quarries

Map Unit Setting

Slope: nearly level to very steep

Landscape: hills, ridges, uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Pits, quarries and similar soils: 90 percent

Minor components: 10 percent

Major Components

Pits, quarries and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

R—0 to 1 inch; unweathered bedrock

Interpretative Groups

Land capability classification (non-irrigated): 8

Minor Components

Included in mapping are small areas of somewhat excessively drained Hollis soils, well drained Holyoke soils and Farmington soils where bedrock is 10 to 20 inches below the surface. bedrock outcrops and cliff faces are also common. Minor components make up 10 percent of this map unit.

Use and Management

This unit is in both active and inactive quarries. Pits and quarries require onsite investigation and evaluation if considered for other uses.

304—Udorthents, loamy, very steep

Map Unit Setting

Slope: steep to very steep

Landscape: terrace escarpments on lake plains

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Udorthents and similar soils: 90 percent

Minor components: 10 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; loam

C1—5 to 21 inches; gravelly loam

C2—21 to 80 inches; very gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: glaciolacustrine deposits
Permeability: very slow to very rapid
Available water capacity: high
Reaction: very strongly acid to slightly alkaline
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 54 to 72 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7e
Hydrologic group: B

Minor Components

Included with this unit in mapping are small areas of very poorly drained Maybid soils and poorly drained Shaker, Scitico, and Raynam soils in depressions and along drainageways. Soils subject to flooding are included along streams in highly dissected areas. Exscesively drained Windsor soils are included on the upper slopes of some map uints. Minor componets make up about 10 percent of the unit.

Use and Management

Most areas are in woodland or brushland. Some areas are in pasture or community developement.

Steep slopes are the main limitation for dwellings with basements and lawns and landscaping. slippage is also a limitation for dwellings with basements. Erosion is a very severe hazard during construction. A site should be selected on a nearby soil that is less sloping.

Slope, slow percolation and slippage are the main limmitations for septic tank adsorption fields. Modifying a conventional system by extending the the length of the distribution lines along the contour and adding fill my allow for onsite sewage disposal. A more suitable site should be considered in a less sloping soil.

Slope, frost action and slippage are the main limitations for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitations. Providing a coarse grained sugrade to the frost depth will reduce the frost limitation.

305—Udorthents-Pits complex, gravelly

Map Unit Setting

Slope: nearly level to steep
Landscape: sand pits, gravel pits
Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Udorthents and similar soils: 65 percent
Pits and similar soils: 25 percent
Minor components: 10 percent

Major Components

Udorthents and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; loam
 C1—5 to 21 inches; gravelly loam

C2—21 to 80 inches; very gravelly sandy loam

Pits and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

C—0 to 65 inches; very gravelly sand

Major Component Properties and Qualities

Udorthents and similar soils

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: gravelly outwash

Permeability: very slow to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 24 to 54 inches

Flooding: none

Interpretative Groups

Udorthents and similar soils

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Pits and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of undisturbed soils. These are excessively drained Hinckley and Windsor soils, somewhat excessively drained Merrimac and Gloucester soils, and moderately well drained Ninigret and Sudbury soils. Also included are small bodies of water in areas that were excavated below the ground water table. Minor components make up 10 percent of the mapunit.

Use and Management

Most areas are in active sand and gravel mines. Some areas are in abandon sand and gravel mines which have reverted back to woodland and or brushland.

The characteristics of this unit are so variable that an onsite investigation is required to determine the suitability for proposed uses.

306—Udorthents-Urban land complex

Map Unit Setting

Slope: nearly level to moderately steep

Landscape: urban lands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Udorthents and similar soils: 50 percent

Urban land and similar soils: 35 percent

Minor components: 15 percent

Major Components

Udorthents and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; loam

C1—5 to 21 inches; gravelly loam

C2—21 to 80 inches; very gravelly sandy loam

Urban land and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Major Component Properties and Qualities

Udorthents and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: drift

Permeability: very slow to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 54 to 72 inches

Flooding: none

Interpretative Groups

Udorthents and similar soils

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Urban land and similar soils

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of udorthents with a wet substratum, which were formerly poorly drained and very poorly drained soils. Also included are areas of undisturbed soils and rock outcrop. Undisturbed soils are in areas between buildings and structures. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in urban and built up areas. Some areas are in adjacent open land.

The characteristics of this unit are so variable that an onsite investigation is required to determine the suitability for proposed uses.

307—Urban Land

Map Unit Setting

Slope: nearly level to steep

Landscape: none assigned

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Urban land and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

H—0 to 6 inches; material

Interpretative Groups

Land capability classification (non-irrigated): 8

Minor Components

Included with this unit in mapping are areas of udorthents with a wet substratum, which were formerly poorly drained and very poorly drained soils. Also included are areas of undisturbed soils and rock outcrop. Undisturbed soils are in areas between buildings and structures. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in urban and built up areas. Some areas are in adjacent open land.

The characteristics of this unit are so variable that an onsite investigation is required to determine the suitability for proposed uses.

308—Udorthents, smoothed

Map Unit Setting

Slope: nearly level to steep

Landscape: leveled lands, fills

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Udorthents and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; loam

C1—5 to 21 inches; gravelly loam

C2—21 to 80 inches; very gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: not specified

Permeability: very slow to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 24 to 54 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of udorthents which have a wet substratum, urban land, rock outcrop, and undisturbed soils. Rock outcrop is in areas which have been cut, exposing bedrock. Undisturbed soils are in areas where the fill is very thin. Inclusions having a wet substratum are filled areas which were formerly poorly drained and very poorly drained. Also included are reclaimed sand and gravel pits and the soil is dominated by sand and gravel in these areas. Minor components make up 20 percent of the unit.

Use and Management

Most areas are used for recreation, some areas are used for cropland, hayland, pasture or community development.

The characteristics of this unit are so variable that an onsite investigation is required to determine the suitability for proposed uses.

309—Udorthents, flood control

Map Unit Setting

Slope: nearly level to steep

Landscape: artificial levees

Size of map unit: Areas commonly range from 3 to 300 acres.

Map Unit Composition

Udorthents and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; loam

C1—5 to 21 inches; gravelly loam

C2—21 to 80 inches; very gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: drift

Permeability: very slow to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 24 to 54 inches

Flooding: very rare

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of udorthents which have a wet substratum, Typic Udifluents, urban land, rock outcrop, and undisturbed soils. Rock outcrop is in areas which have been cut, exposing bedrock. Undisturbed soils are in areas where the fill is very thin. Inclusions having a wet substratum are filled areas which were formerly poorly drained and very poorly drained. Minor components make up 20 percent of the unit.

Use and Management

Most areas are in dams, levees, channels or other flood control structures.

310—Udorthents, periodically flooded

Map Unit Setting

Slope: nearly level to steep

Landscape: river valleys, valleys

Size of map unit: Areas commonly range from 50 to 500 acres in size.

Map Unit Composition

Udorthents and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 5 inches; loam

C1—5 to 21 inches; gravelly loam

C2—21 to 80 inches; very gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: drift

Permeability: very slow to very rapid

Available water capacity: high

Reaction: very strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 24 to 54 inches

Flooding: very rare

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with this unit in mapping are areas of udorthents which have a wet substratum, Typic Udifluents, urban land, rock outcrop, and undisturbed soils. Rock outcrop is in areas which have been cut, exposing bedrock. Undisturbed soils are in areas where the fill is very thin. Inclusions having a wet substratum are filled areas which were formerly poorly drained and very poorly drained. Minor components make up 20 percent of the unit.

Use and Management

Most areas are in flood control impoundments.

The characteristics of this unit are so variable that an onsite investigation is required to determine the suitability for proposed uses.

401C—Macomber-Taconic complex, 3 to 15 percent slopes, very rocky

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock-controlled hills on uplands, bedrock-controlled ridges on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Macomber and similar soils: 55 percent

Taconic and similar soils: 30 percent

Minor components: 15 percent

Major Components

Macomber and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material

A—1 to 2 inches; very channery loam

Bw1—2 to 10 inches; very channery loam

Bw2—10 to 21 inches; very channery loam

C—21 to 30 inches; very channery loam

2R—30 to 80 inches; bedrock

Taconic and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 4 inches; very gravelly loam

Bw—4 to 11 inches; very gravelly loam

2R—11 to 80 inches; bedrock

Major Component Properties and Qualities

Macomber and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: loamy skeletal melt-out till derived from phyllite and/or schist

Permeability: moderate

Available water capacity: low

Reaction: very strongly acid to strongly acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Taconic and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from phyllite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to strongly acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Macomber and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Taconic and similar soils

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of well drained Dummerston and Lanesboro soils, and very poorly drained Wonsqueak and Bucksport soils. Dummerston soils are very deep. Lanesboro soils are very deep and have a dense substratum. Wonsqueak and Bucksport soils are organic soils in depressions. Also included are areas of rock outcrop and steeper slopes. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Shallow depth to bedrock and many rock outcrops in areas of Taconic soils are the main limitations for dwellings with basements and lawns and landscaping. The short, uneven slopes are also a limitation. Erosion is a moderate to severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper, less sloping inclusion or nearby soil. Droughtiness can make establishment and maintenance of lawns difficult. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Shallow depth to bedrock and many rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a deeper inclusion or nearby soil.

Shallow depth to bedrock and many rock outcrops are the main limitations for local roads and streets. Frost action is also a limitation. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

402D—Taconic-Macomber-Rock outcrop complex, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: bedrock-controlled ridges on uplands, bedrock-controlled hills on uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Macomber and similar soils: 50 percent
Taconic and similar soils: 25 percent
Rock outcrop and similar soils: 15 percent
Minor components: 10 percent

Major Components

Macomber and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; highly decomposed plant material
 A—1 to 2 inches; very channery loam
 Bw1—2 to 10 inches; very channery loam
 Bw2—10 to 21 inches; very channery loam
 C—21 to 30 inches; very channery loam
 2R—30 to 80 inches; bedrock

Taconic and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material
 A—1 to 4 inches; very gravelly loam
 Bw—4 to 11 inches; very gravelly loam
 2R—11 to 80 inches; bedrock

Major Component Properties and Qualities

Macomber and similar soils

Depth to bedrock: moderately deep to deep
Drainage class: well drained
Parent material: loamy skeletal melt-out till derived from phyllite and/or schist
Permeability: moderate
Available water capacity: low
Reaction: very strongly acid to strongly acid
Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)
Depth to seasonal water table: greater than 6 feet
Flooding: none

Taconic and similar soils

Depth to bedrock: shallow to moderately deep
Drainage class: somewhat excessively drained
Parent material: loamy melt-out till derived from phyllite and/or schist
Permeability: moderate or moderately rapid
Available water capacity: very low
Reaction: very strongly acid to strongly acid
Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Macomber and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: C

Taconic and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of well drained Dummerston and Lanesboro soils. Dummerston soils are very deep. Lanesboro soils are very deep and have a dense substratum. Also include are small areas of less slope. Minor components make up about 10 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Shallow depth to bedrock and many rock outcrops are also limitations. Erosion is a very severe hazard during construction. Additional fill will reduce the depth limitation. A more suitable site should be selected on a less sloping, deeper portion of the unit or nearby soil.

Slope, shallow depth to bedrock, and many rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a less sloping, deeper inclusion or nearby soil.

Slope and many rock outcrops are the main limitations for local roads and streets. Shallow depth to bedrock is also a limitation. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

403C—Taconic-Rock outcrop complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock-controlled ridges, uplands, bedrock-controlled hills

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Taconic and similar soils: 70 percent

Rock outcrop and similar soils: 25 percent

Minor components: 5 percent

Major Components

Taconic and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 4 inches; very gravelly loam

Bw—4 to 11 inches; very gravelly loam

2R—11 to 80 inches; bedrock

Major Component Properties and Qualities

Taconic and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from phyllite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to strongly acid
Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Taconic and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of well drained Macomber. Macomber soils are moderately deep with bedrock between 20 to 40 inches. Also included are small areas of steeper slopes. areas Minor componets up about 5 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Shallow depth to bedrock and many rock outcrops in areas of Taconic soils are the main limitations for dwellings with basements and lawns and landscaping. The short, uneven slopes are also a limitation. Erosion is a moderate to severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in a deeper, less sloping inclusion or nearby soil. Droughtiness can make establishment and maintenance of lawns difficult. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Shallow depth to bedrock and many rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a deeper inclusion or nearby soil.

Shallow depth to bedrock and many rock outcrops are the main limitations for local roads and streets. Frost action is also a limitation. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

403E—Taconic-Rock outcrop complex, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep to steep
Landscape: uplands, bedrock-controlled hills, bedrock-controlled ridges
Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Taconic and similar soils: 70 percent
Rock outcrop and similar soils: 20 percent
Minor components: 10 percent

Major Components

Taconic and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 4 inches; very gravelly loam

Bw—4 to 11 inches; very gravelly loam

2R—11 to 80 inches; bedrock

Major Component Properties and Qualities

Taconic and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from phyllite and/or schist

Permeability: moderate to moderately rapid

Available water capacity: very low

Reaction: very strongly acid to strongly acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Taconic and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of well drained Macomber and Dummerston soils. Macomber soils are moderately deep with bedrock between 20 and 40 inches and Dummerston soils are very deep. Also included are small areas of less sloping land. Minor components make up about 10 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Shallow depth to bedrock and many rock outcrops are also limitations. Erosion is a very severe hazard during construction. Additional fill will reduce the depth limitation. A more suitable site should be selected on a less sloping, deeper portion of the unit or nearby soil.

Slope, shallow depth to bedrock, and many rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a less sloping, deeper inclusion or nearby soil.

Slope and many rock outcrops are the main limitations for local roads and streets. Shallow depth to bedrock is also a limitation. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

403F—Taconic-Rock outcrop complex, 45 to 70 percent slopes

Map Unit Setting

Slope: very steep

Landscape: bedrock-controlled ridges, bedrock-controlled hills, uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Taconic and similar soils: 70 percent

Rock outcrop and similar soils: 20 percent

Minor components: 10 percent

Major Components

Taconic and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 4 inches; very gravelly loam

Bw—4 to 11 inches; very gravelly loam

2R—11 to 80 inches; bedrock

Major Component Properties and Qualities

Taconic and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from phyllite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: very low

Reaction: very strongly acid to strongly acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Taconic and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of well drained Macomber and Dummerston soils. Macomber soils are moderately deep with bedrock between 20 and 40 inches and Dummerston soils are very deep. Also included are small areas of less sloping land. Minor components make up about 10 percent of the map unit.

Use and Management

Most areas are in woodland.

Many rock outcrops, slope, and shallow depth to bedrock are the main limitations for dwellings with basements and lawns and landscaping. Erosion is a very severe hazard during construction. Addition of fill will reduce the depth limitation. A more suitable site should be selected in a deeper, less sloping inclusion or nearby soil.

Many rock outcrops, slope, and shallow depth to bedrock are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a less sloping, deeper inclusion or nearby soil.

Shallow depth to bedrock, slope, and many rock outcrops are the main limitations for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

405C—Dummerston gravelly loam, 3 to 15 percent slopes, very stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: uplands, hills

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Dummerston and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 2 inches; gravelly loam

E—2 to 3 inches; gravelly loam

Bs—3 to 4 inches; gravelly loam

Bw1—4 to 6 inches; gravelly loam

Bw2—6 to 11 inches; gravelly loam

Bw3—11 to 22 inches; gravelly loam

BC—22 to 27 inches; gravelly loam

C1—27 to 40 inches; very gravelly loam

C2—40 to 64 inches; very stony loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from phyllite and/or schist

Permeability: moderate

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of well drained Lanesboro soils, moderately well drained Fullam soils, and poorly drained Brayton soils. Macomber soils are moderately deep, between 20 and 40 inches to bedrock. Lanesboro, Fullam, and Brayton soils have a dense substratum. Also included are small areas of steeper slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the

land will reduce the slope limitation. Large stones are a limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slope is the main limitation for septic tank absorption fields. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

405E—Dummerston gravelly loam, 15 to 45 percent slopes, very stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: hills, uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Dummerston and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 2 inches; gravelly loam

E—2 to 3 inches; gravelly loam

Bs—3 to 4 inches; gravelly loam

Bw1—4 to 6 inches; gravelly loam

Bw2—6 to 11 inches; gravelly loam

Bw3—11 to 22 inches; gravelly loam

BC—22 to 27 inches; gravelly loam

C1—27 to 40 inches; very gravelly loam

C2—40 to 64 inches; very stony loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from phyllite and/or schist

Permeability: moderate

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of well drained Lanesboro soils and moderately well drained Fullam soils. Lanesboro, Fullam and Brayton soils have a dense substratum. Fullam soils are slight depressions, Brayton soils are in depressions and drainageways. Also included are small areas of lesser slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope is the main limitation for septic tank absorption fields. Placing the distribution lines on the contour increases the efficiency of the system. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

407C—Lanesboro loam, 3 to 15 percent slopes, very stony

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Lanesboro and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 3 inches; highly decomposed plant material

A—3 to 6 inches; loam

Bw1—6 to 8 inches; loam

Bw2—8 to 16 inches; channery loam

Bw3—16 to 22 inches; channery loam

BC—22 to 30 inches; channery loam

Cd—30 to 60 inches; very channery loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from phyllite and/or schist

Permeability: very slow to moderate

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 35 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Dummerston soils, moderately well drained Fullam soils, and poorly drained Brayton soils. Dummerston soils are in areas that lack a dense substratum. Fullam soils are in slight depressions below Lanesboro soils and Brayton soils are in depressions and drainageways. Also included are small areas of steeper slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland or brushland reverting to woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements. Slope is also a limitation. Erosion is a moderate hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Large stones and slope are the main limitations for lawns and landscaping. Removing the stones will reduce this limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation. Constructing roads on the contour will reduce the slope limitation.

407E—Lanesboro loam, 15 to 45 percent slopes, very stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Lanesboro and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 3 inches; highly decomposed plant material

A—3 to 6 inches; loam

Bw1—6 to 8 inches; loam

Bw2—8 to 16 inches; channery loam

Bw3—16 to 22 inches; channery loam
 BC—22 to 30 inches; channery loam
 Cd—30 to 60 inches; very channery loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy lodgment till derived from phyllite and/or schist
Permeability: very slow to moderate
Available water capacity: moderate
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: 20 to 35 inches to densic material
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s
Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Macomber and Dummerston soils, and moderately well drained Fullam soils. Macomber soils are moderately deep, between 20 and 40 inches to bedrock. Dummerston soils are in areas that lack a dense substratum and Fullam soils are in slight depressions below Lanesboro soils. Also included are small areas of lesser slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Designing dwellings to conform to the slope of the land will reduce the slope limitation. A site should be selected on a less sloping portion of the unit or nearby soil.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less dense inclusion or nearby soil.

Slope is the main limitations for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

408C—Fullam silt loam, 3 to 15 percent slopes, very stony

Map Unit Setting

Slope: gently sloping to strongly sloping
Landscape: hills, uplands
Surface cover: 0 to 3 percent stones
Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Fullam and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 2 inches; moderately decomposed plant material

A—2 to 4 inches; silt loam

Bw1—4 to 10 inches; silt loam

Bw2—10 to 20 inches; gravelly loam

Cd1—20 to 49 inches; very channery loam

Cd2—49 to 60 inches; very channery fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from phyllite and/or schist

Permeability: very slow to moderate

Available water capacity: low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Macomber, Lanesboro, and Dummerston soils, and poorly drained Brayton soils. Macomber soils are moderately deep, between 20 and 40 inches to bedrock. Dummerston soils are in areas that lack a dense substratum and Lanesboro soils are higher on the landscape. Brayton soils are in depressions and drainageways. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Slope and large stones are also limitations for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

409B—Brayton mucky silt loam, 0 to 8 percent slopes, very stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Brayton and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 3 inches; moderately decomposed plant material

Oa—3 to 6 inches; highly decomposed plant material

A—6 to 7 inches; mucky silt loam

Bg1—7 to 9 inches; silt loam

Bg2—9 to 13 inches; gravelly loam

Cd1—13 to 18 inches; gravelly fine sandy loam

Cd2—18 to 23 inches; gravelly fine sandy loam

Cd3—23 to 60 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy lodgment till derived from phyllite and/or schist

Permeability: very slow to moderately rapid

Available water capacity: very low

Reaction: extremely acid to neutral

Depth to restrictive feature: 10 to 20 inches to densic material

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately well drained Fullam soils and very poorly drained Wonsqueak and Bucksport soils. Fullam soils are in slightly higher areas on the landscape. Wonsqueak and Bucksport soils are organic soils in depressions. Minor components make up about 15 percent of the map unit.

Use and Management

This soil is mostly in woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil. The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill

materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

412B—Bice fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Bice and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 7 inches; fine sandy loam

Bw1—7 to 16 inches; fine sandy loam

Bw2—16 to 24 inches; gravelly fine sandy loam

C—24 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of somewhat excessively drained Westminster, well drained Millsite and Shelburne soils, and moderately well drained Schroon soils. Schroon soils are in slight depressions and Shelburne soils have a dense substratum. Millsite soils are where bedrock is between 20 to 40 inches below the surface and Westminster soils are where bedrock is 10 to 20 inches below the surface. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland or residential development. Some areas are in pasture.

This unit has few limitations for dwellings with basements. Large stones are a limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Bice soils have few limitations for septic tank absorption fields. This unit has few limitations for local roads and streets.

412C—Bice fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Bice and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 7 inches; fine sandy loam

Bw1—7 to 16 inches; fine sandy loam

Bw2—16 to 24 inches; gravelly fine sandy loam

C—24 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of somewhat excessively drained Westminster, well drained Millsite and Shelburne soils, and moderately well drained Schroon soils. Schroon soils are in slight depressions and Shelburne soils have a dense substratum. Millsite soils are where bedrock is between 20 to 40 inches below the surface and Westminster soils are where bedrock is 10 to 20 inches below the surface. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland or residential development. Some areas are in pasture.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the

land will reduce the slope limitation. Large stones are a limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slope is the main limitation for septic tank absorption fields. Placing the distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

412D—Bice fine sandy loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: hills, uplands

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Bice and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 7 inches; fine sandy loam

Bw1—7 to 16 inches; fine sandy loam

Bw2—16 to 24 inches; gravelly fine sandy loam

C—24 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of somewhat excessively drained Westminster, well drained Millsite and Shelburne soils, and moderately well drained Schroon soils. Schroon soils are in slight depressions and Shelburne soils have a dense substratum. Millsite soils are where bedrock is between 20 to 40 inches below the surface and Westminster soils are where bedrock is 10 to 20 inches below the surface. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or residential development.

Slope is the main limitations for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope is the main limitation for septic tank absorption fields. Placing the distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

413C—Bice-Millsite complex, 3 to 15 percent slopes, very rocky

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: bedrock-controlled hills, bedrock-controlled uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Bice and similar soils: 45 percent

Millsite and similar soils: 40 percent

Minor components: 15 percent

Major Components

Bice and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 7 inches; fine sandy loam

Bw1—7 to 16 inches; fine sandy loam

Bw2—16 to 24 inches; gravelly fine sandy loam

C—24 to 60 inches; gravelly sandy loam

Millsite and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 5 inches; fine sandy loam

Bw1—5 to 13 inches; stony fine sandy loam

Bw2—13 to 24 inches; fine sandy loam

C—24 to 31 inches; sandy loam

2R—31 to 80 inches; bedrock

Major Component Properties and Qualities

Bice and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Millsite and similar soils

Depth to bedrock: moderately deep to deep
Drainage class: well drained
Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss
Permeability: moderate or moderately rapid
Available water capacity: moderate
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Bice and similar soils

Land capability classification (non-irrigated): 6s
Hydrologic group: B

Millsite and similar soils

Land capability classification (non-irrigated): 6s
Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of rock outcrops, somewhat excessively drained Westminster, and moderately well drained Schroon soils. Schroon soils are in slight depressions and areas of shallow Westminster soils are where bedrock is 10 to 20 inches below the surface. Also included are small areas of steeper slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland or residential development. Some areas are in pasture.

Depth to bedrock is the main limitation for dwellings with basements. Slope is also a limitation. Erosion is a moderate hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in areas of very deep Bice soils.

Slope is the main limitation for lawns and landscaping. Large rocks are a limitation in areas of Bice soils, and the thin soil layer is a limitation in areas of Millsite soils. Droughtiness can make establishment and maintenance of lawns difficult. Addition of fill material and removing the rocks will reduce these limitations.

Slope is the main limitation for septic tank absorption fields in areas of Bice soils. Depth to bedrock in areas of Millsite soils and Rock outcrops over portions of the landscape are also limitations. Where possible, septic tank absorption fields should be constructed in areas of very deep Bice soils. Placing distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Depths to bedrock and frost action are limitations in areas of Millsite soils. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock. Providing a coarse grained subgrade will reduce frost action.

413E—Bice-Millsite complex, 15 to 45 percent slopes, very rocky

Map Unit Setting

Slope: moderately steep to steep

Landscape: bedrock-controlled uplands, bedrock-controlled hills

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Bice and similar soils: 45 percent

Millsite and similar soils: 40 percent

Minor components: 15 percent

Major Components

Bice and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 7 inches; fine sandy loam

Bw1—7 to 16 inches; fine sandy loam

Bw2—16 to 24 inches; gravelly fine sandy loam

C—24 to 60 inches; gravelly sandy loam

Millsite and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 5 inches; fine sandy loam

Bw1—5 to 13 inches; stony fine sandy loam

Bw2—13 to 24 inches; fine sandy loam

C—24 to 31 inches; sandy loam

2R—31 to 80 inches; bedrock

Major Component Properties and Qualities

Bice and similar soils

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Millsite and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Bice and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Millsite and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of somewhat excessively drained Westminster and rock outcrops. Areas of shallow Westminster soils are where bedrock is 10 to 20 inches below the surface. Also included are small areas of lesser slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope and depth to bedrock in areas of Millsite soils are the main limitations for dwellings with basements. Erosion is a severe or very severe hazard during construction. Uneven slopes and variable depth to bedrock reduce site selection. Where possible, dwellings with basements should be constructed in areas of very deep Bice soils, a less sloping inclusion, or nearby soil.

Slope is the main limitation for lawns and landscaping. A site should be selected on a less sloping portion of the unit or nearby soil.

Slope is the main limitation for septic tank absorption fields in areas of Bice soils. Depth to bedrock in areas of Millsite soils is also a limitation. Where possible, septic tank absorption fields should be constructed in areas of very deep Bice soils on a less sloping portion of the unit. Placing distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

414—Fredon silt loam, cold

Map Unit Setting

Slope: nearly level

Landscape: depressions on outwash plains, drainageways on outwash plains, terraces on outwash plains

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Fredon and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; silt loam

Bg—8 to 17 inches; fine sandy loam
 Bw—17 to 24 inches; fine sandy loam
 2Cg1—24 to 29 inches; stratified gravelly sand to loamy fine sand
 2C—29 to 48 inches; stratified gravelly sand to loamy fine sand
 2Cg2—48 to 60 inches; stratified gravelly sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: poorly drained
Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist
Permeability: moderate to very rapid
Available water capacity: high
Reaction: moderately acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 0 to 12 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w
Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils, and cold phases of well drained Agawam and moderately well drained Ninigret soils that are higher on the landscape. Cold phases of very poorly drained Halsey soils are included in depressions and along drainageways. Minor components up 15 percent of this map unit.

Use and Management

Most areas are in woodland. Cleared areas are in pasture or cropland. Some cleared areas are drained.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Poor filtering is also a limitation for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Frost action and seasonal highwater table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

415C—Westminster-Millsite-Rock outcrop complex, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping
Landscape: uplands, bedrock controlled ridges, bedrock controlled hills
Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Westminster and similar soils: 40 percent
Millsite and similar soils: 40 percent

Rock outcrop and similar soils: 15 percent

Minor components: 5 percent

Major Components

Westminster and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

A—2 to 5 inches; fine sandy loam

Bw1—5 to 12 inches; fine sandy loam

Bw2—12 to 16 inches; fine sandy loam

2R—16 to 80 inches; bedrock

Millsite and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 5 inches; fine sandy loam

Bw1—5 to 13 inches; stony fine sandy loam

Bw2—13 to 24 inches; fine sandy loam

C—24 to 31 inches; sandy loam

2R—31 to 80 inches; bedrock

Major Component Properties and Qualities

Westminster and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid

Available water capacity: very low

Reaction: extremely acid to strongly acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Millsite and similar soils

Depth to bedrock: moderately deep to deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Westminster and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Millsite and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of well drained Bice soils. Very deep Bice soils are adjacent to Westminster and Millsite soils. Also included are small areas of steeper slopes. Minor components make up about 5 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Shallow depth to bedrock and many rock outcrops are the main limitations for dwellings with basements and lawns and landscaping. Uneven slopes are also a limitation. Erosion is a moderate to severe hazard during construction. Where possible, dwellings with basements should be constructed in a deeper, less sloping inclusion or nearby soil. Droughtiness can make establishment and maintenance of lawns difficult. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Shallow depth to bedrock and many rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a deeper inclusion or nearby soil.

Shallow depth to bedrock and many rock outcrops are the main limitations for local roads and streets. Frost action is also a limitation. Slope is also a limitation in steeper areas. Careful planning of grades and road locations will avoid some removal of rock. Constructing roads on the contour will reduce the slope limitation. Providing a coarse grained subgrade will reduce frost action.

415E—Westminster-Millsite-Rock outcrop complex, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep or steep

Landscape: uplands, bedrock-controlled ridges, bedrock-controlled hills

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Westminster and similar soils: 40 percent

Millsite and similar soils: 40 percent

Rock outcrop and similar soils: 15 percent

Minor components: 5 percent

Major Components

Westminster and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

A—2 to 5 inches; fine sandy loam

Bw1—5 to 12 inches; fine sandy loam

Bw2—12 to 16 inches; fine sandy loam

2R—16 to 80 inches; bedrock

Millsite and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe— 0 to 1 inch; moderately decomposed plant material

A—1 to 5 inches; fine sandy loam

Bw1—5 to 13 inches; stony fine sandy loam

Bw2—13 to 24 inches; fine sandy loam

C—24 to 31 inches; sandy loam

2R—31 to 80 inches; bedrock

Major Component Properties and Qualities**Westminster and similar soils**

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid

Available water capacity: very low

Reaction: extremely acid to strongly acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Millsite and similar soils

Depth to bedrock: moderately deep or deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: 20 to 40 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups**Westminster and similar soils**

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Millsite and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of well drained Bice soils. Very deep Bice soils are adjacent to Westminster and Millsite soils. Also included are small areas of lesser slopes. Minor components make up about 5 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Shallow depth to bedrock and many rock outcrops are also limitations. Erosion is a very severe hazard during construction. Additional fill will reduce the depth limitation. A more suitable site should be selected on a less sloping, deeper portion of the unit, or nearby soil.

Slope, shallow depth to bedrock, and many Rock outcrops are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a less sloping, deeper inclusion, or nearby soil.

Slope and many Rock outcrops are the main limitations for local roads and streets. Shallow depth to bedrock is also a limitation. Constructing roads on the contour will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

416E—Rock outcrop-Westminster complex, 8 to 45 percent slopes

Map Unit Setting

Slope: strongly sloping to steep

Landscape: bedrock-controlled ridges, bedrock-controlled hills, bedrock-controlled uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Rock outcrop and similar soils: 70 percent

Westminster and similar soils: 20 percent

Minor components: 10 percent

Major Components

Westminster and similar soils:

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

A—2 to 5 inches; fine sandy loam

Bw1—5 to 12 inches; fine sandy loam

Bw2—12 to 16 inches; fine sandy loam

2R—16 to 80 inches; bedrock

Major Component Properties and Qualities

Westminster and similar soils:

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid

Available water capacity: very low

Reaction: extremely acid to strongly acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Westminster and similar soils:

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of well drained Millsite soil and well drained Bice soils. Very deep Bice soils and moderately deep Millsite soils are adjacent to Westminster soils. Also included are small areas of lesser slopes. Minor components make up about 10 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Many rock outcrops and shallow depth to bedrock are the main limitations for dwellings with basements and lawns and landscaping. Slope is also a main limitation in steeper areas of the unit. Uneven slopes and variable depth to bedrock reduce site selection. Erosion is a severe to very severe hazard during construction. A more suitable site should be selected in a deeper, less sloping inclusion or nearby soil.

Shallow depth to bedrock and many rock outcrops are the main limitations for septic tank absorption fields. Slope is also a main limitation in steeper areas. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. Placing septic tank absorption field distribution lines on the contour increases the efficiency of the system. A more suitable site should be selected in a less sloping, deeper inclusion or nearby soil.

Shallow depth to bedrock and many rock outcrops are the main limitations for local roads and streets. Slope is also a limitation in steeper areas. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

416F—Rock outcrop-Westminster complex, 45 to 70 percent slopes

Map Unit Setting

Slope: very steep

Landscape: bedrock-controlled ridges, bedrock-controlled hills, bedrock-controlled uplands

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Rock outcrop and similar soils: 70 percent

Westminster and similar soils: 20 percent

Minor components: 10 percent

Major Components

Westminster and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

A—2 to 5 inches; fine sandy loam

Bw1—5 to 12 inches; fine sandy loam

Bw2—12 to 16 inches; fine sandy loam

2R—16 to 80 inches; bedrock

Major Component Properties and Qualities

Westminster and similar soils

Depth to bedrock: shallow to moderately deep

Drainage class: somewhat excessively drained

Parent material: loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderately rapid

Available water capacity: very low

Reaction: extremely acid to strongly acid

Depth to restrictive feature: 10 to 20 inches to bedrock (lithic)

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Westminster and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of well drained Millsite soil and well drained Bice soils. Very deep Bice soils and moderately deep Millsite soils are adjacent to Westminster soils. Also included are small areas of lesser slopes. Minor components make up about 10 percent of the map unit.

Use and Management

Most areas are in woodland.

Many rock outcrops, slope, and shallow depth to bedrock are the main limitations for dwellings with basements and lawns and landscaping. Erosion is a very severe hazard during construction. Addition of fill will reduce the depth limitation. A more suitable site should be selected in a deeper, less sloping inclusion or nearby soil.

Many rock outcrops, slope, and shallow depth to bedrock are the main limitations for septic tank absorption fields. There is the hazard of groundwater pollution because the soil is not thick enough to filter effluent. A more suitable site should be selected in a less sloping, deeper inclusion, or nearby soil.

Shallow depth to bedrock, slope, and many rock outcrops are the main limitations for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation. Careful planning of grades and road locations will avoid some removal of rock.

417B—Bice fine sandy loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: hills, uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Bice and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material
 A—1 to 7 inches; fine sandy loam
 Bw1—7 to 16 inches; fine sandy loam
 Bw2—16 to 24 inches; gravelly fine sandy loam
 C—24 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss
Permeability: moderate or moderately rapid
Available water capacity: high
Reaction: very strongly acid to moderately acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: greater than 6 feet
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s
Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of somewhat excessively drained Westminster, well drained Millsite and Shelburne soils, and moderately well drained Schroon soils. Schroon soils are in slight depressions and Shelburne soils have a dense substratum. Millsite soils are where bedrock is between 20 to 40 inches below the surface and Westminster soils are where bedrock is 10 to 20 inches below the surface. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland or residential development. Some areas are in pasture.

This unit has few limitations for dwellings with basements. Large stones are a limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Bice soils have few limitations for septic tank absorption fields. This unit has few limitations for local roads and streets.

417C—Bice fine sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping
Landscape: hills, uplands
Surface cover: 0 to 3 percent stones
Size of map unit: Areas commonly range from 3 to 500 acres.

Map Unit Composition

Bice and similar soils: 85 percent
Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

O_i—0 to 1 inch; slightly decomposed plant material

A—1 to 7 inches; fine sandy loam

Bw₁—7 to 16 inches; fine sandy loam

Bw₂—16 to 24 inches; gravelly fine sandy loam

C—24 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of somewhat excessively drained Westminster, well drained Millsite and Shelburne soils, and moderately well drained Schroon soils. Schroon soils are in slight depressions and Shelburne soils have a dense substratum. Millsite soils are where bedrock is between 20 to 40 inches below the surface and Westminster soils are where bedrock is 10 to 20 inches below the surface. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland or residential development. Some areas are in pasture.

Slope is the main limitation for dwellings with basements. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Large stones are a limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slope is the main limitation for septic tank absorption fields. Placing the distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

417D—Bice fine sandy loam, 15 to 25 percent slopes, very stony

Map Unit Setting

Slope: moderately steep

Landscape: uplands, hills

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Bice and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

O_i—0 to 1 inch; slightly decomposed plant material

A—1 to 7 inches; fine sandy loam

Bw₁—7 to 16 inches; fine sandy loam

Bw₂—16 to 24 inches; gravelly fine sandy loam

C—24 to 60 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of somewhat excessively drained Westminster, well drained Millsite and Shelburne soils, and moderately well drained Schroon soils. Schroon soils are in slight depressions and Shelburne soils have a dense substratum. Millsite soils are where bedrock is between 20 to 40 inches below the surface and Westminster soils are where bedrock is 10 to 20 inches below the surface. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in residential development.

Slope is the main limitations for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slope is the main limitation for septic tank absorption fields. Placing the distribution lines on the contour increases the efficiency of the system.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

418C—Schroon fine sandy loam, 2 to 15 percent slopes, very stony

Map Unit Setting

Slope: nearly level to strongly sloping

Landscape: hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Schroon and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material

A—3 to 9 inches; fine sandy loam

Bw1—9 to 14 inches; fine sandy loam

Bw2—14 to 23 inches; fine sandy loam

Bw3—23 to 30 inches; sandy loam

C —30 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: extremely acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield, and poorly drained Brayton soils and very poorly drained Loonmeadow soils. Brayton and Loonmeadow soils are in depressions and drainageways. Ashfield soils have a dense substrata and Bice soils are in higher positions on the landscape. Also included are small areas of lesser slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development and pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large stones and slope are also limitations for lawns and landscaping. Removing the stones and designing lawns to conform to the slope of the land will reduce these limitations. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

420A—Schroon fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Schroon and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material

A—3 to 9 inches; fine sandy loam

Bw1—9 to 14 inches; fine sandy loam

Bw2—14 to 23 inches; fine sandy loam

Bw3—23 to 30 inches; sandy loam

C—30 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: extremely acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield, and poorly drained Brayton soils and very poorly drained Loonmeadow soils. Brayton and Loonmeadow soils are in depressions and drainageways. Ashfield soils have a dense substrata and Bice soils are in higher positions on the landscape. Also included are small areas of lesser slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development and pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Slope is also a limitation for lawns and landscaping. Designing lawns to conform to the slope of the land will reduce this limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

420B—Schroon fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: hills on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Schroon and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

O_i—0 to 1 inch; slightly decomposed plant material

O_e—1 to 2 inches; moderately decomposed plant material

O_a—2 to 3 inches; highly decomposed plant material

A—3 to 9 inches; fine sandy loam

Bw₁—9 to 14 inches; fine sandy loam

Bw₂—14 to 23 inches; fine sandy loam

Bw₃—23 to 30 inches; sandy loam

C—30 to 60 inches; sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy melt-out till derived from granite and/or schist and/or gneiss

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: extremely acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w

Hydrologic group: B

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield, and poorly drained Brayton soils and very poorly drained Loonmeadow soils. Brayton and Loonmeadow soils are in depressions and drainageways. Ashfield soils have a dense substrata and Bice soils are in higher positions on the landscape. Also included are small areas of lesser slopes. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development and pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Slope is also a limitation for lawns and landscaping. Designing lawns to conform to the slope of the land will reduce this limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill usually will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

421A—Ninigret fine sandy loam, cold, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ninigret and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 16 inches; fine sandy loam

Bw2—16 to 26 inches; fine sandy loam

2C—26 to 65 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderate to very rapid
Available water capacity: high
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 30 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils, and cold phases of somewhat excessively drained Merrimac and well drained Agawam soils that are higher on the landscape. Also included are cold phases of moderately well drained Sudbury soils. Small areas of poorly drained Moosilauke soils and cold phases of very poorly drained Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are cleared and in cultivated crops, hay, pasture, or brushland. Some areas are in woodland or community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce the wetness.

Poor filtering and the seasonal high water table are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal. There is also a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

423A—Sudbury sandy loam, cold, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level
Landscape: terraces on valleys, outwash plains on valleys
Size of map unit: Areas range from 3 to 40 acres.

Map Unit Composition

Sudbury and similar soils: 85 percent
Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material
 A—1 to 5 inches; sandy loam

Bw1—5 to 17 inches; gravelly sandy loam
Bw2—17 to 25 inches; sandy loam
2C—25 to 60 inches; stratified gravel to sand

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss
Permeability: moderately rapid to very rapid
Available water capacity: moderate
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 18 to 36 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2w
Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils, cold phases of somewhat excessively drained Merrimac and well drained Agawam soils that are higher on the landscape. Also included are cold phases of moderately well drained Ninigret soils. Small areas of poorly drained Moosilauke soils and cold phases of very poorly drained Scarborough soils are included in shallow depressions and drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in cultivated cropland, hay, or pasture. Some areas are in woodland or community development.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Modifying a conventional septic system by extending the length of distribution lines and adding fill usually will allow on site sewage disposal. Specially designed septic systems are necessary in some areas of Sudbury soils.

The seasonal high water table and frost action are the main limitations for local roads and streets. Constructing roads on raised fill materials and installing a drainage system will reduce the wetness limitation. Providing a coarse grained subgrade to frost depth will reduce the limitation.

424B—Shelburne fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping
Landscape: hills on uplands, drumlins on uplands
Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Shelburne and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; fine sandy loam

Bw1—2 to 7 inches; fine sandy loam

Bw2—7 to 21 inches; gravelly fine sandy loam

Bw3—21 to 27 inches; bouldery fine sandy loam

Cd1—27 to 32 inches; gravelly fine sandy loam

Cd2—32 to 43 inches; fine sandy loam

Cd3—43 to 55 inches; fine sandy loam

Cd4—55 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield soils, poorly drained brayton soils and very poorly drained Loonmeadow soils. Bice soils have a more permeable substratum. Ashfield soils are in slightly lower areas and Brayton and Loonmeadow soils are in depressions and drainageways. Also included are small areas of steeper slopes. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or hay and pasture.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

424C—Shelburne fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands, drumlins on uplands

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Shelburne and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; fine sandy loam

Bw1—2 to 7 inches; fine sandy loam

Bw2—7 to 21 inches; gravelly fine sandy loam

Bw3—21 to 27 inches; bouldery fine sandy loam

Cd1—27 to 32 inches; gravelly fine sandy loam

Cd2—32 to 43 inches; fine sandy loam

Cd3—43 to 55 inches; fine sandy loam

Cd4—55 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield soils, poorly drained brayton soils and very poorly drained Loonmeadow soils. Bice soils have a more permeable substratum. Ashfield soils are in slightly lower areas and Brayton and Loonmeadow soils are in depressions and drainageways. Also included are small areas of steeper slopes. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or hay and pasture.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with

gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

424D—Shelburne fine sandy loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: drumlins on uplands, hills on uplands

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Shelburne and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; fine sandy loam

Bw1—2 to 7 inches; fine sandy loam

Bw2—7 to 21 inches; gravelly fine sandy loam

Bw3—21 to 27 inches; bouldery fine sandy loam

Cd1—27 to 32 inches; gravelly fine sandy loam

Cd2—32 to 43 inches; fine sandy loam

Cd3—43 to 55 inches; fine sandy loam

Cd4—55 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield soils, poorly drained brayton soils and very poorly drained Loonmeadow soils. Bice soils have a more permeable substratum. Ashfield soils are in slightly lower areas and Brayton and Loonmeadow soils are in depressions and drainageways. Also included are small areas of steeper slopes. Minor componets make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or hay and pasture.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

425B—Shelburne fine sandy loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Shelburne and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; fine sandy loam

Bw1—2 to 7 inches; fine sandy loam

Bw2— 7 to 21 inches; gravelly fine sandy loam

Bw3—21 to 27 inches; bouldery fine sandy loam

Cd1—27 to 32 inches; gravelly fine sandy loam

Cd2—32 to 43 inches; fine sandy loam

Cd3—43 to 55 inches; fine sandy loam

Cd4—55 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield soils, poorly drained brayton soils and very poorly drained Loonmeadow soils. Bice soils have a more permeable substratum. Ashfield soils are in slightly lower areas and Brayton and Loonmeadow soils are in depressions and drainageways. Also included are small areas of steeper slopes. Minor componets make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Small and large stones are the main limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will reduce the frost action limitation.

425C—Shelburne fine sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: hills on uplands, drumlins on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 75 acres.

Map Unit Composition

Shelburne and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; fine sandy loam

Bw1—2 to 7 inches; fine sandy loam

Bw2—7 to 21 inches; gravelly fine sandy loam

Bw3—21 to 27 inches; bouldery fine sandy loam

Cd1—27 to 32 inches; gravelly fine sandy loam

Cd2—32 to 43 inches; fine sandy loam

Cd3—43 to 55 inches; fine sandy loam

Cd4—55 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Bice soils, moderately well drained Ashfield soils, poorly drained brayton soils and very poorly drained Loonmeadow soils. Bice soils have a more permeable substratum. Ashfield soils are in slightly lower areas and Brayton and Loonmeadow soils are in depressions and drainageways. Also included are small areas of steeper slopes. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements. Slope is also a limitation in areas of Paxton soil. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Large stones and slope are the main limitations for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places.

The seasonal high water table, slope, and frost action are the main limitations for local roads and streets. Construction on raised fill materials and installing a drainage system will reduce wetness. Providing a coarse grained subgrade to frost depth will

reduce the frost action limitation. Constructing roads on the contour will reduce the slope limitation.

426D—Shelburne fine sandy loam, 15 to 35 percent slopes, extremely stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: hills on uplands, drumlins on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Shelburne and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; fine sandy loam

Bw1—2 to 7 inches; fine sandy loam

Bw2—7 to 21 inches; gravelly fine sandy loam

Bw3—21 to 27 inches; bouldery fine sandy loam

Cd1—27 to 32 inches; gravelly fine sandy loam

Cd2—32 to 43 inches; fine sandy loam

Cd3—43 to 55 inches; fine sandy loam

Cd4—55 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: 20 to 30 inches to densic material

Depth to seasonal water table: 18 to 30 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: C

Minor Components

Included with these soils in mapping are areas of well drained Bice soils and moderately well drained Ashfield soils. Bice soils have a more permeable substratum and Ashfield soils are in slightly lower areas. Also included are areas of lesser slopes. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in community development.

Slope is the main limitation for dwellings with basements. Erosion is a severe hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation. A site should be selected on a less sloping portion of the unit or nearby soil.

Slope is the main limitation for lawns and landscaping.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum will allow on site sewage disposal in most places. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less dense inclusion or nearby soil.

Slope is the main limitations for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

427B—Ashfield fine sandy loam, 2 to 8 percent slopes, very stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: drumlins on uplands, hills on uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ashfield and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material

A1—3 to 7 inches; fine sandy loam

A2—7 to 12 inches; fine sandy loam

Bw1—12 to 18 inches; fine sandy loam

Bw2—18 to 24 inches; fine sandy loam

BC—24 to 29 inches; fine sandy loam

Cd1—29 to 44 inches; fine sandy loam

Cd2—44 to 58 inches; sandy loam

Cd3—58 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: high

Reaction: extremely acid to slightly acid

Depth to restrictive feature: 20 to 33 inches to densic material

Depth to seasonal water table: 12 to 24 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Shelburne soils, moderately well drained Schroon soils, poorly drained Brayton soils and very poorly drained Loonmeadow soils. Shelburne soils are higher on the landscape. Schroon soils lack a dense substratum. Brayton and Loonmeadow soils are in depressions and along drainageways. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

427C—Ashfield fine sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: hills, uplands, drumlins

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Ashfield and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material
 A1—3 to 7 inches; fine sandy loam
 A2—7 to 12 inches; fine sandy loam
 Bw1—12 to 18 inches; fine sandy loam
 Bw2—18 to 24 inches; fine sandy loam
 BC—24 to 29 inches; fine sandy loam
 Cd1—29 to 44 inches; fine sandy loam
 Cd2—44 to 58 inches; sandy loam
 Cd3—58 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss
Permeability: very slow to moderately rapid
Available water capacity: moderate
Reaction: extremely acid to slightly acid
Depth to restrictive feature: 20 to 33 inches to densic material
Depth to seasonal water table: 12 to 24 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s
Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Shelburne soils, moderately well drained Schroon soils, poorly drained Brayton soils and very poorly drained Loonmeadow soils. Shelburne soils are higher on the landscape. Schroon soils lack a dense substratum. Brayton and Loonmeadow soils are in depressions and along drainageways. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Slope and large stones are also limitations for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

428A—Ashfield fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: uplands, drumlins, hills

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ashfield and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material

A1—3 to 7 inches; fine sandy loam

A2—7 to 12 inches; fine sandy loam

Bw1—12 to 18 inches; fine sandy loam

Bw2—18 to 24 inches; fine sandy loam

BC—24 to 29 inches; fine sandy loam

Cd1—29 to 44 inches; fine sandy loam

Cd2—44 to 58 inches; sandy loam

Cd3—58 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: extremely acid to slightly acid

Depth to restrictive feature: 20 to 33 inches to densic material

Depth to seasonal water table: 12 to 24 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Shelburne soils, moderately well drained Schroon soils, poorly drained Brayton soils and very poorly drained Loonmeadow soils. Shelburne soils are higher on the landscape. Schroon soils lack a dense substratum. Brayton and Loonmeadow soils are in depressions and along drainageways. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or hay and pasture.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping.

Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

428B—Ashfield fine sandy loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: drumlins, hills, uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Ashfield and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi— 0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material

A1—3 to 7 inches; fine sandy loam

A2—7 to 12 inches; fine sandy loam

Bw1—12 to 18 inches; fine sandy loam

Bw2—18 to 24 inches; fine sandy loam

BC—24 to 29 inches; fine sandy loam

Cd1—29 to 44 inches; fine sandy loam

Cd2—44 to 58 inches; sandy loam

Cd3—58 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: extremely acid to slightly acid

Depth to restrictive feature: 20 to 33 inches to densic material

Depth to seasonal water table: 12 to 24 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Shelburne soils, moderately well drained Schroon soils, poorly drained Brayton soils and very poorly drained Loonmeadow soils. Shelburne soils are higher on the landscape. Schroon soils lack a dense substratum. Brayton and Loonmeadow soils are in depressions and along drainageways. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or hay and pasture.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping.

Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

428C—Ashfield fine sandy loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: hills, drumlins, uplands

Size of map unit: Areas commonly range from 3 to 60 acres.

Map Unit Composition

Ashfield and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material

A1—3 to 7 inches; fine sandy loam

A2—7 to 12 inches; fine sandy loam

Bw1—12 to 18 inches; fine sandy loam

Bw2—18 to 24 inches; fine sandy loam

BC—24 to 29 inches; fine sandy loam

Cd1—29 to 44 inches; fine sandy loam

Cd2—44 to 58 inches; sandy loam
 Cd3—58 to 80 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: moderately well drained
Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss
Permeability: very slow to moderately rapid
Available water capacity: moderate
Reaction: extremely acid to slightly acid
Depth to restrictive feature: 20 to 33 inches to densic material
Depth to seasonal water table: 12 to 24 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e
Hydrologic group: C

Minor Components

Included with this soil in mapping are areas of well drained Shelburne soils, moderately well drained Schroon soils, poorly drained Brayton soils and very poorly drained Loonmeadow soils. Shelburne soils are higher on the landscape. Schroon soils lack a dense substratum. Brayton and Loonmeadow soils are in depressions and along drainageways. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas are in woodland. Some areas are in community development or hay and pasture.

The seasonal high water table is the main limitation if this map unit is used as a site for dwellings with basements or lawns and landscaping. Slope is also a limitation for lawns and landscaping. Locating dwellings in the highest part of the map unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

Wetness and slow percolation are the main limitations if this map unit is used as a site for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill above the impermeable substratum may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation if this map unit is used as a site for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

429A—Agawam fine sandy loam, cold, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level
Landscape: terraces on valleys, outwash plains on valleys
Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Agawam and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 24 inches; fine sandy loam

2C— 24 to 60 inches; stratified very gravelly coarse sand to fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils and cold phases of somewhat excessively drained Merrimac soils. Boscawen Soils are sandy and gravelly throughout; cold phases of Merrimac soils are sandy throughout. Also included are cold phases of moderately well drained Sudbury soils that are sandy and gravelly throughout and cold phases of moderately well drained Ninigret soils that are loamy over sandy and gravelly. Small areas poorly drained Moosilauke soils and cold phases of very poorly drained Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture hayland or woodland. Some areas are in community development or cropland.

This soil has few limitations for dwellings with basements and lawns and landscaping.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

This soil has few limitations for local roads and streets.

429B—Agawam fine sandy loam, cold, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Agawam and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils and cold phases of somewhat excessively drained Merrimac soils. Boscawen Soils are sandy and gravelly through out; cold phases of Merrimac soils are sandy throughout. Also included are cold phases of moderately well drained Sudbury soils that are sandy and gravelly throughout and cold phases of moderately well drained Ninigret soils that are loamy over sandy and gravelly. Small areas poorly drained Moosilauke soils and cold phases of very poorly drained Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture hayland or woodland. Some areas are in community development or cropland.

This soil has few limitations for dwellings with basements and lawns and landscaping. Droughtiness can make establishment and maintenance of lawns difficult.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

This soil has few limitations for local roads and streets.

429C—Agawam fine sandy loam, cold, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: terraces on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Agawam and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 8 inches; fine sandy loam

Bw1—8 to 14 inches; fine sandy loam

Bw2—14 to 24 inches; fine sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy eolian deposits over sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils and cold phases of somewhat excessively drained Merrimac soils. Boscawen Soils are sandy and gravelly through out; cold phases of Merrimac soils are sandy throughout. Also included are cold phases of moderately well drained Sudbury soils that are sandy and gravelly throughout and cold phases of moderately well drained Ninigret soils that are loamy over sandy and gravelly. Small areas poorly drained Moosilauke soils and cold phases of very poorly drained Scarboro soils are included in shallow depressions and drainageways. Minor componets make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture hayland or woodland. Some areas are in community development or cropland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

433—Moosilauke sandy loam

Map Unit Setting

Slope: nearly level

Landscape: valleys, depressions on outwash plains, drainageways on outwash plains, depressions on terraces, drainageways on terraces

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Moosilauke and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 1 inch; slightly decomposed plant material

A—1 to 6 inches; loam

Bg—6 to 16 inches; fine sandy loam

Bw—16 to 24 inches; gravelly fine sandy loam

C1—24 to 39 inches; loamy fine sand

C2—39 to 65 inches; fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils, and cold phases of somewhat excessively drained Merrimac, moderately well drained

Sudbury and Ninigret soils on slightly higher areas. Also included are cold phases of very poorly drained Scarboro soils, and very poorly drained Bucksport and Wonsqueak soils in the depressions. Cold phases of poorly drained Fredon soils are in areas that have a silty surface and subsoil with a higher pH. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland or shrubland. Cleared areas are in pasture.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Poor filtering is also a limitation for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Frost action and seasonal highwater table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

434A—Merrimac sandy loam, cold, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level

Landscape: kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 5 to 75 acres.

Map Unit Composition

Merrimac and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; sandy loam

Bw1—9 to 16 inches; sandy loam

Bw2—16 to 24 inches; gravelly sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 1

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils, and cold phases of well drained Agawam soils, moderately well drained Sudbury and Ninigret soils. Small areas poorly drained Moosilauke soils and cold phases of very poorly drained Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture hayland or woodland. Some areas are in community development or cropland.

This soil has few limitations for dwellings with basements, lawns and landscaping, and local roads and streets.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

434B—Merrimac sandy loam, cold, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: outwash plains on valleys, kames on valleys, terraces on valleys

Size of map unit: Areas commonly range from 5 to 50 acres.

Map Unit Composition

Merrimac and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; sandy loam

Bw1—9 to 16 inches; sandy loam

Bw2—16 to 24 inches; gravelly sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils and cold phases of well drained Agawam, moderately well drained Sudbury and Ninigret soils. Small areas of poorly drained Moosilauke soils and cold phases of very poorly drained Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture hayland or woodland. Some areas are in community development or cropland.

This soil has few limitations for dwellings with basements, lawns and landscaping, and local roads and streets.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

434C—Merrimac sandy loam, cold, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: kames on valleys, outwash plains on valleys, terraces on valleys

Size of map unit: Areas commonly range from 5 to 40 acres.

Map Unit Composition

Merrimac and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 9 inches; sandy loam

Bw1—9 to 16 inches; sandy loam

Bw2—16 to 24 inches; gravelly sandy loam

2C—24 to 60 inches; stratified very gravelly coarse sand to gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: somewhat excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: strongly acid to slightly acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils and cold phases of well drained Agawam, moderately well drained Sudbury and Ninigret soils. Small areas poorly drained Moosilauke soils and cold phases of very poorly drained Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in pasture hayland or woodland. Some areas are in community development or cropland.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the slope of the land will reduce the slope limitation.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

435—Scarboro muck, cold

Map Unit Setting

Slope: nearly level

Landscape: drainageways on outwash plains, terraces on outwash plains, depressions on outwash plains

Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Scarboro and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa—0 to 12 inches; muck

A—12 to 17 inches; loamy sand

Cg1—17 to 31 inches; stratified sand to loamy fine sand

Cg2—31 to 72 inches; stratified very gravelly coarse sand to loamy fine sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: moderate

Reaction: very strongly acid to neutral

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 6 inches above surface

Depth to seasonal water table: 0 to 6 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 5w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils, and cold phases of somewhat excessively drained Merrimac, moderately well drained Sudbury and Ninigret soils in higher areas. Also included are very poorly drained Bucksport and Wonsqeaq in the depressions. Cold phases of poorly drained Fredon soils are in areas that have a silty surface and subsoil with a higher pH. Minor components make up about 20 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are brushland reverting to woodland.

Ponding is the main limitation for dwellings with basements and lawns and landscaping. Ponding and Poor filtering are the main limitations for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Ponding and Frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

436—Halsey silt loam, cold

Map Unit Setting

Slope: nearly level

Landscape: terraces on outwash plains, drainageways on outwash plains, depressions on outwash plains

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Halsey and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 1 inch; moderately decomposed plant material

A—1 to 8 inches; silt loam

Bg1—8 to 16 inches; silt loam

Bg2—16 to 28 inches; fine sandy loam

2Cg1—28 to 38 inches; loamy sand

2Cg2—38 to 60 inches; sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: coarse-loamy over sandy and gravelly glaciofluvial deposits derived from limestone and dolomite and/or schist

Permeability: moderate to very rapid

Available water capacity: high

Reaction: moderately acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Ponding depth: 0 to 6 inches above surface
Depth to seasonal water table: 0 to 6 inches
Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 5w
Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of excessively drained Boscawen soils and cold phases of well drained Agawam and moderately well drained Ninigret soils in slightly higher areas. Cold phases of poorly drained Fredon soils are also included. Very poorly drained Bucksport and Wonsqueak soils are included in depressions and along drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland or marshland.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank adsorption fields. Poor filtering is also a limitation for septic tank adsorption fields. There is a hazard of groundwater pollution because of the rapidly permeable substratum does not adequately filter effluent. A more suitable site should be considered on a drier inclusion or a nearby soil.

Frost action and seasonal highwater table are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarser grained subgrade to frost depth will reduce these limitations.

437—Wonsqueak mucky peat

Map Unit Setting

Slope: nearly level
Landscape: depressions
Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Wonsqueak and similar soils: 85 percent
Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 2 inches; mucky peat
 Oa1—2 to 11 inches; muck
 Oa2—11 to 22 inches; muck
 2Cg1—22 to 25 inches; mucky silt loam
 2Cg2—25 to 45 inches; gravelly fine sandy loam
 2Cg3—45 to 60 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: very poorly drained

Parent material: woody organic material over loamy drift

Permeability: moderately slow to very rapid

Available water capacity: high

Reaction: extremely acid to neutral

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 12 inches above surface

Depth to seasonal water table: 0 to 12 inches

Flooding: rare

Interpretative Groups

Land capability classification (non-irrigated): 5w

Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of poorly drained Brayton soils and very poorly drained Bucksport and Loonmeadow soils. Bucksport soils are where the muck is more than 51 inches thick over mineral substratum. Brayton and Loonmeadow soils formed in coarse-loamy glacial till. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas of this soil are in woodland or wildlife habitat.

Ponding and subsidence are the main limitations for dwellings with basements, septic tank absorption fields, local roads and streets. Excess humus is a limitation for lawns and landscaping. Slow percolation is also a limitation for septic tank absorption fields. Frost action is also a limitation for local roads and streets. A more suitable site for all these uses should be selected on a drier soil.

438—Bucksport muck

Map Unit Setting

Slope: nearly level

Landscape: depressions

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Bucksport and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oa1—0 to 9 inches; muck

Oa2—9 to 33 inches; muck

Oa3—33 to 50 inches; muck

Oa4—50 to 59 inches; muck

2Cg—59 to 63 inches; gravelly sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: woody organic material

Permeability: moderately slow to very rapid
Available water capacity: very high
Reaction: extremely acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Ponding depth: 0 to 12 inches above surface
Depth to seasonal water table: 0 to 12 inches
Flooding: rare

Interpretative Groups

Land capability classification (non-irrigated): 5w
Hydrologic group: D

Minor Components

Included with this soil in mapping are areas of poorly drained Brayton soils, very poorly drained Loonmeadow and Wonsqueak soils. Wonsqueak soils are where the muck is between 16 to 51 inches thick over loamy substratum. Brayton and Loonmeadow soils formed in coarse-loamy glacial till. Minor components make up about 15 percent of the map unit.

Use and Management

Most areas of this soil are in woodland or wildlife habitat.

Ponding and subsidence are the main limitations for dwellings with basements, septic tank absorption fields, local roads and streets. Low strength is also a limitation for dwellings with basements. Excess humus is a limitation for lawns and landscaping. Slow percolation is also a limitation for septic tank absorption fields. Frost action is also a limitation for local roads and streets. A more suitable site for all these uses should be selected on a drier soil.

440A—Boscawen gravelly sandy loam, 0 to 3 percent slopes

Map Unit Setting

Slope: nearly level
Landscape: outwash plains on valleys, kames on valleys, eskers on valleys, terraces on valleys
Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Boscawen and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

O_i—0 to 1 inch; slightly decomposed plant material
 A—1 to 2 inches; gravelly sandy loam
 B_w—2 to 9 inches; very gravelly sandy loam
 B_C—9 to 16 inches; very gravelly loamy sand
 C₁—16 to 29 inches; stratified fine sand to extremely gravelly coarse sand
 C₂—29 to 34 inches; stratified fine sand to extremely gravelly coarse sand
 C₃—34 to 40 inches; stratified fine sand to extremely gravelly coarse sand
 C₄—40 to 44 inches; stratified fine sand to extremely gravelly coarse sand
 C₅—44 to 67 inches; stratified fine sand to extremely gravelly coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3s

Hydrologic group: A

Minor Components

Included in mapping are cold phases of somewhat excessively drained Merrimac soils and cold phases of well drained Agawam soils. Cold phases of Merrimac soils are sandy throughout and cold phases of Agawam soils are loamy over sand and gravel. Small areas of cold phases of moderately well drained Sudbury soils are included in slightly lower areas. Poorly drained Moosilauke soils and very poorly drained Bucksport, Wonsqueak and cold phases Scarborough soils are included in shallow depressions and drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or hayland. Some areas are mined for sand and gravel.

This soil has few limitations for dwellings with basements and local roads and streets. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

440C—Boscawen gravelly sandy loam, 3 to 15 percent slopes

Map Unit Setting

Slope: gently sloping to strongly sloping

Landscape: terraces on valleys, outwash plains on valleys, kames on valleys, eskers on valleys

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Boscawen and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; gravelly sandy loam

Bw—2 to 9 inches; very gravelly sandy loam

BC—9 to 16 inches; very gravelly loamy sand

C1—16 to 29 inches; stratified fine sand to extremely gravelly coarse sand

C2—29 to 34 inches; stratified fine sand to extremely gravelly coarse sand

C3—34 to 40 inches; stratified fine sand to extremely gravelly coarse sand

C4—40 to 44 inches; stratified fine sand to extremely gravelly coarse sand

C5—44 to 67 inches; stratified fine sand to extremely gravelly coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid to very rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: A

Minor Components

Included in mapping are cold phases of somewhat excessively drained Merrimac and well drained Agawam soils. Cold phases of Merrimac soils are sandy throughout and cold phases of Agawam soils are loamy over sand and gravel. Small areas of cold phases of moderately well drained Sudbury soils are included in slightly lower areas. Poorly drained Moosilauke soils and very poorly drained Bucksport, Wonsqueak and cold phases of Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or hayland. Some areas are mined for sand and gravel.

Slope is the main limitation for dwellings with basements. Erosion is a hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Poor filtering is the main limitation for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

440E—Boscawen gravelly sandy loam, 15 to 45 percent slopes

Map Unit Setting

Slope: moderately steep to steep

Landscape: terraces on valleys, eskers on valleys, kames on valleys, outwash plains on valleys

Size of map unit: Areas commonly range from 3 to 200 acres.

Map Unit Composition

Boscawen and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

A—1 to 2 inches; gravelly sandy loam

Bw—2 to 9 inches; very gravelly sandy loam

BC—9 to 16 inches; very gravelly loamy sand

C1—16 to 29 inches; stratified fine sand to extremely gravelly coarse sand

C2—29 to 34 inches; stratified fine sand to extremely gravelly coarse sand

C3—34 to 40 inches; stratified fine sand to extremely gravelly coarse sand

C4—40 to 44 inches; stratified fine sand to extremely gravelly coarse sand

C5—44 to 67 inches; stratified fine sand to extremely gravelly coarse sand

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: excessively drained

Parent material: sandy and gravelly glaciofluvial deposits derived from granite and/or schist and/or gneiss

Permeability: moderately rapid or very rapid

Available water capacity: very low

Reaction: very strongly acid to moderately acid

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6e

Hydrologic group: A

Minor Components

Included in mapping are cold phases of somewhat excessively drained Merrimac soils and well drained Agawam soils. Cold phases of Merrimac soils are sandy throughout and cold phase of Agawam soils are loamy over sand and gravel. Small areas of cold phases of moderately well drained Sudbury soils are included in slightly lower areas. Poorly drained Moosilauke soils and very poorly drained Bucksport, Wonsqueak and cold phases of Scarboro soils are included in shallow depressions and drainageways. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or hayland. Some areas are mined for sand and gravel.

Slope is the main limitation for dwellings with basements. Erosion is a hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation. Droughtiness and slope are the main limitations for lawns and landscaping. Planting early in spring reduces the impact of summer droughtiness and reduces seedling mortality. Lawns need watering in the summer.

Slope and poor filtering are the main limitations for septic tank absorption fields. There is a hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. Specially designed septic systems are necessary in some areas. Locating the septic tank absorption fields in a less sloping inclusion is preferable.

Slope is the main limitation for local roads and streets. Constructing roads on the contour will reduce the slope limitation.

442—Brayton loam

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Brayton and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 2 inches; moderately decomposed plant material

A—2 to 10 inches; loam

Bg1—10 to 17 inches; gravelly sandy loam

Bg2—17 to 22 inches; gravelly sandy loam

Bg3—22 to 27 inches; sandy loam

Cd1—27 to 42 inches; gravelly sandy loam

Cd2—42 to 65 inches; gravelly sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss

Permeability: very slow to moderately rapid

Available water capacity: moderate

Reaction: extremely acid to neutral

Depth to restrictive feature: 20 to 27 inches to densic material

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately well drained Fullam soils and very poorly drained Wonsqueak and Bucksport soils. Fullam soils are in slightly higher areas above Brayton soils. Wonsqueak and Bucksport soils are organic soils in depressions. Minor components make up about 15 percent of the map unit.

Use and Management

This soil is mostly in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil. The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

443—Brayton-Loonmeadow complex, extremely stony

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: drainageways on uplands, depressions on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Brayton and similar soils: 50 percent

Loonmeadow and similar soils: 35 percent

Minor components: 15 percent

Major Components

Brayton and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oe—0 to 2 inches; moderately decomposed plant material

A—2 to 10 inches; loam

Bg1—10 to 17 inches; gravelly sandy loam

Bg2—17 to 22 inches; gravelly sandy loam

Bg3—22 to 27 inches; sandy loam

Cd1—27 to 42 inches; gravelly sandy loam

Cd2—42 to 65 inches; gravelly sandy loam

Loonmeadow and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 2 inches; slightly decomposed plant material

A—2 to 9 inches; mucky fine sandy loam
 Bg—9 to 18 inches; sandy loam
 Cg1—18 to 35 inches; gravelly sandy loam
 Cg2—35 to 80 inches; gravelly sandy loam

Major Component Properties and Qualities

Brayton and similar soils

Depth to bedrock: very deep
Drainage class: poorly drained
Parent material: coarse-loamy lodgment till derived from granite and/or schist and/or gneiss
Permeability: very slow to moderately rapid
Available water capacity: moderate
Reaction: extremely acid to neutral
Depth to restrictive feature: 20 to 27 inches to densic material
Depth to seasonal water table: 0 to 12 inches
Flooding: none

Loonmeadow and similar soils

Depth to bedrock: very deep
Drainage class: very poorly drained
Parent material: coarse-loamy till derived from granite and/or schist and/or gneiss and/or dolomite
Permeability: slow to very rapid
Available water capacity: high
Reaction: strongly acid to moderately alkaline
Depth to restrictive feature: greater than 72 inches
Ponding depth: 0 to 12 inches above surface
Depth to seasonal water table: 0 to 12 inches
Flooding: none

Interpretative Groups

Brayton and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Loonmeadow and similar soils

Land capability classification (non-irrigated): 7s
Hydrologic group: D

Minor Components

Included with these soils in mapping are areas of moderately well drained Fullam and Ashfield soils and very poorly drained Wonsqueak and Bucksport soils. Fullam soils are in slightly higher areas above Brayton soils. Wonsqueak and Bucksport soils are organic soils in depressions. Minor components make up about 15 percent of the map unit.

Use and Management

This soil is mostly in woodland. Some areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil. The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised fill materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

448B—Hogansburg loam, 3 to 8 percent slopes

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: hills, uplands

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Hogansburg and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 12 inches; loam

Bw1—12 to 20 inches; loam

Bw2—20 to 29 inches; loam

Bw3—29 to 43 inches; loam

Cd1—43 to 50 inches; loam

Cd2—50 to 70 inches; loam

Cd3—70 to 84 inches; loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: very slow to moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: 20 to 43 inches to densic material

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this unit in mapping are well drained Pyrites soils in higher areas and cold phases of poorly drained Mudgepond soils. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland or hayland. Some areas are in community development or cropland.

The slow percolation is the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

449B—Hogansburg loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: hills, uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Hogansburg and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 12 inches; loam

Bw1—12 to 20 inches; loam

Bw2—20 to 29 inches; loam

Bw3—29 to 43 inches; loam

Cd1—43 to 50 inches; loam

Cd2—50 to 70 inches; loam

Cd3—70 to 84 inches; loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: very slow to moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: 20 to 43 inches to densic material

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this unit in mapping are well drained Pyrites soils in higher areas. Cold phases of poorly drained Mudgepond soils are included. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are in community development or pasture.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Large and small stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable

outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

449C—Hogansburg loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: uplands, hills

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Hogansburg and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Ap—0 to 12 inches; loam

Bw1—12 to 20 inches; loam

Bw2—20 to 29 inches; loam

Bw3—29 to 43 inches; loam

Cd1—43 to 50 inches; loam

Cd2—50 to 70 inches; loam

Cd3—70 to 84 inches; loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: moderately well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: very slow to moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: 20 to 43 inches to densic material

Depth to seasonal water table: 18 to 36 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this unit in mapping are well drained Pyrites soils in higher areas. Cold phases of poorly drained Mudgepond soils are included. Very poorly drained

Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland. Some areas are in pasture. Other areas are in community development.

The seasonal high water table is the main limitation for dwellings with basements and lawns and landscaping. Slope and large and small stones are also a limitation for lawns and landscaping. Removing the stones will reduce the limitation. Locating dwellings in the highest part of the unit with foundation drains backfilled with gravel to a suitable outlet, waterproofing the outside of basement walls, and diverting runoff from higher areas will reduce wetness.

The seasonal high water table and slow percolation are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines may allow on site sewage disposal. A more suitable site should be considered in a less dense inclusion or nearby soil.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

450B—Pyrities loam, 3 to 8 percent slopes

Map Unit Setting

Slope: gently sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Pyrities and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Ap—1 to 8 inches; loam

Bw1—8 to 13 inches; loam

Bw2—13 to 26 inches; loam

BC—26 to 45 inches; loam

C—45 to 65 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: slow or moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 2e

Hydrologic group: B

Minor Components

Included with this unit in mapping are moderately well drained Hogansburg soils in lower areas. Cold phases of poorly drained Mudgepond soils are included. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland. Other areas are in cropland, pasture or community development.

This soil has few limitations for dwellings with basements and lawns and landscaping.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

450C—Pyrities loam, 8 to 15 percent slopes

Map Unit Setting

Slope: strongly sloping

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Pyrities and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Ap—1 to 8 inches; loam

Bw1—8 to 13 inches; loam

Bw2—13 to 26 inches; loam

BC—26 to 45 inches; loam

C—45 to 65 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: slow or moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 3e

Hydrologic group: B

Minor Components

Included with this unit in mapping are moderately well drained Hogansburg soils in lower areas. Cold phases of poorly drained Mudgepond soils are included. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland. Other areas are in cropland, pasture or community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action and slope are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation. Constructing roads on the contour will reduce the slope limitation.

450D—Pyrities loam, 15 to 25 percent slopes

Map Unit Setting

Slope: moderately steep

Landscape: uplands, hills

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Pyrities and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Ap—1 to 8 inches; loam

Bw1—8 to 13 inches; loam

Bw2—13 to 26 inches; loam

BC—26 to 45 inches; loam

C—45 to 65 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: slow to moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4e

Hydrologic group: B

Minor Components

Included with this unit in mapping are moderately well drained Hogansburg soils in lower areas. Cold phases of poorly drained Mudgepond soils are included. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland. Other areas are in pasture or community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Frost action and slope are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation.

451B—Pyrities loam, 3 to 8 percent slopes, very stony

Map Unit Setting

Slope: gently sloping

Landscape: uplands, hills

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Pyrities and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

O_i—0 to 1 inch; slightly decomposed plant material

A_p—1 to 8 inches; loam

B_w1—8 to 13 inches; loam

B_w2—13 to 26 inches; loam

B_C—26 to 45 inches; loam

C—45 to 65 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: slow to moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this unit in mapping are moderately well drained Hogansburg soils in lower areas. Cold phase of poorly drained Mudgepond soils are included. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland or pasture. Other areas are in community development.

This soil has few limitations for dwellings with basements and lawns and landscaping. Large stones are the main limitation for lawns and landscaping. Removing the stones will reduce the limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action is the main limitation for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation.

451C—Pyrities loam, 8 to 15 percent slopes, very stony

Map Unit Setting

Slope: strongly sloping

Landscape: hills, uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 40 acres.

Map Unit Composition

Pyrities and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Ap—1 to 8 inches; loam

Bw1—8 to 13 inches; loam

Bw2—13 to 26 inches; loam

BC—26 to 45 inches; loam

C—45 to 65 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: slow to moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 6s

Hydrologic group: B

Minor Components

Included with this unit in mapping are moderately well drained Hogansburg soils in lower areas. Cold phases of poorly drained Mudgepond soils are included. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland or pasture. Other areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Large stones are also a limitation for lawns and landscaping. Removing the stones will reduce this limitation. Erosion is a moderate hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation is the main limitation for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution lines and adding fill will allow on site sewage disposal.

Frost action and slope are the main limitations for local roads and streets. Providing a coarse grained subgrade to frost depth will reduce this limitation. Constructing roads on the contour will reduce the slope limitation.

451D—Pyrities loam, 15 to 25 percent slopes, very stony

Map Unit Setting

Slope: moderately steep to steep

Landscape: hills, uplands

Surface cover: 0 to 3 percent stones

Size of map unit: Areas commonly range from 3 to 50 acres.

Map Unit Composition

Pyrities and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inches; slightly decomposed plant material

Ap—1 to 8 inches; loam

Bw1—8 to 13 inches; loam

Bw2—13 to 26 inches; loam

BC—26 to 45 inches; loam

C—45 to 65 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: well drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: slow or moderate

Available water capacity: high

Reaction: moderately acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: greater than 6 feet

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 7s

Hydrologic group: B

Minor Components

Included with this unit in mapping are moderately well drained Hogansburg soils in lower areas. Cold phases of poorly drained Mudgepond soils are included. Very poorly drained Loonmeadow soils are in depressions and along drainageways. Also included are well drained Shelburne soils and moderately well drained Ashfield soils in areas where the soil is more acid. Minor components make up about 15 percent of the unit.

Use and Management

Most areas are in woodland or pasture. Other areas are in community development.

Slope is the main limitation for dwellings with basements and lawns and landscaping. Erosion is a severe hazard during construction. Designing dwellings to conform to the natural slope of the land will reduce the slope limitation.

Slow percolation and slope are the main limitations for septic tank absorption fields. Modifying a conventional system by extending the length of the distribution

lines and adding fill will allow on site sewage disposal. Placing the distribution lines on the contour increases the efficiency of the system. A more suitable site should be considered in a less sloping, less dense inclusion or nearby soil.

Slope is the main limitation for local roads and streets. Constructing roads on the contour or locating them on less sloping inclusions will reduce the slope limitation.

457—Mudgepond silt loam, cold

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Size of map unit: Areas range from 3 to 50 acres.

Map Unit Composition

Mudgepond and similar soils: 80 percent

Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 11 inches; silt loam

Bg—11 to 16 inches; loam

Bw1—16 to 26 inches; fine sandy loam

Bw2—26 to 35 inches; gravelly fine sandy loam

C—35 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: neutral to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are moderately well drained Hogansburg and Ashfield soils upslope from cold phases of poorly drained Mudgepond soils. Also included are cold phases of very poorly drained Alden soils in depressions. Minor components make up about 20 percent of this map unit.

Use and Management

This soil is mostly in woodland. Other areas are in pasture or cropland.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Slow percolation is also a

limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

The seasonal high water table and frost action are the main limitations for local roads and streets. Construction on raised filled materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

458—Mudgepond and Alden soils, extremely stony, cold

Map Unit Setting

Slope: nearly level to gently sloping

Landscape: depressions on uplands, drainageways on uplands

Surface cover: 3 to 15 percent stones

Size of map unit: Areas range from 3 to 50 acres.

Map Unit Composition

Mudgepond and similar soils: 55 percent

Alden and similar soils: 35 percent

Minor components: 10 percent

Major Components

Mudgepond and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 11 inches; silt loam

Bg—11 to 16 inches; loam

Bw1—16 to 26 inches; fine sandy loam

Bw2—26 to 35 inches; gravelly fine sandy loam

C—35 to 65 inches; gravelly fine sandy loam

Alden and similar soils

The typical sequence, depth, and composition of the layers of the soil are as follows—

A1—0 to 4 inches; mucky silt loam

A2—4 to 13 inches; silt loam

Bg1—13 to 23 inches; silt loam

Bg2—23 to 29 inches; silt loam

Cg1—29 to 43 inches; gravelly loam

Cg2—43 to 60 inches; loam

Major Component Properties and Qualities

Mudgepond and similar soils

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy till derived from limestone and dolomite and/or schist

Permeability: moderate or moderately rapid

Available water capacity: high

Reaction: neutral to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Alden and similar soils

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: fine-loamy till derived from limestone and dolomite and/or schist

Permeability: moderately slow or moderate

Available water capacity: high

Reaction: strongly acid to moderately alkaline

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 6 inches above surface

Depth to seasonal water table: 0 to 12 inches

Flooding: none

Interpretative Groups**Mudgepond and similar soils**

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Alden and similar soils

Land capability classification (non-irrigated): 7s

Hydrologic group: D

Minor Components

Included with this soil in mapping are moderately well drained Hogansburg and Ashfield soils upslope from cold phases of poorly drained Mudgepond and very poorly drained Alden soils. Also included are very poorly Bucksport and Wonsqueak soils in depressions. Minor components make up about 10 percent of this map unit.

Use and Management

This soil is mostly in woodland.

The seasonal high water table is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Ponding is a limitation in areas of Alden soils. Slow percolation is also a limitation for septic tank absorption fields. A more suitable site should be selected for these uses in a drier inclusion or nearby soil.

The seasonal high water table and frost action are the main limitations for local roads and streets. Ponding is also a limitation in areas of Alden soils. Construction on raised filled materials, installing a drainage system, and providing a coarse grained subgrade to frost depth will reduce these limitations.

501—Ondawa fine sandy loam***Map Unit Setting***

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 30 acres.

Map Unit Composition

Ondawa and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

Oi—0 to 1 inch; slightly decomposed plant material

Oa—1 to 2 inches; highly decomposed plant material
 Ap—2 to 14 inches; loam
 Bw1—14 to 30 inches; loam
 C1—30 to 33 inches; loam
 C2—33 to 60 inches; fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep
Drainage class: well drained
Parent material: coarse-loamy alluvium
Permeability: slow to moderate
Available water capacity: high
Reaction: very strongly acid to slightly acid
Depth to restrictive feature: greater than 72 inches
Depth to seasonal water table: 60 to 72 inches
Flooding: occasional

Interpretative Groups

Land capability classification (non-irrigated): 1
Hydrologic group: B

Minor Components

Included with this soil in mapping are excessively drained to poorly drained Udifluvents and Fluvaquents on the flood plain. Cold phases of well drained Agawam soils are on nearby outwash plains and terraces. Areas of poorly drained Rumney soils and very poorly drained Medomak soils are included in depressions and channel scars on the flood plain. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or hayland.

Flooding is the main limitation for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Poor filtering is also a limitation for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site for these uses should be selected on a soil that does not flood.

Flooding is the main limitation for local roads and streets. Providing drainage and building on raised fill will reduce this limitation. A more suitable site should be considered on a soil that does not flood.

503—Rumney fine sandy loam

Map Unit Setting

Slope: nearly level
Landscape: depressions on flood plains
Size of map unit: Areas commonly range from 3 to 100 acres.

Map Unit Composition

Rumney and similar soils: 80 percent
Minor components: 20 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 7 inches; silt loam

Bg1—7 to 22 inches; sandy loam

Bg2—22 to 38 inches; sandy loam

Ab—38 to 42 inches; sandy loam

2Cg—42 to 44 inches; extremely gravelly coarse sand

3Cg—44 to 65 inches; gravelly fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: poorly drained

Parent material: coarse-loamy alluvium

Permeability: moderate to very rapid

Available water capacity: high

Reaction: moderately acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Depth to seasonal water table: 0 to 18 inches

Flooding: frequent

Interpretative Groups

Land capability classification (non-irrigated): 4w

Hydrologic group: D

Minor Components

Included with this soil in mapping are excessively drained to poorly drained Udifluvents and Fluvaquents on the flood plain. Well drained Ondawa soils are on higher portions of the flood plain. Areas of very poorly drained Medomak soils are included in depressions and channel scars on the flood plain. Minor components make up about 20 percent of this map unit.

Use and Management

Most areas are in woodland. Some areas are in pasture or hayland.

Flooding and the seasonal high water table are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. Poor filtering is also a limitation for septic tank absorption fields. There is the hazard of groundwater pollution because the rapidly permeable substratum does not adequately filter effluent. A more suitable site for these uses should be selected on a soil that does not flood.

Flooding, wetness, and potential frost action are the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

508—Medomak silt loam

Map Unit Setting

Slope: nearly level

Landscape: flood plains

Size of map unit: Areas commonly range from 3 to 150 acres.

Map Unit Composition

Medomak and similar soils: 85 percent

Minor components: 15 percent

Major Components

The typical sequence, depth, and composition of the layers of the soil are as follows—

A—0 to 7 inches; mucky silt loam

BG—7 to 24 inches; silt loam

Ab—24 to 33 inches; mucky silt loam

2Cg—33 to 46 inches; very gravelly coarse sand

3C—46 to 79 inches; very fine sandy loam

Major Component Properties and Qualities

Depth to bedrock: very deep

Drainage class: very poorly drained

Parent material: coarse-loamy alluvium; over sandy and gravelly alluvium

Permeability: moderate to very rapid

Available water capacity: very high

Reaction: strongly acid to slightly alkaline

Depth to restrictive feature: greater than 72 inches

Ponding depth: 0 to 18 inches above surface

Depth to seasonal water table: 0 to 6 inches

Flooding: frequent

Interpretative Groups

Land capability classification (non-irrigated): not specified

Hydrologic group: D

Minor Components

Included with this soil in mapping are excessively drained to poorly drained Udifluvents and Fluvaquents on the flood plain. Udifluvents and Well drained Ondawa soils are on higher portions of the flood plain. Areas of poorly drained Rumney soils and Fluvaquents are included in depressions and channel scars on the flood plain. Minor components make up about 15 percent of this map unit.

Use and Management

Most areas are in woodland or marsh, and are wetland wildlife habitat.

Flooding and the seasonal high water table are the main limitations for dwellings with basements, lawns and landscaping, and septic tank absorption fields. A more suitable site should be selected on a drier soil that does not flood.

Flooding, potential frost action, and the seasonal high water table are the main limitations for local roads and streets. Providing drainage and building on raised fill with a coarse grained subgrade to frost depth will reduce these limitations. A more suitable site should be considered on a soil that does not flood.

Prime Farmland

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local offices of the Natural Resources Conservation Service.

The U.S. Department of Agriculture defines additional farmland of statewide importance as land, in addition to prime farmland, that is of statewide importance for the production of food, feed, fiber, forage, and oilseed crops. Generally, additional farmlands of statewide importance include those that are nearly prime farmland and that economically produce high yields of crops when treated and managed according to modern farming methods. Some may produce as high a yield as prime farmland if conditions are favorable.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland or additional farmland of statewide importance are listed in [table 5](#). These lists do not constitute a recommendation for a particular land use. On some soils included in the lists, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures.

The extent of each listed map unit is shown in [table 4](#). The location is shown on the detailed soil maps. The soil qualities that affect use and management are described under the heading "Detailed Soil Map Units."

Use and Management of the Soils

This soil survey is an inventory and evaluation of the soils in the survey area. It can be used to adjust land uses to the limitations and potentials of natural resources and the environment. Also, it can help to prevent soil-related failures in land uses.

In preparing a soil survey, soil scientists, conservationists, engineers, and others collect extensive field data about the nature and behavioral characteristics of the soils. They collect data on erosion, droughtiness, flooding, and other factors that affect various soil uses and management. Field experience and collected data on soil properties and performance are used as a basis in predicting soil behavior.

Information in this section can be used to plan the use and management of soils for crops and pasture; as rangeland and forestland; as sites for buildings, sanitary facilities, highways and other transportation systems, and parks and other recreational facilities; and as wildlife habitat. It can be used to identify the potentials and limitations of each soil for specific land uses and to help prevent construction failures caused by unfavorable soil properties.

Planners and others using soil survey information can evaluate the effect of specific land uses on productivity and on the environment in all or part of the survey area. The survey can help planners to maintain or create a land use pattern in harmony with the natural soil.

Contractors can use this survey to locate sources of sand and gravel, roadfill, and topsoil. They can use it to identify areas where bedrock, wetness, or very firm soil layers can cause difficulty in excavation.

Health officials, highway officials, engineers, and others may also find this survey useful. The survey can help them plan the safe disposal of wastes and locate sites for pavements, sidewalks, campgrounds, playgrounds, lawns, and trees and shrubs.

Interpretive Ratings

The interpretive tables in this survey rate the soils in the survey area for various uses. Many of the tables identify the limitations that affect specified uses and indicate the severity of those limitations. The ratings in these tables are both verbal and numerical.

Rating Class Terms

Rating classes are expressed in the tables in terms that indicate the extent to which the soils are limited by all of the soil features that affect a specified use or in terms that indicate the suitability of the soils for the use. Thus, the tables may show limitation classes or suitability classes. Terms for the limitation classes are *not limited*, *slightly limited*, *somewhat limited*, and *very limited*. The suitability ratings are expressed as *well suited*, *moderately well suited*, *poorly suited*, and *unsuited* or as *good*, *fair*, and *poor*.

Numerical Ratings

Numerical ratings in the tables indicate the relative severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact

on the use and the point at which the soil feature is not a limitation. The limitations appear in order from the most limiting to the least limiting. Thus, if more than one limitation is identified, the most severe limitation is listed first and the least severe one is listed last.

Crops and Pasture

According to the 1997 Natural Resources Inventory, approximately 8,100 acres of cultivated crops and 112,000 acres of pasture were in Connecticut (US Department of Agriculture, 2000). The total acreage used for cultivated crops and pasture has been decreasing. The trend is toward the conversion of cropland to urban development, especially in the southern and central parts of the state.

The main crops in northwestern and eastern Connecticut are forage crops and corn for silage. Specialty crops are fruits, vegetables, and nursery stock. The main crops in the Connecticut Valley are nursery stock, tobacco, and vegetables. Specialty crops grown in the Connecticut Valley are fruits, flowers, and turf grass.

Soil erosion is a management concern on about 27 percent of the cropland in Connecticut (1997 NRI, USDA). In areas where the slope is more than 3 percent, erosion is a potential hazard. Stockbridge, Wethersfield, and Paxton are some of the sloping soils that are presently cultivated that are subject to erosion.

Soil erosion can reduce productivity and can result in the pollution of streams. Productivity is reduced as the surface layer of the soil erodes and more of the subsoil is incorporated into the plow layer. Loss of the surface layer is especially damaging on soils that have a silty or clayey subsoil such as Elmridge, Brancroft, or Berlin, or on soils that are shallow over bedrock, such as Hollis, Farmington, Holyoke, or Brimfield. Controlling erosion on farmland minimizes the pollution of streams and improves the quality of water for municipal uses, for recreational uses, and for fish and wildlife.

Erosion control practices provide a protective plant cover, increase the rate of water infiltration, and help to control runoff. A cropping system that keeps plant cover and crop residue on the surface for extended periods can hold soil losses to amounts that will not reduce the productive capacity of the soils. Including grasses and legumes in the cropping system helps to control erosion in sloping areas and improves tilth for the crops that follow in the rotation. The legumes also increase the nitrogen level in the soils.

Applying a system of conservation tillage and leaving crop residue on the surface increase the rate of water infiltration and help to control runoff and erosion. Using a no-till method of planting reduces the hazard of erosion in sloping areas, and this practice is suitable on most of the soils in the state.

The use of buffer strips is effective in controlling erosion on soils with very a slow rate of water infiltration such as Brancroft, Elmridge, and Shaker. Contour farming and strip cropping are very effective erosion control methods in cultivated areas. However, the small size of many fields in the state limits their use. They are best suited to soils that have smooth, uniform slopes, such as Stockbridge and Paxton.

Soil blowing can be a management concern in early spring in areas of Windsor and Berlin soils.

Connecticut has an adequate amount of rainfall for the crops commonly grown. Prolonged periods of drought are rare, but the distribution of rainfall during spring and summer generally results in droughty periods during the growing season in many years. Irrigation may be needed during these periods to reduce drought stress. Soils such as Windsor, Hartford, Hinckley, and Penwood commonly need to be irrigated.

Most of the soils of eastern and western Connecticut have a surface layer of fine sandy loam which is brown in color. Regular additions of crop residue, manure, and other organic material can improve the soil structure and reduce crust formation, thus improving the rate of water infiltration. Some of the soils in the Connecticut Valley

have a loamy sand surface; others have a silt loam surface. Regular additions of crop residue, manure, and other organic material can improve the soil structure and improve the water holding capacity of the sandy soils.

The use of heavy equipment during tillage results in soil compaction in most areas. The compacted areas, sometimes called plowpans, are generally 4 to 10 inches below the soil surface. They restrict the rate of water infiltration and limit the growth of plant roots.

Most of the soils in Connecticut require applications of agricultural limestone to neutralize soil acidity. Some of the soils of northwestern Connecticut, such as Nellis, Stockbridge, Amenia, and Georgia are higher in natural lime content and may not require additional amounts. Crops grown in Connecticut respond well to applications of lime and fertilizer. The level of available phosphorus or potassium is generally low in most of the soils; however, some fields may have a buildup of phosphorus or potassium because of past applications of commercial fertilizer. Therefore, all applications of lime or fertilizer should be based on the results of a soil test. Leaching is a concern in areas of sandy soils, such as Windsor, Hinckley, and Penwood. The Agriculture Experiment Stations or Cooperative Extension can help in determining the kinds and amounts of fertilizer and lime to apply.

Soil wetness is a management concern in areas of Ridgebury, Mudgepond, Walpole, Wilbraham, Leicester, Raypol, Shaker, Scitico, Rippowam, Raynham, Limerick and Lim soils. A drainage system may be needed to minimize the harmful effects of excess wetness. Flooding during the growing season is a concern in areas of Rippowam, Limerick, and Lim soils. Planting dates may be delayed and crops are damaged in some years because of flooding.

Proper stocking rates, control of weeds, proper fertilization, rotation grazing, and proper manure spreading are concerns on land that is used for pasture or hay production. Overgrazing, low rates of fertilization, and acid soils are the main concerns for pasture management. They can result in weak plants and poor stands that are quickly infested with weeds. Maintaining a good, dense cover that has the desired pasture species will prevent weeds from becoming established.

Nursery crops and tobacco grow best in well drained sandy loam soils such as Windsor (fig. 16). These soils promote rapid root development of nursery stock.

General management needed for crops and pasture is suggested in this section. The estimated yields of the main crops and pasture plants are listed, the system of land capability classification used by the Natural Resources Conservation Service is explained, and prime farmland is described.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil under the heading "Detailed Soil Map Units." Specific information can be obtained from the local offices of the Natural Resources Conservation Service or the Cooperative Extension Service.

Yields per Acre

The average yields per acre that can be expected of the principal crops under a high level of management are shown in table 6. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors. The land capability classification of map units in the survey area also is shown in the table.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant



Figure 16.—Tobacco grown on Windsor loamy sand.

diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in table 6 are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local offices of the Natural Resources Conservation Service or of the Cooperative Extension Service can provide information about the management and productivity of the soils for those crops.

Land Capability Classification

Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels—capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

Class 1 soils have slight limitations that restrict their use.

Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.

Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.

Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.

Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral, for example, 2e. The letter *e* shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and *c*, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by *w*, *s*, or *c* because the soils in class 5 are subject to little or no erosion. They have other limitations that restrict their use to pasture, rangeland, forestland, wildlife habitat, or recreation.

The acreage of soils in each capability class or subclass is shown in [table 7](#). The capability classification of map units in this survey area is given in the section "Detailed Soil Map Units" and in the yields table.

Forest Productivity and Management

Stephen Broderick, Extension Forester, University of Connecticut, helped to prepare this section.

Connecticut lies almost entirely within the lower New England section of the eastern broadleaf forest ecological province. A small portion of upper Litchfield County falls within the Berkshire Mountain section of the New England-Adirondack province (Keys, et al, 1995). Forest is the natural vegetative cover in these regions, and despite nearly 400 years since European settlement, about 1.9 million acres of Connecticut (60 percent) was forested in 1998 (USDA Forest Service, 1998). Over 85 percent of that forest belongs to private individuals, families or non-profit organizations.

Connecticut's forest is overwhelmingly deciduous. The oak/hickory forest type occupies just over half of our forestland (51 percent), while northern hardwoods occupy another 29 percent. Forests dominated by conifers (mainly eastern white pine, eastern hemlock, and red pine) represent less than 10 percent of Connecticut's forests.

Prior to European colonization about 95 percent of Connecticut was forested. By the Civil War nearly three-quarters of the state had been cleared for agricultural use. Between the Civil War and the World War I, Connecticut's forests were clearcut repeatedly for charcoal, fuel wood, tanning bark, pine "boxwood" and other products.

With the advent of fossil fuels in the 1920s and 1930s this era of heavy cutting ended abruptly.

Today, the size and age of our forest reflects that land use history. Sawtimber stands, where trees 11 inches in diameter or greater make up at least half the volume, occupy 69 percent of our forest. Poletimber stands occupy 25 percent, while seedling and sapling forests account for only 6 percent of the total.

Trees in the red oak group are Connecticut's most valuable timber resource. Northern red oak represents almost 20 percent of Connecticut's standing sawtimber volume, and high quality lumber finds a worldwide market. These oaks, along with sugar maple, white ash, and Connecticut's other valuable hardwoods, make their best growth on moderately well drained sandy loams, including Paxton and Woodbridge soils which have dense subsurface layers. In general, Connecticut's prime farm soils represent the most productive hardwood forest soils as well. In addition, however, soils in those classes that are too steep or stony to be rated as prime farm soils are often productive forest soils worthy of hardwood timber management investments.

Eastern white pine is Connecticut's most valuable softwood timber species, and its volume of both sawtimber and overall growing stock is increasing. While it also makes its best growth on moist, moderately well drained sandy loams, its ability to tolerate lower moisture and pH levels allows it to compete best on excessively drained soils such as Hinckley.

The fragmentation and parcelization of forests is perhaps the key forest resource issue facing Connecticut today. While the acreage of timberland in Connecticut decreased 5 percent between 1985 and 1998, the amount of urban forests grew by 74,000 acres, or 165 percent (USDA Forest Service). In a densely populated state such as Connecticut, the changes in the spatial pattern of forest cover present new challenges for timber harvest and derivation of other products from the forest.

The tables in this section can help forest owners or managers plan the use of soils for wood crops. They show the potential productivity of the soils for wood crops and rate the soils according to the limitations that affect various aspects of forest management.

Forest Productivity

In [table 8](#), the *potential productivity* of merchantable or *common trees* on a soil is expressed as a site index and as a volume number. The *site index* is the average height, in feet, that dominant and codominant trees of a given species attain in a specified number of years. The site index applies to fully stocked, even-aged, unmanaged stands. Commonly grown trees are those that forest managers generally favor in intermediate or improvement cuttings. They are selected on the basis of growth rate, quality, value, and marketability. More detailed information regarding site index is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet.

The *volume of wood fiber*, a number, is the yield likely to be produced by the most important tree species. This number, expressed as cubic feet per acre per year and calculated at the age of culmination of the mean annual increment (CMAI), indicates the amount of fiber produced in a fully stocked, even-aged, unmanaged stand.

Trees to manage are those that are preferred for planting, seeding, or natural regeneration and those that remain in the stand after thinning or partial harvest.

Forest Management

In [tables 9](#), [10](#), and [11](#) interpretive ratings are given for various aspects of forest management. The ratings are both verbal and numerical.

Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. *Well suited* indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance

can be expected, and little or no maintenance is needed. *Moderately well suited* indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable, and fair performance can be expected. Some maintenance is needed. *Poorly suited* indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. *Unsuited* indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the specified forest management practice (1.00) and the point at which the soil feature is not a limitation (0.00).

Rating class terms for fire damage and seedling mortality are expressed as *low*, *moderate*, and *high*. Where these terms are used, the numerical ratings indicate gradations between the point at which the potential for fire damage or seedling mortality is highest (1.00) and the point at which the potential is lowest (0.00).

The paragraphs that follow indicate the soil properties considered in rating the soils for forest management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service (NRCS) or the NRCS website.

For *limitations affecting construction of haul roads and log landings*, the ratings are based on slope, flooding, permafrost, plasticity index, the hazard of soil slippage, content of sand, the Unified classification, rock fragments on or below the surface, depth to a restrictive layer that is indurated, depth to a water table, and ponding. The limitations are described as slight, moderate, or severe. A rating of *slight* indicates that no significant limitations affect construction activities, *moderate* indicates that one or more limitations can cause some difficulty in construction, and *severe* indicates that one or more limitations can make construction very difficult or very costly.

The ratings of *suitability for log landings* are based on slope, rock fragments on the surface, plasticity index, content of sand, the Unified classification, depth to a water table, ponding, flooding, and the hazard of soil slippage. The soils are described as well suited, moderately suited, or poorly suited to use as log landings.

Ratings in the column *soil rutting hazard* are based on depth to a water table, rock fragments on or below the surface, the Unified classification, depth to a restrictive layer, and slope. Ruts form as a result of the operation of forest equipment. The hazard is described as slight, moderate, or severe. A rating of *slight* indicates that the soil is subject to little or no rutting, *moderate* indicates that rutting is likely, and *severe* indicates that ruts form readily.

Ratings in the column *hazard of off-road or off-trail erosion* are based on slope and on soil erodibility factor K. The soil loss is caused by sheet or rill erosion in off-road or off-trail areas where 50 to 75 percent of the surface has been exposed by logging, grazing, mining, or other kinds of disturbance. The hazard is described as slight, moderate, severe, or very severe. A rating of *slight* indicates that erosion is unlikely under ordinary climatic conditions; *moderate* indicates that some erosion is likely and that erosion-control measures may be needed; *severe* indicates that erosion is very likely and that erosion-control measures, including revegetation of bare areas, are advised; and *very severe* indicates that significant erosion is expected, loss of soil productivity and off-site damage are likely, and erosion-control measures are costly and generally impractical.

Ratings in the column *hazard of erosion on roads and trails* are based on the soil erodibility factor K, slope, and content of rock fragments. The ratings apply to unsurfaced roads and trails. The hazard is described as slight, moderate, or severe. A

rating of *slight* indicates that little or no erosion is likely; *moderate* indicates that some erosion is likely, that the roads or trails may require occasional maintenance; and that simple erosion-control measures are needed; and *severe* indicates that significant erosion is expected, that the roads or trails require frequent maintenance, and that costly erosion-control measures are needed.

Ratings in the column *suitability for roads (natural surface)* are based on slope, rock fragments on the surface, plasticity index, content of sand, the Unified classification, depth to a water table, ponding, flooding, and the hazard of soil slippage. The ratings indicate the suitability for using the natural surface of the soil for roads. The soils are described as well suited, moderately well suited, or poorly suited to this use.

Ratings in the column *potential for damage to soil by fire* are based on texture of the surface layer, content of rock fragments and organic matter in the surface layer, thickness of the surface layer, and slope. The soils are described as having a low, moderate, or high potential for this kind of damage. The ratings indicate an evaluation of the potential impact of prescribed fires or wildfires that are intense enough to remove the duff layer and consume organic matter in the surface layer.

Ratings in the column *potential for seedling mortality* are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality.

Conservation and Environmental Plantings

Conservation and environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Table 12 shows the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates in table 12 are based on measurements and observation of established plantings that have been given adequate care. Additional information on plantings screens and caring for trees and shrubs can be obtained from the local offices of the Natural Resources Conservation Service or the Cooperative Extension Service or from a commercial nursery.

Recreation

The soils of the survey area are rated in tables 13 and 14 according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Slightly limited* indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate

gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in tables 13 and 14 can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Off-road motorcycle trails require little or no site preparation. They are not covered with surfacing material or vegetation. Considerable compaction of the soil material is likely. The ratings are based on the soil properties that influence erodibility, trafficability, dustiness, and the ease of revegetation. These properties are stoniness, slope, depth to a water table, ponding, flooding, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

Wildlife Habitat

Peter Picone, Wildlife Biologist, Connecticut Department of Environmental Protection, helped to prepare this section.

Although Connecticut is the fifth most densely populated state in the United States, a diversity of wildlife may be found throughout the varying landscapes of the Northwest Highlands, the Eastern and Western Uplands, the Central Valley, and the Coastal Slope. With white-tailed deer numbering over 70,000 and wild turkey over 25,000 as of 2002, Connecticut is a state with many wildlife-related recreational opportunities (Kilpatrick et al, 2001). Just over two hundred years ago, the white-tailed deer population was barely a dozen and wild turkeys were extirpated. Land use plays a major role in determining where different species of wildlife live. Today, the fragmentation of forest, grasslands, and other habitat types that has resulted from ex-urban development is creating problems for wildlife. Fragmentation reduces the size of wildlife habitat patches, making forests and grasslands inhospitable to species that require large areas. The interspersed new housing developments within wildlife habitat may result in unwanted human/pet interactions with wildlife. Further, pet cats that are allowed to hunt outside take a tremendous toll on native songbird populations in the vicinity of houses.

Soils affect the kind and amount of vegetation that is available to wildlife as food and cover. Soils also affect the construction of water impoundments. The kind and abundance of wildlife depend largely on the amount and distribution of food, cover, and water. Wildlife habitat can be created or improved by planting appropriate vegetation, by maintaining the existing plant cover, or by promoting the natural establishment of desirable plants.

The better the soil, the better growing conditions for the wildlife habitat. However, even a droughty soil may offer habitat for some of the less abundant wildlife species such as the Eastern spadefoot toad. An area of soils which are seasonally too wet for farming, may have good habitat potential for the declining American woodcock.

As Connecticut's landscape continues to urbanize, the more adaptable wildlife species will persist and the less adaptable species will decline. Bobcats are currently found mostly in northwest and northeast Connecticut. The future of this secretive animal will depend, in a large part, by the degree to which land uses change in those parts of the state. Some animals, such as the highly adaptable Eastern Coyote can be found in every town, including all of the major urban centers.

Wildlife enhancements may be made on a smaller scale on urban and suburban lots and on a larger scale on more rural parcels of land. Landowners may enhance properties for wildlife by discouraging non-native invasive vegetation and by planting

native trees, shrubs, wildflowers, and grasses. Landowners with larger acreages may manipulate forest age and size classes to benefit early successional bird species such as the Eastern Towhee, Chestnut-sided Warbler, and Blue-winged Warbler. Landowners with larger agricultural fields can manage for declining grassland birds such as Bobolink, Eastern Meadowlark and Grasshopper Sparrow.

Wetlands tend to have more diverse vegetative communities and a wider variety of dependent wildlife species. Wetlands serve multiple functions in the life history of some wildlife species. For the less mobile reptiles and amphibians, wetlands serve as breeding habitat, foraging sites, and travel corridors. Even for upland wildlife species, wetlands are utilized for seasonal habitat requirements. Connecticut's wetlands provide habitat for not only resident wildlife species, but also interstate migrants. Vegetation in and adjacent to ponded or flowing water provides migratory songbirds nesting habitat during the breeding season and resting and foraging locations during the migration season.

In [table 15](#), the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

The potential of the soil is rated good, fair, poor, or very poor. A rating of *good* indicates that the element or kind of habitat is easily established, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected. A rating of *fair* indicates that the element or kind of habitat can be established, improved, or maintained in most places. Moderately intensive management is required for satisfactory results. A rating of *poor* indicates that limitations are severe for the designated element or kind of habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and must be intensive. A rating of *very poor* indicates that restrictions for the element or kind of habitat are very severe and that unsatisfactory results can be expected. Creating, improving, or maintaining habitat is impractical or impossible.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, bromegrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing

shrubs that are suitable for planting on soils rated *good* are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, wildrice, saltgrass, cordgrass, rushes, sedges, and reeds.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, and red fox.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, ruffed grouse, woodcock, thrushes, woodpeckers, squirrels, gray fox, raccoon, deer, and bear.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Hydric Soils

In this section, hydric soils are defined and described and the hydric soils in the survey area are listed.

The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are

described in “Soil Taxonomy” (USDA, 1999) and “Keys to Soil Taxonomy” (USDA, 1998) and in the “Soil Survey Manual” (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in “Field Indicators of Hydric Soils in the United States” (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

The following map units are dominated by soils that meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996). The complete list with each map unit component, hydric status, and specific hydric soils criteria status may be accessed through the eFOTG (Electronic Field Office Technical Guide) at the Connecticut NRCS website (www.ct.nrcs.usda.gov).

- 2 Ridgebury fine sandy loam
- 3 Ridgebury, Leicester, and Whitman soils, extremely stony
- 4 Leicester fine sandy loam
- 5 Wilbraham silt loam
- 6 Wilbraham and Menlo soils, extremely stony
- 7 Mudgepond silt loam
- 8 Mudgepond and Alden soils, extremely stony
- 9 Scitico, Shaker, and Maybid soils
- 10 Raynham silt loam
- 12 Raypol silt loam
- 13 Walpole sandy loam
- 14 Fredon silt loam
- 15 Scarboro muck
- 16 Halsey silt loam
- 17 Timakwa and Natchaug soils
- 18 Catden and Freetown soils
- 96 Ipswich mucky peat
- 97 Pawcatuck mucky peat
- 98 Westbrook mucky peat
- 99 Westbrook mucky peat, low salt
- 103 Rippowam fine sandy loam
- 104 Bash silt loam
- 107 Limerick and Lim soils
- 108 Saco silt loam
- 109 Fluvaquents-Udifluvents complex, frequently flooded (Fluvaquents are hydric; Udifluvents are not hydric)
- 409 Brayton mucky silt loam, 0 to 8 percent slopes, very stony
- 414 Fredon silt loam, cold
- 433 Moosilauke sandy loam
- 435 Scarboro muck, cold
- 436 Halsey silt loam, cold

437	Wonsqueak peat
438	Bucksport muck
442	Brayton loam
443	Brayton-Loonmeadow complex, extremely stony
457	Mudgepond silt loam, cold
458	Mudgepond and Alden soils, extremely stony, cold
503	Rumney fine sandy loam
508	Medomak silt loam

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

Map units that are not listed do not meet the definition of hydric soils because the dominant soil component does not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

Connecticut Inland Wetland Soils

The state of Connecticut defines inland wetlands based on soils. The Connecticut Inland Wetlands and Watercourses Act defines wetland soils to include “any of the soil types designated as poorly drained, very poorly drained, alluvial, and floodplain by the National Cooperative Soil Survey, as may be amended from time to time, of the Natural Resources Conservation Service of the United States Department of Agriculture.”

Map units may be dominated by Connecticut inland wetland soils, but have inclusions of non-wetland soils. Non-wetland map units may contain inclusions of Connecticut inland wetland soils. Onsite investigation is necessary to determine the presence or absence of wetland soils in a particular area.

The map units in Connecticut that are dominated by inland wetland are listed in [table 16](#).

Engineering

This section provides information for planning land uses related to urban development and to water management. Soils are rated for various uses, and the most limiting features are identified. Ratings are given for building site development, sanitary facilities, construction materials, and water management. The ratings are based on observed performance of the soils and on the estimated data and test data in the “Soil Properties” section.

Information in this section is intended for land use planning, for evaluating land use alternatives, and for planning site investigations prior to design and construction. The information, however, has limitations. For example, estimates and other data generally apply only to that part of the soil between the surface and a depth of 5 to 7 feet. Because of the map scale, small areas of different soils may be included within the mapped areas of a specific soil.

The information is not site specific and does not eliminate the need for onsite investigation of the soils or for testing and analysis by personnel experienced in the design and construction of engineering works.

Government ordinances and regulations that restrict certain land uses or impose specific design criteria were not considered in preparing the information in this section. Local ordinances and regulations should be considered in planning, in site selection, and in design.

Soil properties, site features, and observed performance were considered in determining the ratings in this section. During the fieldwork for this soil survey, determinations were made about particle-size distribution, liquid limit, plasticity index, soil reaction, depth to bedrock, hardness of bedrock within 5 to 7 feet of the surface, soil wetness, depth to a water table, ponding, slope, likelihood of flooding, natural soil structure aggregation, and soil density. Data were collected about kinds of clay minerals, mineralogy of the sand and silt fractions, and the kinds of adsorbed cations. Estimates were made for erodibility, permeability, corrosivity, shrink-swell potential, available water capacity, and other behavioral characteristics affecting engineering uses.

This information can be used to evaluate the potential of areas for residential, commercial, industrial, and recreational uses; make preliminary estimates of construction conditions; evaluate alternative routes for roads, streets, highways, pipelines, and underground cables; evaluate alternative sites for septic tank absorption fields and sewage lagoons; plan detailed onsite investigations of soils and geology; locate potential sources of gravel, sand, earthfill, and topsoil; plan drainage systems, irrigation systems, ponds, terraces, and other structures for soil and water conservation; and predict performance of proposed small structures and pavements by comparing the performance of existing similar structures on the same or similar soils.

The information in the tables, along with the soil maps, the soil descriptions, and other data provided in this survey, can be used to make additional interpretations.

Some of the terms used in this soil survey have a special meaning in soil science and are defined in the Glossary.

Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. [Tables 17](#) and [18](#) show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Slightly limited* indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties

that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

Sanitary Facilities

Table 19 shows the degree and kind of soil limitations that affect septic tank absorption fields and sewage lagoons. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil

features that affect these uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Slightly limited* indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Soil potential ratings for septic tank absorption fields for single family residences are provided at www.ct.nrcs.usda.gov/soils.html. Soil potential ratings are interpretive ratings that stress the suitability of use.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

Construction Materials

Tables 20 and 21 give information about the soils as potential sources of gravel, sand, roadfill, and topsoil. Normal compaction, minor processing, and other standard construction practices are assumed.

Sand and *gravel* are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In table 20, only the likelihood of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the bottom layer of the soil contains sand or gravel, the soil is considered a likely source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

The soils are rated *good*, *fair*, or *poor* as potential sources of sand and gravel. A rating of *good* or *fair* means that the source material is likely to be in or below the soil. The bottom layer and the thickest layer of the soils are assigned numerical ratings. These ratings indicate the likelihood that the layer is a source of sand or gravel. The number 0.00 indicates that the layer is a poor source. The number 0.99 indicates that the layer is a good source. A number between 0.00 and 0.99 indicates the degree to which the layer is a likely source.

The soils are rated *good*, *fair*, or *poor* as potential sources of roadfill and topsoil. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of roadfill or topsoil. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Water Management

Table 22 gives information on the soil properties and site features that affect water management. The degree and kind of soil limitations are given for pond reservoir areas; embankments, dikes, and levees; and aquifer-fed excavated ponds.

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Slightly limited* indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects trafficability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Soil Properties

Data relating to soil properties are collected during the course of the soil survey.

Soil properties are ascertained by field examination of the soils and by laboratory index testing of some benchmark soils. Established standard procedures are followed. During the survey, many shallow borings are made and examined to identify and classify the soils and to delineate them on the soil maps. Samples are taken from some typical profiles and tested in the laboratory to determine particle-size distribution, plasticity, and compaction characteristics.

Estimates of soil properties are based on field examinations, on laboratory tests of samples from the survey area, and on laboratory tests of samples of similar soils in nearby areas. Tests verify field observations, verify properties that cannot be estimated accurately by field observation, and help to characterize key soils.

The estimates of soil properties are shown in tables. They include engineering index properties, physical and chemical properties, and pertinent soil and water features.

Engineering Index Properties

[Table 23](#) gives the engineering classifications and the range of index properties for the layers of each soil in the survey area.

Depth to the upper and lower boundaries of each layer is indicated.

Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. "Loam," for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, "gravelly." Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998).

The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection.

If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional

refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in [table 23](#).

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage.

Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination.

The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

Physical Properties

[Table 24](#) shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In [table 24](#), the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In [table 24](#), the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In [table 24](#), the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earthmoving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at $1/3$ - or $1/10$ -bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Permeability (K_{sat}) refers to the ability of a soil to transmit water or air. The term “permeability,” as used in soil surveys, indicates saturated hydraulic conductivity (K_{sat}). The estimates in the table indicate the rate of water movement, in both inches per hour and micrometers per second (um/sec), when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at $1/3$ - or $1/10$ -bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In table 24, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in [table 24](#) as the K factor (K_w and K_f) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of several factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor Kw indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor Kf indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Chemical Properties

Table 25 shows estimates of some chemical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. The ability to retain cations reduces the hazard of ground-water pollution.

Effective cation-exchange capacity refers to the sum of extractable bases plus aluminum expressed in terms of milliequivalents per 100 grams of soil. It is determined for soils that have pH of less than 5.5.

Soil reaction is a measure of acidity or alkalinity. The pH of each soil horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Extractable aluminum is a measure of the active acidity present in soils. It is important for soil classification and for certain evaluations of soil nutrient availability and of toxicities. It may be a useful measurement for assessing potential lime needs for acid soils.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Water Features

Table 26 gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations.

Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The *months* in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. Table 26 indicates, by month, depth to the top (*upper limit*) and base (*lower limit*) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table.

Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. Table 26 indicates *surface water depth* and the *duration* and *frequency* of ponding. Duration is expressed as *very brief* if less than 2 days, *brief* if 2 to 7 days, *long* if 7 to 30 days, and *very long* if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. *None* means that ponding is not probable; *rare* that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); *occasional* that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and *frequent* that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding is the temporary inundation of an area caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and *frequency* are estimated. Duration is expressed as *extremely brief* if 0.1 hour to 4 hours, *very brief* if 4 hours to 2 days, *brief* if 2 to 7 days, *long* if 7 to 30 days, and *very long* if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. *None* means that flooding is not probable; *very rare* that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); *rare* that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); *occasional* that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); *frequent* that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and *very frequent* that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

Soil Features

[Table 27](#) gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A *restrictive layer* is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. *Depth to top* is the vertical distance from the soil surface to the upper boundary of the restrictive layer. If there is no restrictive layer identified, it is assumed that the depth to bedrock is greater than 6 feet.

Subsidence is the settlement of organic soils or of saturated mineral soils of very low density. Subsidence generally results from either desiccation and shrinkage or oxidation of organic material, or both, following drainage. Subsidence takes place gradually, usually over a period of several years. The table shows the expected initial subsidence, which usually is a result of drainage, and total subsidence, which results from a combination of factors.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as *low*, *moderate*, or *high*, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as *low*, *moderate*, or *high*. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

Storm Water Runoff Management Systems

Soil properties influence storm water runoff management, including the design, installation, and maintenance of the system, and especially the selection of the site and type of system to be used. [Table 28](#) shows the degree of limitation and limiting soil factors for four commonly used storm water runoff management systems. The practices are detention basin type, infiltration systems, perennial water type, and intermittent water type.

This interpretation was developed using criteria in the NRCS Connecticut/Rhode Island Runoff Management System Standard number 570. These ratings apply to storm water runoff from residential and commercial areas including driveways, roads, roofs, parking areas, and lawns. They are not intended for the placement of treatment systems for animal waste, domestic or industrial wastewater, or other highly concentrated waste material. The information in this table indicates interpretation of the dominant soil condition but does not eliminate the need for onsite investigation.

The table includes rating classes and numerical ratings. Rating classes indicate the extent to which the soils are limited by the soil properties that affect the management system. *Unlimited* indicates that the soil is very favorable for the specified system. Good performance and relatively low installation and maintenance costs can be expected. *Somewhat limited* indicates that the soil is moderately favorable for the system. The limitations can be overcome by or minimized by special planning, design, installation, and maintenance. Increased installation costs and maintenance will be required to sustain performance. *Very limited* indicates that one or more soil feature is unfavorable for the specified system. The limitation generally cannot be overcome. Sometimes expensive design, installation, and maintenance may be employed, but performance may still be poor.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which the soil is not a limitation (0.00) and at which the soil feature(s) has the greatest negative impact (1.00). The larger the value, the greater the limitation.

Detention basins are the most commonly used system for storm water management and include retention basins. Runoff from each rain event is directed into a basin or pond, where it is detained and treated until it infiltrates into the ground or is displaced by runoff from the next storm. These basins detain the storm water runoff to allow particles and associated pollutants to settle. They may reduce peak flows, remove particulate pollutants, allow groundwater recharge, and provide flood control. Of all the practices evaluated, detention/retention basins are practical on the widest range of soils. Only soils with shallow bedrock, ponding, or very steep slopes cannot be used. Soil properties that influence performance and were used to develop this interpretation are: depth to bedrock, hydrologic soil group, flooding, and slope.

Infiltration systems receive storm water runoff and allow it to seep into the soil below. Runoff is stored in either void space between the stones in a stone filled trench or stored in underground chambers made of plastic or concrete. The stored runoff infiltrates through the bottom and sides of these systems into the soil matrix. An advantage of these systems is that they provide ground water recharge as well as pollutant control. The primary pollutant removal mechanism of these systems is filtering through the soil. Infiltration systems are used to remove suspended solids, particulate pollutants, coliform bacteria, organics, and some soluble forms of metals and nutrients. Runoff should be clean or pretreated for infiltration systems in soils with permeability greater than six inches per hour in the substratum.

Specific practices included in this interpretation are infiltration trenches, which are excavated trenches at or below grade that are backfilled with stone and lined with filter fabric; and underground galleries (or galleys or leaching galleys), which are

similar to infiltration trenches except that underground chambers made of plastic and concrete are used to store runoff and allow infiltration. Infiltration systems perform best in deep, permeable soils, with a deep or absent water table on a level area. Properties that influence performance and were used to develop this interpretation are depth to restrictive layer (bedrock, dense till), depth to seasonal high water table and ponding, saturated hydraulic conductivity in the top 40 inches or above a restrictive layer, bulk density, flooding, and slope.

Perennial water systems include ponds, shallow marsh systems, and constructed wetlands. Storm water runoff flows into the system and is held there for long periods. Sedimentation processes remove particulates, organic matter, and metals, while dissolved metals and nutrients are removed through biological uptake. These systems require deep soils with slow hydraulic conductivity, permanent water tables and a level site. Soil properties that influence performance and were used to develop this interpretation are: seasonal high water table, saturated hydraulic conductivity, hydrologic soil group, and slope.

Intermittent water systems are similar to perennial systems except that they may dry out for short periods during the year. These systems include pond/wetland systems, extended detention wetlands, and pocket wetlands. Soils may have slightly more rapid hydraulic conductivity or lower water tables than those required for perennial systems. Soil properties that influence performance and were used to develop this interpretation are: seasonal high water table, saturated hydraulic conductivity, hydrologic soil group, and slope.

Low Impact Development (LID) Management Practices are designed to maintain or replicate predevelopment hydrology through the use of small-scale controls integrated throughout the site. These systems are generally designed to accept runoff from a single or part of a single residential lot. LID practices that depend on site and soil conditions include vegetated swales, buffers, and filter strips; rain gardens; dry wells and leaching trenches; and the incorporation of pervious constructing materials such as porous pavers for driveways, walkways, and parking areas.

Use the interpretation in table 28 that best addresses these practices for small residential scale systems. For vegetated swales, buffers, filter strips, and rain gardens* use the ratings for detention/retention basins. For dry wells, leaching trenches, and pervious constructing materials for driveways, walkways, and parking areas use the ratings for infiltration systems.

***For large scale bio-retention systems (“rain gardens”) use the ratings for infiltration systems.**

Classification of the Soils

The system of soil classification used by the National Cooperative Soil Survey has six categories (USDA, 1998 and 1999). Beginning with the broadest, these categories are the order, suborder, great group, subgroup, family, and series. Classification is based on soil properties observed in the field or inferred from those observations or from laboratory measurements. [Table 29](#) shows the classification of the soils in the survey area. The categories are defined in the following paragraphs.

ORDER. Twelve soil orders are recognized. The differences among orders reflect the dominant soil-forming processes and the degree of soil formation. Each order is identified by a word ending in *sol*. An example is Inceptisol.

SUBORDER. Each order is divided into suborders primarily on the basis of properties that influence soil genesis and are important to plant growth or properties that reflect the most important variables within the orders. The last syllable in the name of a suborder indicates the order. An example is Udept (*Ud*, meaning humid, plus *ept*, from Inceptisol).

GREAT GROUP. Each suborder is divided into great groups on the basis of close similarities in kind, arrangement, and degree of development of pedogenic horizons; soil moisture and temperature regimes; type of saturation; and base status. Each great group is identified by the name of a suborder and by a prefix that indicates a property of the soil. An example is Dystrudepts (*Dystr*, meaning low base saturation, plus *udept*, the suborder of the Inceptisols that has an udic moisture regime).

SUBGROUP. Each great group has a typic subgroup. Other subgroups are intergrades or extragrades. The typic subgroup is the central concept of the great group; it is not necessarily the most extensive. Intergrades are transitions to other orders, suborders, or great groups. Extragrades have some properties that are not representative of the great group but do not indicate transitions to any other taxonomic class. Each subgroup is identified by one or more adjectives preceding the name of the great group. The adjective *Typic* identifies the subgroup that typifies the great group. An example is Typic Dystrudepts.

FAMILY. Families are established within a subgroup on the basis of physical and chemical properties and other characteristics that affect management. Generally, the properties are those of horizons below plow depth where there is much biological activity. Among the properties and characteristics considered are particle size, mineral content, soil temperature regime, soil depth, and reaction. A family name consists of the name of a subgroup preceded by terms that indicate soil properties. An example is coarse-loamy, mixed, mesic Typic Dystrudepts.

SERIES. The series consists of soils within a family that have horizons similar in color, texture, structure, reaction, consistence, mineral and chemical composition, and arrangement in the profile.

Soil Series and Their Morphology

In this section, each soil series recognized in the survey area is described. Characteristics of the soil and the material in which it formed are identified for each series. A pedon, a small three-dimensional area of soil that is typical of the series in

the survey area is described. The detailed description of each soil horizon follows standards in the "Soil Survey Manual" (USDA, 1993). Many of the technical terms used in the descriptions are defined in "Soil Taxonomy" (USDA, 1999) and in "Keys to Soil Taxonomy" (USDA, 1998). Unless otherwise indicated, colors in the descriptions are for moist soil. Following the pedon description is the range of important characteristics of the soils in the series.

Agawam Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Eolian deposits over glaciofluvial deposits derived from schist, granite, and gneiss

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Ninigret (moderately well drained)

Walpole (poorly drained)

Scarboro (very poorly drained)

Associated soils:

Enfield (coarse-silty over sandy or sandy-skeletal)

Haven (finer texture in the subsoil)

Hinckley (excessively drained, sandy-skeletal)

Merrimac (somewhat excessively drained, sandy)

Windsor (excessively drained, sandy)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts

Typical Pedon

Agawam fine sandy loam, 3 to 8 percent slopes, located in the town of Essex, 1,900 feet northeast of the intersection of Bokum Road and Plains Road, on the Essex USGS topographic quadrangle, lat. 41 degrees 20 minutes 36 seconds N., long. 72 degrees 24 minutes 29 seconds W., NAD 27:

Ap—0 to 8 inches; dark brown (10YR 3/3) fine sandy loam, pale brown (10YR 6/3) dry; weak medium granular structure; friable; common fine roots; neutral; clear smooth boundary.

Bw1—8 to 14 inches; brown (7.5YR 4/4) fine sandy loam; weak, medium subangular blocky structure; friable; few fine roots; moderately acid; gradual wavy boundary.

Bw2—14 to 24 inches; strong brown (7.5YR 5/6) fine sandy loam; weak, medium subangular blocky structure; friable; moderately acid; clear wavy boundary.

2C—24 to 60 inches; brown (7.5YR 4/4) and grayish brown (10YR 5/2) stratified fine sand to very gravelly coarse sand; single grain; loose; 5 percent rock fragments; moderately acid.

Range in Characteristics

Solum thickness: 15 to 35 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid

Ap horizon:

Hue—7.5YR to 2.5Y

Value—3 or 4 (6 or more dry) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)
 Content of rock fragments—0 to 10 percent

Bw1 horizon:

Hue—5YR to 10YR
 Value—4 to 7
 Chroma—3 to 8
 Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, or loam
 Content of rock fragments—0 to 10 percent

Bw2 horizon:

Hue—7.5YR to 5Y
 Value—4 to 7
 Chroma—3 to 8
 Texture of the fine earth fraction—fine sandy loam or very fine sandy loam
 Content of rock fragments—0 to 14 percent

BC horizon (where present)

Hue—7.5YR to 5Y
 Value—4 to 7
 Chroma—3 to 8
 Texture of the fine earth fraction—sandy loam or loamy sand
 Content of rock fragments—0 to 14 percent

2C horizon:

Hue—7.5YR to 5Y
 Value—3 to 7
 Chroma—1 to 4
 Texture—stratified fine sand to very gravelly coarse sand
 Content of rock fragments—0 to 30 percent above a depth of 40 inches, 0 to 60 percent below

Some of the Agawam soils in this survey area have a mean annual soil temperature which is colder than typical of the series. These map units (429A, 429B, and 429C) are identified as cold phases of the Agawam series.

Alden Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate in the surface layer and subsoil, and moderately slow in the substratum

Landform: Depressions and drainageways on hills

Parent material: Till derived from limestone, dolomite, and schist

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Amenia (moderately well drained, coarse-loamy)

Georgia (moderately well drained, coarse-loamy)

Mudgepond (poorly drained, coarse-loamy)

Associated other soil:

Natchaug (16 to 51 inches of organic materials, over loamy deposits)

Taxonomic class: Fine-loamy, mixed, active, nonacid, mesic Mollic Endoaquepts

Typical Pedon

Alden mucky silt loam, in an area of Mudgepond and Alden soils, extremely stony, located in the town of Sharon, 1.5 miles northeast on Indian Mountain Road from the

intersection of Indian Mountain Road and Route 4, 200 feet east of Indian Mountain Road, on the Ellsworth USGS topographic quadrangle, lat. 41 degrees 54 minutes 42 seconds N., long. 73 degrees 28 minutes 18 seconds W., NAD 27, in a swamp:

- A1—0 to 4 inches; very dark grayish brown (2.5Y 3/2) mucky silt loam, grayish brown (2.5Y 5/2) dry; weak fine granular structure; very friable; many fine to medium roots; neutral; clear wavy boundary.
- A2—4 to 13 inches; very dark gray (2.5Y 3/1) silt loam; weak medium granular structure; friable; many fine to medium roots; neutral; clear wavy boundary.
- Bg1—13 to 23 inches; dark gray (5Y 4/1) silt loam; massive; friable; common fine to medium roots; common very fine prominent yellowish red (5YR 5/8) soft masses of iron accumulation; 5 percent rock fragments; neutral; clear wavy boundary.
- Bg2—23 to 29 inches; gray (5Y 5/1) silt loam; massive; friable; common fine roots; few medium prominent gray (10YR 6/1) iron depletions and many fine prominent brownish yellow (10YR 6/8) soft masses of iron accumulation; 5 percent rock fragments; neutral; clear wavy boundary.
- Cg1—29 to 43 inches; olive gray (5Y 4/2) gravelly loam; massive; friable; few fine roots; few medium prominent brownish yellow (10YR 6/8) soft masses of iron accumulation; 18 percent rock fragments; neutral; clear wavy boundary.
- Cg2—43 to 60 inches; dark olive gray (5Y 3/2) loam; massive; friable; few fine roots; common medium distinct pale olive (5Y 6/3) iron depletions and many medium prominent very dark brown (7.5YR 2.5/2) soft masses of iron accumulation; 8 percent rock fragments; neutral.

Range in Characteristics

Solum thickness: 19 to 48 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer, moderately acid to neutral in the subsoil, slightly acid to neutral to a depth of 40 inches, and slightly acid to moderately alkaline at depths greater than 40 inches

A horizons:

Hue—10YR or 2.5Y

Value—2 or 3

Chroma—0 to 2

Content of rock fragments—0 to 14 percent

Bg horizons:

Hue—5YR to 5Y or 5GY, or neutral

Value—4 to 6

Chroma—0 to 2

Texture of the fine earth fraction—very fine sandy loam or silt loam

Content of rock fragments—0 to 14 percent

Cg horizons:

Hue—5YR to 5Y

Value—3 to 6

Chroma—0 to 2

Texture of the fine earth fraction—loam or silt loam

Content of rock fragments—5 to 34 percent

Some of the Alden soils in this survey area have a mean annual soil temperature which is colder than typical of the series. This map unit (458) is identified as a cold phase of the Alden series.

Amenia Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, moderately slow in the substratum

Landform: Hills

Parent material: Till derived from limestone, dolomite and schist

Slope range: 2 to 15 percent

Associated soils in a drainage sequence:

Nellis (well drained)

Mudgepond (poorly drained)

Alden (very poorly drained, fine-loamy)

Associated similar soil:

Georgia (carbonates at greater depth)

Associated other soils:

Farmington (shallow, well drained)

Stockbridge (well drained, carbonates at greater depth)

Taxonomic class: Coarse-loamy, mixed, active, mesic Aquic Eutrudepts

Typical Pedon

Amenia silt loam, in an area of Georgia and Amenias silt loams, 2 to 8 percent slopes, located in the town of Salisbury, 2,300 feet east/southeast of the intersection of Long Pond Road and Indian Mountain Road, on the Sharon USGS topographic quadrangle, lat. 41 degrees 55 minutes 35 seconds N., long. 73 degrees 27 minutes 40 seconds W., NAD 27, in a cultivated field:

Ap—0 to 9 inches; brown (10YR 4/3) silt loam, light brownish gray (10YR 6/2) dry; weak medium granular structure; very friable; many fine and very fine roots; 10 percent gravel, 2 percent cobbles; neutral; clear smooth boundary.

Bw1—9 to 16 inches; olive brown (2.5Y 4/4) silt loam; weak medium subangular blocky structure; friable; common fine and very fine roots; few fine faint light olive brown (2.5Y 5/4) soft masses of iron accumulation; 10 percent gravel, 2 percent cobbles; neutral; gradual wavy boundary.

Bw2—16 to 25 inches; light olive brown (2.5Y 5/4) silt loam; weak medium subangular blocky structure; friable; few fine and very fine roots; common fine faint dark yellowish brown (10YR 4/4), distinct (10YR 4/6) soft masses of iron accumulation and distinct grayish brown (2.5Y 5/2) iron depletions; 10 percent gravel, 2 percent cobbles; neutral; clear smooth boundary.

C—25 to 60 inches; olive (5Y 4/3) gravelly loam; massive; firm; few fine distinct gray (5Y 5/1) iron depletions, distinct dark yellowish brown (10YR 4/4 and 10YR 4/6), faint olive brown (2.5Y 4/4), and prominent red (2.5YR 4/6) soft masses of iron accumulation; 20 percent gravel, 5 percent cobbles; slight effervescence; slightly alkaline.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to slightly alkaline in the surface layer and subsoil, slightly alkaline or moderately alkaline in the substratum

Depth to carbonates: 10 to 34 inches

Ap horizon:

Hue—or 2.5Y

Value—3 or 4

Chroma—2 or 3 (If A horizon present instead of Ap, chroma 2)
 Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—10YR or 2.5Y
 Value—4 to 6
 Chroma—2 to 6
 Texture of the fine earth fraction—loam or silt loam
 Content of rock fragments—5 to 30 percent

C horizon:

Hue—10YR to 5Y
 Value—4 or 5
 Chrom—1 to 3
 Texture of the fine earth fraction—loam or silt loam
 Content of rock fragments—5 to 30 percent

Ashfield Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderately rapid in the surface layer, moderate to moderately rapid in the subsoil, and very slow or slow in the substratum

Landform: Hills and drumlins

Parent material: Lodgement till derived from granite, schist, and gneiss

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Shelburne (well drained)
 Loonmeadow (very poorly drained)

Associated similar soil:

Schroon (friable substratum)

Associated other soils:

Westminster (shallow, somewhat excessively drained)
 Millsite (moderately deep, well drained)

Taxonomic class: Coarse-loamy, mixed, active, frigid Aquic Dystrudepts

Typical Pedon

Ashfield fine sandy loam, 2 to 8 percent slopes, very stony, located 2,000 feet north along a private gravel road from the intersection with Loon Meadow Drive and 1,400 feet southwest of Doolittle Lake, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees 00 minutes 35 seconds N., long. 73 degrees 09 minutes 53 seconds W., NAD 27, in a wooded area:

Oi—0 to 1 inch; slightly decomposed plant material

Oe—1 to 2 inches; moderately decomposed plant material

Oa—2 to 3 inches; highly decomposed plant material

A1—3 to 7 inches; very dark grayish brown (10YR 3/2) fine sandy loam, brown (10YR 5/3) dry; weak very fine and fine granular structure; very friable; many fine and very fine roots; 4 percent gravel, 1 percent cobbles; extremely acid; clear smooth boundary.

A2—7 to 12 inches; brown (10YR 4/3) fine sandy loam, brown (10YR 5/3) dry; weak fine and medium granular structure; very friable; many medium and very coarse roots; 5 percent gravel, 1 percent cobbles; very strongly acid; clear smooth boundary.

Bw1—12 to 18 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak fine and medium subangular blocky structure; friable; many fine to medium roots; few

- fine faint brown (10YR 5/3) soft masses of iron accumulation and few fine distinct grayish brown (10YR 5/2) iron depletions; 7 percent gravel, 1 percent cobbles; very strongly acid; abrupt wavy boundary.
- Bw2—18 to 24 inches; yellowish brown (10YR 5/4) fine sandy loam; moderate medium subangular blocky structure; friable; common fine to medium roots; few fine and medium faint brown (10YR 5/3) and common coarse and very coarse distinct strong brown (7.5YR 4/6) soft masses of iron accumulation and few fine and medium distinct grayish brown (10YR 5/2) iron depletions; 10 percent gravel, 5 percent cobbles; very strongly acid; clear wavy boundary.
- BC—24 to 29 inches; brown (10YR 5/3) fine sandy loam; weak fine and medium subangular blocky structure; friable to firm; few very fine and fine roots and common medium roots in cracks; common medium and coarse distinct strong brown (7.5YR 4/6) soft masses of iron accumulation and few fine and medium faint grayish brown (2.5Y 5/2) iron depletions; 8 percent gravel, 5 percent cobbles; strongly acid; clear wavy boundary.
- Cd1—29 to 44 inches; brown (10YR 5/3) fine sandy loam; massive; firm to very firm; few fine and medium roots; many medium prominent strong brown (7.5YR 4/6) soft masses of iron accumulation, common medium faint light brownish gray (2.5Y 6/2) iron depletions, and few fine prominent dark reddish brown (5YR 2.5/1) soft masses of iron and manganese accumulation; 12 percent gravel, 1 percent cobbles; strongly acid; gradual smooth boundary.
- Cd2—44 to 58 inches; light olive brown (2.5Y 5/3) sandy loam; massive; very firm to firm; few fine and medium roots in cracks; common medium prominent strong brown (7.5YR 4/6) soft masses of iron accumulation and few medium distinct gray (2.5Y 6/1) and common medium and coarse faint light brownish gray (2.5Y 6/2) iron depletions; 4 percent gravel, 5 percent cobbles; moderately acid; gradual smooth boundary.
- Cd3—58 to 80 inches; olive brown (2.5Y 4/3) fine sandy loam; pockets of loamy sand; massive; friable; few fine faint olive brown (2.5Y 4/4) and few fine distinct brown (7.5YR 4/4) soft masses of iron accumulation; 8 percent gravel, 5 percent cobbles; slightly acid.

Range in Characteristics

Solum thickness: 20 to 33 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to strongly acid in the surface layer and subsoil, strongly acid to slightly acid in the substratum

A horizons:

Hue—10YR

Value—2 or 3

Chroma—1 to 4

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—10YR to 5Y

Value—3 to 5

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam, silt loam, or loam

Content of rock fragments—5 to 25 percent

BC horizon:

Hue—10YR to 5Y

Value—3 to 5

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam, silt loam or loam
Content of rock fragments—5 to 25 percent

Cd horizons:

Hue—2.5Y or 5Y

Value—3 to 5

Chroma—1 to 3

Texture of the fine earth fraction—fine sandy loam, sandy loam, or loam (pockets of loamy sand possible in lower substratum)

Content of rock fragments—5 to 25 percent

Bash Series

Depth class: Very deep

Drainage class: Somewhat poorly drained

Permeability: Moderate in the surface layer and subsoil, and moderate or moderately slow in the substratum

Landform: Flood plains

Parent material: Alluvium derived from sandstone and shale

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Hadley (well drained)

Winooski (moderately well drained)

Lim (poorly drained)

Limerick (coarse-silty, poorly drained)

Saco (very poorly drained)

Associated similar soil:

Pootatuck (moderately well drained, browner)

Taxonomic class: Coarse-loamy, mixed, semiactive, mesic Fluvaquentic Dystrudepts

Typical Pedon

Bash silt loam, located in the town of Middlefield, 1,800 feet southeast along Route 147 from the intersection with Miller Road, 100 feet south of Route 147 and 200 feet west of Coginchaug River, on the Durham USGS topographic quadrangle, lat. 41 degrees 29 minutes 42 seconds N., long. 71 degrees 42 minutes 18 seconds W., NAD 27, in a grass field:

Ap—0 to 11 inches; brown (7.5YR 4/2) silt loam, light brown (7.5 YR 6/3) dry; weak fine granular structure; very friable; few very fine and fine roots; moderately acid; abrupt smooth boundary.

Bw1—11 to 21 inches; reddish brown (5YR 4/3) silt loam; weak medium subangular blocky structure; friable; few fine roots; common, medium distinct yellowish red (5YR 5/6) soft masses of iron accumulation and common medium distinct grayish brown (10YR 5/2) iron depletions; strongly acid; clear wavy boundary.

Bw2—21 to 28 inches; reddish brown (5YR 4/3) silt loam; weak medium subangular blocky structure; friable; common fine prominent strong brown (7.5 YR 5/6) and few fine prominent dark red (2.5YR 3/6) soft masses of iron accumulation; strongly acid; clear wavy boundary.

C—28 to 60 inches; reddish brown (5YR 4/3) silt loam; massive; friable; common medium distinct yellowish red (5YR 4/6) soft masses of iron accumulation and common medium prominent gray (5 Y 5/1) iron depletions; strongly acid.

Range in Characteristics

Solum thickness: 16 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—2.5 YR to 10 YR

Value—3 or 4

Chroma—2 to 4

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—2.5 YR to 7.5 YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, silt loam, or loam

Content of rock fragments—0 to 34 percent

C horizon:

Hue—5 YR to 10YR

Value—3 to 5

Chroma—1 to 4

Texture of the fine earth fraction—fine sandy loam, silt loam, or loam

Content of rock fragments—0 to 34 percent

Belgrade Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate

Landform: Terraces on lake plains

Parent material: Glaciolacustrine deposits

Slope range: 0 to 5 percent

Associated soil in a drainage sequence:

Raynham (poorly drained)

Associated similar soils:

Berlin (fine-silty)

Brancroft (fine-silty)

Elmridge (coarse-loamy over clayey)

Associated other soils:

Scitico (poorly drained, fine)

Shaker (poorly drained, coarse-loamy over clayey)

Maybid (very poorly drained, fine)

Taxonomic class: Coarse-silty, mixed, active, mesic Aquic Dystric Eutrudepts

Typical Pedon

Belgrade silt loam, 0 to 5 percent slopes, located in the town of Suffield, 1,000 feet west northwest of the junction of Hill Street and Connecticut Route 190, on the Windsor Locks USGS topographic quadrangle, lat. 41 degrees 58 minutes 50 seconds N., long. 72 degrees 40 minutes 01 seconds W., NAD 27, in a wooded area:

Ap—0 to 8 inches; dark brown (10YR 3/3) silt loam, pale brown (10YR 6/3) dry; moderate medium granular structure; very friable; few very fine, fine and medium roots; strongly acid; clear smooth boundary.

Bw1—8 to 16 inches; dark yellowish brown (10YR 4/4) silt loam; weak fine and medium subangular blocky structure; friable; few fine medium and coarse roots; moderately acid; gradual wavy boundary.

Bw2—16 to 27 inches; yellowish brown (10YR 5/4) silt loam; weak fine and medium subangular blocky structure; friable; few fine, medium and coarse roots; common

fine faint pale brown (10YR 6/3) and distinct grayish brown (10YR 5/2) iron depletions; moderately acid; clear smooth boundary.

C1—27 to 45 inches; light olive brown (2.5Y 5/4) varved silt loam and very fine sandy loam; massive parting to thin platy structure; friable; few fine medium and coarse roots that decrease with depth; common fine and medium distinct light brownish gray (2.5Y 6/2) iron depletions and distinct yellowish brown (10YR 5/6) and prominent strong brown (7.5YR 5/6) soft masses of iron accumulations; moderately acid; clear smooth boundary.

C2—45 to 60 inches; olive brown (2.5Y 4/4) varved silt loam and very fine sandy loam (few thin varves of silty clay loam); massive parting to thin platy structure; friable; few fine and medium distinct light brownish gray (2.5Y 6/2) iron depletions and distinct yellowish brown (10YR 5/6) and prominent strong brown (7.5YR 5/6) soft masses of iron accumulations; moderately acid.

Range in Characteristics

Solum thickness: 20 to 44 inches.

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral in the surface layer and subsoil, and moderately acid to neutral in the substratum (some subhorizon between depths of 10 and 30 inches moderately acid to neutral)

Ap horizon:

Hue—10YR

Value—3 or 4 (dry value 6 or more)

Chroma—2 to 4

Content of rock fragments—less than 1 percent

Bw1 horizon:

Hue—10YR to 5Y

Value—4 or 5

Chroma—4 to 6

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—less than 1 percent

Bw2 horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 to 6

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—less than 1 percent

C horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 to 4

Texture of the fine earth fraction—very fine sandy loam or silt loam

Content of rock fragments—less than 1 percent

Berlin Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer, moderately slow or moderate in the upper subsoil, slow or very slow in the lower subsoil, and very slow in the substratum

Landform: Terraces on lake plains

Parent material: Glaciolacustrine deposits

Slope range: 0 to 8 percent

Associated similar soils:

Belgrade (coarse-silty)

Brancroft (7.5YR or yellower in subsoil and substratum)

Elmridge (coarse-loamy over clayey)

Ludlow (coarse-loamy)

Associated other soils:

Wethersfield (well drained, coarse-loamy)

Scitico (poorly drained, fine)

Maybid (very poorly drained, fine)

Taxonomic class: Fine-silty, mixed, semiactive, mesic Aquic Dystric Eutrudepts

Typical Pedon

Berlin silt loam, 3 to 8 percent slopes, located in the town of Rocky Hill, 450 feet northwest of the intersection of Orchard Street and Raymond Road, 50 feet west of Interstate 91, on the Hartford South USGS topographic quadrangle, lat. 41 degrees 40 minutes 37 seconds N., long. 72 degrees 39 minutes 30 seconds W., NAD 27, in a wooded area:

Ap—0 to 6 inches; dark reddish brown (5YR 3/2) silt loam, light brownish gray (10YR 6/2) dry; weak medium granular structure; friable; many very fine and fine roots; 5 percent fine gravel; strongly acid; abrupt smooth boundary.

Bw1—6 to 12 inches; reddish brown (5YR 4/3) silt loam; weak medium subangular blocky structure; friable; common very fine and fine roots; 5 percent fine gravel; strongly acid; clear wavy boundary.

Bw2—12 to 20 inches; reddish brown (5YR 4/4) silty clay loam; weak coarse subangular blocky structure; friable; few fine roots; 3 percent gravel; strongly acid; clear wavy boundary.

Bw3—20 to 34 inches; dark reddish brown (5YR 3/4) silty clay loam; weak very coarse prismatic structure parting to moderate coarse subangular blocky structure; firm; few fine roots along prism faces; discontinuous black coatings on ped faces; common fine and medium distinct pinkish gray (5YR 6/2), prominent pale brown (10YR 6/3) and distinct light reddish brown (5YR 6/3) iron depletions; 2 percent fine gravel; moderately acid; gradual wavy boundary.

C1—34 to 48 inches; dark reddish brown (5YR 3/4) silty clay loam; weak very coarse prismatic structure parting to thin platy structure along varve bedding planes; firm; discontinuous black coatings on plate surfaces; few fine prominent reddish yellow (7.5YR 6/6, 6/8) soft masses of iron accumulations and distinct reddish gray (5YR 5/2) iron depletions; slightly acid; clear wavy boundary.

C2—48 to 65 inches; dark reddish brown (5YR 3/3) varved silt and clay; silty clay loam weighted average texture; massive separating into thin platy structure along varve bedding planes; very firm; few thin strata of very fine sand; slightly acid.

Range in Characteristics

Solum thickness 24 to 45 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral in the surface layer, strongly acid to neutral in the subsoil, and moderately acid to neutral in the substratum

Ap horizon:

Hue—5YR or 7.5YR

Value—3 or 4 (Dry value 6 or more) (If A horizon present instead of Ap, value 2 or 3)

Chroma—2 or 3 (If A horizon present instead of Ap, chroma 1 to 3)

Content of rock fragments—0 to 10 percent

Upper Bw horizons:

Hue—5YR or 2.5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam or silty clay loam

Content of rock fragments—0 to 10 percent

Lower Bw horizons:

Hue—5YR or 2.5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, silty clay loam, or silty clay

Content of rock fragments—0 to 10 percent

C horizons:

Hue—5YR or 2.5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam to silty clay (very fine sand to clay in individual varves)

Content of rock fragments—0 to 5 percent

Bernardston Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderate in the surface layer and subsoil, slow or very slow in the substratum*Landform:* Hills*Parent material:* Lodgement till derived from phyllite or schist*Slope range:* 3 to 25 percent*Associated similar soils:*

Dummerston (friable substratum, higher elevations)

Fullam (moderately well drained, higher elevations)

Lanesboro (higher elevations)

Paxton (less silty)

Associated other soils:

Taconic (shallow, somewhat excessively drained, higher elevations)

Macomber (moderately deep, higher elevations)

Woodbridge (moderately well drained)

Taxonomic class: Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts**Typical Pedon**

Bernardston silt loam, 3 to 8 percent slopes, located in the town of Salisbury, 50 feet north of Sellick Hill Road, 0.25 miles east of the intersection of Sellick Hill Road and Lincoln City Road, on the Sharon USGS topographic quadrangle, lat. 41 degrees 58 minutes 49 seconds N., long. 73 degrees 26 minutes 47 seconds W., NAD 27, in a hayfield:

Ap—0 to 8 inches, dark brown (10YR 3/3) silt loam, pale brown (10 YR 6/3) dry; weak fine granular structure; friable; many fine roots; 13 percent rock fragments; moderately acid; clear smooth boundary.

Bw1—8 to 14 inches, yellowish brown (10YR 5/4) channery silt loam; weak medium subangular blocky structure; friable; few fine roots; 15 percent rock fragments; strongly acid; gradual wavy boundary.

Bw2—14 to 24 inches, light olive brown (2.5Y 5/4) channery silt loam; weak medium subangular blocky structure; friable; few fine roots; 25 percent rock fragments; very strongly acid; gradual wavy boundary.

BC—24 to 26 inches, light olive brown (2.5Y 5/4) to olive (5Y 5/3) channery silt loam; weak thick platy structure; firm; 25 percent rock fragments; very strongly acid; clear wavy boundary.

Cd—26 to 60 inches, olive brown (2.5Y 4/4) to light olive brown (2.5Y 5/4) channery silt loam; weak thick platy structure; very firm; 30 percent rock fragments, primarily schist; very strongly acid.

Range in Characteristics

Solum thickness: 20 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap or A horizon:

Hue—7.5YR to 2.5Y

Value—3 or 4

Chroma—2 or 3

Content of rock fragments—0 to 14 percent

Bw1 horizon:

Hue—7.5YR to 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—5 to 25 percent

Bw2 horizon:

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—5 to 25 percent

BC horizon:

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—5 to 25 percent

Cd horizon:

Hue—2.5Y or 5Y

Value—4 or 5

Chroma—2 to 4

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—5 to 30 percent

Bice Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid

Landform: Hills

Parent material: Melt-out till derived from granite, schist, and/or gneiss

Slope range: 3 to 45 percent

Associated soils in a drainage sequence:

- Schroon (moderately well drained)
- Loonmeadow (very poorly drained)

Associated other soils:

- Westminster (shallow, somewhat excessively drained)
- Millsite (moderately deep)
- Shelburne (dense substratum)

Taxonomic class: Coarse-loamy, mixed, active, frigid Typic Dystrudepts

Typical Pedon

Bice fine sandy loam, 3 to 8 percent slopes, very stony, located in the town of Norfolk, 2,000 feet north of the intersection of Barry Hill Road and Wheeler Street, 30 feet East of Barry Hill Road, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees 02 minutes 18 seconds N., long. 73 degrees 10 minutes 33 seconds W. NAD 27, in a wooded area:

- Oi—0 to 1 inch; slightly decomposed plant materials derived from leaf and twig litter matted with roots.
- Ap—1 to 7 inches, dark brown (10YR 3/3) fine sandy loam, brown (10YR 5/3) dry; weak fine granular structure; very friable; common fine to coarse roots; 5 percent rock fragments; strongly acid; clear smooth boundary.
- Bw1—7 to 16 inches, dark yellowish brown (10Y 4/6) fine sandy loam; weak medium subangular blocky structure; very friable; common fine to coarse roots; 10 percent rock fragments; strongly acid; gradual smooth boundary.
- Bw2—16 to 24 inches, dark yellowish brown (10Y 4/4) gravelly fine sandy loam; weak medium subangular blocky structure; very friable; common fine to coarse roots; 25 percent rock fragments; moderately acid; clear smooth boundary.
- C—24 to 60 inches; light olive brown (2.5Y 5/3) gravelly sandy loam; massive; friable; few fine roots in the upper part; 25 percent rock fragments; moderately acid.

Range in Characteristics

Solum thickness: 20 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

- Hue—7.5YR or 10YR
- Value—3 or 4 (If A horizon present instead of Ap, value 2 or 3)
- Chroma—2 to 4 (If A horizon present instead of Ap, chroma 1 or 2)
- Content of rock fragments—5 to 14 percent

Bw1 horizon:

- Hue—7.5YR or 10YR
- Value—3 to 6
- Chroma—3 to 6
- Texture of the fine earth fraction—coarse sandy loam, sandy loam, fine sandy loam, or silt loam
- Content of rock fragments—5 to 25 percent

Bw2 horizon:

- Hue—2.5Y or 10YR
- Value—4 to 6
- Chroma—3 to 6
- Texture of the fine earth fraction—fine sandy loam, sandy loam, or loam
- Content of rock fragments—5 to 25 percent

C horizon:

Hue—2.5Y or 10YR

Value—4 to 6

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam, sandy loam or loam

Content of rock fragments—5 to 25 percent

Boscawen Series*Depth class:* Very deep*Drainage class:* Excessively drained*Permeability:* Moderately rapid or rapid in the surface layer and subsoil, and rapid or very rapid in the substratum*Landform:* Outwash plains, terraces, kames, and eskers*Parent material:* Glaciofluvial deposits derived from granite, schist, and/or gneiss*Slope range:* 0 to 45 percent*Associated soils in a drainage sequence:*

Merrimac, cold (somewhat excessively drained)

Sudbury, cold (moderately well drained)

Moosilauke (poorly drained)

Associated similar soils:

Agawam, cold (coarse-loamy over sandy or sandy-skeletal)

Taxonomic class: Sandy-skeletal, mixed, frigid Typic Udorthents**Typical Pedon**

Boscawen gravelly sandy loam, 3 to 15 percent slopes, located in the town of Norfolk, 800 feet west along Windrow Road from the intersection with Westside Road, and 1,000 feet north of Windrow Road, on the Norfolk USGS topographic quadrangle, lat. 41 degrees 58 minutes 46 seconds N., long. 73 degrees 12 minutes 50 seconds W., NAD 27, in a wooded area:

Oi—0 to 1 inch; slightly decomposed plant materials

A—1 to 2 inches; very dark brown (10YR 2/2) gravelly sandy loam; weak fine granular structure; friable; many very fine and medium roots, common coarse and very coarse roots; 20 percent gravel; very strongly acid; abrupt wavy boundary.

Bw—2 to 9 inches; brown (7.5YR 4/4) very gravelly sandy loam; weak, medium subangular blocky structure; friable; common very fine, fine and medium roots, few coarse and very coarse roots; 40 percent gravel, 5 percent cobbles; strongly acid; clear wavy boundary.

BC—9 to 16 inches; dark yellowish brown (10YR 4/6) very gravelly loamy sand; single grain; loose; few very fine, fine, and medium roots, few coarse and very coarse roots; 50 percent gravel, 5 percent cobbles; strongly acid; clear wavy boundary.

C1—16 to 29 inches; dark yellowish brown (10YR 4/6) extremely gravelly sand; single grain; loose; few very fine, fine and medium roots; 60 percent gravel, 10 percent cobbles; strongly acid; clear wavy boundary.

C2—29 to 34 inches; yellowish brown (10YR 5/6) very gravelly coarse sand; single grain; loose; 40 percent gravel, 10 percent cobbles; strongly acid; clear smooth boundary.

C3—34 to 40 inches; yellowish brown (10YR 5/6) and light olive brown (2.5Y 5/6) gravelly sand; single grain; loose; 20 percent gravel; moderately acid; clear smooth boundary.

C4—40 to 44 inches; light olive brown (2.5Y 5/6) fine sand; single grain; loose; 5 percent gravel; moderately acid; clear smooth boundary.

C5—44 to 67 inches; light olive brown (2.5Y 5/6) and yellowish brown (10YR 5/6) sand; single grain; loose; 10 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: 6 to 24 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to strongly acid in the surface layer and subsoil, and very strongly acid to moderately acid in the substratum

A horizon:

Hue—7.5YR or 10YR

Value—2 to 4

Chroma—2 to 4

Content of rock fragments—15 to 20 percent

Bw horizon:

Hue—7.5YR or 10YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—loamy sand or sandy loam

Content of rock fragments—20 to 50 percent

BC horizon (where present)

Hue—7.5YR or 10YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—loamy sand

Content of rock fragments—35 to 60 percent

C horizon:

Hue—7.5YR to 2.5Y

Value—4 or 5

Chroma—4 to 6

Texture—stratified fine sand to very gravelly coarse sand

Content of rock fragments—40 to 75 percent (weighted average, individual subhorizons may be lower)

Brancroft Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer, moderate or moderately slow in the upper subsoil, slow or very slow in the lower subsoil, and very slow in the substratum

Landform: Terraces on lake plains

Parent material: Glaciolacustrine deposits

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Scitico (poorly drained)

Maybid (very poorly drained)

Associated similar soils:

Belgrade (coarse-silty)

Berlin (redder in the subsoil and substratum)

Elmridge (coarse-loamy over clayey)

Taxonomic class: Fine-silty, mixed, active, mesic Aquic Dystric Eutrudepts

Typical Pedon

Brancroft silt loam, 3 to 8 percent slopes, located in the town of Suffield, 1,800 feet west along Halladay Avenue from the intersection of Mapleton Avenue and Halladay Avenue, 600 feet north of Halladay Avenue, on the West Springfield USGS topographic quadrangle, lat. 42 degrees 00 minutes 35 seconds N., long. 72 degrees 38 minutes 08 seconds W., NAD 27, in a grassy field:

- Ap—0 to 6 inches; dark grayish brown (10YR 4/2) silt loam; light brownish gray (2.5Y 6/2) dry; moderate fine and medium granular structure; friable; many fine and very fine roots; strongly acid; clear smooth boundary.
- Bw1—6 to 17 inches; brown (10YR 4/3) silt loam; moderate medium subangular blocky structure; friable; few fine and very fine roots; strongly acid; clear smooth boundary.
- Bw2—17 to 22 inches; brown (10YR 4/3) silty clay loam; weak very coarse prismatic structure parting to weak, thin platy structure; firm; few fine roots along prism faces; common fine and medium prominent strong brown (7.5YR 5/6, 5/8) soft masses of iron accumulations and faint grayish brown (10YR 5/2) iron depletions on prism faces and plate surfaces; strongly acid; clear smooth boundary.
- Bw3—22 to 32 inches; light olive brown (2.5Y 5/4) silt loam; weak very coarse prismatic structure parting to weak, thin platy structure; firm; few fine roots along prism faces; common light olive gray (5Y 6/2) coatings on prism faces; many fine to coarse prominent strong brown (7.5YR 5/6) soft masses of iron accumulations and faint light olive gray (5Y 6/2) iron depletions on prism faces and plate surfaces; strongly acid; clear smooth boundary.
- C1—32 to 43 inches; olive (5Y 5/3) silty clay loam; massive; firm; many light olive gray (5Y 6/2) coatings on prism faces; many fine to coarse prominent strong brown (7.5YR 5/6) soft masses of iron accumulations and faint light olive gray (5Y 6/2) iron depletions on prism faces and plate surfaces; moderately acid; clear smooth boundary.
- C2—43 to 66 inches; olive (5Y 5/3) varved silt and clay (silt loam weighted average); massive; firm; few fine and medium prominent strong brown (7.5YR 5/6) soft masses of iron accumulations and faint light olive gray (5Y 6/2) depletions on varve surfaces; neutral.

Range in Characteristics

Solum thickness: 24 to 45 inches

Depth to bedrock: More than 80 inches

Reaction: *Very strongly acid to slightly acid in the surface layer, strongly acid to neutral in the subsoil, and moderately acid to neutral in the substratum*

Ap horizon:

Hue—10YR or 2.5Y

Value—2 to 4 (dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 or 3 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—0 to 3 percent

Bw1 horizon:

Hue—7.5YR to 5Y

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam or silty clay loam

Content of rock fragments—0 to 3 percent

Lower Bw horizons:

Hue—10YR to 5Y

Value—3 to 6
 Chroma—3 or 4
 Texture of the fine earth fraction—silt loam or silty clay loam
 Content of rock fragments—0 to 3 percent

C horizons:

Hue—7.5YR to 5Y
 Value—3 to 6
 Chroma—2 to 4
 Texture of the fine earth fraction—silt loam or silty clay loam (silt loam, silty clay loam, or silty clay in individual varves)
 Content of rock fragments—0 to 3 percent

Branford Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Eolian deposits over glaciofluvial deposits derived from sandstone, shale, and basalt

Slope range: 0 to 15 percent

Associated soil in a drainage sequence:

Ellington (moderately well drained)

Associated similar soils:

Manchester (excessively drained, sandy and gravelly)

Penwood (excessively drained, sandy)

Hartford (somewhat excessively drained, sandy)

Associated other soils:

Raypol (poorly drained)

Walpole (poorly drained, sandy)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts

Typical Pedon

Branford silt loam, 0 to 3 percent slopes, located in the town of Branford, 600 feet west of the intersection of Connecticut Route 139 and the New Haven Trap Rock Company railroad, on the Branford USGS topographic quadrangle, lat. 41 degrees 18 minutes 07 seconds N., long. 72 degrees 46 minutes 42 seconds W., NAD 27:

Ap—0 to 8 inches; dark reddish brown (5YR 3/3) silt loam, light reddish brown (5YR 6/3) dry; weak medium granular structure; friable; common very fine and fine roots; 10 percent gravel; moderately acid; clear smooth boundary.

Bw1—8 to 18 inches; reddish brown (5YR 4/4) loam; weak medium subangular blocky structure; friable; few fine roots; common earthworm holes and worm casts; 10 percent gravel; strongly acid; gradual wavy boundary.

Bw2—to 24 inches; reddish brown (5YR 4/4) gravelly loam; weak coarse subangular blocky structure; very friable; few fine roots; 15 percent gravel; strongly acid; clear wavy boundary.

2C—24 to 65 inches; reddish brown (5YR 4/3) stratified loamy fine sand to very gravelly coarse sand; single grain; loose; 25 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—5YR to 10YR

Value—3 or 4 (dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon present instead of Ap, chroma 1 to 3)

Content of rock fragments—0 to 14 percent

Bw1 horizon:

Hue—5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam, loam, or fine sandy loam

Content of rock fragments—0 to 30 percent

Bw2 horizon:

Hue—5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam, loam, or fine sandy loam

Content of rock fragments—0 to 30 percent

2C horizon:

Hue—2.5YR or 5YR

Value—3 to 6

Chroma—3 to 6

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—10 to 50 percent

Brayton Series Taxadjunct

This taxonomic description is for the Brayton loam of map units 442 and 443.

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Depressions and drainageways on hills

Parent material: Lodgement till derived from granite, schist and gneiss

Slope range: 0 to 8 percent

Associated soils in a drainage sequence:

Shelburne (well drained)

Ashfield (moderately well drained)

Associated other soils:

Loonmeadow (very poorly drained, fine-loamy)

Bucksport (very poorly drained, more than 51 inches of organic material)

Wonsqueak (very poorly drained, 16 to 51 inches of organic material over loamy deposits)

Taxonomic class: Coarse-loamy, mixed, active, nonacid, frigid Typic Humaquepts

Typical Pedon

Brayton loam, in an area of Brayton-Loonmeadow complex, extremely stony, located in the town of Norfolk, 600 feet east along North Colebrook Road from the

intersection with State Line Hill Road, and 300 feet North of North Colebrook Road, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees 01 minutes 41 seconds N., long. 73 degrees 08 minutes 17 seconds W., NAD 27, in a wooded area:

- Oe—0 to 2 inches; very dark gray (2.5Y 3/1) moderately decomposed plant materials; many very fine to medium roots, few coarse and very coarse roots; very strongly acid; clear wavy boundary.
- A—2 to 10 inches; very dark grayish brown (10YR 3/2) loam, grayish brown (10 YR 5/2) dry; moderate medium granular structure; very friable, slightly sticky, non-plastic; many very fine to medium roots, few coarse and very coarse roots; common medium distinct brown (7.5 YR 4/4) masses of iron accumulation, few fine faint dark grayish brown (2.5 Y 4/2) iron depletions; 10 percent gravel, 1 percent cobbles; strongly acid; clear wavy boundary.
- Bg1—10 to 17 inches; dark grayish brown (2.5Y 4/2) gravelly sandy loam; weak medium subangular blocky structure; very friable, non-sticky, non-plastic; few very fine to coarse roots; few medium faint grayish brown (2.5Y 5/2) iron depletions, common fine prominent brown (7.5YR 4/4) masses of iron accumulation; 12 percent gravel, 2 percent cobbles, 1 percent stones; moderately acid; gradual smooth boundary.
- Bg2—17 to 22 inches; grayish brown (2.5Y 5/2) gravelly sandy loam; weak coarse subangular blocky structure; very friable, non-sticky, non-plastic; few fine faint gray (2.5Y 5/1) iron depletions and common medium prominent brown (7.5YR 4/4) masses of iron accumulation; 12 percent gravel, 2 percent cobbles, 1 percent stones; moderately acid; clear smooth boundary.
- Bg3—22 to 28 inches; olive gray (5Y 5/2) sandy loam; weak coarse subangular blocky structure; friable, non-sticky, non-plastic; few fine faint gray (5Y 5/1) iron depletions, common medium distinct olive brown (2.5Y 4/4) masses of iron accumulation; 10 percent gravel, 2 percent cobbles, 1 percent stones; neutral; clear smooth boundary.
- Cd1—28 to 42 inches; olive brown (2.5Y 4/3) gravelly sandy loam; massive; firm, non-sticky, non-plastic; common medium faint grayish brown (2.5Y 5/2) iron depletions, common medium prominent yellowish brown (10 YR 5/6) masses of iron accumulation; 15 percent gravel, 2 percent cobbles, 2 percent stones; neutral; gradual smooth boundary.
- Cd2—42 to 65 inches; olive (5Y 5/3) gravelly sandy loam; massive; firm, non-sticky, non-plastic; 15 percent gravel, 2 percent cobbles, 2 percent stones; neutral.

Range in Characteristics

Solum thickness: 20 to 28 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to moderately acid in the surface layers, strongly acid to slightly acid in the upper subsoil, strongly acid to neutral in the lower subsoil, and moderately acid to neutral in the substratum (some subhorizon has a pH value greater than 5.5)

A horizon:

Hue—10YR to 5Y

Value—2 or 3 (dry value is 5 or less)

Chroma—1 to 3

Content of rock fragments—5 to 14 percent

Bg horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam, loam, very fine sandy loam, fine sandy loam, or sandy loam

Content of rock fragments—5 to 34 percent

Cd horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 to 4

Texture of the fine earth fraction— loam, fine sandy loam, or sandy loam

Content of rock fragments—5 to 34 percent

The Brayton soils in eastern Litchfield County are taxadjuncts because the soils are very deep and the surface layers are thicker and darker in color than the range for the series. This difference, however, does not significantly affect the use, management, or interpretation of the soils. In the eastern area of Litchfield County, the Brayton soils are coarse-loamy, mixed, active, nonacid frigid Typic Humaquepts.

Brayton Series Taxadjunct

This taxonomic description is for the Brayton mucky silt loam of map unit 409B.

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate in the surface layer, moderate or moderately rapid in the subsoil, and slow or very slow in the substratum

Landform: Depressions and drainageways on hills

Parent material: Lodgement till derived from phyllite or schist

Slope range: 0 to 8 percent

Associated soils in a drainage sequence:

Lanesboro (well drained)

Fullam (moderately well drained)

Associated other soils:

Taconic (shallow, somewhat excessively drained)

Dummerston (well drained, friable substratum)

Macomber (moderately deep, well drained)

Bucksport (very poorly drained, more than 51 inches of organic material)

Wonsqueak (very poorly drained, 16 to 51 inches of organic material over loamy deposits)

Taxonomic class: Coarse-loamy, mixed, active, nonacid, frigid shallow Typic Endoaquepts

Typical Pedon

Brayton mucky silt loam, 0 to 8 percent slopes, very stony, located in the town of Salisbury, 6,000 feet south of the Connecticut/Massachusetts state line on Mt. Washington Road, 2500 feet northwest following the jeep trail, 1250 feet northeast of jeep trail and stream crossing, on the Bash Bish USGS topographic quadrangle, lat. 42 degrees 02 minutes 30 seconds N., long. 73 degrees 28 minutes 29 seconds W., NAD 27, in a wooded area:

Oe—0 to 3 inches; moderately decomposed plant materials derived from hemlock needles; many very fine to very coarse roots; abrupt smooth boundary.

Oa—3 to 6 inches; highly decomposed plant materials derived from hemlock needles; many very fine to very coarse roots; extremely acid; abrupt smooth boundary.

A—6 to 7 inches; black (10YR 2/1) mucky silt loam, dark gray (10 YR 4/1) dry; weak fine granular structure; very friable; many very fine to very coarse roots; many

very fine irregular pores; 10 percent gravel; extremely acid; very abrupt wavy boundary.

Bg1—7 to 9 inches; light brownish gray (2.5Y 6/2) silt loam; weak medium subangular blocky structure; friable; few very fine roots; few medium faint light gray (2.5Y 7/1) iron depletions; 10 percent gravel; very strongly acid; abrupt wavy boundary.

Bg2—9 to 13 inches; olive gray (5Y 5/2) gravelly loam; weak medium subangular blocky structure; friable; few very fine to fine roots; common fine to medium distinct gray (2.5Y 6/1) iron depletions and many fine to medium prominent brownish yellow (10YR 6/8) soft masses of iron accumulation; 15 percent gravel, 2 percent cobbles; slightly acid; clear smooth boundary.

Cd1—13 to 18 inches; olive brown (2.5Y 4/3) gravelly fine sandy loam; weak coarse platy structure; very firm; common fine to medium faint light brownish gray (2.5Y 6/2) iron depletions and common fine to medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; 20 percent gravel, 5 percent cobbles; slightly acid; gradual smooth boundary.

Cd2—18 to 23 inches; olive brown (2.5Y 4/3) gravelly fine sandy loam; massive; firm; few fine to medium distinct light olive gray (5Y 6/2) iron depletions and few fine to medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; 20 percent gravel, 5 percent cobbles; slightly acid; gradual smooth boundary.

Cd3—23 to 60 inches; olive brown (2.5Y 4/4) gravelly fine sandy loam; massive; firm; few fine to medium distinct light olive gray (5Y 6/2) iron depletions and few fine to medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; 20 percent gravel, 5 percent cobbles; neutral.

Range in Characteristics

Solum thickness: 10 to 20 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to moderately acid in the surface layers, strongly acid to slightly acid in the subsoil, and moderately acid to neutral in the substratum (some subhorizon has a pH value greater than 5.5)

A horizon:

Hue—10YR to 5Y

Value—2 to 4

Chroma—1 to 4

Content of rock fragments—5 to 14 percent

Bg1 horizon:

Hue—10YR to 5Y

Value—5 or 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam or loam

Content of rock fragments—5 to 34 percent

Bg2 horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 or less

Texture of the fine earth fraction—loam or fine sandy loam

Content of rock fragments—5 to 34 percent

Cd horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 to 4

Texture of the fine earth fraction—loam or fine sandy loam

Content of rock fragments—5 to 34 percent

The Brayton soils in western Litchfield County are taxadjuncts because the surface layers are darker in color than the range for the series. This difference, however, does not significantly affect the use, management, or interpretation of the soils. In the western area of Litchfield County, the Brayton soils are coarse-loamy, mixed, active, nonacid frigid shallow Typic Endoaquepts.

Brimfield Series

Depth class: Shallow

Drainage class: Somewhat excessively drained

Permeability: Moderate or moderately rapid

Landform: Bedrock-controlled hills and ridges

Parent material: Melt-out till derived from schist

Slope range: 3 to 45 percent

Associated soil in a drainage sequence:

Brookfield (very deep, well drained, coarse-loamy)

Associated similar soil:

Hollis (shallow over schist, granite, and gneiss)

Associated other soils:

Leicester (very deep, poorly drained, coarse-loamy)

Whitman (very deep, very poorly drained, coarse-loamy)

Taxonomic class: Loamy, mixed, active, mesic Lithic Dystrudepts

Typical Pedon

Brimfield fine sandy loam, in an area of Brookfield-Brimfield-Rock outcrop complex, 3 to 15 percent slopes, very rocky, located in the town of Stafford, 50 feet north of the intersection of Connecticut Route 190 and Collette Road, on the Stafford Springs USGS topographic quadrangle, lat. 41 degrees 57 minutes 47 seconds N., long. 72 degrees 15 minutes 05 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials

A—1 to 3 inches; very dark grayish brown (10YR 3/2) fine sandy loam, grayish brown (10 YR 5/2) dry; weak fine granular structure; very friable; many fine and medium roots; 5 percent gravel and channers; very strongly acid; abrupt smooth boundary.

Bw1—3 to 6 inches; dark reddish brown (5YR 3/4) fine sandy loam; weak medium subangular blocky structure; very friable; many fine and medium roots; 10 percent gravel and channers; very strongly acid, clear wavy boundary.

Bw2—6 to 17 inches; yellowish red (5YR 5/6) gravelly fine sandy loam; weak coarse subangular blocky structure; few fine, medium and coarse roots; 20 percent gravel and channers; strongly acid; abrupt wavy boundary.

2R—17 inches; brown mica schist bedrock.

Range in Characteristics

Solum thickness: 10 to 20 inches

Depth to bedrock: 10 to 20 inches

Reaction: Very strongly acid to moderately acid

In some pedons, the Oe horizon is absent.

A horizon:

Hue—5YR to 10YR

Value—2 to 4

Chroma—1 to 3

Content of rock fragments—5 to 14 percent

Bw1 horizon:

Hue—2.5 YR to 7.5YR

Value—3 to 6

Chroma—4 to 8

Texture of the fine earth fraction—fine sandy loam, loam or sandy loam

Content of rock fragments—5 to 14 percent

Bw2 horizon:

Hue—2.5YR to 7.5YR (7.5YR cannot be in both Bw1 and Bw2 horizons)

Value—3 to 6

Chroma—4 to 8

Texture of the fine earth fraction—fine sandy loam, sandy loam, or loam

Content of rock fragments—5 to 34 percent

Broadbrook Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderate in the surface layer and subsoil, and slow or very slow in the substratum*Landform:* Hills, drumlins and till plains*Parent material:* Eolian deposits over lodgement till derived from gneiss, basalt, schist, and shale*Slope range:* 0 to 25 percent*Associated soil in a drainage sequence:*

Rainbow (moderately well drained)

Associated similar soil:

Narragansett (coarse loamy over sandy or sandy-skeletal)

Taxonomic class: Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts**Typical Pedon**

Broadbrook silt loam, 3 to 8 percent slopes, located in the town of Ledyard, 2,500 feet east northeast of the junction of Connecticut Route 117 and Thomas Road, 1,000 feet north of Thomas Road, on the Uncasville USGS topographic quadrangle, lat. 41 degrees 28 minutes 24 seconds N., long. 72 degrees 01 minutes 09 seconds W., NAD 27, in a cultivated field:

Ap—0 to 8 inches; very dark brown (10YR 2/2) silt loam, pale brown (10YR 6/3) dry; moderate medium granular structure; very friable; few fine roots; 5 percent gravel; moderately acid; clear wavy boundary.

Bw1—8 to 14 inches; dark yellowish brown (10YR 4/6) silt loam; moderate medium subangular blocky structure; very friable; few fine roots; 5 percent gravel; moderately acid; clear wavy boundary.

Bw2—14 to 25 inches; dark yellowish brown (10YR 4/4) silt loam; moderate medium subangular blocky structure; very friable; 5 percent gravel; moderately acid; gradual wavy boundary.

2Cd—25 to 65 inches; olive brown (2.5Y 4/4) gravelly fine sandy loam; weak medium platy structure; very firm, brittle; 15 percent gravel and cobbles; moderately acid.

Range in Characteristics*Solum thickness:* 20 to 40 inches*Depth to bedrock:* More than 80 inches*Reaction:* Very strongly acid to moderately acid

Ap horizon:

Hue—7.5YR or 10YR

Value—2 to 4 (Dry value is 6 or more) (If A horizon present instead of Ap, value 2 or 3)

Chroma—2 to 4 (If A horizon present instead of Ap, chroma 1 to 3)

Content of rock fragments—0 to 14 percent

Bw1 horizon:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—4 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam or loam

Content of rock fragments—0 to 20 percent

Bw2 horizon:

Hue—7.5YR to 2.5Y

Value—4 or 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam or loam

Content of rock fragments—0 to 20 percent

2Cd horizon:

Hue—.5YR to 5Y

Value—4 to 6

Chroma—2 to 6

Texture of the fine earth fraction—fine sandy loam or sandy loam

Content of rock fragments—5 to 34 percent

Brookfield Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderate or moderately rapid*Landform:* Bedrock-controlled hills and ridges*Parent material:* Melt-out till derived from schist*Slope range:* 3 to 45 percent*Associated soil in a drainage sequence:*

Brimfield (shallow, somewhat excessively drained, loamy)

Associated similar soils:

Canton (yellower, coarse-loamy over sandy or sandy-skeletal)

Charlton (yellower)

Taxonomic class: Coarse-loamy, mixed, active, mesic Typic Dystrudepts**Typical Pedon**

Brookfield fine sandy loam, in an area of Brookfield-Brimfield-Rock outcrop complex, 3 to 15 percent slopes, very rocky, located in the town of Union, 50 feet east along Szych Road from the intersection with Route 190, 200 feet north of Szych Road, and 200 feet south of Buckley Pond, on the Westford USGS topographic quadrangle, lat. 41 degrees 59 minutes 19 seconds N., long. 72 degrees 10 minutes 05 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials

A—1 to 3 inches; dark brown (10YR 3/3) fine sandy loam, brown (10 YR 5/3) dry;

weak fine granular structure; very friable; many fine roots; 5 percent gravel; very strongly acid; abrupt smooth boundary.

Bw1—3 to 13 inches; yellowish red (5YR 4/6) gravelly fine sandy loam; weak medium granular structure; very friable; common fine and medium roots; common mica flakes; 15 percent gravel and cobbles; very strongly acid; gradual wavy boundary.

Bw2—13 to 27 inches; strong brown (7.5YR 5/8) gravelly fine sandy loam; massive; very friable; few fine and medium roots; common mica flakes; 15 percent gravel and cobbles; strongly acid; clear wavy boundary.

C—27 to 60 inches; strong brown (7.5YR 5/6) gravelly sandy loam; grayish brown (2.5Y 5/2) streaks; massive; very friable; few large roots; many mica flakes; 25 percent gravel and cobbles; strongly acid; clear wavy boundary.

Range in Characteristics

Solum thickness: 20 to 38 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—7.5YR or 10YR

Value—2 or 3 (If Ap horizon is present instead of A, value is 3 or 4)

Chroma—1 to 3 (If Ap horizon is present instead of A, chroma is 2 to 4)

Content of rock fragments—5 to 14 percent

Bw1 horizon:

Hue—2.5YR or 5YR

Value—3 to 6

Chroma—4 to 8

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—15 to 34 percent

Bw2 horizon:

Hue—5YR to 10YR

Value—4 or 5

Chroma—4 to 8

Texture of fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—15 to 34 percent

C horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—2 to 6

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—5 to 34 percent

Bucksport Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderately slow to moderately rapid (may range to very rapid in the underlying mineral soil)

Landform: Depressions

Parent material: Organic materials

Slope range: 0 to 2 percent

Associated similar soil:

Wonsqueak (16 to 51 inches of organic material over loamy deposits)

Associated other soils:

Brayton (poorly drained, coarse-loamy)

Loonmeadow (very poorly drained, coarse-loamy)

Taxonomic class: Euic, frigid Typic Haplosaprists

Typical Pedon

Bucksport muck, located in the town of Norfolk, 3,000 feet northwest along Wheeler Street from the intersection with Barry Hill Rd and 1,000 feet west of Wheeler Street, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees 02 minutes 08 seconds N., long. 73 degrees 11 minutes 18 seconds W., NAD 27, in a bog:

Oa1—0 to 9 inches; very dark brown (7.5YR 2.5/2) muck; 15 percent fiber, 5 percent rubbed; massive; 50 percent woody and 50 percent herbaceous fibers; strongly acid; clear smooth boundary.

Oa2—9 to 33 inches; black (7.5YR 2.5/1) muck; 20 percent fiber, 10 percent rubbed; massive; 60 percent woody and 40 percent herbaceous fibers; moderately acid; clear smooth boundary.

Oa3—33 to 50 inches; black (7.5YR 2.5/1) muck; 10 percent fiber, 2 percent rubbed; massive; 80 percent woody and 20 percent herbaceous fibers; moderately acid; clear smooth boundary.

Oa4—50 to 59 inches; black (10YR 2/1) muck; 2 percent fiber, 0 percent rubbed; massive; 80 percent woody and 20 percent herbaceous fibers; very strongly acid; gradual smooth boundary.

2Cg—59 to 63 inches; olive gray (5Y4/2) gravelly sand; single grain; loose; 25 percent gravel; slightly acid.

Range in Characteristics

Organic material thickness: More than 51 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to strongly acid in the surface tier, very strongly acid to slightly acid in the subsurface tier and bottom tiers, and very strongly acid to neutral in the mineral substratum.

Surface tier:

Hue—neutral or 2.5YR to 10YR

Value—2 to 4

Chroma—0 to 2

Subsurface and bottom tiers:

Hue—2.5YR to 10YR

Value—2 to 4

Chroma—1 to 3

2Cg horizon: (present in some pedons)

Hue—neutral or 5YR to 5GY

Value—3 to 6

Chroma—0 to 4

Texture of the fine earth fraction—sand, sandy loam, fine sandy loam, or loam

Content of rock fragments—0 to 25 percent

Canton Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderately rapid in the surface layer and subsoil, and rapid in the substratum

Landform: Hills

Parent material: Melt-out till derived from granite, gneiss, and schist

Slope range: 3 to 35 percent

Associated soils in a drainage sequence:

Sutton (moderately well drained)
Leicester (poorly drained)

Associated similar soils:

Charlton (coarse-loamy)
Cheshire (redder, coarse-loamy)
Montauk (dense substratum, coarse-loamy)
Narragansett (finer texture in the surface layer and subsoil)
Paxton (dense substratum, coarse-loamy)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, semi-active, mesic Typic Dystrudepts

Typical Pedon

Canton fine sandy loam, in an area of Canton and Charlton fine sandy loams, 3 to 8 percent slopes, very stony, located in the town of Chester, in Cockaponset State Forest, 6,000 feet west of the intersection of Connecticut Route 148 and Connecticut Route 145 and 2800 feet north, on the Haddam USGS topographic quadrangle, lat. 41 degrees 24 minutes 14 seconds N., long. 72 degrees 31 minutes 24 seconds W., NAD 27, in a wooded area:

- Oe—0 to 1 inch, very dark brown (10YR 2/2) moderately decomposed plant materials; weak fine granular structure; very friable; many fine roots; extremely acid; abrupt smooth boundary.
- A—1 to 3 inches, dark brown (10YR 3/3) gravelly fine sandy loam, pale brown (10 YR 6/3) dry; weak fine granular structure; very friable, many fine and few medium and coarse roots; 10 percent gravel, 5 percent cobbles, 2 percent stones; very strongly acid; abrupt smooth boundary.
- Bw1—3 to 15 inches, strong brown (7.5YR 5/6) gravelly loam; weak fine subangular blocky structure; very friable; many fine and few medium and coarse roots; 10 percent gravel, 5 percent cobbles, 2 percent stones; very strongly acid; gradual wavy boundary.
- Bw2—15 to 24 inches, yellowish brown (10YR 5/6) gravelly loam; weak fine subangular blocky structure; very friable; few fine medium and coarse roots; 10 percent gravel, 5 percent cobbles, 2 percent stones; very strongly acid; gradual wavy boundary.
- Bw3—24 to 30 inches, brownish yellow (10YR 6/6) gravelly loam; weak fine subangular blocky structure; very friable; few fine and medium roots; 10 percent gravel, 5 percent cobbles, 2 percent stones; very strongly acid; abrupt smooth boundary.
- 2C—30 to 60 inches, light brownish gray (2.5Y 6/2) and olive (5Y 5/3) very gravelly loamy sand; massive; very friable; few fine roots to 40 inches; 30 percent gravel, 15 percent cobbles; strongly acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to moderately acid

In some pedons, the Oe horizon may be absent.

A horizon:

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Content of rock fragments—15 to 25 percent

Bw1 horizon:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—4 to 6

Texture of the fine earth fraction— loam, very fine sandy loam, or fine sandy loam

Content of rock fragments—5 to 25 percent

Lower Bw horizons:

Hue—10YR or 2.5Y

Value—4 to 7

Chroma—4 to 6

Texture of the fine earth fraction— loam, very fine sandy loam, or fine sandy loam

Content of rock fragments—5 to 25 percent

C horizon:

Hue—2.5Y or 5Y (10YR with chroma of 4 to 6 is allowed)

Value—5 to 7

Chroma—2 or 3 (if hue is 10YR, chroma is 4 to 6)

Texture of the fine earth fraction—loamy sand, loamy fine sand, or loamy coarse sand

Content of rock fragments—15 to 50 percent

Catden Series*Depth class:* Very deep*Drainage class:* Very poorly drained*Permeability:* Moderate or moderately rapid*Landform:* Depressions*Parent Material:* Organic materials*Slope range:* 0 to 2 percent*Associated similar soils:*

Freetown (dysic reaction class)

Natchaug (16 to 51 inches of organic materials, over loamy deposits)

Timakwa (16 to 51 inches of organic materials, over sandy or sandy-skeletal)

Taxonomic class: Euic, mesic Typic Haplosaprists**Typical Pedon**

Catden muck, in an area of Catden and Freetown soils, located in the town of Ashford, 3,500 feet west on Horse Hill Road from the intersection of North Road and Horse Hill Road, 75 feet north of Horse Hill Road, on the Westford USGS topographic quadrangle, lat. 41 degrees 54 minutes 05 seconds N., long. 72 degrees 08 minutes 41 seconds W., NAD 27, in a swamp:

Oa1—0 to 2 inches; black (10YR 2/1) muck; 25 percent fibers, 15 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; many very fine to medium roots; 10 percent woody fragments; less than 5 percent mineral material; very strongly acid (pH 4.6 in 0.01 M calcium chloride); clear smooth boundary.

Oa2—2 to 18 inches; black (7.5YR 2.5/1) muck; 15 percent fibers, 10 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; many very fine to fine roots; less than 5 percent mineral material; very strongly acid (pH 4.7 in 0.01 M calcium chloride); clear smooth boundary.

Oa3—18 to 47 inches; black (7.5YR 2.5/1) muck; 40 percent fibers, 10 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; less than 5 percent mineral material; very strongly acid (pH 4.8 in 0.01 M calcium chloride); clear smooth boundary. Oa4—47 to 49 inches; very dark brown (7.5YR 2.5/2) muck; 5

percent fibers, 0 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; less than 5 percent mineral material; very strongly acid (pH 4.9 in 0.01 M calcium chloride); clear smooth boundary.

Oa5—49 to 61 inches; very dark brown (10YR 2/2) muck; 2 percent fibers, 0 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; less than 5 percent mineral material; very strongly acid (pH 4.8 in 0.01 M calcium chloride).

Range in Characteristics

Solum thickness: Organic layers more than 51 inches thick

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral (in 0.01 M calcium chloride); some layers may range to extremely acid

Woody fragments: 0 to 30 percent

Mineral material: Less than 5 percent

Oa1 horizon:

Hue—5YR to 2.5Y, or is neutral

Value—1 to 4

Chroma—0 to 6

Lower Oa horizons:

Hue—5YR to 2.5Y, or is neutral

Value—2 or 3

Chroma—0 to 4

Charlton Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid

Landform: Hills

Parent material: Melt-out till derived from granite, gneiss and/or schist

Slope range: 3 to 45 percent

Associated soils in a drainage sequence:

Sutton (moderately well drained)

Leicester (poorly drained)

Associated similar soils:

Canton (coarse-loamy over sandy or sandy-skeletal)

Cheshire (redder)

Montauk (dense substratum)

Narragansett (coarse loamy over sandy or sandy-skeletal)

Paxton (dense substratum)

Taxonomic class: Coarse-loamy, mixed, active, mesic Typic Dystrudepts

Typical Pedon

Charlton fine sandy loam, in an area of Charlton-Chatfield complex, 3 to 15 percent slopes, very rocky, located in the town of Middlebury, 3,800 feet along Long Meadow Road from the intersection with South Street, 450 feet southeast along a gravel road, 50 feet west of gravel road, 400 feet northeast of Long Meadow Pond and 700 feet east of power transmission line, on the Naugatuck USGS topographic quadrangle, lat. 41 degrees 29 minutes 50 seconds N., long. 73 degrees 06 minutes 29 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials

A—1 to 4 inches; dark brown (10YR 3/3) fine sandy loam, pale brown (10 YR 6/3);

weak fine granular structure; very friable; many fine roots; 5 percent gravel; very strongly acid; clear wavy boundary.

Bw1—4 to 7 inches; brown (7.5YR 4/4) fine sandy loam; weak coarse granular structure; very friable; many fine and medium roots; 5 percent gravel; very strongly acid; clear wavy boundary.

Bw2—7 to 19 inches; yellowish brown (10YR 5/6) fine sandy loam; weak medium subangular blocky structure; very friable; common fine and medium roots; 10 percent gravel and cobbles; very strongly acid; abrupt wavy boundary.

Bw3—19 to 27 inches; light olive brown (2.5Y 5/4) gravelly fine sandy loam; massive; very friable; few medium roots; 15 percent gravel and cobbles; very strongly acid; abrupt wavy boundary.

C—27 to 65 inches; grayish brown (2.5Y 5/2) gravelly fine sandy loam, thin lenses of loamy sand; massive; friable, some lenses firm; few medium roots; 25 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 20 to 38 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

In some pedons, the Oe horizon may be absent.

A horizon

Hue—7.5YR or 10YR

Value—2 or 3 (If Ap horizon is present instead of A, value is 3 or 4)

Chroma—1 to 3 (If Ap horizon is present instead of A, chroma is 2 to 4)

Content of rock fragments—5 to 14 percent

Bw1 and Bw2 horizons:

Hue—7.5YR or 10YR

Value—4 to 6

Chroma—4 to 6

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—5 to 34 percent

Bw3 horizon:

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—4 to 6

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—5 to 34 percent

C horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 to 6

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—5 to 34 percent

Chatfield Series

Depth class: Moderately deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid

Landform: Bedrock controlled hills and ridges

Parent material: Melt-out till derived from granite, gneiss and schist

Slope range: 3 to 45 percent

Associated other soils:

Canton (very deep, coarse-loamy over sandy or sandy-skeletal)

Charlton (very deep)

Hollis (shallow, loamy)

Taxonomic class: Coarse-loamy, mixed, superactive, mesic Typic Dystrudepts

Typical Pedon

Chatfield gravelly fine sandy loam, in an area of Charlton-Chatfield complex, 3 to 15 percent slopes, very rocky, located in the town of East Hampton, 1,450 feet west of the intersection of Connecticut Route 196 and Connecticut Route 151, and 3000 feet north of Connecticut Route 151, on the Moodus USGS topographic quadrangle, lat. 41 degrees 31 minutes 01 seconds N., long. 72 degrees 29 minutes 24 seconds W., NAD 27, in a wooded area:

Oa—0 to 1 inch; very dark brown (10YR 2/2) highly decomposed plant material; weak fine granular structure; very friable; many fine and very fine roots; very strongly acid; abrupt smooth boundary.

A—1 to 6 inches; very dark grayish brown (10YR 3/2) gravelly fine sandy loam, light brownish gray (10YR 6/2) dry; weak fine and medium granular structure; very friable; common fine, very fine and medium roots, few coarse roots; 10 percent gravel, 5 percent channers, 2 percent stones; very strongly acid; clear smooth boundary.

Bw1—6 to 15 inches; dark yellowish brown (10YR 4/6) gravelly fine sandy loam; moderate medium subangular blocky structure; friable; few very fine and fine roots, common medium roots; 10 percent gravel, 5 percent channers, 2 percent stones; strongly acid; clear smooth boundary.

Bw2—15 to 29 inches; yellowish brown (10YR 5/6) gravelly fine sandy loam; moderate medium and coarse subangular blocky structure; friable; few fine and very fine roots; 10 percent gravel, 5 percent channers, 2 percent stones; strongly acid; abrupt smooth boundary.

2R—29+ inches; schist bedrock

Range in Characteristics

Solum thickness: 16 to 36 inches

Depth to bedrock: 20 to 40 inches

Reaction: Very strongly acid to moderately acid

In some pedons, the Oe horizon may be absent.

A horizon:

Hue—7.5YR to 2.5Y

Value—2 to 4

Chroma—1 to 4

Content of rock fragments—15 to 34 percent

Bw horizons:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—4 or 6

Texture of the fine earth fraction—fine sandy loam, sandy loam, or loam

Content of rock fragments—5 to 30 percent

BC horizon:

Some pedons have a BC horizon with color and texture similar to the C horizon.

C horizon (where present):

Hue—7.5YR to 5Y

Value—4 or 5

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam or sandy loam

Content of rock fragments—5 to 30 percent

Cheshire Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid

Landform: Hills or till plains

Parent material: Melt-out till derived from sandstone, shale, and basalt

Slope range: 3 to 35 percent

Associated soil in a drainage sequence:

Watchaug (moderately well drained)

Associated similar soils:

Canton (yellower, coarse-loamy over sandy or sandy-skeletal)

Charlton (yellower)

Wethersfield (dense substratum)

Yalesville (moderately deep)

Taxonomic class: Coarse-loamy, mixed, semiactive, mesic Typic Dystrudepts

Typical Pedon

Cheshire fine sandy loam, 3 to 8 percent slopes, located in the town of Wallingford, 50 feet east of Northford Road and 500 feet north of the junction of Northford and Anderson Roads, on the Wallingford USGS topographic quadrangle, lat. 41 degrees 24 minutes 57 seconds N., long. 72 degrees 46 minutes 23 seconds W., NAD 27, in a cultivated field:

Ap—0 to 8 inches; dark brown (7.5YR 3/2) fine sandy loam, pinkish gray (7.5YR 6/2) dry; weak medium granular structure; friable; common fine roots; 5 percent gravel; strongly acid; clear wavy boundary.

Bw1—8 to 16 inches; reddish brown (5YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; few fine roots; 10 percent gravel; strongly acid; gradual wavy boundary

Bw2—16 to 26 inches; reddish brown (5YR 5/4) fine sandy loam, weak medium subangular blocky structure; very friable; few fine roots; 10 percent gravel; strongly acid; clear wavy boundary.

C—26 to 65 inches; reddish brown (2.5YR 4/4) gravelly sandy loam; massive; very friable with firm lenses; 20 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 20 to 38 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—5YR to 10YR

Value—3 or 4 (Dry value is 6 or more) (If Ap horizon present instead of A, value 2 or 3)

Chroma—2 to 4 (If Ap horizon present instead of A, chroma 1 to 3)

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—2.5YR or 5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, silt loam or sandy loam

Content of rock fragments—5 to 34 percent

C horizon:

Hue—10R to 5YR

Value—3 or 4

Chroma—3 to 6

Texture of the fine earth fraction—sandy loam, fine sandy loam

Content of rock fragments—5 to 34 percent

Copake Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderately rapid in the surface layer, moderate or moderately rapid in the subsoil, and rapid or very rapid in the substratum*Landform:* Outwash plains, terraces, and kames*Parent material:* Glaciofluvial deposits derived from schist, limestone, and dolomite*Slope range:* 0 to 15 percent*Associated soils in a drainage sequence:*

Hero (moderately well drained)

Fredon (poorly drained)

Halsey (very poorly drained)

Associated similar soil:

Groton (excessively drained, sandy skeletal)

Associated other soil:

Farmington (shallow, loamy)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, semiactive, mesic Dystric Eutrudepts**Typical Pedon**

Copake fine sandy loam, 0 to 3 percent slopes, located in the town of Salisbury, 1.5 miles north along Route 41 from the New York state line at Amenia Union, NY, 500 feet northeast of Bollen District Cemetery, and 150 feet east of Route 41, on the Ellsworth USGS topographic quadrangle, lat. 41 degrees 50 minutes 31 seconds N., long. 73 degrees 29 minutes 27 seconds W., NAD 27, in a corn field:

Ap—0 to 6 inches, dark brown (10YR 3/3) fine sandy loam, pale brown (10YR 6/3) dry; weak fine granular structure; friable; common very fine and fine roots; 10 percent gravel; neutral; gradual wavy boundary.

AB—6 to 13 inches, dark olive brown (2.5 Y 3/3) gravelly fine sandy loam, pale brown (10 YR 6/3) dry; weak medium subangular blocky structure; friable; common fine and very fine roots; 15 percent gravel; slightly acid; clear smooth boundary.

Bw1—13 to 21 inches, dark yellowish brown (10YR 4/4) gravelly fine sandy loam; weak medium subangular blocky structure; friable; common very fine and fine roots; 25 percent gravel; slightly acid; clear smooth boundary.

Bw2—21 to 31 inches, dark yellowish brown (10YR 4/4) gravelly fine sandy loam; weak medium subangular blocky structure; friable; few very fine and fine roots; 20 percent gravel; neutral; clear wavy boundary.

2C1—31 to 56 inches, dark brown (10 YR 3/3) very gravelly coarse sand; single grain; loose; 40 percent fine gravel; neutral; abrupt smooth boundary

- 2C2—56 to 65 inches, light olive brown (2.5 Y 5/3) fine sand; single grain; loose; 5 percent gravel; slight effervescence; slightly alkaline; abrupt smooth boundary
- 2C3—65 to 75 inches, olive brown (2.5 Y 4/3) gravelly sand; single grain; loose; 15 percent gravel; slight effervescence; slightly alkaline; abrupt smooth boundary.
- 2C4—75 to 80 inches, olive brown (2.5Y 4/3) gravelly sand; single grain; loose; 15 percent gravel; slight effervescence; moderately alkaline.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer, strongly acid to neutral in the subsoil, and slightly acid to moderately alkaline in the substratum

Ap horizon:

Hue—7.5YR to 2.5Y

Value—3 to 5 (Dry value is 6 or more)

Chroma—2 to 4

Content of rock fragments—5 to 14 percent

A horizon (where present)

Hue—7.5 YR to 2.5Y

Value—2 or 3 (Dry value is 6 or more)

Chroma—1 to 3

Content of rock fragments—5 to 25 percent

AB horizon (where present)

Hue—7.5 YR to 2.5 Y

Value—3

Chroma—2 to 4

Content of rock fragments—5 to 25 percent

Bw horizons:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—3 to 8

Texture of the fine earth fraction—silt loam, loam, or fine sandy loam

Content of rock fragments—5 to 34 percent

2C horizons:

Hue—7.5YR to 5Y

Value—3 to 6

Chroma—3 to 6

Texture of the fine earth fraction—loamy fine sand to coarse sand

Content of rock fragments—5 to 70 percent (substratum average more than 20 percent)

Deerfield Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Glaciofluvial deposits derived from granite, gneiss, and/or schist

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

- Windsor (excessively drained)
- Scarboro (very poorly drained)

Associated similar soils:

- Ellington (coarse-silty over sandy or sandy-skeletal, redder in the subsoil and substratum)
- Ninigret (coarse-loamy over sandy or sandy-skeletal)
- Sudbury (sandy)
- Tisbury (coarse-silty over sandy or sandy-skeletal)

Taxonomic class: mixed, mesic Aquic Udipsamments

Typical Pedon

Deerfield loamy fine sand, 0 to 3 percent slopes, located in the town of North Haven, 100 feet west of Manor Drive, 600 feet north of the intersection of Montowese Avenue and Fitch Street, on the Branford USGS topographic quadrangle, lat. 41 degrees 21 minutes 10 seconds N., long. 72 degrees 51 minutes 29 seconds W., NAD 27:

- Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) loamy fine sand, light brownish gray (10 YR 6/2) dry; weak medium granular structure; very friable; common fine roots; very strongly acid; clear smooth boundary.
- Bw1—8 to 16 inches; dark yellowish brown (10YR 4/4) loamy sand; massive; very friable; few fine roots; strongly acid; gradual wavy boundary.
- Bw2—16 to 28 inches; yellowish brown (10YR 5/4) loamy sand; single grain; loose; few fine roots; common fine prominent yellowish red (5YR 5/8) soft masses of iron accumulations and distinct pinkish gray (7.5YR 6/2) iron depletions; strongly acid; gradual wavy boundary.
- C1—28 to 34 inches; brown (7.5YR 4/4) fine sand; single grain; loose; few fine prominent yellowish red (5YR 5/8) soft masses of iron accumulations and distinct pinkish gray (7.5YR 6/2) iron depletions; moderately acid.
- C2—34 to 60 inches; brown (7.5YR 5/4) fine sand; single grain; loose; few fine distinct pinkish gray (7.5YR 6/2) iron depletions; moderately acid.

Range in Characteristics

Solum thickness: 15 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid

Ap horizon:

- Hue—10YR
- Value—2 to 4
- Chroma—1 to 3
- Content of rock fragments—0 to 14 percent

Bw horizons:

- Hue—7.5YR to 2.5Y
- Value—4 to 6
- Chroma—3 to 6
- Texture of the fine earth fraction—loamy fine sand, loamy sand or fine sand
- Content of rock fragments—0 to 14 percent

C1 horizon:

- Hue—7.5YR to 5Y
- Value—4 to 6
- Chroma—1 to 4
- Texture of the fine earth fraction—loamy sand, fine sand or sand
- Content of rock fragments—0 to 14 percent

C2 horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—1 to 4

Texture of the fine earth fraction—loamy sand, fine sand or coarse sand

Content of rock fragments—0 to 14 percent

Dummerston Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderate*Landform:* Hills*Parent material:* Melt-out till derived from phyllite, and/or schist*Slope range:* 3 to 45 percent*Associated soils in a drainage sequence:*

Fullam (moderately well drained)

Brayton (poorly drained)

Associated similar soil:

Lanesboro (dense substratum)

Associated other soils:

Taconic (shallow, somewhat excessively drained)

Macomber (moderately deep, well drained)

Taxonomic class: Coarse-loamy, mixed, active, frigid Typic Dystrudepts**Typical Pedon**

Dummerston gravelly loam, 3 to 15 percent slopes, very stony, located in the town of Salisbury, 6,000 feet south of the Massachusetts/Connecticut state line on Mt. Washington Road, 25 feet northwest following jeep trail, 10 feet on the northwest side of jeep trail, on the Bash Bish USGS topographic quadrangle, lat. 42 degrees 02 minutes 03 seconds N., long. 73 degrees 28 minutes 20 seconds W., NAD 27, in an old dug pit, in a wooded area:

Oe—0 to 1 inch; moderately decomposed plant materials derived from leaf litter; many very fine to medium roots; extremely acid; abrupt smooth boundary.

A—1 to 2 inches; black (7.5YR 2.5/1) gravelly loam, dark gray (7.5 YR 4/1) dry; weak fine granular structure; very friable; many very fine to medium roots; 20 percent gravel; very strongly acid; abrupt broken boundary.

E—2 to 3 inches; brown (7.5YR 4/2) gravelly loam; weak fine granular structure; very friable; many very fine to medium roots; 20 percent gravel; very strongly acid; abrupt broken boundary.

Bs—3 to 4 inches; dark brown (7.5YR 3/4) gravelly loam; weak medium subangular blocky structure; very friable; many very fine to medium roots; 20 percent gravel; very strongly acid; abrupt wavy boundary.

Bw1—4 to 6 inches; yellowish brown (10YR 5/6) gravelly loam; weak medium subangular blocky structure; very friable; common fine to very coarse roots; 20 percent gravel; very strongly acid; abrupt wavy boundary.

Bw2—6 to 11 inches; light olive brown (2.5Y 5/4) gravelly loam; weak medium subangular blocky structure; very friable; common fine to very coarse roots; 20 percent gravel; very strongly acid; abrupt wavy boundary.

Bw3—11 to 22 inches; light olive brown (2.5Y 5/4) gravelly loam; weak medium subangular blocky structure; very friable; common fine to very coarse roots; 25 percent gravel; strongly acid; clear smooth boundary.

BC—22 to 27 inches; light olive brown (2.5Y 5/4) gravelly loam; weak medium subangular blocky structure; very friable; common fine to very coarse roots; 37 percent gravel; strongly acid; clear smooth boundary.

C1—27 to 40 inches; olive brown (2.5Y 4/3) very gravelly loam; massive; firm; few fine to medium roots; 40 percent gravel; strongly acid; clear wavy boundary.

C2—40 to 64 inches; olive brown (2.5Y 4/3) very stony loam; massive; firm; 20 percent gravel, 20 percent stones; strongly acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Content of rock fragments: 15 to 40 percent, but averages less than 35 percent

A horizon:

Hue—10YR or 2.5Y

Value—2 to 4

Chroma—1 to 3

E horizon (if present)

Hue—2.5YR to 10YR

Value—4 to 6

Chroma—0 to 2

Bs horizon (if present):

Hue—7.5YR to 10YR

Value—3 to 5

Chroma—4 to 6

Bw horizons:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam or loam

Content of rock fragments—15 to 40 percent

BC horizon:

Hue—10YR to 5Y

Value—3 to 6

Chroma—2 to 6

Texture of the fine earth fraction—silt loam or loam

Content of rock fragments—15 to 40 percent

C horizons:

Hue—10YR to 5Y

Value—3 to 5

Chroma—2 to 4

Texture of the fine earth fraction—loam or fine sandy loam

Ellington Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Eolian deposits over glaciofluvial deposits derived from sandstone, shale, and/or basalt

Slope range: 0 to 5 percent

Associated soil in a drainage sequence:

Branford (well drained)

Associated similar soils:

Deerfield (sandy)

Ninigret (yellower)

Sudbury (sandy)

Tisbury (coarse-silty over sandy or sandy-skeletal)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, subactive, mesic Aquic Dystrudepts

Typical Pedon

Ellington silt loam, 0 to 5 percent slopes, located in the town of Cheshire, 0.5 miles south on Cheshire Street from the intersection of Cheshire Street and East Johnson Avenue, 75 feet west of Cheshire Street, on the Meriden USGS topographic quadrangle, lat. 41 degrees 32 minutes 32 seconds N., long. 72 degrees 52 minutes 06 seconds W., NAD 27, in a cultivated field (limed):

Ap—0 to 8 inches; dark reddish brown (5YR 3/2) silt loam; pinkish gray (5YR 6/2) dry; weak medium granular structure; friable; few fine roots; 5 percent gravel; slightly acid; clear smooth boundary.

Bw1—8 to 18 inches; reddish brown (5YR 4/4) silt loam; weak medium subangular blocky structure; friable; few fine roots; 5 percent gravel; moderately acid; gradual wavy boundary.

Bw2—18 to 26 inches; reddish brown (5YR 4/4) very fine sandy loam; massive; friable; common medium distinct reddish gray (5YR 5/2) iron depletions and common medium distinct dark red (2.5YR 3/6) soft masses of iron accumulation; 10 percent gravel; strongly acid; abrupt smooth boundary.

2C—26 to 65 inches; dark reddish brown (5YR 3/4) stratified sand and gravel with a few thin lenses of sandy loam; single grain; loose; few fine faint reddish gray (5YR 5/2) iron depletions and few fine distinct yellowish red (5YR 4/6) soft masses of iron accumulation; 50 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 18 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—5YR to 10YR

Value—3 or 4 (If A horizon present instead of Ap, value 2 or 3)

Chroma—2 to 4 (If A horizon present instead of Ap, chroma 1 to 3)

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—2.5YR or 5YR

Value—4 or 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam or fine sandy loam

Content of rock fragments—0 to 30 percent

2C horizon:

Hue—2.5YR or 5YR

Value—3 to 6
 Chroma—3 to 6
 Texture of the fine earth fraction—stratified loamy fine sand to coarse sand
 Content of rock fragments—0 to 50 percent

Elmridge Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderately rapid in the surface layer and subsoil, and very slow in the substratum

Landform: Terraces on lake plains

Parent material: Eolian deposits over glaciolacustrine deposits

Slope range: 0 to 8 percent

Associated soil in a drainage sequence:

Shaker (poorly drained)

Associated similar soils:

Belgrade (coarse-silty)

Berlin (fine-silty)

Brancroft (fine-silty)

Ninigret (coarse-loamy over sandy or sandy skeletal)

Sudbury (sandy)

Taxonomic class: Coarse-loamy over clayey, mixed, semiactive, mesic Aquic Dystric Eutrudepts

Typical Pedon

Elmridge fine sandy loam, 0 to 3 percent slopes, located in the town of Windsor Locks, 300 feet south of Connecticut Route 140 and 500 feet west of South Street, on the Windsor Locks USGS topographic quadrangle, lat. 41 degrees 55 minutes 30 seconds N., long. 72 degrees 38 minutes 07 seconds W., NAD 27:

Ap—0 to 6 inches; very dark grayish brown (10YR 3/2) fine sandy loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; friable; many fine roots; moderately acid; clear wavy boundary.

Bw1—6 to 10 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; many fine roots; moderately acid; clear wavy boundary.

Bw2—10 to 18 inches; brown (10YR 4/3) fine sandy loam; weak medium subangular blocky structure; friable; common fine roots; slightly acid; gradual wavy boundary.

Bw3—18 to 25 inches; yellowish brown (10YR 5/4) sandy loam; weak medium subangular blocky structure; very friable; few fine roots; common medium distinct grayish brown (2.5Y 5/2) iron depletions and common medium prominent strong brown (7.5YR 5/8) soft masses of iron accumulation; slightly acid; clear wavy boundary.

2C—25 to 65 inches; olive brown (2.5Y 4/4) varved silt and clay (silty clay weighted average texture); massive parting to weak thick plates along varved bedding planes; firm, very sticky, plastic; common medium distinct grayish brown (10YR 5/2) iron depletions and yellowish brown (10YR 5/6) soft masses of iron accumulation; slightly acid.

Range in Characteristics

Solum thickness: 18 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral in the surface layer, strongly acid to neutral in

the subsoil, (at least one subhorizon is moderately acid, slightly acid, or neutral), and moderately acid to slightly alkaline in the substratum

Ap horizon:

Hue—7.5YR or 10YR

Value—2 to 4 (Dry value 6 or more) (If A horizon present instead of Ap, value 2 or 3)

Chroma—1 to 3

Content of rock fragments—0 to 5 percent

Upper Bw horizons:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—sandy loam, fine sandy loam, or loam

Content of rock fragments—0 to 5 percent

Lower Bw horizons:

Hue—7.5YR to 5Y

Value—3 to 6

Chroma—2 to 6

Texture of the fine earth fraction—sandy loam, fine sandy loam, or loam

Content of rock fragments—0 to 5 percent

2C horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—2 to 4

Texture of the fine earth fraction—silty clay, silty clay loam, or clay

Content of rock fragments—0 to 2 percent

Enfield Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Eolian deposits over glaciofluvial deposits derived from granite, gneiss, and/or schist

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Tisbury (moderately well drained)

Raypol (poorly drained)

Associated similar soils:

Agawam (coarse-loamy over sandy or sandy-skeletal)

Haven (coarse-loamy over sandy or sandy skeletal)

Taxonomic class: Coarse-silty over sandy or sandy-skeletal, mixed, active, mesic
Typic Dystrudepts

Typical Pedon

Enfield silt loam, in an area of Haven and Enfield silt loams, 3 to 8 percent slopes, located in the town of North Stonington, 2,800 feet southeast along Route 2 from the intersection with Route 201, 1400 feet northwest along an unnamed gravel road, 45 feet south of old cemetery, and 200 feet west of Shunock River, on the Old Mystic

USGS topographic quadrangle, lat. 41 degrees 26 minutes 23 seconds N., long. 71 degrees 54 minutes 07 seconds W., NAD 27, in a wooded area.

Oi—0 to 3 inches slightly decomposed plant material

Oe—3 to 4 inches moderately decomposed plant material

Ap—4 to 12 inches, dark yellowish brown (10YR 3/4) silt loam, pale brown (10YR 6/3) dry; weak medium granular structure; very friable; common fine and medium roots; 2 percent gravel; very strongly acid; abrupt wavy boundary.

Bw1—12 to 20 inches, yellowish brown (10YR 5/6) silt loam; weak medium subangular blocky structure; friable; few fine roots; 1 percent gravel; strongly acid, clear smooth boundary.

Bw2—20 to 26 inches, yellowish brown (10YR 5/6) silt loam; weak medium subangular blocky structure; friable; few fine roots; lenses of very fine sandy loam in the lower part; moderately acid; clear wavy boundary.

Bw3—26 to 30 inches, light olive brown (2.5Y 5/4) silt loam; massive; friable; few fine roots; 2 percent gravel; moderately acid; clear wavy boundary.

2C—30 to 37 inches, light olive brown (2.5Y 5/4) very gravelly loamy sand; single grain; loose; 35 percent gravel, 5 percent cobbles; moderately acid; gradual wavy boundary.

3C—37 to 65 inches, olive gray (5Y 5/2) very gravelly coarse sand; single grain; loose; 50 percent gravel, 7 percent cobbles; moderately acid.

Range in Characteristics

Solum thickness: 16 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—7.5YR or 10YR

Value—3 or 4 (Dry value is 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 through 3)

Content of rock fragments—0 to 5 percent

Bw1 and Bw2 horizons:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—3 to 8

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 5 percent

Bw3 horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 5 percent

2C horizons:

Hue—2.5YR to 2.5Y

Value—4 to 6

Chroma—0 to 6

Texture of the fine earth fraction—stratified loamy sand to coarse sand

Content of rock fragments—0 to 59 percent

Farmington Series

Depth class: Shallow

Drainage class: Well drained

Permeability: Moderate or moderately rapid

Landform: Bedrock-controlled hills and ridges

Parent material: Melt-out till derived from limestone, dolomite, and schist

Slope range: 3 to 45 percent

Associated similar soil:

Hollis (somewhat excessively drained, over schist, granite, or gneiss bedrock)

Associated other soils:

Copake (very deep, coarse-loamy over sandy or sandy-skeletal)

Nellis (very deep)

Stockbridge (very deep)

Amenia (very deep, moderately well drained)

Georgia (very deep, moderately well drained)

Mudgepond (very deep, poorly drained)

Alden (very deep, very poorly drained)

Taxonomic class: Loamy, mixed, active, mesic Lithic Eutrudepts

Typical Pedon

Farmington fine sandy loam, in an area of Farmington-Rock outcrop complex, 3 to 15 percent slopes, located in the town of North Canaan, 0.6 miles south on US Route 7 from the junction of Route 7 and Sand Road, 150 feet east of Route 7, on the Ashley Falls USGS topographic quadrangle, lat. 42 degrees 00 minutes 33 seconds N., long. 73 degrees 19 minutes 47 seconds W., in a wooded area:

A—0 to 3 inches, very dark grayish brown (10YR 3/2) fine sandy loam, light brownish gray (10 YR 6/2) dry; weak fine and medium granular structure; very friable; common very fine, fine, and medium roots, few coarse roots; 7 percent gravel; neutral; clear smooth boundary.

Bw1—3 to 8 inches, dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; common very fine, fine, and medium roots, few coarse roots; 7 percent gravel; neutral; gradual wavy boundary.

Bw2—8 to 17 inches, yellowish brown (10YR 5/4) fine sandy loam; weak medium subangular blocky structure; friable; common fine and medium roots, few coarse roots; 10 percent gravel; neutral; abrupt smooth boundary.

2R—17 inches, siliceous limestone bedrock, weathered in the first inch.

Range in Characteristics

Solum thickness: 10 to 20 inches

Depth to bedrock: 10 to 20 inches

Reaction: Strongly acid to neutral in the surface layer, moderately acid to slightly alkaline in the subsoil

A horizon:

Hue—10YR

Value—3 to 5 (Dry value is 6 or more)

Chroma—1 to 3

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—7.5YR to 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, loam, or silt loam

Content of rock fragments—5 to 34 percent

Fluvaquents

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate to very rapid

Landform: Flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soil in a drainage sequence:

Udifluvents (moderately well drained to somewhat excessively drained)

Associated similar soils:

Occum (well drained)

Pootatuck (moderately well drained)

Rippowam (poorly drained, coarse loamy)

Saco (very poorly drained, thick dark surface layer)

Taxonomic class: Fluvaquents

Sample Pedon

Fluvaquents, in an area of Fluvaquents-Udifluvents complex, frequently flooded, located in the town of Cornwall, 3,000 feet southwest of the intersection of Dawn Hill Road and River Road, and 50 feet west of the Housatonic River, on the USGS Ellsworth topographic quadrangle, lat. 41 degrees 48 minutes 05 seconds N., long. 73 degrees 23 minutes 54 seconds W., NAD 27, in a wooded area:

A—0 to 4 inches; black (2.5Y 2.5/1) silt loam, light brownish gray (2.5Y 6/2) dry; weak fine to medium granular structure; very friable; many fine to coarse roots; slightly alkaline; abrupt wavy boundary.

Cg1—4 to 14 inches; dark grayish brown (2.5Y 4/2) fine sand; single grain; loose; many fine to coarse roots; 10 percent light olive gray (5Y 6/2) lenses of stratified loamy fine sand to sand; common fine to coarse prominent strong brown (7.5YR 5/8) soft masses of iron accumulation and few fine to coarse faint gray (2.5Y 5/1) iron depletions; slightly alkaline; gradual wavy boundary.

Cg2—14 to 21 inches; very dark grayish brown (2.5Y 4/2) very fine sand; single grain; loose; common fine to medium roots; many fine to coarse prominent strong brown (7.5YR 5/8) soft masses of iron accumulation; slightly alkaline; abrupt wavy boundary.

Ab1—21 to 38 inches; very dark gray (2.5Y 3/1) silt loam; massive; very friable; few fine to medium roots; 1 inch thick lense of medium sand; common partially decomposed wood fragments; common fine prominent yellowish red (5YR 4/6) soft masses of iron accumulation; slightly alkaline; clear wavy boundary.

Ab2—38 to 45 inches; very dark gray (10YR 3/1) fine sandy loam; massive; very friable; many charcoal fragments; common fine prominent yellowish red (5YR 4/6) soft masses of iron accumulation; slightly alkaline; clear smooth boundary.

Cgb—45 to 55 inches; very dark gray (10YR 3/1) sand; single grain; loose; slightly alkaline; clear smooth boundary.

A'b—55 to 60 inches; black (2.5Y 2.5/1) fine sandy loam; massive; very friable; neutral.

Range in Characteristics

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly alkaline

Content of rock fragments: 0 to 59 percent

A or Ab horizons:

Hue—5YR to 2.5Y

Value—2 to 4

Chroma—1 to 3

Texture of the fine earth fraction—loamy sand, loamy fine sand, loamy very fine sand, sandy loam, fine sandy loam, very fine sandy loam, loam, or silt loam

C horizons:

Hue—5 YR to 5Y

Value—3 to 6

Chroma—1 or 2

Texture of the fine earth fraction—coarse sand, sand, fine sand, very fine sand, loamy fine sand, loamy very fine sand, sandy loam, fine sandy loam, very fine sandy loam, loam, or silt loam

Fredon Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate in the surface layer and subsoil, and moderately rapid to very rapid in the substratum

Landform: Depressions and drainageways on outwash plains and terraces

Parent material: Glaciofluvial deposits derived from limestone, dolomite, and schist

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Copake (well drained)

Halsey (very poorly drained)

Hero (moderately well drained)

Associated other soil:

Groton (excessively drained, sandy-skeletal)

Taxonomic class: Coarse-loamy over sandy or sandy skeletal, mixed, active, nonacid, mesic Aeric Endoaquepts

Typical Pedon

Fredon silt loam, located in the town of Canaan, 1,600 feet northeast of the intersection of Sand Road and Belden Street, on the South Canaan USGS topographic quadrangle, lat. 41 degrees 58 minutes 43 seconds N., long. 73 degrees 21 minutes 07 seconds W., NAD 27:

Ap—0 to 8 inches; very dark grayish brown (2.5Y 3/2) silt loam, light brownish gray (2.5Y6/2) dry; moderate medium granular structure; friable; few very fine and fine roots; neutral; clear smooth boundary.

Bg—8 to 17 inches; olive gray (5Y5/2) fine sandy loam; weak medium subangular blocky structure; friable; few very fine roots; few fine and medium prominent yellowish brown (10YR 5/6, 5/8) and dark yellowish brown (10YR 4/4, 4/6) soft masses of iron accumulation; neutral; clear smooth boundary.

Bw—17 to 24 inches; olive brown (2.5Y 4/4) fine sandy loam; weak coarse subangular blocky structure; friable; many coarse faint yellowish brown (10YR 4/4) and distinct yellowish brown (10 YR 4/6) soft masses of iron accumulation and many coarse distinct light brownish gray (2.5Y 6/2) iron depletions; neutral; abrupt smooth boundary.

2Cg1—24 to 29 inches; light olive gray (5Y 6/2) loamy sand; single grain; loose; few fine prominent yellowish brown (10YR 5/8) and dark yellowish brown (10YR 4/4) soft masses of iron accumulation; neutral; abrupt smooth boundary.

2C—29 to 48 inches; olive (5Y 5/3) loamy sand; single grain; loose; many coarse faint light olive brown (2.5Y 5/4) and distinct dark yellowish brown (10YR 4/4) and prominent light olive brown (2.5 Y 5/6) soft masses of iron accumulation and many coarse distinct gray (5Y 5/1) iron depletions; 10 percent fine gravel; neutral; clear smooth boundary.

2Cg2—48 to 60 inches; light olive gray (5Y 6/2) sand; single grain; loose; 10 percent fine gravel; neutral.

Range in Characteristics

Solum thickness: 22 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Slightly acid to neutral in the surface layer and subsoil, and moderately acid to moderately alkaline in the substratum

Ap horizon:

Hue—10YR or 2.5Y

Value—2 to 4

Chroma—1 or 2

Content of rock fragments—0 to 14 percent

Bg horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—loam, fine sandy loam, very fine sandy loam or silt loam

Content of rock fragments—0 to 14 percent

Bw horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—loam, fine sandy loam, very fine sandy loam or silt loam

Content of rock fragments—0 to 14 percent

2Cg horizons:

Hue—5YR to 5Y

Value—2 to 4

Chroma—0 to 2

Texture of the fine earth fraction—sand, loamy fine sand, or loamy sand

Content of rock fragments—10 to 30 percent

2C horizons:

Hue—5YR to 5Y

Value—3 to 6

Chroma—3 or 4

Texture of the fine earth fraction—sand, loamy fine sand, or loamy sand

Content of rock fragments—10 to 30 percent

Some of the Fredon soils in this survey area have a mean annual soil temperature which is colder than typical of the series. This map unit (414) is identified as a cold phase of the Fredon series.

Freetown Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate or moderately rapid

Landform: Depressions

Parent Material: Organic materials

Slope range: 0 to 2 percent

Associated similar soils:

Catden (euic reaction class)

Natchaug (16 to 51 inches of organic materials, over loamy)

Timakwa (16 to 51 inches of organic materials, over sandy or sandy-skeletal)

Taxonomic class: Dysic, mesic Typic Haplosaprists

Typical Pedon

Freetown peat, in an area of Catden and Freetown soils, located in the town of Eastford, 500 feet west of the intersection of Pifershire Road and Fayette Wright Road, on the Hampton USGS topographic quadrangle, lat. 41 degrees 50 minutes 37 seconds N., long. 72 degrees 03 minutes 13 seconds W., NAD 27, in Catden Swamp:

Oi1—0 to 4 inches; dark yellowish brown (10YR 4/6) peat; 95 percent fibers, 90 percent rubbed; massive; soft, friable; ultra acid; abrupt smooth boundary.

Oi2—4 to 10 inches; dark yellowish brown (10YR 3/4) peat; 90 percent fibers, 85 percent rubbed; massive; soft, friable; extremely acid; abrupt smooth boundary.

Oa1—10 to 22 inches; very dark brown (7.5YR 2.5/2) muck; 10 percent fibers, 3 percent rubbed; massive; soft, friable; extremely acid; clear smooth boundary.

Oa2—22 to 35 inches; black (10YR 2/1) muck; 5 percent fibers, 0 percent rubbed; massive; soft, friable; extremely acid; clear smooth boundary.

Oa3—35 to 41 inches; black (7.5YR 2.5/1) muck; 5 percent fibers, 0 percent rubbed; massive; soft, friable; extremely acid; clear smooth boundary.

Oa4—41 to 55 inches; black (10YR 2/1) muck; 15 percent fibers, 7 percent rubbed; massive; soft, friable; extremely acid; clear smooth boundary.

Oa5—55 to 71 inches; black (10YR 2/1) muck; 5 percent fibers, 0 percent rubbed; massive; soft, friable; extremely acid; clear smooth boundary.

Oa6—71 to 91 inches; black (7.5YR 2.5/1) muck; 5 percent fibers, 0 percent rubbed; massive; soft, friable; extremely acid.

Range in Characteristics

Solum thickness: Organic layers more than 51 inches thick

Depth to bedrock: More than 80 inches

Reaction: Ultra acid to extremely acid (in 0.01 M calcium chloride)

Woody fragments: 0 to 25 percent

Mineral material: Less than 5 percent

Oi horizons:

Hue—5YR to 10YR

Value—2 to 4

Chroma—0 to 6

Oa horizons:

Hue—5YR to 10YR, or is neutral

Value—2 or 3

Chroma—0 to 4

Fullam Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Hills

Parent material: Lodgement till derived from schist and phyllite

Slope range: 3 to 15 percent

Associated soils in a drainage sequence:

Lanesboro (well drained)

Brayton (poorly drained)

Associated other soils:

Taconic (shallow, somewhat excessively drained)

Dummerston (well drained, friable substratum)

Macomber (moderately deep, well drained)

Taxonomic class: Coarse-loamy, mixed, active, frigid Aquic Dystrudepts

Typical Pedon

Fullam silt loam, 3 to 15 percent slopes, located in the town of Salisbury, 1,800 feet east along Mt. Riga Road from the intersection with Mt. Washington Road, and 600 feet south of Mt. Riga Road, on the Bash Bish Falls USGS topographic quadrangle, lat. 42 degrees 00 minutes 10 seconds and long. 73 degrees 27 minutes 45 seconds, NAD 27, in a wooded area:

Oe—0 to 2 inches; dusky red (2.5YR 3/2); moderately decomposed plant materials derived from leaf litter; many very fine and fine roots.

A—2 to 4 inches; very dark grayish brown (10YR 3/2) silt loam, light brownish gray (10 YR 6/2) dry; weak medium subangular blocky structure parting to weak fine granular structure; very friable; many very fine and medium roots; 8 percent gravel; very strongly acid; abrupt smooth boundary.

Bw1—4 to 10 inches; light olive brown (2.5Y 5/3) silt loam; weak fine and medium subangular blocky structure; friable; common very fine to very coarse roots; 6 percent gravel; very strongly acid; clear smooth boundary.

Bw2—10 to 20 inches; light yellowish brown (2.5Y 6/3) gravelly loam; weak fine to medium subangular blocky structure; friable; few very fine to coarse roots; few coarse faint light olive gray (5Y 6/2) iron depletions and few coarse prominent strong brown (7.5YR 5/8) soft masses of iron accumulation; 15 percent gravel, 2 percent cobbles, 5 percent stones; strongly acid; gradual wavy boundary.

Cd1—20 to 49 inches; light olive brown (2.5Y 5/3) very channery loam; massive; firm; few very fine roots; few fine distinct gray (5Y 5/1) iron depletions and few fine prominent strong brown (7.5YR 5/8) soft masses of iron accumulation; 20 percent gravel, 20 percent channers; strongly acid.

Cd2—49 to 60 inches; olive brown (2.5Y 4/3) very channery fine sandy loam; massive; firm; 20 percent gravel, 20 percent channers; strongly acid.

Range in Characteristics

Solum thickness: 20 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—10YR or 2.5Y

Value—2 to 4

Chroma—2 or 3

Content of rock fragments—0 to 14 percent

Bw1 horizon:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam or loam
Content of rock fragments—5 to 30 percent

Bw2 horizon:

Hue—2.5Y or 5Y
Value—3 to 5
Chroma—2 to 4
Texture of the fine earth fraction—silt loam or loam
Content of rock fragments—15 to 30 percent

Cd horizon:

Hue—2.5Y or 5Y
Value—3 to 5
Chroma—2 to 4
Texture of the fine earth fraction—loam or fine sandy loam
Content of rock fragments—15 to 40 percent

Georgia Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, moderately slow in the substratum

Landform: Hills

Parent material: Till derived from limestone, dolomite, and schist

Slope range: 2 to 15 percent

Associated soils in a drainage sequence:

Stockbridge (well drained)
Mudgepond (poorly drained)
Alden (very poorly drained, fine-loamy)

Associated similar soil:

Amenia (carbonates within 40 inches)

Associated other soils:

Farmington (well drained, shallow, loamy)
Nellis (well drained, carbonates within 40 inches)

Taxonomic class: Coarse-loamy, mixed, semiactive, mesic Aquic Dystric Eutrudepts

Typical Pedon

Georgia silt loam, in an area of Georgia and Ameniam silt loams, 2 to 8 percent slopes, located in the town of Ridgefield, 1600 feet west of the intersection of George Washington Highway and North Ridgebury Road, on the Peach Lake USGS topographic quadrangle, lat. 41 degrees 21 minutes 39 seconds N., long. 73 degrees 31 minutes 48 seconds W., NAD 27:

Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) silt loam, light brownish gray (10 YR 6/2) dry; weak medium granular structure; friable; common fine and medium roots; 5 percent rock fragments; strongly acid; clear smooth boundary.

Bw1—8 to 14 inches; yellowish brown (10YR 5/4) loam; weak medium subangular blocky structure; friable; few fine and medium roots; 5 percent rock fragments; strongly acid; gradual wavy boundary.

Bw2—14 to 24 inches; olive brown (2.5Y 4/4) loam; moderate medium subangular blocky structure; firm; few fine roots; common medium distinct light olive gray (5Y 6/2) iron depletions and prominent strong brown (7.5YR 5/6) soft masses of iron accumulation; 5 percent rock fragments; moderately acid; gradual wavy boundary.

C—24 to 60 inches, dark grayish brown (2.5Y 4/2) gravelly fine sandy loam; moderate thick platy structure; firm; common medium prominent strong brown (7.5YR 5/6) soft masses of iron accumulations and faint olive gray (5Y 5/2) iron depletions; 15 percent rock fragments; slightly acid.

Range in Characteristics

Solum thickness: 16 to 32 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral

Ap horizon:

Hue—10YR or 2.5Y

Value—3 or 4

Chroma—2 or 3

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, silt loam or loam

Content of rock fragments—5 to 30 percent

C horizon:

Hue—10YR to 5Y

Value—3 to 5

Chroma—1 to 4

Texture of the fine earth fraction—fine sandy loam, silt loam, or loam

Content of rock fragments—5 to 30 percent

Gloucester Series

Depth class: Very deep

Drainage class: Somewhat excessively drained

Permeability: Rapid

Landform: Hills

Parent material: Melt-out till derived from granite, gneiss, and schist

Slope range: 3 to 35 percent

Associated similar soils:

Canton (coarse-loamy over sandy or sandy-skeletal, well drained)

Charlton (coarse-loamy, well drained)

Hinckley (outwash)

Merrimac (sandy, well drained, outwash)

Paxton (coarse-loamy, well drained, dense substratum)

Associated other soils:

Sutton (moderately well drained)

Woodbridge (moderately well drained, dense substratum)

Taxonomic class: Sandy-skeletal, mixed, mesic Typic Dystrudepts

Typical Pedon

Gloucester gravelly sandy loam, 8 to 15 percent slopes, located in the town of Sterling, 2,000 feet south along Gibson Hill Road from the intersection with River Road, 1,200 feet west of Gibson Hill Road, on the Oneco USGS topographic quadrangle, lat. 41 degrees 43 minutes 33 seconds N., long. 71 degrees 48 minutes 14 seconds W., NAD 27, in a wooded area:

- A—0 to 4 inches, very dark grayish brown (10YR 3/2) gravelly sandy loam, light brownish gray (10 YR 6/2) dry; weak fine granular structure; very friable; many fine and medium roots; 15 percent rock fragments; strongly acid; abrupt smooth boundary.
- Bw1—4 to 12 inches, dark yellowish brown (10YR 4/6) gravelly sandy loam; weak medium granular structure; very friable; common fine and medium roots; 20 percent rock fragments; strongly acid; clear wavy boundary.
- Bw2—12 to 25 inches, yellowish brown (10YR 5/6) very gravelly loamy sand; weak medium granular structure; very friable; few fine and medium roots; 35 percent rock fragments; strongly acid; clear wavy boundary.
- C1—25 to 35 inches, light olive brown (2.5Y 5/4) very gravelly loamy coarse sand; single grain; loose; few medium roots; 35 percent rock fragments; strongly acid; gradual wavy boundary.
- C2—35 to 60 inches, light brownish gray (2.5Y 6/2) very gravelly loamy coarse sand; single grain; loose; 50 percent rock fragments: strongly acid.

Range in Characteristics

Solum thickness: 20 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to moderately acid in the surface, very strongly acid to moderately acid in the subsoil and substratum

A horizon:

Hue—10YR

Value—2 to 4

Chroma—1 to 3

Content of rock fragments—15 to 30 percent

Bw1 horizon:

Hue—7.5YR or 10YR

Value—4 to 6

Chroma—3 to 8

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—20 to 34 percent

Bw2 horizon:

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture of the fine earth fraction—loamy sand, loamy coarse sand or loamy fine sand

Content of rock fragments—35 to 59 percent

C horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 to 4

Texture of the fine earth fraction—loamy coarse sand, loamy sand, or loamy fine sand

Content of rock fragments—35 to 59 percent

Groton Series

Depth class: Very deep

Drainage class: Excessively drained

Permeability: Moderately rapid in the surface layer and upper subsoil, moderately

rapid or rapid in the middle subsoil, rapid in the lower subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains, terraces, kames, and eskers

Parent material: Glaciofluvial deposits derived from limestone, dolomite, and schist

Slope range: 0 to 45 percent

Associated soils in a drainage sequence:

Hero (moderately well drained)

Fredon (poorly drained)

Halsey (very poorly drained)

Associated similar soil:

Copake (well drained, coarse-loamy over sandy or sandy skeletal)

Taxonomic class: Sandy-skeletal, mixed, mesic Typic Eutrudepts

Typical Pedon

Groton gravelly sandy loam, 3 to 15 percent slopes, located in the town of Canaan, 2,600 feet northwest along Connecticut Route 63 from the intersection with Connecticut Route 126, 800 feet southwest of Route 63, on the South Canaan USGS topographic quadrangle, lat. 41 degrees 51 minutes 00 seconds N., long. 73 degrees 19 minutes 36 seconds W., NAD 27, in a hayfield:

Ap—0 to 8 inches, brown (10YR 4/3) gravelly sandy loam, pale brown (10 YR 6/3) dry; weak fine granular structure; very friable; many very fine and very fine roots; 30 percent gravel; neutral; abrupt smooth boundary.

Bw1—8 to 18 inches, dark yellowish brown (10YR 4/6) very gravelly sandy loam; weak fine and medium granular structure; very friable; common very fine and fine, and few medium roots; 40 percent gravel; neutral; clear wavy boundary.

Bw2—18 to 24 inches, light olive brown (2.5Y 5/4) very gravelly loamy sand; very weak fine granular structure; very friable; few very fine and fine roots; 50 percent gravel; neutral; gradual wavy boundary.

Bw3—24 to 30 inches, olive brown (2.5Y 4/4) very gravelly loamy sand; single grain; loose; few very fine roots; 50 percent gravel; neutral; abrupt wavy boundary.

C1—30 to 52 inches, light olive gray (5Y 6/2) extremely gravelly sand; single grain; loose; very few very fine roots; 65 percent gravel; slightly effervescent; slightly alkaline; abrupt wavy boundary.

C2—52 to 72 inches, pale olive (5Y 6/3) gravelly sand; single grain; loose; 30 percent gravel; slightly effervescent; slightly alkaline.

Range in Characteristics

Solum thickness: 12 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to neutral in the surface layer and upper subsoil, moderately acid to slightly alkaline in the lower subsoil, and neutral to moderately alkaline in the substratum (calcareous within 40 inches)

Ap horizon:

Hue—7.5YR to 2.5Y

Value—3 or 4 (If A is horizon present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A is horizon present instead of Ap, chroma is 1 to 3)

Content of rock fragments—15 to 34 percent

Upper Bw horizons:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam or sandy loam

Content of rock fragments—20 to 59 percent

Lower Bw horizons:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—sandy loam, loamy fine sand, or loamy sand

Content of rock fragments—20 to 59 percent

C horizons:

Hue—7.5YR to 5Y

Value—3 to 6

Chroma—2 to 6

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—25 to 70 percent

Hadley Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderate in the surface layer, moderate or moderately rapid in the upper substratum, and moderate to rapid in the lower substratum*Landform:* Flood plains*Parent material:* Alluvium*Slope range:* 0 to 3 percent*Associated soils in a drainage sequence:*

Winooski (moderately well drained)

Bash (somewhat poorly drained, coarse-loamy)

Lim (poorly drained, coarse-loamy)

Limerick (poorly drained)

Saco (very poorly drained)

Taxonomic class: Coarse-silty, mixed, superactive, nonacid, mesic, Typic Udifluvents**Typical Pedon**

Hadley silt loam, located in the town of Windsor, 4,200 feet east-southeast of the intersection of Connecticut routes 178 and 159, 3,600 feet southeast of the Island Road railroad crossing, on the Hartford North USGS topographic quadrangle, lat. 41 degrees 50 minutes 06 seconds N., long. 72 degrees 38 minutes 20 seconds W., NAD 27:

Ap—0 to 12 inches, very dark grayish brown (10YR 3/2) silt loam; light brownish gray (10YR 6/2) dry; moderate fine and medium granular structure; friable; few very fine, fine and medium roots; slightly acid; clear smooth boundary.

C1—12 to 29 inches, dark brown (10YR 3/3) and very dark grayish brown (10YR 3/2) silt loam; massive; friable; few very fine and fine roots; slightly acid; clear smooth boundary.

C2—29 to 40 inches, dark brown (10YR 3/3) silt loam; massive; friable; slightly acid; abrupt smooth boundary.

C3—40 to 45 inches, very dark grayish brown (2.5Y 3/2) silt loam; massive; friable; slightly acid; clear smooth boundary.

C4—45 to 60 inches, dark brown (10YR 3/3) silt loam; massive; friable; slightly acid.

Range in Characteristics*Solum thickness:* 6 to 14 inches*Depth to bedrock:* More than 80 inches*Reaction:* Very strongly acid to neutral to a depth of 40 inches and moderately acid to slightly alkaline below a depth of 40 inches

Ap horizon:

Hue—10YR to 5Y
 Value—3 or 4 (Dry value 6 or 7)
 Chroma—2 to 4
 Content of rock fragments—less than 1 percent

C1 and C2 horizons:

Hue—10YR to 5Y
 Value—3 to 6
 Chroma—2 to 6
 Texture of the fine earth fraction—stratified silt loam to very fine sand
 Content of rock fragments—less than 1 percent

C3 and C4 horizons:

Hue—10YR to 5Y
 Value—3 to 6
 Chroma—2 to 6
 Texture of the fine earth fraction—stratified silt loam to sand
 Content of rock fragments—less than 1 percent

Halsey Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate in the surface layer and upper subsoil, moderately rapid in the lower subsoil, and rapid or very rapid in the substratum

Landform: Depressions and drainageways on outwash plains and terraces

Parent material: Glaciofluvial deposits derived from limestone, dolomite, and schist

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Groton (excessively drained, sandy-skeletal)
 Copake (well drained)
 Hero (moderately well drained)
 Fredon (poorly drained)

Associated similar soil:

Timakwa (16 to 51 inches of organic materials, over sandy deposits)

Taxonomic class: Coarse-loamy over sandy or sandy skeletal, mixed, active, nonacid, mesic Typic Humaquepts

Typical Pedon

Halsey silt loam, located in the town of Canaan, 2,100 feet northeast of the intersection of Sand Road and Belden Street, on the South Canaan USGS topographic quadrangle, lat. 41 degrees 58 minutes 53 seconds N., long. 73 degrees 21 minutes 09 seconds W., NAD 27:

Oe—0 to 1 inch; black (5YR 2.5/1) moderately decomposed plant materials; moderate medium granular structure; friable; slightly acid; abrupt smooth boundary.

A—1 to 8 inches; very dark brown (10YR 2/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine and medium granular structure; friable; common very fine, fine, and medium roots; slightly acid; clear smooth boundary.

Bg1—8 to 16 inches; dark gray (10YR 4/1) silt loam; weak medium subangular blocky structure; friable; few very fine, fine, and medium roots; neutral; clear smooth boundary.

Bg2—16 to 28 inches; gray (5Y 5/1) and olive gray (5Y 5/2) fine sandy loam; weak coarse subangular blocky structure; friable; slightly plastic and slightly sticky; few

very fine and fine roots in upper part; few fine prominent yellowish brown (10YR 5/6) and dark yellowish brown (10YR 4/4, 4/6) soft masses of iron accumulation; neutral; abrupt smooth boundary.

2Cg1—28 to 38 inches; gray (5Y 5/1) loamy sand; single grain; loose; nonplastic, nonsticky; neutral; clear smooth boundary.

2Cg2—38 to 60 inches; gray (5Y 5/1) sand; single grain; loose; nonplastic, nonsticky; 10 percent fine gravel; neutral.

Range in Characteristics

Solum thickness: 20 to 39 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to neutral in the surface layer and subsoil, and slightly acid to moderately alkaline in the substratum

A horizon:

Hue—10YR or 2.5Y

Value—2 or 3

Chroma—0 to 2

Content of rock fragments—0 to 14 percent

Bg1 horizon:

Hue—10YR to 5Y or 5BG

Value—4 to 6

Chroma—0 to 2

Texture of the fine earth fraction—very fine sandy loam, loam, or silt loam

Content of rock fragments—0 to 20 percent

Bg2 horizon:

Hue—10YR to 5Y or 5BG

Value—4 to 6

Chroma—0 to 2

Texture of the fine earth fraction—fine sandy loam or very fine sandy loam

Content of rock fragments—0 to 20 percent

2Cg horizons

Hue—2.5Y or 5Y

Value—3 to 6

Chroma—0 to 2

Texture of the fine earth fraction—loamy sand or sand

Content of rock fragments—10 to 50 percent

2C horizons (where present, below 30 inch depth)

Hue—2.5 Y or 5 Y

Value—3 to 6

Chroma—3 or 4

Texture of the fine earth fraction—loamy sand or sand

Content of rock fragments—10 to 50 percent

Some of the Halsey soils in this survey area have a mean annual soil temperature which is colder than typical of the series. This map unit (436) is identified as a cold phase of the Halsey series.

Hartford Series

Depth class: Very deep

Drainage class: Somewhat excessively drained

Permeability: Moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Glaciofluvial deposits derived from sandstone and basalt

Slope range: 0 to 8 percent

Associated soil in a drainage sequence:

Manchester (excessively drained)

Associated similar soils:

Branford (coarse-loamy over sandy or sandy-skeletal)

Penwood (loamy fine sand or coarser in the surface layer and subsoil)

Taxonomic class: Sandy, mixed, mesic Typic Dystrudepts

Typical Pedon

Hartford sandy loam, 0 to 3 percent slopes, located in the town of Rocky Hill, 0.95 miles west of the intersection of Brook Street and Connecticut Route 9, and 150 feet south of Brook Street, on the Hartford South USGS topographic quadrangle, lat. 41 degrees 38 minutes 34 seconds N., long. 72 degrees 39 minutes 53 seconds W., NAD 27:

Ap—0 to 8 inches; dark reddish brown (5YR 3/4) sandy loam, reddish brown (5 YR 5/4) dry; weak coarse granular structure; very friable; many fine roots; 5 percent gravel; strongly acid; clear smooth boundary.

Bw1—8 to 20 inches; yellowish red (5YR 4/6) sandy loam; weak fine granular structure; very friable; few fine roots; 5 percent gravel; strongly acid; clear wavy boundary.

Bw2—20 to 26 inches; reddish brown (5YR 4/4) loamy sand; single grain; loose; 10 percent gravel; strongly acid; clear wavy boundary.

2C—26 to 65 inches; reddish brown (5YR 4/4) stratified very gravelly coarse sand to loamy fine sand; single grain; loose; 35 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 18 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—5YR to 10YR

Value—3 or 4 (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—5 to 14 percent

Bw1 horizon:

Hue—5YR or 2.5YR

Value—3 to 5

Chroma—3 to 8

Texture of the fine earth fraction—sandy loam

Content of rock fragments—5 to 30 percent

Bw2 horizon:

Hue—5YR or 2.5YR

Value—3 to 5

Chroma—3 to 8

Texture of the fine earth fraction—loamy sand or sandy loam

Content of rock fragments—5 to 30 percent

2C horizon:

Hue—2.5YR or 5YR

Value—3 to 6

Chroma—3 to 6

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—10 to 50 percent

Haven Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate in the surface layer and subsoil, and very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Eolian deposits over glaciofluvial deposits derived from granite, gneiss, and/or schist

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Tisbury (moderately well drained, coarse-silty over sandy or sandy-skeletal)

Raypol (poorly drained)

Associated similar soils:

Agawam (coarser texture in subsoil)

Enfield (coarse-silty over sandy or sandy-skeletal)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic, Typic Dystrudepts

Typical Pedon

Haven silt loam, in an area of Haven and Enfield silt loams, 0 to 3 percent slopes, located in the town of Fairfield, 300 feet north along Orchard Hill Lane from the intersection with Ross Hill Road, and 100 feet west of Orchard Hill Lane, on the Westport USGS topographic quadrangle, lat. 41 degrees 10 minutes 16 seconds N., long. 73 degrees 15 minutes 52 seconds W., NAD 27, in a lawn:

- Ap—0 to 7 inches, very dark grayish brown (10YR 3/2) silt loam, light brownish gray (10 YR 6/2) dry; weak fine granular structure; very friable; many fine roots; 5 percent rock fragments; very strongly acid; clear smooth boundary.
- Bw1—7 to 14 inches, brown (7.5YR 4/4) silt loam; weak medium subangular blocky structure; friable; common fine roots; 5 percent rock fragments; very strongly acid; gradual wavy boundary.
- Bw2—14 to 20 inches, dark yellowish brown (10YR 4/4) silt loam; weak medium subangular blocky structure; very strongly acid; gradual wavy boundary.
- BC—20 to 24 inches, strong brown (7.5YR 5/6) fine sandy loam; weak medium subangular blocky structure; very friable; few fine roots; 10 percent rock fragments; strongly acid; abrupt smooth boundary.
- 2C—24 to 60 inches, yellowish brown (10YR 5/4) gravelly sand; single grain; loose; 30 percent rock fragments; moderately acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—7.5YR or 10YR

Value—2 to 4 (If Ap horizon is present instead of A, value is 3 or 4)

Chroma—1 to 3 (if Ap horizon is present instead of A, chroma is 2 or 3)

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—5YR to 2.5Y

Value—4 to 6

Chroma—3 to 8

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 14 percent

BC horizon (where present):

Color and rock fragment content—similar to Bw horizons

Texture of the fine earth fraction—fine sandy loam or very fine sandy loam

2C horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—2 to 6

Texture of the fine earth fraction—stratified sand and fine sand

Content of rock fragments—15 to 59 percent

Hero Series*Depth class:* Very deep*Drainage class:* Moderately well drained*Permeability:* Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum*Landform:* Outwash plains and terraces*Parent material:* Glaciofluvial deposits derived from limestone, schist, and dolomite*Slope range:* 0 to 8 percent*Associated soils in a drainage sequence:*

Groton (excessively drained, sandy-skeletal)

Copake (well drained)

Fredon (poorly drained)

Halsey (very poorly drained)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, semiactive, mesic Aquic Eutrudepts**Typical Pedon**

Hero gravelly loam, 0 to 3 percent slopes, located in the town of Sharon, 500 feet southwest of the intersection of Connecticut Route 361 and Indian Mountain Road, on the Sharon USGS topographic quadrangle, lat. 41 degrees 53 minutes 25 seconds N., long. 73 degrees 28 minutes 56 seconds W., in a pasture:

Ap—0 to 9 inches; very dark grayish brown (10YR 3/2) gravelly loam, pale brown (10YR 6/3) dry; weak medium granular structure; very friable; many very fine and fine roots; 15 percent gravel; slightly acid; clear smooth boundary.

Bw1—9 to 18 inches; olive brown (2.5Y 4/4) gravelly silt loam; weak medium subangular blocky structure; friable; few fine and very fine roots; 20 percent gravel; neutral; gradual wavy boundary.

Bw2—18 to 24 inches; olive brown (2.5Y 4/4) gravelly silt loam; weak medium subangular blocky structure; friable; few medium and fine distinct grayish brown (2.5Y 5/2) and distinct olive gray (5Y 5/2) iron depletions and few fine and medium faint brown (7.5YR 4/4) soft masses of iron accumulation; 25 percent gravel; neutral; clear wavy boundary.

Bw3—24 to 27 inches; dark grayish brown (2.5Y 4/2) gravelly sandy loam; massive; very friable; common fine and medium distinct dark yellowish brown (10YR 4/4) soft masses of iron accumulation and common fine and medium faint grayish

brown (2.5Y 5/2) iron depletions; 30 percent gravel; slightly alkaline; slightly effervescent; clear smooth boundary.

2C—27 to 60 inches; grayish brown (2.5Y 5/2) and dark grayish brown (2.5Y 4/2) very gravelly sand; single grain; loose; few thin lenses of gravelly sandy loam; few fine and medium prominent light olive brown (2.5Y 5/6) soft masses of iron accumulation; few strong brown (7.5YR 5/6) and gray (N6/) weathered limestone pebbles; 50 percent gravel, 10 percent cobbles; slightly effervescent; slightly alkaline.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to neutral in the surface layer, moderately acid to slightly alkaline in the subsoil, and neutral to moderately alkaline in the substratum

A horizon:

Hue—10YR or 2.5Y

Value—2 to 4 (dry value 6 or more)

Chroma—1 to 3

Content of rock fragments—15 to 34 percent

Bw1 horizon:

Hue—10YR or 2.5Y

Value—3 to 6

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, loam, or silt loam

Content of rock fragments—5 to 34 percent

Bw2 horizon:

Hue—10YR or 2.5Y

Value—3 to 6

Chroma—2 to 6

Texture of the fine earth fraction—fine sandy loam, loam, or silt loam

Content of rock fragments—5 to 34 percent

Bw3 horizon:

Hue—10YR or 2.5Y

Value—3 to 6

Chroma—2 to 6

Texture of the fine earth fraction—sandy loam, fine sandy loam, loam, or silt loam

Content of rock fragments—5 to 34 percent

2C horizon:

Hue—10YR to 5Y

Value—3 to 6

Chroma—2 to 4

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—15 to 65 percent

Hinckley Series

Depth class: Very deep

Drainage class: Excessively drained

Permeability: Rapid in the surface layer and subsoil, and very rapid in the substratum

Landform: Outwash plains, terraces, kames, and eskers

Parent material: Glaciofluvial deposits derived from granite, gneiss, and schist

Slope range: 0 to 45 percent

Associated soils in a drainage sequence:

Merrimac (somewhat excessively drained, sandy)
 Sudbury (moderately well drained, sandy)
 Walpole (poorly drained, sandy)

Associated similar soils:

Agawam (coarse-loamy over sandy or sandy-skeletal)
 Windsor (sandy)

Taxonomic class: Sandy-skeletal, mixed, mesic Typic Udorthents

Typical Pedon

Hinckley gravelly sandy loam, 0 to 3 percent slopes, located in the town of East Haddam, 600 feet west southwest of the intersection of Connecticut Routes 151 and 149, 200 feet north of Dykas Brook, on the Deep River USGS topographic quadrangle, lat. 41 degrees 29 minutes 39 seconds N., long. 72 degrees 27 minutes 41 seconds W., NAD 27:

- Ap—0 to 8 inches, dark grayish brown (10YR 4/2) gravelly sandy loam, light brownish gray (10 YR 6/2) dry; weak coarse granular structure; friable; many fine and medium roots; 20 percent rock fragments; strongly acid; abrupt smooth boundary.
 Bw1—8 to 20 inches, brown (7.5YR 5/4) very gravelly loamy sand; weak fine granular structure; very friable; common fine and medium roots; 40 percent rock fragments; strongly acid; clear wavy boundary.
 Bw2—20 to 27 inches, yellowish brown (10YR 5/4) very gravelly sand; single grain; loose; few fine roots; 45 percent rock fragments; strongly acid; clear wavy boundary.
 C1—27 to 42 inches, brown (10YR 5/3) very gravelly sand; single grain; loose; 50 percent rock fragments; strongly acid; clear wavy boundary.
 C2—42 to 60 inches, light brownish gray (10YR 6/2) extremely gravelly sand; single grain; loose; 60 percent rock fragments; strongly acid.

Range in Characteristics

Solum thickness: 12 to 30 inches
Depth to bedrock: More than 80 inches
Reaction: Extremely acid to moderately acid
Rock fragments: Overall, more than 35 percent

Ap horizon:

Hue—10YR
 Value—2 to 4 (If A horizon is present instead of Ap, value is 2)
 Chroma—1 to 3
 Content of rock fragments—15 to 34 percent

Bw1 horizon:

Hue—7.5YR or 10YR
 Value—3 to 5
 Chroma—4 to 8
 Texture of the fine earth fraction—loamy sand, loamy coarse sand, or loamy fine sand
 Content of rock fragments—15 to 50 percent

Bw2 horizon:

Hue—7.5YR to 2.5Y
 Value—3 to 6
 Chroma—4 to 8
 Texture of the fine earth fraction—sand
 Content of rock fragments—15 to 50 percent

C horizon:

Hue—7.5YR to 5Y

Value—4 to 7

Chroma—2 to 8

Texture of the fine earth fraction—stratified sand or coarse sand

Content of rock fragments—30 to 70

Hogansburg Series*Depth class:* Very deep*Drainage class:* Moderately well drained*Permeability:* Moderate in the surface layer and subsoil, slow or very slow in the substratum*Landform:* Hills*Parent material:* Till derived from limestone, dolomite, and schist*Slope range:* 2 to 15 percent*Associated soils in a drainage sequence:*

Pyrities (well drained)

Mudgepond, cold (poorly drained)

Loonmeadow (very poorly drained, fine-loamy)

Taxonomic class: Coarse-loamy, mixed, semiactive, frigid Aquic Eutrudepts**Typical Pedon**

Hogansburg loam, located in the town of Canaan, 2,500 feet southwest of the south end of Wangum Lake, on the South Canaan USGS topographic quadrangle, lat. 41 degrees 58 minutes 19 seconds N., long. 73 degrees 16 minutes 25 seconds W., in an old field:

Ap—0 to 12 inches; very dark grayish brown (10YR 3/2) loam, light brownish gray (10YR 6/2 dry); weak very fine and medium subangular blocky structure; very friable; common very fine to coarse roots; 5 percent gravel; neutral; clear wavy boundary.

Bw1—12 to 20 inches; dark yellowish brown (10YR 4/4) loam; weak very fine and medium subangular blocky structure; very friable; common very fine and fine roots; 10 percent gravel; slightly alkaline; clear wavy boundary.

Bw2—20 to 29 inches; olive brown (2.5Y 4/4) loam; weak coarse subangular blocky structure; very friable; common very fine and fine roots; 13 percent gravel; common medium and coarse distinct grayish brown (2.5Y 5/2), faint light olive brown (2.5Y 5/3), and distinct dark grayish brown (2.5Y 4/2) iron depletions; very slightly alkaline; gradual wavy boundary.

Bw3—29 to 43 inches; olive brown (2.5Y 4/3) loam; weak coarse subangular blocky structure; friable; common fine roots; 13 percent gravel; common coarse faint grayish brown (2.5Y 5/2) iron depletions; neutral; very slightly effervescent; clear wavy boundary.

Cd1—43 to 50 inches; olive brown (2.5Y 4/3) loam; massive; firm; 10 percent gravel; common coarse faint grayish brown (2.5Y 5/2) iron depletions and few medium prominent strong brown (7.5YR 4/6) masses of iron accumulation; slightly alkaline; slightly effervescent; gradual wavy boundary.

Cd2—50 to 70 inches; olive brown (2.5Y 4/3) fine sandy loam; massive; firm; 9 percent gravel, 1 percent cobbles; many coarse faint grayish brown (2.5Y 5/2) iron depletions; slightly alkaline; slightly effervescent; gradual wavy boundary.

Cd3—70 to 84 inches; olive brown (2.5Y 4/3) loam; massive; firm; 13 percent gravel; common coarse faint grayish brown (2.5Y 5/2), common coarse faint dark grayish brown (2.5Y 4/2) iron depletions and few fine prominent strong brown (7.5YR 4/6) masses of iron accumulation; slightly alkaline; slightly effervescent.

Range in Characteristics

Solum thickness: 20 to 43 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer, strongly acid to slightly alkaline in the subsoil, and slightly alkaline or moderately alkaline in the substratum (depth to carbonates 10 to 33 inches)

Ap horizon:

Hue—7.5YR or 10YR

Value—3 or 4

Chroma—2 or 3

Content of rock fragments—3 to 14 percent

Bw horizons:

Hue—5YR to 2.5Y

Value—4 or 5

Chroma—3 or 4

Texture of the fine earth fraction—fine sandy loam, silt loam or loam

Content of rock fragments—5 to 34 percent

Cd horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, or loam

Content of rock fragments—5 to 40 percent

Hollis Series

Depth class: Shallow

Drainage class: Somewhat excessively drained

Permeability: Moderate or moderately rapid

Landform: Bedrock-controlled hills and ridges

Parent material: Melt-out till derived from granite, gneiss, and schist

Slope range: 3 to 60 percent

Associated similar soils:

Brimfield (redder subsoil)

Farmington (well drained, over limestone bedrock)

Holyoke (well drained, over basalt or sandstone)

Associated other soils:

Canton (well drained, very deep, coarse-loamy over sandy or sandy-skeletal)

Charlton (well drained, very deep, coarse-loamy)

Chatfield (well drained, moderately deep, coarse-loamy)

Taxonomic class: Loamy, mixed, active, mesic Lithic Dystrudepts

Typical Pedon

Hollis gravelly fine sandy loam, extremely stony, in an area of Hollis-Chatfield Rock outcrop complex, 3 to 15 percent slopes, located in the town of East Hampton, 1,000 feet west of CT Route 196 and 3,200 feet north of CT Route 151, on the Moodus USGS topographic quadrangle, lat. 41 degrees 31 minutes 28 seconds N., long. 72 degrees 29 minutes 48 seconds W., NAD 27, in a wooded area:

Oa—0 to 1 inch; black (10YR 2/1) highly decomposed plant materials; moderate fine granular structure; very friable; many fine and very fine roots; abrupt smooth boundary.

A—1 to 6 inches; very dark grayish brown (10YR 3/2) gravelly fine sandy loam, light brownish gray (10YR 6/2) dry; weak fine granular structure; very friable; common fine, very fine, medium and coarse roots; 10 percent gravel; 5 percent channers; very strongly acid; clear smooth boundary.

Bw1—6 to 9 inches; dark yellowish brown (10YR 4/4) channery fine sandy loam; weak medium subangular blocky structure; friable; few very fine and fine roots, common medium roots; 10 percent gravel, 10 percent channers; strongly acid; clear wavy boundary.

Bw2—9 to 15 inches; yellowish brown (10YR 5/6) gravelly fine sandy loam; weak medium and coarse subangular blocky structure; friable; few fine and very fine roots, common medium roots; 10 percent gravel, 5 percent channers; strongly acid; abrupt smooth boundary.

2R—15+ inches; schist bedrock.

Range in Characteristics

Solum thickness: 10 to 20 inches

Depth to bedrock: 10 to 20 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—7.5YR or 10YR

Value—2 to 4

Chroma—1 to 3

Content of rock fragments—15 to 34 percent

Bw horizons:

Hue—7.5YR to 2.5Y

Value—4 or 5

Chroma—4 to 8

Texture of the fine earth fraction—fine sandy loam

Content of rock fragments—5 to 34 percent

BC or C horizon (where present):

Hue—7.5YR to 5Y

Value—4 or 5

Chroma—4 to 8

Texture of the fine earth fraction—fine sandy loam

Content of rock fragments—5 to 34 percent

Holyoke Series

Depth class: Shallow

Drainage class: Well drained

Permeability: Moderate

Landform: Bedrock-controlled hills and ridges

Parent material: Melt-out till or eolian deposits over basalt, sandstone or shale

Slope range: 3 to 45 percent

Associated similar soils:

Brimfield (somewhat excessively drained, dominantly mica schist rock fragments)

Hollis (somewhat excessively drained, 5YR or yellower in B horizon)

Associated other soils:

Cheshire (very deep)

Wethersfield (very deep, dense substratum)

Yalesville (moderately deep)

Taxonomic class: Loamy, mixed, superactive, mesic Lithic Dystrudepts

Typical Pedon

Holyoke silt loam, in an area of Holyoke-Rock outcrop complex, 3 to 15 percent slopes, located in the town of West Hartford just east of Ely Pond, 200 feet south of Route 44, on the Avon USGS topographic quadrangle, lat. 41 degrees 47 minutes 44 seconds W., long. 72 degrees 47 minutes 55 seconds W., NAD 27, in a wooded area:

- Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials
 A—1 to 3 inches; dark brown (10YR 3/3) silt loam, pale brown (10 YR 6/3) dry; weak medium granular structure; very friable; many fine roots; 10 percent rock fragments; very strongly acid; abrupt wavy boundary.
 Bw1—3 to 8 inches; brown (7.5YR 4/4) silt loam; weak coarse granular structure; very friable; many fine roots; 10 percent rock fragments; very strongly acid; clear wavy boundary.
 Bw2—8 to 18 inches; yellowish red (5YR 4/6) gravelly silt loam; weak medium subangular blocky structure; friable; common fine roots; 15 percent rock fragments; very strongly acid; abrupt wavy boundary.
 2R—18 inches; basalt bedrock.

Range in Characteristics

Solum thickness: 10 to 20 inches

Depth to bedrock: 10 to 20 inches

Reaction: Extremely acid to moderately acid

A horizon:

Hue—5YR to 10YR

Value—3 or 4

Chroma—1 to 3

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—2.5YR to 7.5YR

Value—3 to 6

Chroma—4 to 6

Texture of the fine earth fraction—silt loam, loam, very fine sandy loam, or fine sandy loam

Content of rock fragments—5 to 34 percent

BC or C horizon (where present)—Similar to Bw horizons

Ipswich Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate to rapid

Landform: Tidal marshes and salt marshes

Parent material: Grassy organic materials

Slope range: 0 to 2 percent

Associated similar soils:

Pawcatuck (16 to 51 inches organic materials over sandy deposits)

Westbrook (16 to 51 inches organic materials over loamy deposits)

Taxonomic class: Euic, mesic Typic Sulfihemists

Typical Pedon

Ipswich mucky peat, located in the town of Old Lyme, 1,000 feet west-northwest of the junction of Connecticut Route 156 and Button Ball Road, on the Old Lyme USGS

topographic quadrangle, lat. 41 degrees 17 minutes 19 seconds N., long. 72 degrees 18 minutes 38 seconds W., NAD 27, in a tidal marsh:

- Oe1—0 to 16 inches; very dark grayish brown (10YR 3/2) mucky peat; 85 percent fiber, 35 percent rubbed; massive; friable; many fine and medium roots; 5 percent mineral content; slightly acid; moderately saline; clear wavy boundary.
- Oe2—16 to 23 inches; very dark brown (10YR 2/2) mucky peat; 75 percent fiber, 30 percent rubbed; massive; friable; few fine and medium roots; 5 percent mineral content; neutral; moderately saline; clear wavy boundary.
- Oe3—23 to 64 inches; very dark grayish brown (10YR 3/2) and very dark gray (10YR 3/1) mucky peat (crushed); 70 percent fiber, 25 percent rubbed; massive; friable; 10 percent mineral content; neutral; strongly saline; clear wavy boundary.
- Oa—64 to 80 inches; very dark grayish brown (10YR 3/2) muck; 35 percent fiber, 10 percent rubbed; massive; friable; 15 percent mineral content; neutral; strongly saline.

Range in Characteristics

Thickness of organic materials: More than 51 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to slightly alkaline

Salinity: Moderately saline in the surface tier, moderately saline to strongly saline in the upper subsurface and bottom tiers

Surface tier:

Hue—7.5YR to 5Y

Value—2 to 5

Chroma—0 to 3

Content of rock fragments—none

Subsurface tiers:

Hue—5YR to 5Y

Value—2 to 5

Chroma—0 to 3

Texture of the fine earth fraction—mucky peat

Content of rock fragments—none

Bottom tier:

Hue—5YR to 5Y

Value—2 to 4

Chroma—0 to 3

Texture of the fine earth fraction—muck

Content of rock fragments—none

Lanesboro Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate in the surface layer and subsoil, slow or very slow in the substratum

Landform: Hills

Parent material: Lodgement till derived from phyllite and schist

Slope range: 3 to 45 percent

Associated soils in a drainage sequence:

Fullam (moderately well drained)

Brayton (poorly drained)

Associated similar soil:

Dummerston (well drained, friable substratum)

Associated other soils:

Taconic (shallow, somewhat excessively drained)

Macomber (moderately deep, well drained)

Taxonomic class: Coarse-loamy, mixed, active, frigid Oxyaquic Dystrudepts

Typical Pedon

Lanesboro loam, 15 to 45 percent slopes, located in the town of Salisbury, 3,500 feet east along Mt. Riga Road from the intersection of Mt. Washington Road and Mt. Riga Road, 50 feet southeast of Mt. Riga Road, on the Bash Bish Falls USGS topographic quadrangle, lat. 42 degrees, 00 minutes, 25 seconds N., long. 73 degrees, 27 minutes, 10 seconds W., NAD 27, in a wooded area:

- Oa—0 to 3 inches; highly decomposed plant materials derived from leaf litter; many fine to medium roots; very strongly acid; clear wavy boundary.
- A—3 to 6 inches; black (10YR 2/1) loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; very friable; many fine to medium roots; 5 percent channers, 2 percent cobbles; very strongly acid; abrupt wavy boundary.
- Bw1—6 to 8 inches; dark yellowish brown (10YR 4/6) loam; weak medium subangular blocky structure; friable; many fine to medium roots; 10 percent channers, 2 percent cobbles; very strongly acid; clear wavy boundary.
- Bw2—8 to 16 inches; olive brown (2.5Y 4/4) channery loam; weak medium subangular blocky structure; friable; many fine to very coarse roots; 20 percent channers, 5 percent stones; strongly acid; clear wavy boundary.
- Bw3—16 to 22 inches; light olive brown (2.5Y 5/4) channery loam; weak medium subangular blocky structure; friable; common fine to coarse roots; 25 percent channers, 5 percent stones; strongly acid; gradual wavy boundary.
- BC—22 to 30 inches; olive brown (2.5Y 4/3) channery loam; massive; friable; few fine roots; few medium prominent dark reddish brown (5YR 3/4) soft masses of iron accumulation; 25 percent channers, 5 percent stones; strongly acid; clear smooth boundary.
- Cd—30 to 60 inches; olive brown (2.5Y 4/3) very channery loam; massive; firm; few medium prominent dark reddish brown (5YR 3/4) masses of iron accumulation; 40 percent channers, 5 percent stones; strongly acid

Range in Characteristics

Solum thickness: 20 to 35 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—7.5YR to 2.5Y

Value—2 to 4

Chroma—1 to 3

Content of rock fragments—0 to 14 percent

Bw1 horizon:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—10 to 34 percent

Bw2 horizon:

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—15 to 50 percent

Bw3 horizon:

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—loam, silt loam, or very fine sandy loam

Content of rock fragments—15 to 50 percent

BC horizon:

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—loam, silt loam, or very fine sandy loam

Content of rock fragments—15 to 50 percent

Cd horizon:

Hue—2.5Y or 5Y

Value—4 to 6

Chroma—2 or 3

Texture of the fine earth fraction—loam or very fine sandy loam

Content of rock fragments—15 to 50 percent

Leicester Series*Depth class:* Very deep*Drainage class:* Poorly drained*Permeability:* Moderate or moderately rapid in the surface layer and subsoil, and moderate to rapid in the substratum*Landform:* Depressions and drainageways on hills and drumlins*Parent material:* Melt-out till derived from granite, gneiss, and schist*Slope range:* 0 to 5 percent*Associated soils in a drainage sequence:*

Canton (well drained, coarse-loamy over sandy or sandy-skeletal)

Charlton (well drained)

Sutton (moderately well drained)

Associated similar soils:

Ridgebury Taxadjunct (dense substratum)

Whitman (very poorly drained, dense substratum)

Taxonomic class: Coarse-loamy, mixed, active, acid, mesic Aeric Endoaquepts**Typical Pedon**

Leicester fine sandy loam, in an area of Ridgebury, Leicester, and Whitman soils, extremely stony, located in the town of Prospect, 4,500 feet north of the Prospect-Bethany town line and 300 feet east of Route 69, on the Mount Carmel USGS topographic quadrangle, lat. 41 degrees 28 minutes 49 seconds N., long. 72 degrees 58 minutes 49 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials

A—1 to 7 inches; black (10YR 2/1) fine sandy loam, gray (10 YR 5/1) dry; moderate

medium granular structure; friable; common fine and medium roots; 10 percent gravel and cobbles; strongly acid; clear wavy boundary.

- Bg1—7 to 10 inches; grayish brown (2.5Y 5/2) fine sandy loam; weak medium subangular blocky structure; friable; common fine and medium roots; common medium prominent yellowish red (5YR 5/6) soft masses of iron accumulation; 10 percent gravel and cobbles; strongly acid; gradual wavy boundary.
- Bg2—10 to 18 inches: light brownish gray (2.5Y 6/2) fine sandy loam; weak medium subangular blocky structure; friable; few fine and medium roots; common fine prominent yellowish brown (10YR 5/6) soft masses of iron accumulation; 10 percent gravel and cobbles; strongly acid; gradual wavy boundary.
- BC—18 to 24 inches; pale brown (10YR 6/3) fine sandy loam; massive; friable; few fine roots; many medium distinct yellowish brown (10YR 5/6) and prominent yellowish red (5YR 4/6) soft masses of iron accumulation; 10 percent gravel and cobbles; strongly acid; clear wavy boundary.
- C1—24 to 43 inches; dark yellowish brown (10YR 4/4) gravelly fine sandy loam; massive; friable; many medium distinct yellowish brown (10YR 5/6) soft masses of iron accumulation and distinct pinkish gray (7.5YR 6/2) iron depletions; 15 percent gravel and cobbles; strongly acid; gradual wavy boundary.
- C2—43 to 65 inches; dark yellowish brown (10YR 4/4) gravelly fine sandy loam; massive; friable; few fine distinct yellowish brown (10YR 5/6) soft masses of iron accumulation; 15 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 18 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid or strongly acid in the upper 40 inches and very strongly acid to moderately acid below

A horizon:

Hue—10YR

Value—2 or 3

Chroma—1 or 2 (If Ap horizon is present instead of A, chroma is 1 to 3)

Content of rock fragments—5 to 14 percent

Bg horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 to 4

Texture of the fine earth fraction—fine sandy loam, loam, or sandy loam

Content of rock fragments—5 to 34 percent

BC horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—fine sandy loam, loam, or sandy loam

Content of rock fragments—5 to 34 percent

C1 horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—1 to 4

Texture of the fine earth fraction—fine sandy loam or sandy loam

Content of rock fragments—5 to 34 percent

C2 horizon:

Hue—7.5YR to 5Y

Value—4 to 6
 Chroma—1 to 4
 Texture of the fine earth fraction—fine sandy loam or sandy loam
 Content of rock fragments—5 to 50 percent

Lim Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate in the loamy layers and rapid or very rapid in the underlying sandy materials

Landform: Depressions on flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Hadley (well drained, coarse-silty)

Winooski (moderately well drained, coarse-silty)

Bash (somewhat poorly drained)

Saco (very poorly drained, coarse-silty)

Associated similar soils:

Limerick (coarse-silty)

Rippowam (fine sandy loam or coarser in the substratum)

Taxonomic class: Coarse-loamy, mixed, superactive, nonacid, mesic Fluvaquentic Endoaquepts

Typical Pedon

Lim very fine sandy loam, in an area of Limerick and Lim soils, located in the town of Canaan, 1,425 feet southwest of the junction of Connecticut Route 126 and Sand Road, on the east side of Hollenbeck River, on the South Canaan USGS topographic quadrangle, lat. 41 degrees 58 minutes 25 seconds N., long. 73 degrees 21 minutes 42 seconds W., NAD 27, in a brushy area:

A—0 to 6 inches; very dark grayish brown (2.5Y 3/2) very fine sandy loam; light brownish gray (10YR 6/2) dry; weak medium granular structure; very friable; many very fine and common fine and medium roots; neutral; clear smooth boundary.

Bg1—6 to 11 inches; dark grayish brown (10YR 4/2) very fine sandy loam; massive; friable; common fine and medium roots; common fine prominent yellowish brown (10YR 5/6) and distinct dark yellowish brown (10YR 4/4) soft masses of iron accumulation and prominent light olive gray (5Y 6/2) iron depletions; neutral; clear smooth boundary.

Bg2—11 to 15 inches; dark grayish brown (2.5Y 4/2) very fine sandy loam; massive; friable; few fine, medium and coarse roots; common pieces of slightly decomposed herbaceous plant materials; few fine prominent yellowish brown (10YR 5/6) and distinct dark yellowish brown (10YR 4/4) soft masses of iron accumulation and distinct light olive gray (5Y 6/2) iron depletions; neutral; clear smooth boundary.

Bg3—15 to 22 inches; dark gray (10YR 4/1) silt loam; massive; friable; few fine roots; common pieces of slightly decomposed herbaceous plant materials; common fine prominent strong brown (7.5YR 5/6) and yellowish red (5YR 4/6) soft masses of iron accumulation; neutral; abrupt smooth boundary.

Bg4—22 to 29 inches; gray (5Y 5/1) and olive gray (5Y 5/2) fine sandy loam; massive; friable; common pieces of slightly decomposed herbaceous plant materials; many coarse prominent yellowish brown (10YR 5/6, 5/8) and dark yellowish brown (10YR 4/4) soft masses of iron accumulation; neutral; clear smooth boundary.

- CBg—29 to 42 inches; gray (5Y 5/1 and 5Y 6/1) loamy fine sand; massive; very friable; few pieces of slightly decomposed herbaceous plant materials; few medium prominent dark brown (7.5YR 4/4) soft masses of iron accumulation; neutral; clear smooth boundary.
- Cg1—42 to 50 inches; olive gray (5Y 5/2) sand; single grain; loose; few pieces of slightly decomposed herbaceous and woody plant materials; many fine prominent strong brown (7.5YR 5/6) and yellowish brown (10YR 5/6) soft masses of iron accumulation; neutral; abrupt smooth boundary.
- Cg2—50 to 57 inches; dark gray (10YR 4/1) loamy sand; single grain; loose; few thin black (10YR 2/1) silt loam lenses high in organic matter; few pieces of slightly decomposed herbaceous and woody plant materials; neutral; abrupt smooth boundary.
- Cg3—57 to 65 inches; dark gray (5Y 4/1) sand; single grain; loose; few slightly decomposed pieces of herbaceous and woody plant materials; neutral.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral (Some sub-horizon is moderately acid, slightly acid, or neutral within a depth of 40 inches)

A horizon:

Hue—2.5Y or 10YR

Value—2 to 4

Chroma—1 or 2

Content of rock fragments—0 to 5 percent

Bg horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 or 2

Texture of the fine earth fraction—very fine sandy loam, silt loam, or loam in the upper part, very fine sandy loam or silt loam in the lower part

Content of rock fragments—0 to 5 percent

BC or CB horizons (where present):

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 or 2

Texture of the fine earth fraction—very fine sandy loam, sandy loam, fine sandy loam, loamy fine sand, loamy sand, fine sand, sand, or coarse sand

Content of rock fragments—0 to 14 percent

Cg horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 or 2

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—0 to 50 percent

Limerick Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate

Landform: Depressions on flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Hadley (well drained)

Winooski (moderately well drained)

Bash (somewhat poorly drained, coarse-loamy)

Saco (very poorly drained)

Associated similar soils:

Lim (coarse-loamy)

Rippowam (coarse-loamy)

Taxonomic class: Coarse-silty, mixed, active, nonacid, mesic Fluvaquentic Endoaquepts

Typical Pedon

Limerick silt loam, in an area of Limerick and Lim soils, located in the town of Wethersfield, 1,200 feet east on Second Lane Road from Interstate 91 underpass, 50 feet south of Second Lane Road, on the Hartford South USGS topographic quadrangle, lat. 41 degrees 41 minutes 52 seconds N., long. 72 degrees 38 minutes 24 seconds W., NAD 27, in a hayfield on the floodplain of the Connecticut River:

Ap—0 to 8 inches; dark grayish brown (10YR 4/2) silt loam, light grayish brown (10YR 6/2) dry; moderate medium granular structure; friable; common very fine and fine and few medium roots; moderately acid; clear smooth boundary.

BCg1—8 to 20 inches; olive gray (5Y 4/2) silt loam; massive; friable (moist), slightly sticky (wet); few very fine and fine roots; common medium prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/4, 5/6) soft masses of iron accumulation; moderately acid; clear smooth boundary.

BCg2—20 to 36 inches; olive gray (5Y 4/2) silt loam; massive; slightly sticky; slightly plastic; common medium prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/4, 5/6) soft masses of iron accumulation; moderately acid; clear smooth boundary.

BCg3—36 to 54 inches; dark gray (5Y 4/1) silt loam; massive; slightly sticky; slightly plastic; common medium prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/4, 5/6) soft masses of iron accumulation; moderately acid; clear smooth boundary.

Cg—54 to 65 inches; dark greenish gray (5GY 4/1) silt loam; massive; slightly sticky; slightly plastic; few, fine prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/4, 5/6) soft masses of iron accumulation; neutral.

Range in Characteristics

Solum thickness: 17 to 60 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer, moderately acid to neutral in the subsoil and substratum

A horizon:

Hue—10YR to 5Y

Value—3 or 4

Chroma—1 or 2

Content of rock fragments—none

BCg horizons:

Hue—2.5Y or 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam or very fine sandy loam
Content of rock fragments—none

Cg horizon:

Hue—2.5Y to 5GY

Value—4

Chroma—0 to 2

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—none

Loonmeadow series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderately rapid in the surface layer and subsoil, moderately rapid to moderately slow in the substratum to a depth of 40 inches, and slow to rapid below 40 inches

Landform: Depressions and drainageways on hills

Parent material: Till derived from dolomite, granite, schist, and gneiss

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Bice (well drained)

Shelburne (well drained, dense substratum)

Ashfield (moderately well drained, dense substratum)

Schroon (moderately well drained)

Associated other soils:

Bucksport (organic materials greater than 51 inches deep)

Wonsqueak (16 to 51 inches of organic materials over loamy deposits)

Taxonomic class: Coarse-loamy, mixed, active, nonacid, frigid Mollic Endoaquepts

Typical Pedon

Loonmeadow mucky fine sandy loam, in an area of Brayton-Loonmeadow complex, extremely stony, located in the town of Norfolk, 1,600 feet south along Doolittle Drive from the intersection with North Colebrook Road and 50 feet east of Doolittle Drive, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees, 01 minutes, 18 seconds N., long. 73 degrees, 09 minutes, 33 seconds W. NAD 27; in a wooded area:

Oi—0 to 2 inches; very dusky red (2.5 YR 2.5/2) slightly decomposed plant materials derived from forest leaf litter

A—2 to 9 inches; black (10YR 2/1) mucky fine sandy loam, dark gray (2.5Y 4/1) dry; weak fine to medium granular structure; friable; many fine to coarse roots; 2 percent gravel, 1 percent cobbles, 1 percent stones; slightly acid; clear wavy boundary.

Bg—9 to 18 inches; dark grayish brown (2.5Y 4/2) sandy loam; weak fine to medium subangular blocky structure; friable; common fine to coarse roots; 8 percent gravel, 2 percent cobbles, 1 percent stones; few fine faint gray (2.5Y 6/1) iron depletions and few medium distinct dark yellowish brown (10YR 3/4) soft masses of iron accumulation; neutral; gradual smooth boundary.

Cg1—18 to 35 inches; dark grayish brown (2.5Y 4/2) gravelly sandy loam; massive; friable; few fine faint olive (5Y 4/3) soft masses of iron accumulation; 10 percent gravel, 5 percent cobbles, 5 percent stones; neutral; gradual smooth boundary.

Cg2—35 to 80 inches; gray (10 YR 5/1) gravelly sandy loam; massive; friable; few fine prominent olive (5Y 4/3) soft masses of iron accumulation; 20 percent gravel, 5 percent cobbles, 5 percent stones; neutral.

Range in Characteristics

Solum thickness: 14 to 25 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer and subsoil, neutral to moderately alkaline in the substratum

Depth to carbonates: a trace may be present above a depth of 40 inches

O horizon:

Hue—2.5YR to 10YR

Value—2 or 2.5

Chroma—2 or 3

A or Ap horizon:

Hue—7.5YR or 10YR

Value—2 or 3 (5 or less dry)

Chroma—1

Content of rock fragments—0 to 14 percent

Bg horizon:

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—5 to 30 percent

Cg or Cd_g horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—sandy loam or fine sandy loam (or fine sandy loam to coarse sand below a depth of 40 inches)

Content of rock fragments—5 to 30 percent

Ludlow Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Hills and drumlins

Parent material: Lodgement till derived from sandstone, shale and basalt

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Wethersfield (well drained)

Wilbraham (poorly drained)

Menlo (very poorly drained)

Associated similar soils:

Rainbow (friable substratum)

Watchaug (friable substratum)

Taxonomic class: Coarse-loamy, mixed, semiactive, mesic Aquic Dystrudepts

Typical Pedon

Ludlow silt loam, 3 to 8 percent slopes, located in the town of Middletown, 400 feet southwest of the intersection of Bush Hill Road and Laurel Grove Road, on the Middletown USGS topographic quadrangle, lat. 41 degrees 30 minutes 54 seconds N., long. 72 degrees 40 minutes 42 seconds W., NAD 27, in a cultivated field:

- Ap—0 to 8 inches; dark brown (7.5YR 3/2) silt loam, pinkish gray (7.5YR 6/2) dry; weak coarse granular structure; friable; many fine roots; 8 percent gravel; strongly acid; clear wavy boundary.
- Bw1—8 to 20 inches; reddish brown (5YR 4/4) silt loam; weak coarse subangular blocky structure; friable; few fine roots; 10 percent gravel; strongly acid; gradual wavy boundary.
- Bw2—20 to 26 inches; dark reddish brown (5YR 3/4) silt loam; weak coarse subangular blocky structure; friable; few fine roots; common medium distinct pinkish gray (5YR 6/2) iron depletions and prominent strong brown (7.5YR 5/8) soft masses of iron accumulation; 12 percent gravel; strongly acid; clear wavy boundary.
- Cd—26 to 65 inches; dark reddish brown (2.5YR 3/4) gravelly loam; weak thick platy structure; very firm; brittle; thin patchy silt films and black (10YR 2/1) manganese coatings on some plates; few fine distinct reddish gray (5YR 5/2) iron depletions; 20 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—5YR to 10R

Value—3 or 4 (Dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—2.5YR or 5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, fine sandy loam or loam

Content of rock fragments—5 to 25 percent

Cd horizon:

Hue—2.5YR or 5YR

Value—3 to 5

Chroma—2 to 6

Texture of the fine earth fraction—loam, silt loam, or fine sandy loam

Content of rock fragments—5 to 34 percent

Macomber Series

Depth class: Moderately deep

Drainage class: Well drained

Permeability: Moderate

Landform: Bedrock-controlled hills and ridges

Parent material: Melt-out till derived from phyllite and schist

Slope range: 3 to 25 percent

Associated similar soil:

Taconic (shallow, somewhat excessively drained)

Associated other soils:

Dummerston (very deep, well drained)

Fullam (very deep, moderately well drained)

Brayton (very deep, poorly drained)

Taxonomic class: Loamy-skeletal, mixed, active, frigid Typic Dystrudepts

Typical Pedon

Macomber very channery loam, in an area of Macomber Taconic complex, 3 to 15 percent slopes, very rocky, located in the town of Salisbury, 2,400 feet southwest of the South Pond dam and 2,200 feet west of an old cemetery, on the Bash Bish USGS topographic quadrangle, lat. 42 degrees 00 minutes 11 seconds N., long. 73 degrees 28 minutes 40 seconds W., NAD 27, in a wooded area:

- Oa—0 to 1 inch, highly decomposed plant materials derived from leaf litter
 A—1 to 2 inches, dark brown (10YR 3/3) very channery loam, brown (10 YR 5/3) dry; weak fine granular structure; friable; many fine to medium roots; 30 percent channers, 5 percent gravel; very strongly acid; abrupt wavy boundary.
 Bw1—2 to 10 inches, yellowish brown (10YR 5/8) very channery loam; weak very fine to medium subangular blocky structure; friable; many fine to coarse roots; 30 percent channers, 5 percent gravel; strongly acid; clear smooth boundary.
 Bw2—10 to 21 inches, light olive brown (2.5Y 5/4) very channery loam; weak medium subangular blocky structure; friable; common fine to coarse roots; 30 percent channers, 2 percent gravel; 3 percent cobbles; strongly acid; gradual smooth boundary.
 C—21 to 30 inches, olive (5Y 5/3) very channery loam; massive; friable to firm; few fine roots; 30 percent channers, 10 percent gravel; strongly acid; very abrupt irregular boundary.
 2R—30 inches, phyllite bedrock

Range in Characteristics

Solum thickness: 15 to 30 inches
Depth to bedrock: 20 to 40 inches
Reaction: Very strongly acid or strongly acid

A horizon:

Hue—10YR or 2.5Y
 Value—2 to 4
 Chroma—1 to 4
 Content of rock fragments—10 to 34 percent

Bw horizons:

Hue—7.5YR to 2.5Y
 Value—3 to 5
 Chroma—3 to 8
 Texture of the fine earth fraction—silt loam or loam
 Content of rock fragments—30 to 59 percent

C horizon (where present)

Hue—10YR to 5Y
 Value—4 or 5
 Chroma—2 to 6
 Texture of the fine earth fraction—silt loam or loam
 Content of rock fragments—40 to 65 percent

Manchester Series

Depth class: Very deep
Drainage class: Excessively drained
Permeability: Rapid in the surface layer, rapid or very rapid in the subsoil and substratum

Landform: Outwash plains, terraces, kames, and eskers

Parent material: Glaciofluvial deposits derived from sandstone, shale, and basalt

Slope range: 0 to 45 percent

Associated soil in a drainage sequence:

Hartford (somewhat excessively drained, sandy)

Associated similar soils:

Penwood (sandy)

Windsor (sandy)

Taxonomic class: Sandy-skeletal, mixed, mesic Typic Udorthents

Typical Pedon

Manchester gravelly sandy loam, 0 to 3 percent slopes, located in the town of Portland, 2,300 feet west-southwest of the intersection of Isinglass Hill Road and Connecticut Route 17, and 2,200 feet south of the Middlesex-Hartford County line, on the Glastonbury USGS topographic quadrangle, lat. 41 degrees 37 minutes 44 seconds N., long. 72 degrees 36 minutes 48 seconds W., NAD 27, in a cultivated field:

Ap—0 to 9 inches; dark brown (7.5YR 3/2) gravelly sandy loam, pinkish gray (7.5 YR 6/2) dry; weak medium granular structure; very friable; many fine and common medium roots; 20 percent gravel; strongly acid; clear smooth boundary.

Bw—9 to 18 inches; reddish brown (5YR 4/3) gravelly loamy sand; very weak fine and medium granular structure; very friable; few fine roots; 25 percent gravel; strongly acid; clear wavy boundary.

C—18 to 65 inches; reddish brown (5YR 4/4) stratified extremely gravelly coarse sand to very gravelly loamy sand; single grain; loose; 50 percent gravel; very strongly acid.

Range in Characteristics

Solum thickness: 12 to 24 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—5YR to 10YR

Value—3 or 4 (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—15 to 34 percent

Bw horizon:

Hue—5YR or 2.5YR

Value—4 or 5

Chroma—3 to 8

Texture of the fine earth fraction—loamy sand or sand

Content of rock fragments—15 to 50 percent

C horizon:

Hue—5YR or 2.5YR

Value—3 to 6

Chroma—3 to 6

Texture of the fine earth fraction—stratified loamy sand to coarse sand

Content of rock fragments—35 to 70 percent

Maybid Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate in the surface layer, and slow or very slow in the subsoil and substratum

Landform: Depressions and drainageways on lake plains and terraces

Parent material: Glaciolacustrine deposits

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Brancroft (moderately well drained, fine silty)

Scitico (poorly drained)

Associated other soils:

Raynham (coarse-silty)

Shaker (coarse-loamy over clayey)

Taxonomic class: Fine, mixed, semiactive, nonacid, mesic Typic Humaquepts

Typical Pedon

Maybid silt loam, in an area of Scitico, Shaker, and Maybid soils, located in the town of East Windsor, 2,100 feet west of the intersection of Newberry and Winkler Roads, 250 feet north of Newberry Road, on the Broad Brook USGS topographic quadrangle, lat. 41 degrees 55 minutes 20 seconds N., long. 72 degrees 35 minutes 33 seconds W., NAD 27, in a brushy area:

A—0 to 9 inches; very dark grayish brown (10YR 3/2) silt loam, dark gray (10YR 4/1) dry; moderate medium granular structure; friable; slightly sticky and slightly plastic; many very fine and fine and common medium roots; strongly acid; clear smooth boundary.

Bg1—9 to 18 inches; gray (5Y 5/1) silty clay loam; weak very coarse prismatic structure parting to moderate coarse subangular blocky structure; prism surfaces are a continuous gray (5Y 5/1); firm; sticky and plastic; few very fine and fine roots along prism faces; common coarse prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/6 and 10YR 5/8) soft masses of iron accumulation in the interior of peds; moderately acid; clear smooth boundary.

Bg2—18 to 26 inches; gray (5Y 5/1) silty clay loam; weak very coarse prismatic structure parting to moderate coarse subangular blocky; prism surfaces are a continuous gray (5Y 5/1) and dark gray (5Y 4/1); firm; sticky and plastic; common coarse prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/6 and 10YR 5/8) soft masses of iron accumulation in the interiors of peds; slightly acid; clear smooth boundary.

Cg1—26 to 36 inches; gray (5Y 5/1) and dark gray (5Y 4/1) varved silt loam, silty clay loam, and clay (silty clay loam weighted average texture); common light olive brown (2.5Y 5/4, 5/6) and dark gray (5Y 4/1) variegations; weak thin platy structure; firm; sticky and plastic; neutral.

Cg2—36 to 60 inches; grayish brown (2.5Y 5/2) and olive gray (5Y 5/2) varved silt loam, silty clay loam, and clay (silty clay loam weighted average texture); common light olive brown (2.5Y 5/4, 5/6) and dark gray (5Y 4/1) variegations; weak thin platy structure; firm; sticky and plastic; neutral.

Range in Characteristics

Solum thickness: 18 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to moderately acid in the surface layer and strongly acid to neutral in the subsoil and substratum

A horizon:

Hue—10YR to 5Y

Value—2 or 3 (dry value 4 or 5)

Chroma—0 to 2
Content of rock fragments—none

Bg1 horizon:

Hue—5Y
Value—4 or 5
Chroma—1 or 2
Texture of the fine earth fraction—silty clay loam, or silty clay
Content of rock fragments—none

Bg2 horizon:

Hue—5Y
Value—4 to 6
Chroma—0 to 2
Texture of the fine earth fraction—silty clay loam, silty clay or clay
Content of rock fragments—none

Cg horizons:

Hue—2.5Y, 5Y, or 5GY
Value—4 or 5
Chroma—0 or 1
Texture of the fine earth fraction—silty clay loam, silty clay, or clay.
Content of rock fragments—none

Medomak Series Taxadjunct

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate in the surface layer and upper substratum and rapid or very rapid in the lower substratum

Landform: Depressions and drainageways on flood plains

Parent material: Alluvium

Slope range: 0 to 2 percent

Associated other soils:

Ondawa (well drained, coarse-loamy)

Rumney (poorly drained, coarse-loamy)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, nonacid, superactive, frigid Fluvaquentic Endoaquepts

Typical Pedon

Medomak mucky silt loam, located in the town of Canaan, 2,000 feet south along Wangum Lake Brook from the outlet of Wangum Lake and 30 feet east of the brook, on the South Canaan USGS topographic quadrangle, lat. 41 degrees 58 minutes 08 seconds N., long. 73 degrees 16 minutes 09 seconds W., NAD 27, in a brushy area:

A—0 to 7 inches; very dark grayish brown (10YR 3/2) mucky silt loam, light brownish gray (10YR 6/2) dry; moderate medium granular structure; very friable; many very fine and fine roots; 2 percent gravel; slightly acid; gradual wavy boundary.

Bg—7 to 23 inches; dark gray (2.5Y 4/1) silt loam; massive; very friable; few very fine and fine roots; common very coarse prominent (5YR 5/8) masses of iron accumulation; 2 percent gravel; neutral; clear wavy boundary.

Ab—23 to 33 inches; very dark brown (10YR 2/2) mucky silt loam; massive; very friable; few medium distinct dark gray (2.5Y 4/1) iron depletions; 2 percent gravel; neutral; abrupt smooth boundary.

2Cg—33 to 46 inches; grayish brown (2.5Y 5/2) very gravelly coarse sand; single grain; loose; common very coarse prominent strong brown (7.5YR 4/6) masses of iron accumulation; 40 percent gravel; slightly alkaline; abrupt smooth boundary.

3C—46 to 78 inches; light olive brown (2.5Y 5/3) very fine sandy loam; massive; firm; common medium faint grayish brown (2.5Y 5/2) iron depletions; 5 percent gravel; slightly alkaline.

Range in Characteristics

Solum thickness: 6 to 15 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral to a depth of about 30 inches, moderately acid to slightly alkaline below

O horizon (where present):

Hue—5YR to 10YR

Value—2 or 3

Chroma—1 or 2

A horizon:

Hue—7.5YR to 2.5Y

Value—2 or 3

Chroma—0 to 2

Content of rock fragments—0 to 5 percent

Cg horizons:

Hue—10YR to 5GY

Value—3 to 6

Chroma—0 to 2

Texture of the fine earth fraction between 6 and 20 inches—silt loam, very fine sandy loam, or loamy very fine sand

Texture of the fine earth fraction between 20 and 40 inches—loamy very fine sand to very gravelly coarse sand

Texture below 40 inches—very fine sandy loam to fine gravel

Content of rock fragments—0 to 5 percent above 40 inches, 0 to 50 percent below 40 inches

The Medomak soils in this survey area are taxadjuncts because the soil is coarser textured than is defined as the range for the series. This difference, however, does not significantly affect the use, management, or interpretations of the soils. In this survey, the Medomak soils are coarse-loamy, mixed, superactive, frigid Fluvaquentic Humaquepts.

Menlo Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderately rapid in the organic surface layer, moderate in the mineral surface layer and subsoil, and slow or very slow in the substratum

Landform: Depressions and drainageways on hills and drumlins

Parent material: Lodgement till derived from sandstone, shale, and basalt

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Wethersfield (well drained)

Ludlow (moderately well drained)

Wilbraham (poorly drained)

Taxonomic class: Coarse-loamy, mixed, active, mesic Typic Endoaquolls

Typical Pedon

Menlo mucky silt loam, in an area of Wilbraham and Menlo soils, extremely stony, located in the town of Middletown, 700 feet northwest of the intersection of Long Hill Road and Pine Street, on the Middletown USGS topographic quadrangle, lat. 41 degrees 32 minutes 21 seconds N., long. 72 degrees 39 minutes 51 seconds W., NAD 27, in a wooded swamp:

- Oa—0 to 5 inches; black (10YR 2/1) highly decomposed plant materials; 5 percent fiber; massive; very friable; many very fine and fine roots, few medium to very coarse roots; moderately acid; clear wavy boundary.
- A—5 to 16 inches; black (10YR 2/1) mucky silt loam, gray (7.5 YR 5/1) dry; weak coarse subangular blocky structure; friable; common very fine and fine roots; common fine prominent strong brown (7.5 YR 4/6) soft masses of iron accumulation; 2 percent quartz and basalt gravel; neutral; clear wavy boundary.
- Bg1—16 to 22 inches; gray (7.5YR 5/1) flaggy very fine sandy loam; weak coarse subangular blocky structure; friable; common medium prominent strong brown (7.5 YR 5/6) and yellowish brown (10YR 5/6) soft masses of iron accumulation; 15 percent red sandstone flagstones, 10 percent quartz and basalt gravel, 5 percent red sandstone channers; neutral; clear smooth boundary.
- Bg2—22 to 27 inches; grayish brown (10YR 5/2) flaggy fine sandy loam; weak coarse subangular blocky structure; friable; many fine to coarse prominent strong brown (7.5 YR 5/6) soft masses of iron accumulation and common medium faint gray (10YR 5/1) iron depletions; 10 percent red sandstone flagstones, 10 percent red sandstone channers, 5 percent quartz and basalt gravel; neutral; abrupt smooth boundary.
- Cd1—27 to 40 inches; reddish brown (5YR 4/3) fine sandy loam; massive; firm; 1 inch lenses of brown (7.5YR 4/3) loamy sand on top of the horizon; common medium prominent strong brown (7.5YR 4/6) soft masses of iron accumulation, common medium faint reddish gray (5YR 5/2) and few medium distinct pinkish gray (7.5YR 6/2) iron depletions; 8 percent quartz and basalt gravel, 2 percent red sandstone channers and 2 percent red sandstone flagstones; neutral; gradual smooth boundary.
- Cd2—40 to 60 inches; reddish brown (5YR 4/3) fine sandy loam; massive; firm; common fine distinct yellowish red (5YR 4/6) soft masses of iron accumulation; 8 percent quartz and red sandstone gravel, 2 percent red sandstone channers and 2 percent red sandstone flagstones; neutral.

Range in Characteristics

Solum thickness: 20 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid in the organic layers, very strongly acid to neutral in the mineral surface layers, and strongly acid to slightly alkaline in the subsoil and substratum (ranges to moderately alkaline below 40 inches)

O horizon:

Hue—5YR to 10YR

Value—2 to 3

Chroma—0 to 2

Content of rock fragments—0 to 25 percent

A horizon:

Hue—5YR to 10YR

Value—2 to 3

Chroma—1 or 2

Content of rock fragments—2 to 14 percent

Bg horizons:

Hue—2.5YR to 10YR

Value—3 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam, loam, very fine sandy loam or fine sandy loam

Content of rock fragments—2 to 25 percent

Cd horizons:

Hue—10R to 5YR

Value—3 to 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam, loam, very fine sandy loam or fine sandy loam

Content of rock fragments—5 to 34 percent

Merrimac Series*Depth class:* Very deep*Drainage class:* Somewhat excessively drained*Permeability:* Moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum*Landform:* Outwash plains, terraces, and kames*Parent material:* Glaciofluvial deposits derived from granite, gneiss, and schist*Slope range:* 0 to 15 percent*Associated soils in a drainage sequence:*

Hinckley (excessively drained, sandy skeletal)

Sudbury (moderately well drained)

Walpole (poorly drained)

Associated similar soils:

Agawam (coarse-loamy over sandy or sandy-skeletal)

Enfield (coarse-silty over sandy or sandy-skeletal)

Taxonomic class: Sandy, mixed, mesic Typic Dystrudepts**Typical Pedon**

Merrimac sandy loam, 3 to 8 percent slopes, located in the town of Newtown, 0.5 miles north of the intersection of Glen Road and Walnut Tree Hill Road and 0.1 miles east of Walnut Tree Road, on the Newtown USGS topographic quadrangle, lat. 41 degrees 26 minutes 37 seconds N., long. 73 degrees 16 minutes 20 seconds W., NAD 27:

Ap—0 to 9 inches; very dark grayish brown (10YR 3/2) sandy loam, light brownish gray (10 YR 6/2) dry; weak medium granular structure; very friable; common fine roots; 5 percent rock fragments; slightly acid; clear wavy boundary.

Bw1—9 to 16 inches; brown (7.5YR 4/4) sandy loam; weak medium granular structure; very friable; few fine roots; 5 percent rock fragments; moderately acid; gradual wavy boundary.

Bw2—16 to 24 inches; brown (7.5YR 4/4) gravelly sandy loam; weak medium granular structure; very friable; few fine roots; 15 percent rock fragments; moderately acid; clear wavy boundary.

2C—24 to 60 inches; yellowish brown (10YR 5/4) stratified very gravelly coarse sand to very gravelly sand; single grain; loose; 40 percent rock fragments; moderately acid.

Range in Characteristics

Solum thickness: 18 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to slightly acid in the surface layer, moderately acid in the subsoil and substratum.

Ap horizon:

Hue—7.5YR or 10YR

Value—3 or 4 (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 or 2)

Content of rock fragments—5 to 14 percent

Bw1 horizon:

Hue—7.5YR or 10YR

Value—3 to 6

Chroma—3 to 8

Texture of the fine earth fraction—sandy loam or fine sandy loam,

Content of rock fragments—5 to 14 percent

Bw2 horizon:

Hue—7.5YR to 2.5Y

Value—3 to 6

Chroma—3 to 8

Texture of the fine earth fraction—sandy loam

Content of rock fragments—5 to 30 percent

2C horizon:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 to 6

Texture—stratified gravelly sand to very gravelly coarse sand

Content of rock fragments—25 to 45 percent gravel and 5 to 10 percent cobbles

Some of the Merrimac soils in this survey area have a mean annual soil temperature which is colder than typical of the series. These map units (434A, 434B, and 434C) are identified as cold phases of the Merrimac series.

Millsite Series

Depth class: Moderately deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid

Landform: Bedrock-controlled hills and ridges

Parent material: Melt-out till derived from granite, gneiss, and schist

Slope range: 3 to 45 percent

Associated other soils:

Westminster (shallow, somewhat excessively drained)

Bice (very deep)

Schroon (very deep, moderately well drained)

Loonmeadow (very deep, very poorly drained)

Taxonomic class: Coarse-loamy, mixed, active, frigid Typic Dystrudepts

Typical Pedon

Millsite fine sandy loam, in an area of Bice-Millsite complex, 3 to 15 percent slopes, very rocky, located in the town of Norfolk, on Dutton Mountain, on the Norfolk USGS

topographic quadrangle, lat. 41 degrees 58 minutes 20 seconds N., long. 73 degrees 11 minutes 32 seconds W., NAD 27, in a wooded area:

- Oe—0 to 1 inch moderately decomposed plant materials derived from leaf litter
 A—1 to 5 inches, dark brown (7.5YR 3/2) fine sandy loam, brown (7.5 YR 4/2) dry; weak very fine and fine granular structure; friable; many fine and medium roots; very strongly acid; clear smooth boundary.
 Bw1—5 to 13 inches, dark yellowish brown (10YR 3/4) stony fine sandy loam; weak medium subangular blocky structure; friable; few fine and common medium roots; 5 percent gravel, 5 percent cobbles, 10 percent stones; very strongly acid; gradual irregular boundary.
 Bw2—13 to 24 inches, dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; few very fine, fine and medium roots; 5 percent gravel; strongly acid; clear wavy boundary.
 C—24 to 31 inches, yellowish brown (10YR 5/4) sandy loam; massive; firm; 5 percent gravel; strongly acid; abrupt irregular boundary.
 2R—31 inches, gneiss bedrock

Range in Characteristics

Solum thickness: 16 to 36 inches

Depth to bedrock: 20 to 40 inches

Reaction: Very strongly acid to slightly acid

Ap or A horizon:

Hue—7.5YR to 2.5Y

Value—3 or 4

Chroma—2 to 4

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—4 to 6

Texture of the fine earth fraction—loam, fine sandy loam, or sandy loam

Content of rock fragments—5 to 34 percent

C horizon:

Hue—7.5YR to 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture of the fine earth fraction—loam, fine sandy loam, or sandy loam

Content of rock fragments—5 to 34 percent

Montauk Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Hills and drumlins

Parent material: Lodgement till derived from granite, gneiss, and schist

Slope range: 3 to 35 percent

Associated soils in a drainage sequence:

Woodbridge (moderately well drained)

Ridgebury taxadjunct (poorly drained)

Whitman (very poorly drained)

Associated similar soils:

- Broadbrook (finer textured substratum)
- Canton (coarse-loamy over sandy or sandy-skeletal, friable substratum)
- Charlton (friable substratum)
- Paxton (finer textured substratum)

Taxonomic class: Coarse-loamy, mixed, subactive, mesic Oxisol Dystrudepts

Typical Pedon

Montauk fine sandy loam, in an area of Paxton and Montauk fine sandy loams, 15 to 35 percent slopes, extremely stony, located in the town of Killingworth, Chatfield Hollow State Park, 400 feet west along Abner Lane from the crossing with Chatfield Hollow Brook, 400 feet south of Abner Lane and 600 feet west of Old Mill Pond, on the Haddam USGS topographic quadrangle, lat. 41 degrees 22 minutes 40 seconds N., long. 72 degrees 35 minutes 50 seconds W., NAD 27, in a wooded area:

- A—0 to 4 inches; dark brown (10YR 3/3) fine sandy loam, pale brown (10 YR 6/3) dry; weak medium granular structure; friable; common very fine and fine roots; 5 percent gravel; strongly acid; abrupt smooth boundary.
- Bw1—4 to 14 inches; dark yellowish brown (10YR 4/6) fine sandy loam; weak medium subangular blocky structure; friable; many very fine and common fine and medium roots; 5 percent gravel; strongly acid; clear smooth boundary.
- Bw2—14 to 25 inches; dark yellowish brown (10YR 4/4) sandy loam; weak medium subangular blocky structure; friable; common very fine, fine, and medium roots; 10 percent gravel; strongly acid; abrupt smooth boundary.
- 2Cd1—25 to 39 inches; dark grayish brown (2.5Y 4/2) gravelly loamy coarse sand; massive; firm; few fine roots; 20 percent gravel; strongly acid; clear smooth boundary.
- 2Cd2—39 to 60 inches; dark grayish brown (2.5Y 4/2) gravelly sandy loam; massive; firm; 25 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 20 to 38 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

- Hue—7.5YR or 10YR
- Value—2 to 5
- Chroma—1 to 4
- Content of rock fragments—3 to 14 percent

Bw1 horizon:

- Hue—7.5YR or 10YR
- Value—4 or 5
- Chroma—3 to 8
- Texture of the fine earth fraction—loam, fine sandy loam or sandy loam
- Content of rock fragments—3 to 34 percent

Bw2 horizon:

- Hue—7.5YR to 2.5Y
- Value—4 to 6
- Chroma—3 to 6
- Texture of the fine earth fraction—loam, fine sandy loam or sandy loam
- Content of rock fragments—3 to 34 percent

2Cd horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 to 6

Texture of the fine earth fraction—loamy coarse sand, sandy loam, or loamy sand

Content of rock fragments—5 to 34 percent

Moosilauke Series*Depth class:* Very deep*Drainage class:* Poorly drained*Permeability:* Moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum*Landform:* Depressions and drainageways on outwash plains and terraces*Parent material:* Glaciofluvial deposits derived from granite, schist, and gneiss*Slope range:* 0 to 3 percent*Associated soils in a drainage sequence:*

Boscawen (excessively drained, sandy-skeletal)

Merrimac, cold (somewhat excessively drained)

Sudbury, cold (moderately well drained)

Associated similar soils:

Scarboro, cold (very poorly drained)

Taxonomic class: Sandy, mixed, frigid Aeric Endoaquepts**Typical Pedon**

Moosilauke loam, located in the town of Norfolk, 1,800 feet northwest along Gamefields Road from the intersection with Windrow Road, 50 feet east of Gamefields Road and 100 feet west of Tobey Pond, on the Norfolk USGS topographic quadrangle, lat. 41 degrees 58 minutes 33 seconds N., long. 73 degrees 13 minutes 35 seconds W., NAD 27, in a wooded area:

Oa—0 to 1 inch; very dark gray (5YR 3/1) highly decomposed plant materials

A—1 to 6 inches; very dark gray (10YR 3/1) loam; moderate fine granular structure; very friable; common fine distinct brown (7.5YR 4/3) masses of iron accumulation and few medium faint dark gray (10YR 4/1) iron depletions; many fine and very fine roots, common medium roots, and few very coarse and coarse roots; strongly acid; clear wavy boundary.

Bg—6 to 16 inches; dark grayish brown (10YR 4/2) fine sandy loam; weak medium subangular blocky structure; very friable; common very fine and fine roots; common fine faint brown (7.5YR 4/3) mass of iron accumulation and common fine faint gray (7.5YR 5/1) iron depletions; 12 percent gravel; strongly acid; clear smooth boundary.

Bw—16 to 24 inches; dark yellowish brown (10YR 4/4) gravelly fine sandy loam; weak medium subangular blocky structure; friable; common medium distinct dark yellowish brown (10YR 4/6) and few fine prominent strong brown (7.5YR 5/8) soft masses of iron accumulation, few fine distinct light brownish gray (10YR 6/2) iron depletions; 20 percent gravel; strongly acid; gradual smooth boundary.

C1—24 to 39 inches; dark yellowish brown (10YR 4/4) loamy fine sand; massive; very friable; common coarse prominent yellowish red (5YR 4/6) soft masses of iron accumulation; 10 percent gravel; moderately acid; gradual smooth boundary.

C2—39 to 65 inches; light olive brown (2.5Y 5/4) fine sand; single grain; loose; 5 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: 18 to 28 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—10YR or 2.5Y

Value—2 to 4

Chroma—1 or 2

Content of rock fragments—0 to 14 percent

Bg horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—0 to 25 percent

Bw horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—0 to 25 percent

C horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 to 4

Texture of the fine earth fraction—loamy sand to sand

Content of rock fragments—0 to 50 percent

Mudgepond Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate in the surface layer, moderate or moderately rapid in the subsoil, and moderate in the substratum

Landform: Depressions and drainageways on hills and plains

Parent material: Till from limestone, dolomite, and schist

Slope range: 0 to 8 percent

Associated soils in a drainage sequence:

Nellis (well drained)

Amenia (moderately well drained)

Alden (very poorly drained)

Associated other soils:

Stockbridge (well drained, carbonates at greater depth)

Georgia (moderately well drained, carbonates at greater depth)

Taxonomic class: Coarse-loamy, mixed, superactive, mesic Typic Endoaquolls

Typical Pedon

Mudgepond silt loam, located in the town of Sharon, 4,000 feet southeast along Norton Hill Road from the intersection with Race Track Road and 800 feet southwest of Norton Hill Road, on the Sharon USGS topographic quadrangle, lat. 41 degrees, 56 minutes, 29 seconds N. and 73 degrees, 24 minutes, 33 seconds W. , NAD 27, in a wooded area:

- A—0 to 11 inches; very dark gray (10YR 3/1) silt loam; gray (10YR 5/1) dry; weak fine and medium granular structure; very friable; slightly sticky, slightly plastic; many very fine to very coarse roots; 5 percent subrounded gravel; slightly alkaline; clear wavy boundary.
- Bg—11 to 16 inches; olive gray (5Y 5/2) loam; weak medium subangular blocky structure; friable; slightly sticky, slightly plastic; few very fine and fine roots; common medium faint light olive brown (2.5Y 5/3) soft masses of iron accumulation and common coarse faint gray (5Y 5/1) iron depletions; 10 percent subrounded gravel; slightly alkaline; gradual smooth boundary.
- Bw1—16 to 26 inches; light olive brown (2.5Y 5/3) fine sandy loam; weak medium subangular blocky structure; friable; slightly sticky, slightly plastic; few very fine and fine roots; common medium prominent olive yellow (2.5Y 6/8) soft masses of iron accumulation and common coarse faint grayish brown (2.5Y 5/2) iron depletions; 10 percent subrounded gravel; slightly alkaline; gradual wavy boundary.
- Bw2—26 to 35 inches; light olive brown (2.5Y 5/4) gravelly fine sandy loam; weak medium subangular blocky structure; friable; non sticky, non plastic; common medium prominent yellowish brown (10 YR 5/8) soft masses of iron accumulation and common medium prominent gray (10YR 6/1) iron depletions; 15 percent subrounded gravel; slightly effervescent; slightly alkaline; gradual smooth boundary.
- C—35 to 65 inches; olive brown (2.5Y 4/4) gravelly fine sandy loam; massive; firm; non sticky, non plastic; common medium distinct yellowish brown (10 YR 5/6) masses of iron accumulation and few fine distinct light olive gray (5 Y 6/2) iron depletions; 15 percent subrounded gravel; slightly effervescent; slightly alkaline.

Range in Characteristics

Solum thickness: 20 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Neutral to slightly alkaline in the surface layer and subsoil, neutral to moderately alkaline in the substratum

Depth to carbonates: Mainly from 12 to 40 inches, but may be deeper than 40 inches

A or Ap horizon:

Hue—10YR to 2.5Y

Value—2 or 3 (5 or less dry)

Chroma—1 to 3

Content of rock fragments—5 to 14 percent

Bg horizon:

Hue—10 YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, loam, or silt loam

Content of rock fragments—5 to 34 percent

Bw Horizons:

Hue—2.5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, loam or silt loam

Content of rock fragments—5 to 34 percent

C or Cg horizon:

Hue—10YR to 5Y

Value—4 or 5

Chroma—1 to 6

Texture of the fine earth fraction—sandy loam, fine sandy loam, very fine sandy loam or loam

Content of rock fragments—5 to 34 percent

Some of the Mudgepond soils in this survey area have a mean annual soil temperature which is colder than typical of the series. These map units (457 and 458) are identified as cold phases of the Mudgepond series.

Narragansett Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate in the surface layer and subsoil, and moderately rapid or rapid in the substratum

Landform: Hills or till plains

Parent material: Eolian deposits over melt-out till derived from gneiss, schist, sandstone and shale

Slope range: 2 to 25 percent

Associated soil in a drainage sequence:

Wapping (moderately well drained, coarse-loamy)

Associated similar soils:

Broadbrook (coarse-loamy, dense substratum)

Canton (coarser textured)

Charlton (coarse-loamy)

Montauk (coarse-loamy, dense substratum)

Paxton (coarse-loamy, dense substratum)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic
Typic Dystrudepts

Typical Pedon

Narragansett silt loam, in an area of Narragansett-Hollis complex, 3 to 15 percent slopes, very rocky, located in the town of Montville, 0.6 miles north of the intersection of Cherry Lane and Raymond Hill Road and 30 feet west of Cherry Lane, on the Montville USGS topographic quadrangle, lat. 41 degrees 29 minutes 28 seconds N., long. 72 degrees 09 minutes 23 seconds W., NAD 27, in a wooded area:

Ap—0 to 6 inches; dark brown (10YR 3/3) silt loam, pale brown (10 YR 6/3) dry; weak medium granular structure; very friable; common medium roots; very strongly acid; clear wavy boundary.

Bw1—6 to 15 inches; dark yellowish brown (10YR 4/6) silt loam; weak medium subangular blocky structure; very friable; common medium roots; very strongly acid; gradual wavy boundary.

Bw2—15 to 24 inches; yellowish brown (10YR 5/6) silt loam; weak medium subangular blocky structure; very friable; common medium roots; strongly acid; clear wavy boundary.

Bw3—24 to 28 inches; yellowish brown (10YR 5/6) gravelly silt loam; weak medium subangular blocky structure; very friable; few fine roots; 15 percent gravel; strongly acid; clear wavy boundary.

2C—28 to 60 inches; light olive brown (2.5Y 5/4) very gravelly loamy coarse sand; single grain; loose; 45 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 18 to 38 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—7.5YR or 10YR

Value—3 or 4 (Dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—0 to 14 percent

Upper Bw horizons:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—4 to 8

Texture of the fine earth fraction—silt loam, very fine sandy loam, or loam

Content of rock fragments—0 to 25 percent

Lower Bw horizons:

Hue—7.5YR to 2.5Y

Value—4 to 7

Chroma—4 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam, or loam

Content of rock fragments—0 to 25 percent

2C horizon:

Hue—2.5YR to 5Y

Value—3 to 5

Chroma—2 to 6

Texture of the fine earth fraction—loamy coarse sand, loamy sand, or sand

Content of rock fragments—10 to 50 percent

Natchaug Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate to very rapid in the organic material and moderate or moderately slow in the loamy substratum

Landform: Depressions

Parent Material: Organic materials over alluvium, till, glaciolacustrine deposits, or glaciofluvial deposits

Slope range: 0 to 2 percent

Associated similar soils:

Catden (organic materials greater than 51 inches deep)

Freetown (dysic reaction class, organic materials greater than 51 inches deep)

Timakwa (16 to 51 inches of organic materials, over sandy or sandy-skeletal)

Taxonomic class: Loamy, mixed, euic, mesic Terric Haplosaprists

Typical Pedon

Natchaug peat, in an area of Timakwa and Natchaug soils, located in the town of Eastford, 1,500 feet west on Pifershire Road from the intersection of Pifershire Road and Fayette Wright Road, 500 feet south of Pifershire Road, on the Hampton USGS topographic quadrangle, lat. 41 degrees 50 minutes 35 seconds N., long. 72 degrees 03 minutes 36 seconds W., NAD 27, in a swamp:

- Oi1—0 to 2 inches; dark yellowish brown (10YR 4/6) peat; 90 percent fibers, 90 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; common fine to medium roots; 5 percent coarse woody fragments; less than 5 percent mineral material; ultra acid; abrupt smooth boundary.
- Oi2—2 to 4 inches; dark brown (7.5YR 3/4) peat; 80 percent fibers, 60 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; common fine to medium roots; 5 percent coarse woody fragments; less than 5 percent mineral material; extremely acid; clear smooth boundary.
- Oa1—4 to 6 inches; dark reddish brown (5YR 3/2) muck; 25 percent fibers, 7 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; common fine to medium roots; 5 percent coarse woody fragments; less than 5 percent mineral material; extremely acid; clear smooth boundary.
- Oa2—6 to 11 inches; dark reddish brown (5YR 2.5/2) muck; 20 percent fibers, 2 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; few fine and medium roots; 2 percent coarse woody fragments; less than 5 percent mineral material; extremely acid; clear smooth boundary.
- Oa3—11 to 18 inches; black (5YR 2.5/1) muck; 5 percent fibers, 0 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; 2 percent coarse woody fragments; less than 5 percent mineral material; extremely acid; clear smooth boundary.
- Oa4—18 to 24 inches; black (10YR 2/1) muck; 0 percent fibers, 0 percent rubbed; massive; soft, very friable, nonsticky and nonplastic; less than 5 percent mineral material; very strongly acid; abrupt wavy boundary.
- 2Cg1—24 to 33 inches; grayish brown (10YR 5/2) fine sandy loam; massive; slightly hard, friable, nonsticky and nonplastic; common medium faint gray (10YR 6/1) iron depletions, common coarse prominent strong brown (7.5YR 4/6) and common medium distinct yellowish brown (10YR 5/4) soft masses of iron accumulation; strongly acid; clear smooth boundary.
- 2Cg2—33 to 36 inches; gray (2.5Y 6/1) fine sandy loam; massive; slightly hard, friable, nonsticky and nonplastic; common percent medium prominent yellowish brown (10YR 5/6) soft masses of iron accumulation; moderately acid; clear smooth boundary.
- 2Cg3—36 to 80 inches; gray (2.5Y 6/1) and yellowish brown (10YR 5/6) loam; massive; slightly hard, friable, nonsticky and nonplastic; 1 percent fine gravel; moderately acid

Range in Characteristics

Solum thickness: Organic layers 16 to 51 inches deep

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to moderately acid (in 0.01 M calcium chloride) in the organic layers and strongly acid to neutral in the loamy substratum (some layers may range to ultra acid)

Woody fragments in organic soil materials: 2 to 15 percent

Oi horizons: (absent in some pedons)

Hue—10YR to 5YR, or is neutral

Value—2 to 4

Chroma—0 to 6

Oa horizons:

Hue—10YR to 5YR, or is neutral

Value—2 to 3

Chroma—0 to 4

2Cg horizons:

Hue—10YR to 5Y, or is neutral

Value—4 to 6
 Chroma—1 or 2
 Texture of the fine earth fraction—very fine sandy loam, fine sandy loam, sandy loam, loam, or silt loam
 Content of rock fragments—0 to 20 percent

Nellis Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid

Landform: Hills or bedrock-controlled ridges

Parent material: Melt-out till derived from limestone, dolomite, and schist

Slope range: 3 to 35 percent

Associated soils in a drainage sequence:

Amenia (moderately well drained)

Mudgepond (poorly drained)

Alden (very poorly drained, fine-loamy)

Associated similar soils:

Charlton (no carbonates)

Stockbridge (deeper to carbonates)

Associated other soils:

Copake (coarse-loamy over sandy or sandy skeletal)

Farmington (shallow to limestone bedrock)

Taxonomic class: Coarse-loamy, mixed, superactive, mesic Typic Eutrudepts

Typical Pedon

Nellis fine sandy loam, 3 to 8 percent slopes, located in the town of Ridgefield, 0.2 miles south of Haviland Road and 400 feet east of Limekiln Road, on the Bethel USGS topographic quadrangle, lat. 41 degrees 18 minutes 19 seconds N., 73 degrees 28 minutes 37 seconds W., NAD 27:

Ap—0 to 8 inches, very dark grayish brown (10YR 3/2) fine sandy loam, pale brown (10 YR 6/3) dry; moderate medium granular structure; friable; many fine roots; 10 percent rock fragments; neutral, clear wavy boundary.

Bw1—8 to 14 inches, dark yellowish brown (10YR 3/4) fine sandy loam; weak medium subangular blocky structure; very friable; common fine roots; 10 percent rock fragments; neutral; gradual wavy boundary.

Bw2—14 to 25 inches, dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; very friable; few fine roots; 5 percent rock fragments; neutral; gradual wavy boundary.

BC—25 to 27 inches, dark yellowish brown (10YR 4/4) loam; massive; friable; few fine roots; 5 percent rock fragments; slightly alkaline; calcareous; clear wavy boundary.

C—27 to 60 inches, very pale brown (10YR 7/4) sandy loam; massive; very friable; 10 percent rock fragments; slightly alkaline; calcareous.

Range in Characteristics

Solum thickness: 15 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to neutral in the surface layer and upper subsoil, moderately acid to slightly alkaline in the lower subsoil, and neutral to moderately alkaline in the substratum (calcareous within 40 inches)

Ap horizon:

Hue—10YR or 2.5Y
 Value—3 or 4
 Chroma—2 to 4
 Content of rock fragments—3 to 14 percent

Bw horizons:

Hue—7.5YR to 2.5Y
 Value—3 to 5
 Chroma—2 to 4
 Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, loam, or silt loam
 Content of rock fragments—5 to 34 percent

BC horizon:

Hue—7.5YR to 2.5Y
 Value—3 to 5
 Chroma—2 to 6
 Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, loam, or silt loam
 Content of rock fragments—5 to 34 percent

C horizon:

Hue—10YR to 5Y
 Value—3 to 7
 Chroma—2 to 6
 Texture of the fine earth fraction—sandy loam, fine sandy loam, or loam
 Content of rock fragments—5 to 50 percent

Ninigret Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Eolian deposits over glaciofluvial deposits derived from granite, schist, and gneiss

Slope range: 0 to 5 percent

Associated soil in a drainage sequence:

Agawam (well drained)

Associated similar soil:

Tisbury (coarse-silty over sandy or sandy-skeletal)

Associated other soils:

Enfield (well drained, coarse-silty over sandy or sandy-skeletal)

Haven (well drained)

Raypol (poorly drained)

Walpole (poorly drained, sandy)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Aquic Dystrudepts

Typical Pedon

Ninigret fine sandy loam, in an area of Ninigret and Tisbury soils, 0 to 5 percent slopes, located in the town of Stonington, 1,000 feet southwest along Riverside Drive from the intersection with Green Haven Road; 300 feet east of Riverside Drive and 350 feet west of the Pawcatuck River, on the Watch Hill USGS topographic

quadrangle, lat. 41 degrees 20 minutes 03 seconds N., long. 71 degrees 50 minutes 36 seconds W., NAD 27, in a grassy field:

- Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) fine sandy loam; pale brown (10YR 6/3) dry; weak medium granular structure; very friable; many fine roots; strongly acid; abrupt smooth boundary.
- Bw1—8 to 16 inches; yellowish brown (10YR 5/6) fine sandy loam; weak coarse granular structure; very friable; few fine roots; strongly acid; clear wavy boundary.
- Bw2—16 to 26 inches; yellowish brown (10YR 5/4) fine sandy loam; very weak coarse granular structure; very friable; very few fine roots; common medium distinct light brownish gray (10YR 6/2) iron depletions and brownish yellow (10YR 6/6) soft masses of iron accumulation; strongly acid; clear wavy boundary.
- 2C—26 to 65 inches; pale brown (10YR 6/3) loamy sand and few lenses of loamy fine sand; single grain; loose; many medium distinct light olive gray (5Y 6/2) iron depletions and prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; strongly acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid to a depth of 30 inches and very strongly acid to slightly acid below 30 inches

Ap horizon:

Hue—7.5YR to 2.5Y

Value—2 to 4 (dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—0 to 14 percent

Bw1 horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—4 to 8

Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, or silt loam

Content of rock fragments—0 to 14 percent

Bw2 horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—2 to 6

Texture of the fine earth fraction—fine sandy loam, very fine sandy loam, or silt loam

Content of rock fragments—0 to 14 percent

2C horizon:

Hue—10YR to 5Y

Value—4 to 7

Chroma—1 to 6

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—0 to 30 percent above a depth of 40 inches, 0 to 60 percent below

Some of the Ninigret soils in this survey area have a mean annual soil temperature which is colder than typical of the series. This map unit (421A) is identified as a cold phase of the Ninigret series.

Occum Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Pootatuck (moderately well drained)

Rippowam (poorly drained)

Associated similar soil:

Hadley (coarse-silty)

Associated other soils:

Suncook (excessively drained, sandy)

Agawam (doesn't flood, coarse-loamy over sandy or sandy-skeletal)

Lim (poorly drained)

Limerick (poorly drained, coarse-silty)

Saco (very poorly drained, coarse-silty)

Taxonomic class: Coarse-loamy, mixed, superactive, mesic Fluventic Dystrudepts

Typical Pedon

Occum fine sandy loam, located in the town of Granby, 1,400 feet west of the junction of Connecticut Route 10 and Mechanicsville Road, on the east side of the East Branch Salmon Brook, on the Tariffville USGS topographic quadrangle, lat. 41 degrees 58 minutes 15 seconds N., long. 72 degrees 48 minutes 11 seconds W., NAD 27, in a hayfield:

Ap—0 to 10 inches; dark brown (10YR 3/3) fine sandy loam, pale brown (10YR 6/3) dry; weak fine and medium granular structure; very friable; many very fine and fine roots; moderately acid; clear smooth boundary.

Bw1—10 to 17 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; very friable; common very fine and fine roots; moderately acid; clear smooth boundary.

Bw2—17 to 28 inches; dark yellowish brown (10YR 4/6) sandy loam; weak medium subangular blocky structure; very friable; few very fine and fine roots; moderately acid; clear smooth boundary.

C1—28 to 32 inches; yellowish brown (10YR 5/4) loamy sand; single grain; loose; moderately acid; clear smooth boundary.

C2—32 to 42 inches; brown (10YR 5/3) and light olive brown (2.5Y 5/4) sand; single grain; loose; 10 percent gravel; moderately acid; clear smooth boundary.

C3—42 to 65 inches; brown (10YR 5/3) and light olive brown (2.5Y 5/4) very gravelly coarse sand; single grain; loose; 35 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid

Ap horizon:

Hue—10YR or 2.5Y

Value—3 to 5

Chroma—1 to 4

Content of rock fragments—0 to 14 percent

Bw1 horizon:

Hue—7.5YR to 2.5Y

Value—3 to 8

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, sandy loam, or loam

Content of rock fragments—0 to 14 percent

Bw2 horizon:

Hue—7.5YR to 2.5Y

Value—3 to 8

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam or sandy loam

Content of rock fragments—0 to 14 percent

C horizons:

Hue—7.5YR to 5Y

Value—3 to 7

Chroma—2 to 6

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—0 to 59 percent

Ondawa Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum*Landform:* Flood plains*Parent material:* Alluvium*Slope range:* 0 to 3 percent*Associated soils in a drainage sequence:*

Rumney (poorly drained)

Associated other soils:

Medomak Taxadjunct (very poorly drained, coarse-silty)

Taxonomic class: Coarse-loamy, mixed, active, frigid Fluventic Dystrudepts**Typical Pedon**

Ondawa fine sandy loam, located in the town of Colebrook, 1,000 feet northwest along Egler Road from the intersection with Connecticut Route 183 and 1,000 feet south of Egler Road, on the Tolland Center USGS topographic quadrangle, lat. 42 degrees 01 minutes 03 seconds N., long. 73 degrees 07 minutes 17 seconds W., NAD 27, in a wooded area:

Oi—0 to 1 inch; strong brown (7.5YR 4/6) slightly decomposed plant materials

Oa—1 to 2 inches; very dark brown (7.5YR 2.5/2) highly decomposed plant materials

Ap—2 to 14 inches; dark brown (10YR 3/3) fine sandy loam, pale brown (10YR 6/3) dry; weak fine granular structure; very friable; many very fine and fine roots; very strongly acid; clear smooth boundary.

Bw—14 to 30 inches; dark yellowish brown (10YR 4/6) fine sandy loam; moderate medium subangular blocky structure; very friable; common very fine to very coarse roots; strongly acid; clear smooth boundary.

C1—30 to 33 inches; light olive brown (2.5Y 5/4) sandy loam; massive; very friable; few very fine to coarse roots; 5 percent gravel; strongly acid; clear smooth boundary.

C2—33 to 60 inches; light olive brown (2.5Y 5/4) gravelly sandy loam; massive; 25 percent gravel; strongly acid

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid

A horizon:

Hue—10YR or 2.5Y

Value—3 to 5 (Dry value 6 or more)

Chroma—1 to 4

Content of rock fragments—0 to 14 percent

Bw horizon:

Hue—10YR or 2.5Y

Value—3 to 8

Chroma—2 to 8

Texture of the fine earth fraction—fine sandy loam, sandy loam, or loam

Content of rock fragments—0 to 14 percent

C horizons:

Hue—10YR to 5Y

Value—3 to 7

Chroma—2 to 6

Texture of the fine earth fraction—loamy fine sand to coarse sand (upper C horizon may be sandy loam)

Content of rock fragments—0 to 40 percent

Pawcatuck Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate to rapid in the organic layers and very rapid in the substratum

Landform: Tidal marshes and salt marshes

Parent material: Grassy organic materials over glaciofluvial deposits

Slope range: 0 to 2 percent

Associated similar soils:

Ipswich (organic material thicker than 51 inches)

Westbrook (organic over loamy)

Taxonomic class: Sandy or sandy-skeletal, mixed, euic, mesic Terric Sulfihemists

Typical Pedon

Pawcatuck mucky peat, located in the town of Stonington, Barn Island area, 2,600 feet northeast of the elevation benchmark on Barn Island, 4,000 feet north of the elevation benchmark on Pawcatuck Point, and 1,000 feet south of a gravel road, on the Watch Hill USGS topographic quadrangle, lat. 41 degrees 20 minutes 00 seconds N., long. 72 degrees 38 minutes 07 seconds W., NAD 27, in a tidal marsh:

Oe1—0 to 12 inches; very dark gray (10YR 3/1) mucky peat, dark grayish brown (10YR 4/2) dry; 65 percent fiber, 30 percent rubbed; dense mat of roots; stems, and leaves; massive; slightly sticky; many very fine, fine, and medium roots; fibers herbaceous; thin lenses and coatings of silt; 57 percent organic matter; strongly saline; slightly acid; clear wavy boundary.

Oe2—12 to 40 inches; black (10YR 2/1) mucky peat, very dark grayish brown (10YR 3/1) dry; 50 percent fiber, 25 percent rubbed; massive; slightly sticky; few very

- fine, fine, and medium roots; fibers herbaceous; 54 percent organic matter; strongly saline; slightly acid; gradual wavy boundary.
- Oe3—40 to 46 inches; black (10YR 2/1) moist and dry mucky peat; 40 percent fiber, 25 percent rubbed; massive; slightly sticky; fibers herbaceous; 27 percent organic matter; strongly saline; slightly acid; clear wavy boundary.
- 2C1—46 to 50 inches; gray (N 5/0) very fine sandy loam, gray (10YR 5/1) dry; 10 percent organic matter; massive; slightly sticky; strongly saline; slightly acid; clear wavy boundary.
- 2C2—50 to 60 inches; black (10YR 2/1) loamy sand, dark gray (10YR 4/1) dry; single grain; loose; 10 percent rock fragments; strongly saline; slightly acid.

Range in Characteristics

Solum thickness: 16 to 51 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to slightly alkaline

Salinity: Moderately saline to strongly saline

Surface tier:

Hue—5YR to 5Y

Value—2 to 5

Chroma—0 to 3

Content of rock fragments—none

Subsurface and Bottom tiers:

Hue—5YR to 5Y

Value—2 to 5

Chroma—0 to 3

Organic materials—dominantly hemic, some pedons contain fibric or sapric materials

Content of rock fragments—none

2C1 horizon:

Hue—10YR to 5BG

Value—2 to 7

Chroma—0 to 3

Texture of the fine earth fraction—very fine sandy loam, silt loam or sandy loam

Content of rock fragments—0 to 14 percent

2C2 horizon:

Hue—10YR to 5BG

Value—2 to 7

Chroma—0 to 3

Texture of the fine earth fraction—loamy fine sand, loamy sand or sand

Content of rock fragments—0 to 25 percent

Paxton Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Hills and drumlins

Parent material: Lodgement till derived from granite, gneiss and schist

Slope range: 3 to 35 percent

Associated soils in a drainage sequence:

Woodbridge (moderately well drained)

Ridgebury taxadjunct (poorly drained)

Whitman (very poorly drained)

Associated similar soils:

Broadbrook (siltier in surface layer and subsoil)

Canton (coarse-loamy over sandy or sandy-skeletal)

Charlton (friable substratum)

Montauk (sandy substratum)

Narragansett (coarse-loamy over sandy or sandy-skeletal)

Wethersfield (redder)

Taxonomic class: Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts

Typical Pedon

Paxton fine sandy loam, in an area of Paxton and Montauk fine sandy loams, 3 to 8 percent slopes, very stony, located in the town of Prospect. 0.4 miles east of Straitsville Road and 0.5 miles north of the Bethany–Prospect line, on the Mount Carmel USGS topographic quadrangle, lat. 41 degrees 28 minutes 34 seconds N., long. 72 degrees 59 minutes 16 seconds W., NAD 27, in a brushy field:

Ap—0 to 8 inches; dark brown (10YR 3/3) fine sandy loam, pale brown (10YR 6/3) dry; moderate medium granular structure; friable; many fine roots; 5 percent gravel; strongly acid; abrupt smooth boundary.

Bw1—8 to 15 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; common fine roots; 5 percent gravel; few earthworm casts; strongly acid; gradual wavy boundary.

Bw2—15 to 26 inches; olive brown (2.5Y 4/4) fine sandy loam; weak medium subangular blocky structure; friable; few fine roots; 10 percent gravel; strongly acid; clear wavy boundary.

Cd—26 to 65 inches; olive (5Y 5/3) gravelly fine sandy loam; moderate thick platy structure; very firm, brittle; 25 percent gravel; many dark coatings on plates; strongly acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid in the surface layer, very strongly acid to moderately acid in the subsoil and substratum

Ap horizon:

Hue—10YR

Value—3 or 4 (dry value 6 or more) (If A horizon present instead of Ap, value 2 or 3)

Chroma—2 to 4 (If A horizon present instead of Ap, chroma 1 or 2)

Content of rock fragments—5 to 14 percent

Bw1 horizon:

Hue—7.5YR or 10YR

Value—4 to 6

Chroma—4 to 8

Texture of the fine earth fraction—loam, fine sandy loam, or sandy loam

Content of rock fragments—5 to 34 percent

Bw2 horizon:

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture of the fine earth fraction—loam, fine sandy loam, or sandy loam

Content of rock fragments—5 to 34 percent

Cd horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 to 4

Texture of the fine earth fraction—loam, fine sandy loam, or sandy loam

Content of rock fragments—5 to 34 percent

Penwood Series*Depth class:* Very deep*Drainage class:* Excessively drained*Permeability:* Rapid or very rapid*Landform:* Outwash plains and terraces*Parent material:* Glaciofluvial deposits derived from sandstone and shale*Slope range:* 0 to 8 percent*Associated similar soils:*

Manchester (sandy-skeletal)

Hartford (somewhat excessively drained, sandy)

Branford (well drained, coarse-loamy over sandy or sandy-skeletal)

Windsor (brownier)

Taxonomic class: Mixed, mesic Typic Udipsamments**Typical Pedon**

Penwood loamy sand, 3 to 8 percent slopes, located in the town of East Haven, 1 mile north on Connecticut Route 100 from the interchange of Connecticut Route 100 and Interstate 95, and 250 feet east of Connecticut Route 100 just south of the State Rifle Range, on the Branford USGS topographic quadrangle, lat. 49 degrees 17 minutes 48 seconds N., long. 72 degrees 51 minutes 45 seconds W., NAD 27, in a brushy field:

Ap—0 to 8 inches; dark brown (7.5YR 3/2) loamy sand; pinkish gray (7.5YR 6/2) dry; weak medium granular structure; very friable; many fine roots; strongly acid; clear smooth boundary.

Bw1—8 to 18 inches; yellowish red (5YR 4/6) loamy sand; single grain, loose; common fine roots; strongly acid; gradual wavy boundary.

Bw2—18 to 30 inches; reddish brown (5YR 4/4) sand; single grain, loose; few fine roots; strongly acid; gradual wavy boundary.

C—30 to 60 inches; reddish brown (5YR 4/3) sand with thin strata of fine sand; single grain; loose; few fine roots in the upper part; strongly acid.

Range in Characteristics*Solum thickness:* 20 to 36 inches*Depth to bedrock:* More than 80 inches*Reaction:* Very strongly acid to moderately acid**Ap horizon:**

Hue—10YR to 5YR

Value—3 or 4 (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—0 to 10 percent

Bw1 horizon:

Hue—5YR or 2.5YR

Value—4 or 5

Chroma—4 to 8

Texture of the fine earth fraction—loamy sand or loamy fine sand
Content of rock fragments—0 to 10 percent

Bw2 horizon:

Hue—5YR or 2.5YR
Value—4 or 5
Chroma—4 to 8
Texture of the fine earth fraction—sand or fine sand
Content of rock fragments—0 to 10 percent

C horizon:

Hue—2.5YR to 7.5YR
Value—4 to 6
Chroma—3 to 6
Texture of the fine earth fraction—sand or fine sand
Content of rock fragments—0 to 14 percent

Pootatuck Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Occum (well drained)
Rippowam (poorly drained)

Associated similar soil:

Winooski (coarse-silty)

Associated other soils:

Suncook (excessively drained, sandy)
Hadley (well drained, coarse-silty)
Limerick (poorly drained, coarse-silty)
Lim (poorly drained)
Saco (very poorly drained, coarse-silty)

Taxonomic class: Coarse-loamy, mixed, active, mesic Fluvaquentic Dystrudepts

Typical Pedon

Pootatuck fine sandy loam, located in the town of Easton, 200 feet east of Connecticut Route 58, 0.1 mile north of Silver Hill Road, and 80 feet west of the Aspetuck River, on the Botsford USGS topographic quadrangle, lat. 41 degrees 16 minutes 40 seconds N., long. 73 degrees 19 minutes 32 seconds W., NAD 27, in a wooded area:

A—0 to 4 inches; very dark grayish brown (10YR 3/2) fine sandy loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; friable; common fine and medium roots; strongly acid; gradual wavy boundary.

Bw1—4 to 16 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; common fine and medium roots; strongly acid; clear wavy boundary.

Bw2—16 to 21 inches; brown (10YR 4/3) fine sandy loam; weak medium subangular blocky structure; friable; common fine medium roots; few medium prominent

- strong brown (7.5YR 5/6) soft masses of iron accumulation and faint grayish brown (10YR 5/2) iron depletions; moderately acid; gradual wavy boundary.
- Bw3—21 to 29 inches; dark brown (10YR 3/3) sandy loam; weak medium subangular blocky structure; friable; common fine roots; common medium faint grayish brown (10YR 5/2) iron depletions and prominent strong brown (7.5YR 5/6) soft masses of iron accumulation; moderately acid; clear wavy boundary.
- C1—29 to 35 inches; brown (10YR 4/3) sand; single grain; loose; few fine roots; common medium distinct grayish brown (10YR 5/2) iron depletions and prominent strong brown (7.5YR 5/6) soft masses of iron accumulation; moderately acid; clear wavy boundary;
- C2—35 to 40 inches; grayish brown (2.5Y 5/2) sand; single grain; loose; few fine faint pale brown (10YR 6/3) soft masses of iron accumulation; 5 percent gravel; moderately acid; clear wavy boundary.
- C3—40 to 65 inches; grayish brown (10YR 5/2) gravelly sand; single grain; loose; 25 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid

A horizon:

Hue—10YR or 2.5Y

Value—3 to 5

Chroma—1 to 4

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam or sandy loam

Content of rock fragments—0 to 14 percent

C horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 to 6

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—0 to 40 percent

Pyrities Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate in the surface layer and subsoil, moderately slow or slow in the substratum

Landform: Hills

Parent material: Till derived from limestone, schist, and dolomite

Slope range: 3 to 25 percent

Associated soils in a drainage sequence:

Hogansburg (moderately well drained)

Alden, cold (very poorly drained)

Mudgepond, cold (poorly drained)

Associated similar soil:

Bice (no carbonates)

Taxonomic class: Coarse-loamy, mixed, active, frigid Dystric Eutrudepts**Typical Pedon**

Pyrities loam, 8 to 15 percent slopes, located in the town of Canaan, 300 feet south of Canaan Mountain Road and 1600 feet southwest of the south end of Wangum Lake, on the South Canaan USGS topographic quadrangle, lat. 41 degrees 58 minutes 22 seconds N., long. 73 degrees 16 minutes 14 seconds W., NAD 27, in a wooded area:

Oi—0 to 1 inch, slightly decomposed plant materials

Ap—1 to 8 inches, very dark grayish brown (10YR 3/2) loam, light brownish gray (10YR 6/2) dry; weak medium granular structure; very friable, slightly sticky, slightly plastic; many very fine roots, common medium to coarse roots; 5 percent gravel, 2 percent cobbles; neutral; abrupt wavy boundary.

Bw1—8 to 13 inches, brown (10 YR 4/3) loam; moderate medium subangular blocky structure; friable, slightly sticky, slightly plastic; common very fine to coarse roots; 5 percent gravel, 2 percent cobbles; neutral; clear smooth boundary.

Bw2—13 to 26 inches, dark yellowish brown (10YR 4/4) loam; weak coarse subangular blocky structure; friable, slightly sticky, slightly plastic; common very fine and fine roots, few medium and coarse roots; 5 percent gravel, 1 percent cobbles; neutral; clear smooth boundary.

BC—26 to 45 inches, light olive brown (2.5Y 5/3) loam; weak thin platy structure; friable, slightly sticky, slightly plastic; few very fine to coarse roots; 5 percent gravel, 1 percent cobbles; neutral; clear smooth boundary.

C—45 to 60 inches; light olive brown (2.5YR 4/4) fine sandy loam; massive; friable, slightly sticky, slightly plastic; 10 percent, 2 percent cobbles; slightly alkaline; slightly effervescent.

Range in Characteristics

Solum thickness: 25 to 50 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to neutral in the surface layer, slightly acid to slightly alkaline in the subsoil, and slightly acid to moderately alkaline in the substratum

Ap horizon:

Hue—7.5YR or 10YR

Value—5 or 6 (dry value 6 or more)

Chroma—1 to 3

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—7.5YR or 10YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam or loam

Content of rock fragments—5 to 34 percent

BC horizon:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam or loam

Content of rock fragments—5 to 34 percent

C horizon:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam or loam

Content of rock fragments—5 to 50 percent

Rainbow Series*Depth class:* Very deep*Drainage class:* Moderately well drained*Permeability:* Moderate in the surface layer and subsoil, and slow or very slow in the substratum*Landform:* Hills and drumlins*Parent material:* Eolian deposits over lodgement till derived from gneiss, schist, sandstone and basalt*Slope range:* 0 to 8 percent*Associated soil in a drainage sequence:*

Broadbrook (well drained)

Associated similar soils:

Ludlow (redder in the subsoil)

Wapping (friable substratum)

Watchaug (friable substratum)

Taxonomic class: Coarse-loamy, mixed, active, mesic Aquic Dystrudepts**Typical Pedon**

Rainbow silt loam, 2 to 8 percent slopes, very stony, located in the town of Montville, 3,000 feet west of Connecticut Route 32 and 1,000 feet north of the Montville-Waterford town line, on the Uncasville USGS topographic quadrangle, lat. 41 degrees 25 minutes 33 seconds N., long. 72 degrees 07 minutes 08 seconds W., NAD 27, in an old field:

Ap—0 to 6 inches; dark brown (10YR 3/3) silt loam, pale brown (10YR 6/3) dry; weak fine granular structure; very friable; many fine roots; few pebbles; strongly acid; clear wavy boundary

Bw1—6 to 18 inches; yellowish brown (10YR 5/6) silt loam; weak medium subangular blocky structure; very friable; few fine roots; few pebbles; strongly acid; clear wavy boundary.

Bw2—18 to 26 inches; light yellowish brown (10YR 6/4) silt loam; weak fine and medium subangular blocky structure; very friable; few fine roots; common medium prominent light gray (5Y 7/1) iron depletions and distinct strong brown (7.5YR 5/6) soft masses of iron accumulation; few pebbles; strongly acid; clear wavy boundary.

2Cd—26 to 65 inches; pale brown (10YR 6/3) gravelly fine sandy loam; weak thick platy structure; very firm, brittle; common silt films on rock fragments; common medium faint light olive brown (2.5Y 5/4) and distinct brownish yellow (10YR 6/6) soft masses of iron accumulation; 15 percent gravel; strongly acid.

Range in Characteristics*Solum thickness:* 20 to 40 inches*Depth to bedrock:* More than 80 inches*Reaction:* Very strongly acid to moderately acid**Ap horizon:**

Hue—7.5YR or 10YR

Value—2 to 4 (Dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 or 2)

Content of rock fragments—0 to 14 percent

Bw1 Horizon:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—4 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam or loam

Content of rock fragments—0 to 20 percent

Bw2 Horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, very fine sandy loam or loam

Content of rock fragments—0 to 20 percent

2Cd Horizon:

Hue—2.5 YR to 5Y

Value—2 to 6

Chroma—2 to 6

Texture of the fine earth fraction—fine sandy loam, sandy loam or loam

Content of rock fragments—5 to 34 percent

Raynham Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate or moderately slow in the surface layer and subsoil, and moderate to slow in the substratum

Landform: Depressions and drainageways on lake plains and terraces

Parent material: Glaciolacustrine deposits

Slope range: 0 to 3 percent

Associated soil in a drainage sequence:

Belgrade (moderately well drained)

Associated similar soils:

Ellington (moderately well drained, coarse-loamy over sandy or sandy-skeletal)

Raypol (coarse-loamy over sandy or sandy-skeletal)

Scitico (fine)

Shaker (coarse-loamy over clayey)

Maybid (very poorly drained, fine)

Taxonomic class: Coarse-silty, mixed, active, nonacid, mesic Aeric Epiaquepts

Typical Pedon

Raynham silt loam, located in the town of Suffield, 2,500 feet south of the intersection of Mountain and Remington roads, 600 feet west of Remington Road, on the Windsor Locks USGS topographic quadrangle, lat. 41 degrees 58 minutes 35 seconds N., 72 degrees 40 minutes 06 seconds W., NAD 27, in a field:

Ap—0 to 10 inches; brown (10YR 4/3) silt loam, pale brown (10YR 6/3) dry; moderate medium and coarse granular structure; friable; common very fine and few fine and medium roots; slightly acid; abrupt smooth boundary.

Bg1—10 to 16 inches; variegated grayish brown (10YR 5/2), yellowish brown (10YR 5/4, 5/6), strong brown (7.5YR 5/6), and yellowish red (5YR 4/6) silt loam, 50

percent of the horizon is chroma 2; weak thin platy structure; friable; few very fine, fine, and medium roots; moderately acid; clear smooth boundary.

Bg₂—16 to 26 inches; grayish brown (10YR 5/2) silt loam; weak very coarse prismatic structure parting to weak thin platy structure; firm; few very fine and fine roots along prism faces; common medium distinct yellowish brown (10YR 5/4), prominent yellowish brown (10YR 5/6) and strong brown (7.5 YR 5/6), and distinct yellowish red (5YR 4/6) variegations and soft masses of iron accumulation; dark grayish brown (10YR 4/2) coatings on prism faces; moderately acid; clear smooth boundary.

Bw—26 to 34 inches; yellowish brown (10YR 5/4) varved silt loam and very fine sandy loam (very fine sandy loam weighted average texture); weak very coarse prismatic structure parting to weak thin platy structure; friable; few very fine and fine roots along prism faces; common medium faint strong brown (7.5YR 5/4) and distinct (7.5YR 5/6) variegations and soft masses of iron accumulation, and distinct light brownish gray (10YR 6/2) iron depletions; discontinuous dark grayish brown (10YR 4/2) coating on prism faces; moderately acid; clear smooth boundary.

Cg—34 to 47 inches; gray (10YR 5/1) and grayish brown (10YR 5/2) varved silt loam, very fine sandy loam, silt, and silty clay loam (weighted average silt loam texture); weak thin platy structure; firm; common medium distinct yellowish brown (10YR 5/4), prominent yellowish brown (10YR 5/6), and prominent strong brown (7.5YR 5/6) variegations and soft masses of iron accumulation; moderately acid; abrupt smooth boundary.

C—47 to 60 inches; brown (10YR 4/3) varved silt loam, very fine sandy loam, silt, and silty clay loam (weighted average silt loam texture); weak thin platy structure; firm; common medium faint grayish brown (10YR 5/2) iron depletions and faint yellowish brown (10YR 5/4) and distinct yellowish brown (10YR 5/6) soft masses of iron accumulation; moderately acid.

Range in Characteristics

Solum thickness: 16 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer and subsoil, moderately acid to slightly alkaline in the substratum

Ap horizon:

Hue—10YR or 2.5Y

Value—2 to 4

Chroma—1 to 3

Content of rock fragments—0 to 2 percent

Bg horizons:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—2

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 2 percent

Bw horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 2 percent

Cg horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 2 percent

C horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 2 percent

Raypol Series*Depth class:* Very deep*Drainage class:* Poorly drained*Permeability:* Moderate in the surface layer and subsoil, and rapid or very rapid in the substratum*Landform:* Depressions and drainageways on outwash plains and terraces*Parent material:* Eolian deposits over glaciofluvial deposits*Slope range:* 0 to 3 percent*Associated soils in a drainage sequence:*

Enfield (well drained, coarse-silty over sandy or sandy-skeletal)

Haven (well drained)

Tisbury (moderately well drained)

Associated similar soil:

Walpole (sandy)

Taxonomic class: Coarse-loamy over sandy or sandy-skeletal, mixed, active, acid, mesic Aeric Endoaquepts**Typical Pedon**

Raypol silt loam, located in the town of Orange, 1,800 feet north of the intersection of the Derby-Milford Road and the Wilbur Cross Parkway overpass, 800 feet east of Derby-Milford Road, on the Ansonia USGS topographic quadrangle, lat. 41 degrees 16 minutes 56 seconds N., long. 73 degrees 02 minutes 56 seconds W., NAD 27, in a grass field:

Ap—0 to 8 inches; very dark brown (10YR 2/2) silt loam, pale brown (10YR 6/3) dry; weak medium granular structure; friable; common very fine, fine, and medium roots; strongly acid; clear smooth boundary.

Bg1—8 to 12 inches; grayish brown (10YR 5/2) very fine sandy loam; weak medium subangular blocky structure; friable; common very fine, fine, and medium roots; common medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; strongly acid; clear wavy boundary.

Bg2—12 to 20 inches; grayish brown (10YR 5/2) silt loam; weak medium subangular blocky structure; friable; common fine and medium roots; common medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; strongly acid; clear wavy boundary.

Bw1—20 to 26 inches; dark yellowish brown (10YR 4/4) silt loam; weak medium subangular blocky structure; friable; few fine roots; common medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation and distinct light brownish gray (10YR 6/2) iron depletions; strongly acid; gradual wavy boundary.

Bw2—26 to 29 inches; olive brown (2.5Y 4/4) very fine sandy loam; massive; friable; common medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation and distinct light brownish gray (10YR 6/2) iron depletions; 5 percent gravel; strongly acid; clear wavy boundary.

2C1—29 to 52 inches; light olive brown (2.5Y 5/4) gravelly sand; single grain; loose; few medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; 25 percent gravel; strongly acid; gradual wavy boundary.

2C2—52 to 65 inches; dark grayish brown (2.5Y 4/2) very gravelly sand; single grain; loose; few medium prominent yellowish brown (10YR 5/6) masses of iron accumulation; 35 percent gravel and 5 percent cobbles; strongly acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid or strongly acid above 40 inches and very strongly acid to slightly acid below 40 inches

Ap horizon:

Hue—7.5YR or 10YR

Value—2 to 4 (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 or 3 (If A horizon is present instead of Ap, chroma is 1 or 2)

Content of rock fragments—0 to 10 percent

Bg horizons:

Hue—5YR to 2.5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam, very fine sandy loam, or loam

Content of rock fragments—0 to 10 percent

Bw horizons:

Hue—5YR to 2.5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—silt loam, very fine sandy loam, or loam

Content of rock fragments—0 to 10 percent

2C horizons:

Hue—5YR to 5Y

Value—4 to 6

Chroma—1 to 4

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—0 to 59 percent

Ridgebury Series Taxadjunct

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, slow or very slow in the substratum

Landform: Depressions and drainageways on hills and drumlins

Parent material: Lodgement till derived from granite, gneiss and schist

Slope range: 0 to 5 percent

Associated soils in a drainage sequence:

Paxton (well drained)

Woodbridge (moderately well drained)

Whitman (very poorly drained)

Associated similar soils:

Leicester (friable substratum)

Mudgepond (calcareous)

Wilbraham (redder)

Taxonomic class: Coarse-loamy, mixed, active, nonacid, mesic Aeric Endoaquepts

Typical Pedon

Ridgebury fine sandy loam, in an area of Ridgebury taxadjunct, Leicester, and Whitman soils, extremely stony, located in the town of Colchester, 2,500 feet south of the intersection of Connecticut Routes 149 and 16, and 100 feet west of Connecticut Route 149, on the Moodus USGS topographic quadrangle, lat. 41 degrees 32 minutes 18 seconds N., long. 72 degrees 24 minutes 49 seconds W., NAD 27, in a wooded area:

Oi—0 to 1 inch; slightly decomposed plant materials

A—1 to 5 inches; black (10YR 2/1) fine sandy loam, dark gray (10 YR 4/1) dry; weak medium granular structure; friable; common fine roots; 5 percent rock fragments; strongly acid; clear wavy boundary.

Bg—5 to 14 inches; gray (10YR 5/1) fine sandy loam; massive; friable; common medium prominent strong brown (7.5YR 5/8) and yellowish brown (10YR 5/8) soft masses of iron accumulation; 5 percent rock fragments; strongly acid; gradual wavy boundary.

Bw—14 to 21 inches; brown (10YR 5/3) fine sandy loam; massive; friable to firm; many medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation and few fine faint grayish brown (10YR 5/2) iron depletions; 10 percent rock fragments; moderately acid; clear wavy boundary.

Cd—21 to 60 inches; grayish brown (10YR 5/2) sandy loam; massive; very firm, brittle; few fine prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; 5 percent rock fragments; moderately acid.

Range in Characteristics

Solum thickness: 20 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—10YR or 2.5Y

Value—2 or 3

Chroma—1 or 2

Content of rock fragments—5 to 14 percent

Bg horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—fine sandy loam, sandy loam or loam

Content of rock fragments—5 to 34 percent

Bw horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—3

Texture of the fine earth fraction—fine sandy loam, sandy loam or loam

Content of rock fragments—5 to 34 percent

Cd horizons:

Hue—10YR to 2.5Y

Value—3 to 6

Chroma—1 to 4

Texture of the fine earth fraction—sandy loam or loam

Content of rock fragments—5 to 34 percent

The Ridgebury soils in this survey area are taxadjuncts because the soil is deeper than is defined as the range for the series. This difference, however, does not significantly affect the use, management, or interpretations of the soils. In this survey, the Ridgebury soils are coarse-loamy, mixed, active, nonacid mesic Aeric Epiaquepts.

Rippowam Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Depressions on flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Occum (well drained)

Pootatuck (moderately well drained)

Associated similar soils:

Lim (finer texture in substratum)

Limerick (coarse-silty)

Taxonomic class: Coarse-loamy, mixed, superactive, nonacid, mesic Fluvaquentic Endoaquepts

Typical Pedon

Rippowam fine sandy loam, located in the town of Redding, 100 feet south of Cross Highway and 100 feet east of the Little River, on the Botsford USGS topographic quadrangle, lat. 41 degrees 18 minutes 32 seconds N., long. 73 degrees 21 minutes 57 seconds W., NAD 27, in a wooded area:

A—0 to 5 inches; very dark grayish brown (10YR 3/2) fine sandy loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; friable; common fine and medium roots; very strongly acid; clear wavy boundary.

Bg1—5 to 12 inches; dark grayish brown (10YR 4/2) fine sandy loam; weak medium subangular blocky structure; friable; common fine and medium roots; common medium prominent strong brown (7.5YR 5/6) soft masses of iron accumulation; very strongly acid; clear wavy boundary.

Bg2—12 to 19 inches; dark gray (10YR 4/1) fine sandy loam; weak medium subangular blocky structure; friable; few fine and medium roots; many medium prominent yellowish red (5YR 4/6) soft masses of iron accumulation; strongly acid; clear wavy boundary.

BCg1—19 to 24 inches; grayish brown (10YR 5/2) sandy loam; massive; friable; few fine and medium roots; common medium prominent strong brown (7.5YR 5/6) soft masses of iron accumulation; strongly acid; clear wavy boundary.

BCg2—24 to 27 inches; very dark gray (10YR 3/1) sandy loam; massive; friable; few fine and medium roots; moderately acid; clear wavy boundary.

Cg1—27 to 31 inches; dark gray (10YR 4/1) loamy sand; single grain; loose; moderately acid; clear wavy boundary.

Cg2—31 to 65 inches; grayish brown (10YR 5/2) very gravelly sand; single grain; loose; 35 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral above 24 inches, moderately acid to neutral below 24 inches (some subhorizon is moderately acid, slightly acid, or neutral within a depth of 40 inches)

A horizon:

Hue—10YR or 2.5Y

Value—2 to 4

Chroma—1 or 2

Content of rock fragments—0 to 14 percent

Bg horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—0 to 14 percent

BCg horizons (where present):

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 to 4

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—0 to 14 percent

C horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 to 4

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—0 to 40 percent

Rumney Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soil in a drainage sequence:

Ondawa (well drained)

Associated similar soils:

Udifluvents (well drained)

Fluvaquents

Medomak Taxadjunct (finer texture)

Taxonomic class: Coarse-loamy, mixed, active, nonacid, frigid Fluvaquentic Endoaquents

Typical Pedon

Rumney silt loam, located in the town of Canaan, 1,600 feet south along Wangum Lake Brook from the outlet of Wangum Lake, and 30 feet east of brook, on the South Canaan USGS topographic quadrangle, lat. 41 degrees, 58 minutes, 12 seconds N., long. 73 degrees 16 minutes 09 seconds W., NAD 27, in a brushy area:

- A—0 to 7 inches; very dark grayish brown (10YR 3/2) silt loam, light brownish gray (10 YR 6/2) dry; moderate medium granular structure; very friable; many very fine and fine roots, few medium roots; common fine distinct dark yellowish brown (10 YR 4/4) masses of iron accumulation; 2 percent gravel; slightly acid; clear smooth boundary.
- Bg1—7 to 22 inches; grayish brown (2.5Y 5/2) sandy loam; weak medium subangular blocky structure; very friable; few very fine and fine roots; common coarse prominent yellowish brown (10 YR 5/6) masses of iron accumulation and few fine faint dark gray (2.5 Y 4/1) iron depletions; 5 percent gravel; neutral; clear smooth boundary.
- Bg2—22 to 38 inches; dark grayish brown (10 YR 4/2) sandy loam; weak medium subangular blocky structure; very friable; common fine prominent dark yellowish brown (10 YR 4/6) masses of iron accumulation; 5 percent gravel; neutral; abrupt smooth boundary.
- Ab—38 to 42 inches; black (10 YR 2/1) sandy loam; weak medium subangular blocky structure; very friable; common fine prominent dark yellowish brown (10 YR 4/6) masses of iron accumulation; 5 percent gravel; neutral; abrupt smooth boundary.
- 2Cg—42 to 44 inches; dark grayish brown (2.5Y 4/2) extremely gravelly coarse sand; single grain; loose; 65 percent gravel; neutral; abrupt smooth boundary.
- 3Cg—44 to 65 inches; dark greenish gray (5 GY 4/1) gravelly fine sandy loam; massive; friable; 20 percent gravel; slightly alkaline.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to neutral (may range to slightly acid below a depth of 40 inches, some subhorizon is moderately acid, slightly acid, or neutral within a depth of 40 inches)

A horizon:

Hue—10YR or 2.5Y

Value—2 to 4 (dry value is 6 or more)

Chroma—1 to 3

Content of rock fragments—0 to 14 percent

B horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 to 4 (at least one subhorizon has hue of 10YR or 2.5Y, value 3 to 5, and chroma 2)

Texture of the fine earth fraction—sandy loam, fine sandy loam, or loam

Content of rock fragments—0 to 14

C horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—1 to 4

Texture of the fine earth fraction—loamy fine sand to coarse sand (some pedons have a loamy layer)

Content of rock fragments—0 to 34 percent (weighted average in the substratum)

Saco Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate in the surface layer and upper substratum and rapid or very rapid in the lower substratum

Landform: Depressions and drainageways on flood plains

Parent material: Alluvium

Slope range: 0 to 2 percent

Associated soils in a drainage sequence:

Hadley (well drained)

Winooski (moderately well drained)

Bash (somewhat poorly drained, coarse-loamy)

Lim (poorly drained, coarse-loamy)

Limerick (poorly drained)

Associated other soils:

Occum (well drained, coarse-loamy)

Pootatuck (moderately well drained, coarse-loamy)

Rippowam (poorly drained, coarse-loamy)

Taxonomic class: Coarse-silty, mixed, active, nonacid, mesic Fluvaquentic Humaquepts

Typical Pedon

Saco silt loam, located in the town of South Windsor, 1200 feet west along Newbury Road from the intersection with Main Street, and 270 feet south of Newbury Road, on the Manchester USGS topographic quadrangle, lat. 41 degrees 49 minutes 49 seconds N., long. 72 degrees 37 minutes 23 seconds W., NAD 27, in a marsh:

A—0 to 12 inches; very dark gray (10YR 3/1) silt loam, gray (10YR 5/1) dry; weak coarse granular structure; very friable, many fine roots; moderately acid; clear wavy boundary.

Cg1—12 to 32 inches; gray (10YR 5/1) silt loam; massive; friable; few fine roots; common medium faint light brownish gray (10YR 6/2) iron depletions and prominent strong brown (7.5YR 5/8) soft masses of iron accumulation; moderately acid; clear wavy boundary.

Cg2—32 to 48 inches; gray (5Y 5/1) silt loam with thin strata of very dark gray (5Y 3/1) silt loam; massive; friable; moderately acid; clear wavy boundary.

2Cg3—48 to 60 inches; gray (10YR 6/1 and 5/1) stratified coarse sand and medium sand; single grain; loose; moderately acid.

Range in Characteristics

Solum thickness: 10 to 15 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral to a depth of about 30 inches, moderately acid to neutral below

A horizon:

Hue—7.5YR to 2.5Y

Value—2 or 3

Chroma—1 to 3

Content of rock fragments—0 to 5 percent

Cg horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—0 to 2
 Texture of the fine earth fraction—silt loam
 Texture below 40 inches—silt loam to fine gravel
 Content of rock fragments—0 to 5 percent

2Cg horizon:

Hue—10YR to 5Y
 Value—3 to 6
 Chroma—0 to 2
 Texture of the fine earth fraction—stratified loamy fine sand to coarse sand
 Content of rock fragments—0 to 40 percent

Scarboro Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderately rapid in the organic layer, moderately rapid or rapid in the surface layer, and rapid or very rapid in the substratum

Landform: Depressions and drainageways on outwash plains and terraces

Parent material: Organic materials over glaciofluvial deposits derived from granite, schist, and gneiss

Slope range: 0 to 2 percent

Associated soils in a drainage sequence:

Windsor (excessively drained)

Deerfield (moderately well drained)

Associated other soils:

Hinckley (excessively drained, sandy-skeletal)

Merrimac (somewhat excessively drained)

Agawam (well drained, coarse-loamy over sandy or sandy-skeletal)

Ninigret (moderately well drained, coarse-loamy over sandy or sandy-skeletal)

Sudbury (moderately well drained)

Walpole (poorly drained)

Taxonomic class: Sandy, mixed, mesic Histic Humaquepts

Typical Pedon

Scarboro muck, located in the town of Wolcott, 2,800 feet northeast along Long Swamp Road from the intersection of Route 69 and Long Swamp Road, 150 feet south of Long Swamp Road, and 100 feet west of Roaring Brook, on the Bristol USGS topographic quadrangle, lat. 41 degrees 37 minutes 59 seconds N., long. 72 degrees 57 minutes 14 seconds W., NAD 27, in a wooded area:

Oa—0 to 12 inches; black (10YR 2/1) muck; weak medium granular structure; very friable; many fine roots; strongly acid; clear wavy boundary.

A—12 to 17 inches; very dark gray (10YR 3/1) loamy sand, dark gray (10 YR 4/1) dry; weak medium granular structure; very friable; few roots; strongly acid; clear wavy boundary.

Cg1—17 to 31 inches; gray (N 6/) fine sand; single grain; loose; strongly acid; gradual wavy boundary.

Cg2—31 to 72 inches; grayish brown (2.5Y 5/2) sand; single grain; loose; few medium prominent yellowish brown (10YR5/6) soft masses of iron accumulation; 5 percent gravel; moderately acid

Range in Characteristics

Solum thickness: 13 to 27 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid in the surface layer and upper substratum, very strongly acid to neutral in the lower substratum

Oa horizon:

Hue—10YR
Value—2 to 4
Chroma—1 or 2

A horizon:

Hue—5YR to 2.5Y
Value—2 or 3
Chroma—0 to 2
Texture of the fine earth fraction—fine sandy loam, sandy loam, loamy sand, loamy fine sand
Content of rock fragments—0 to 10 percent

Cg1 horizon:

Hue—10YR to 5Y
Value—4 to 7
Chroma—0 to 2
Texture of the fine earth fraction—stratified loamy sand to sand
Content of rock fragments—0 to 14 percent

Cg2 horizon:

Hue—10YR to 5Y or N to 5GY
Value—4 to 6
Chroma—0 to 2
Texture of the fine earth fraction—stratified loamy fine sand to coarse sand
Content of rock fragments—0 to 50 percent

Some of the Scarborough soils in this survey area have a mean annual soil temperature which is colder than typical of the series. This map unit (435) is identified as a cold phase of the Scarborough series.

Schroon Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate to moderately rapid in the surface layer and subsoil, and moderately rapid in the substratum

Landform: Hills

Parent material: Melt-out till derived from granite, schist, and gneiss

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Bice (well drained)
Loonmeadow (very poorly drained)

Associated other soils:

Westminster (shallow, somewhat excessively drained)
Millsite (moderately deep, well drained)

Taxonomic class: Coarse-loamy, mixed, superactive, frigid Aquic Dystrudepts

Typical Pedon

Schroon fine sandy loam, 3 to 15 percent slopes, very stony, located in the town of Norfolk, 500 feet south of the intersection of Wheeler Street and Barry Hill Road, 300 feet east of Barry Hill Road, on the South Sandisfield USGS topographic quadrangle,

lat. 42 degrees 01 minutes 54 second N., long. 73 degrees 10 minutes 38 seconds W., NAD 27, in a wooded area:

- Oi—0 to 1 inch; slightly decomposed plant materials derived from leaves
- Oe—1 to 2 inches; moderately decomposed plant materials derived from leaf litter
- Oa—2 to 3 inches; highly decomposed plant materials
- A—3 to 9 inches; very dark grayish brown (10YR 3/2) fine sandy loam, brown (10YR 5/3) dry; weak fine and medium granular structure; very friable; many fine to coarse roots; 3 percent gravel; very strongly acid; clear smooth boundary.
- Bw1—9 to 14 inches; dark brown (7.5YR 3/4) fine sandy loam; weak very fine to medium subangular blocky structure; very friable; many very fine to coarse roots; 3 percent gravel; strongly acid; clear smooth boundary.
- Bw2—14 to 23 inches; brown (10YR 4/4) fine sandy loam; weak very fine to medium subangular blocky structure; very friable; common very fine to medium roots; 3 percent gravel; strongly acid; clear smooth boundary.
- Bw3—23 to 30 inches light olive brown (2.5Y 5/4) sandy loam; weak medium platy structure; friable; few fine roots; few fine prominent strong brown (7.5YR 4/6) soft masses of iron accumulation and common fine distinct light brownish gray (2.5Y 6/2) iron depletions; 3 percent gravel; strongly acid; gradual smooth boundary.
- C—30 to 60 inches pale olive (5Y 6/3) sandy loam; massive; friable; 5 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to moderately acid in the surface layer, very strongly acid to moderately acid in the subsoil, and strongly acid to slightly acid in the substratum

A horizon:

Hue—7.5YR or 10YR

Value—2 or 3

Chroma—1 or 2

Content of rock fragments—3 to 14 percent

Upper Bw horizon:

Hue—5YR to 10YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, sandy loam or loam

Content of rock fragments—3 to 34 percent

Lower Bw horizon:

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, sandy loam or loam

Content of rock fragments—3 to 34 percent

C horizon:

Hue—10YR to 5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—fine sandy loam, sandy loam or coarse sandy loam

Content of rock fragments—3 to 34 percent

Scitico Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate in the surface layer, moderately slow or slow in the upper subsoil, slow or very slow in the lower subsoil, and very slow in the substratum

Landform: Depressions and drainageways on lake plains and terraces

Parent material: Glaciolacustrine deposits

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Brancroft (moderately well drained)

Maybid (very poorly drained)

Associated similar soils:

Raynham (coarse-silty)

Shaker (coarse-loamy over clayey)

Taxonomic class: Fine, illitic, semiactive, nonacid, mesic Typic Endoaquepts

Typical Pedon

Scitico silt loam, in an area of Scitico, Shaker, and Maybid soils, located in the town of East Windsor, 2,000 feet west of the intersection of Newberry Road and Winkler Road, 100 feet north of Newberry Road, on the Broad Brook USGS topographic quadrangle, lat. 41 degrees 55 minutes 20 seconds N., long. 72 degrees 35 minutes 22 seconds W., NAD 27:

- Ap—0 to 8 inches; very dark grayish brown (2.5Y 3/2) silt loam, light gray (5Y 7/1) dry; moderate fine and medium granular structure; friable, sticky, plastic; few very fine, fine, and medium roots; slightly acid; clear smooth boundary.
- Eg—8 to 11 inches; olive gray (5Y 5/2) silt loam; moderate medium blocky structure; friable, sticky, plastic; few very fine and fine roots; common fine distinct light olive brown (2.5Y 5/4), prominent yellowish brown (10YR 5/4), and prominent dark yellowish brown (10YR 4/4) soft masses of iron accumulation; slightly acid; clear smooth boundary.
- Bg1—11 to 18 inches; olive gray (5Y 5/2, 5Y 4/2) silty clay loam; moderate coarse blocky structure; firm, very sticky, plastic; few fine roots between peds; common fine prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/6) soft masses of iron accumulation; continuous distinct gray (5Y 5/1) coatings on ped faces; neutral; clear smooth boundary.
- Bg2—18 to 30 inches; dark gray (5Y 4/1) silty clay loam; moderate coarse prismatic structure parting to coarse blocky structure; firm, very sticky, plastic; few fine roots between peds; many fine prominent dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/6) soft masses of iron accumulation; continuous distinct gray (5Y 5/1) coatings on vertical structure faces; few worm casts along prism faces; neutral; clear smooth boundary.
- Bg3—30 to 38 inches; olive gray (5Y 5/2) and grayish brown (2.5Y 5/2) silty clay; weak coarse prismatic structure; firm, very sticky, plastic; few fine roots between prisms; many fine prominent yellowish brown (10YR 5/6) and dark yellowish brown (10YR 4/4) soft masses of iron accumulation; continuous distinct gray (5Y 5/1) coatings on vertical structure faces; few worm casts along prism faces; neutral; clear smooth boundary.
- Cg1—38 to 52 inches; olive gray (5Y 5/2), dark gray (5Y 4/1), grayish brown (2.5Y 5/2), and light olive brown (2.5Y 5/4) varved silt and clay; silty clay loam weighted average texture; massive parting to weak thin platy structure along varved bedding planes; firm, very sticky, plastic; few fine prominent yellowish brown (10YR 5/6, 5/8) soft masses of iron accumulation; neutral; clear smooth boundary.

Cg2—52 to 65 inches; olive gray (5Y 5/2), gray (5Y 5/1), grayish brown (2.5Y 5/2), and light olive brown (2.5Y 5/4) varved silt and clay; silty clay weighted average texture; weak thin platy structure along varved bedding planes; firm, very sticky, plastic; few prominent dark yellowish brown (10YR 4/4) soft masses of iron accumulation; massive; neutral.

Range in Characteristics

Solum thickness: 20 to 45 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral in the surface layer, strongly acid to neutral in the upper subsoil, and moderately acid to slightly alkaline in the lower subsoil and substratum

Ap horizon:

Hue—10YR to 5Y

Value—3 to 5 (Dry value 6 or more) (If A horizon present instead of Ap, value 2 or 3)

Chroma—1 to 3 (If A horizon present instead of Ap, chroma 1 or 2)

Content of rock fragments—0 to 3 percent

Eg horizon (where present):

Hue—2.5Y or 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam, silty clay loam or silty clay

Content of rock fragments—0 to 3 percent

Upper Bg horizons:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silt loam, silty clay loam, or silty clay

Content of rock fragments—0 to 3 percent

Lower Bg horizons:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—silty clay, clay or silty clay loam

Content of rock fragments—0 to 3 percent

Cg horizons:

Hue—7.5YR to 5Y

Value—3 to 6

Chroma—0 to 2

Texture of the fine earth fraction—weighted average of silty clay, clay, or silty clay loam

Content of rock fragments—0 to 3 percent

Shaker Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderately rapid in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Depressions and drainageways on lake plains and terraces

Parent material: Eolian deposits over glaciolacustrine deposits

Slope range: 0 to 3 percent

Associated soil in a drainage sequence:

Elmridge (moderately well drained)

Associated similar soils:

Raynham (coarse-silty)

Scitico (fine)

Maybid (very poorly drained, fine)

Taxonomic class: Coarse-loamy over clayey, mixed, semiactive, nonacid, mesic
Aeric Epiaquepts

Typical Pedon

Shaker fine sandy loam, in an area of Scitico, Shaker, and Maybid soils, located in the town of Windsor, 2000 feet west of Basswood Road and the Conrail Railroad Crossing, 330 feet north of Basswood Road, on the Windsor Locks USGS topographic quadrangle, lat. 41 degrees 53 minutes 04 seconds N., long. 72 degrees 38 minutes 02 seconds W., NAD 27, in a wooded area:

Oe—0 to 2 inches; black (10YR 2/1) moderately decomposed plant materials

A—2 to 6 inches; very dark brown (10YR 2/2) fine sandy loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; very friable; many fine and medium roots; strongly acid; clear wavy boundary.

Bg—6 to 20 inches; light brownish gray (10YR 6/2) sandy loam weak medium subangular blocky structure; very friable; common fine and medium roots; common medium prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; strongly acid; clear wavy boundary.

Bw—20 to 30 inches; brown (10YR 4/3) sandy loam; weak medium subangular blocky structure; very friable; few fine and medium roots; common medium prominent light gray (2.5Y 7/2) iron depletions and strong brown (7.5YR 5/8) soft masses of iron accumulation; moderately acid; abrupt wavy boundary.

2C—30 to 65 inches; dark yellowish brown (10YR 4/4) varved silt and clay; silty clay weighted average texture; massive structure, parting to weak thin platy structure along varved bedding planes; firm, very sticky, plastic; thin films of very fine sand on plate surfaces; common medium prominent light gray (2.5Y 7/2) iron depletions; slightly acid.

Range in Characteristics

Solum thickness: 18 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer and subsoil, moderately acid to slightly alkaline in the substratum

In some pedons, the Oe horizon is absent.

Oe horizon (where present):

Hue—10YR

Value—2

Chroma—1

Content of rock fragments—none

A horizon:

Hue—7.5YR to 2.5Y

Value—2 to 4 (Dry value 6 or more) (If Ap horizon is present instead of A, value is 3 to 5)

Chroma—1 to 3 (If Ap horizon is present instead of A, chroma is 2 to 4)

Content of rock fragments—0 to 5 percent

Bg horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—1 or 2

Texture of the fine earth fraction—sandy loam, fine sandy loam, or loam

Content of rock fragments—0 to 5 percent

Bw horizon:

Hue—7.5YR to 5Y

Value—4 to 6

Chroma—3 or 4

Texture of the fine earth fraction—sandy loam, fine sandy loam, or loam

Content of rock fragments—0 to 5 percent

2C horizon:

Hue—7.5YR to 5Y

Value—3 to 6

Chroma—0 to 4

Texture of the fine earth fraction—weighted average of silty clay loam, silty clay, or clay

Content of rock fragments—0 to 3 percent

Shelburne Series*Depth class:* Very deep*Drainage class:* Well drained*Permeability:* Moderate or moderately rapid in the surface layer and subsoil, slow or very slow in the substratum*Landform:* Hills and drumlins*Parent material:* Lodgement till derived from granite, schist and gneiss*Slope range:* 3 to 35 percent*Associated soil in a drainage sequence:*

Ashfield (moderately well drained)

Associated other soils:

Westminster (shallow, somewhat excessively drained)

Bice (friable substratum)

Millsite (moderately deep, well drained)

Taxonomic class: Coarse-loamy, mixed, active, frigid Oxyaquic Dystrudepts**Typical Pedon**

Shelburne fine sandy loam, 8 to 15 percent slopes, very stony, located in the town of Norfolk, 700 feet west of Doolittle Lake, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees 00 minutes 57 seconds N., long. 73 degrees 09 minutes 56 seconds W., NAD 27, in a wooded area:

Oi – 0 to 1 inch; slightly decomposed plant materials derived from leaf and twig litter
 A – 1 to 2 inches; black (10YR 2/1) fine sandy loam; dark gray (10YR 4/1) dry; weak medium granular structure; friable; many fine to very coarse roots; 2 percent gravel; strongly acid; abrupt, wavy boundary

Bw1 – 2 to 7 inches; dark yellowish brown (10YR 3/4) fine sandy loam; weak medium subangular blocky structure; friable; common medium and coarse roots; 9 percent gravel; very strongly acid; clear wavy boundary

Bw2 – 7 to 21 inches; olive brown (2.5Y 4/4) gravelly fine sandy loam; moderate medium and coarse subangular blocky structure; friable; common fine to very coarse roots; 12 percent gravel and 5 percent cobbles; very strongly acid; clear wavy boundary.

Bw3 – 21 to 27 inches; olive brown (2.5Y 4/4) bouldery fine sandy loam; moderate medium subangular blocky structure; friable; common fine and medium roots; 12 percent gravel, 5 percent cobbles, and 17 percent boulders; very strongly acid; clear wavy boundary.

Cd1 – 27 to 32 inches; brown (10YR 4/3) gravelly fine sandy loam; moderate coarse and medium subangular blocky structure; firm; common very fine and fine roots; 13 percent gravel and 2 percent cobbles; very strongly acid; clear wavy boundary

Cd2 – 32 to 43 inches; olive brown (2.5Y 4/3) fine sandy loam; massive; very firm; common very fine to medium roots in cracks; few medium distinct light brownish gray (2.5Y 6/2) iron depletions and few medium prominent strong brown (7.5YR 5/8) soft masses of iron accumulation; 8 percent gravel and 2 percent cobbles; strongly acid; clear wavy boundary.

Cd3 – 43 to 55 inches; olive brown (2.5Y 4/4) fine sandy loam; massive; very firm; common very fine to medium roots in cracks; common medium distinct light brownish gray (2.5Y 6/2) iron depletions and common coarse prominent strong brown (7.5YR 4/6) soft masses of iron accumulation; 10 percent gravel; strongly acid; gradual wavy boundary.

Cd4 – 55 to 80 inches; olive brown (2.5Y 4/3) fine sandy loam; massive; very firm; common very fine roots in cracks; common medium distinct gray (2.5Y 5/1) iron depletions and common medium prominent (7.5YR 4/6) masses of iron accumulation; 8 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 20 to 30 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—10YR

Value—3 or 4

Chroma—2 or 3

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—10YR or 2.5Y

Value—3 to 5

Chroma—2 to 4

Texture of the fine earth fraction— fine sandy loam, sandy loam or loam

Content of rock fragments—5 to 34 percent

Cd horizon:

Hue—2.5Y or 5Y

Value—3 to 5

Chroma—2 to 4

Texture of the fine earth fraction— fine sandy loam, sandy loam or loam

Content of rock fragments—5 to 25 percent

Stockbridge Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate in the surface layer and subsoil, moderately slow in the substratum

Landform: Hills

Parent material: Till derived from limestone, schist, and dolomite

Slope range: 3 to 35 percent

Associated soils in a drainage sequence:

Georgia (moderately well drained)

Mudgepond (poorly drained)

Alden (very poorly drained)

Associated similar soils:

Charlton (no carbonates)

Nellis (carbonates within 40 inches)

Associated other soil:

Farmington (shallow to limestone bedrock)

Taxonomic class: Coarse-loamy, mixed, semiactive, mesic Dystric Eutrudepts

Typical Pedon

Stockbridge loam, 3 to 8 percent slopes, located in the town of Salisbury, 1800 feet south of the intersection of Race Track Road and Farnum Road, 400 feet east of Race Track Road, on Sharon USGS topographic quadrangle, lat. 41 degrees 57 minutes 27 seconds N., long. 73 degrees 25 minutes 03 seconds W., NAD 27, in a cultivated field:

Ap—0 to 10 inches, dark brown (10YR 3/3) loam, light brownish gray (2.5Y 6/2) dry; weak coarse granular structure; friable; many fine and very fine roots; 10 percent gravel; moderately acid; clear smooth boundary.

Bw1—10 to 20 inches, olive brown (2.5Y 4/4) loam; weak coarse subangular blocky structure; friable; common fine roots; 10 percent gravel; neutral; clear wavy boundary.

Bw2—20 to 28 inches, light olive brown (2.5Y 5/4) loam; weak coarse subangular blocky structure; firm; few fine roots; few weathered limestone fragments in lower part; 10 percent gravel; neutral; gradual wavy boundary.

C1—28 to 42 inches, olive (5Y 4/3) gravelly loam; weak thick platy structure; firm; few fine roots; many brown (10YR 4/3) weathered limestone fragments and few grayish brown (2.5Y 5/2) streaks; 15 percent gravel and 2 percent cobbles; neutral; gradual wavy boundary.

C2—42 to 48 inches, olive (5Y 4/3) gravelly loam; weak thick platy structure; firm; few brown (10YR 4/3) and light gray (10YR 7/1) streaks from weathered and partially weathered limestone and quartzite fragments; 15 percent gravel and 2 percent cobbles; slightly effervescent; slightly alkaline; gradual wavy boundary.

C3—48 to 65 inches, olive (5Y 4/3) gravelly loam; weak thick platy structure; firm; few brown (10YR 4/3) and light gray (10YR 7/1) streaks from weathered and partially weathered limestone and quartzite fragments; 15 percent gravel and 2 percent cobbles; slightly effervescent; moderately alkaline.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to neutral in the surface layer, moderately acid to neutral to a depth of 40 inches, and moderately acid to moderately alkaline below 40 inches

Ap horizon:

Hue—10YR or 2.5Y

Value—2 to 4 (dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—1 to 3

Content of rock fragments—5 to 14 percent

Bw1 horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6
 Chroma—3 to 6
 Texture of the fine earth fraction—loam or silt loam
 Content of rock fragments—5 to 34 percent

Bw2 horizon:

Hue—10YR to 5Y
 Value—4 to 6
 Chroma—3 to 6
 Texture of the fine earth fraction—loam or silt loam
 Content of rock fragments—5 to 34 percent

C horizon:

Hue—10YR to 5Y
 Value—3 to 6
 Chroma—2 to 4
 Texture of the fine earth fraction—loam or silt loam (may be fine sandy loam below a depth of 40 inches)
 Content of rock fragments—5 to 34 percent above a depth of 40 inches, up to 50 percent below 40 inches

Sudbury Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Glaciofluvial deposits derived from granite, gneiss, and schist

Slope range: 0 to 5 percent

Associated soils in a drainage sequence:

Hinckley (excessively drained, sandy-skeletal)

Merrimac (somewhat excessively drained)

Walpole (poorly drained)

Associated similar soils:

Deerfield (coarser texture in the subsoil)

Ninigret (coarse-loamy over sandy or sandy-skeletal)

Tisbury (coarse-silty over sandy or sandy-skeletal)

Taxonomic class: Sandy, mixed, mesic Aquic Dystrudepts

Typical Pedon

Sudbury sandy loam, 0 to 5 percent slopes, located in the town of East Lyme, 800 feet southeast along Plants Dam Road from the intersection with Boston Post Road, 700 feet south of Plants Dam Road, and 100 feet north of the power transmission lines, on the Old Lyme USGS topographic quadrangle, lat. 41 degrees 20 minutes 30 seconds N., long. 72 degrees 15 minutes 30 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch; moderately decomposed plant materials

A—1 to 5 inches; very dark brown (10YR 2/2) sandy loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; friable; many roots; 5 percent rock fragments; strongly acid; clear wavy boundary.

Bw1—5 to 17 inches; dark yellowish brown (10YR 4/4) gravelly sandy loam; weak medium subangular blocky structure; friable; common fine and medium roots; 15 percent rock fragments; strongly acid; gradual wavy boundary.

Bw2—17 to 25 inches; yellowish brown (10YR 5/6) sandy loam; weak medium

subangular blocky structure; friable; few fine and medium roots; common medium distinct strong brown (7.5YR 5/8) soft masses of iron accumulation and prominent light brownish gray (10YR 6/2) iron depletions; 5 percent rock fragments; strongly acid; clear wavy boundary.

2C—25 to 60 inches; dark yellowish brown (10YR 4/6) gravelly sand; single grain; loose; common medium distinct strong brown (7.5YR 5/8) soft masses of iron accumulation; 15 percent rock fragments; strongly acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid

In some pedons, the Oe horizon is absent.

A or Ap horizon:

Hue—7.5 YR or 10YR

Value—2 to 4

Chroma—1 to 4

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—7.5YR to 2.5Y

Value—3 to 5

Chroma—2 to 8

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—0 to 30 percent

2C horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—2 to 8

Texture —stratified sand and gravel

Content of rock fragments—5 to 75 percent

Some of the Sudbury soils in this survey area have a mean annual soil temperature which is colder than typical of the series. This map unit (423A) is identified as a cold phase of the Sudbury series.

Suncook Series

Depth class: Very deep

Drainage class: Excessively drained

Permeability: Rapid or very rapid

Landform: Flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soils:

Hadley (well drained, coarse-silty)

Occum (well drained, coarse-loamy)

Pootatuck (moderately well drained, coarse-loamy)

Winooski (moderately well drained, coarse-silty)

Rippowam (poorly drained, coarse-loamy)

Lim (poorly drained, coarse-loamy)

Limerick (poorly drained, coarse-silty)

Saco (very poorly drained, coarse-silty)

Taxonomic class: Mixed, mesic Typic Udipsamments

Typical Pedon

Suncook loamy fine sand, located in the town of Granby, 1000 feet east along Mechanicsville Road from the intersection with Connecticut Route 189, and 1200 feet north of Mechanicsville Road and 50 feet east of the East Branch Salmon Brook, on the Tariffville USGS topographic quadrangle, lat. 41 degrees 58 minutes 26 seconds N., long. 72 degrees 48 minutes 12 seconds W., NAD 27, in a wooded area:

- Ap—0 to 7 inches; very dark grayish brown (10YR 3/2) loamy fine sand, pale brown (10 YR 6/3) dry; very weak coarse granular structure; very friable; many fine roots; strongly acid; abrupt smooth boundary.
- C1—7 to 15 inches; dark grayish brown (10YR 4/2) and brown (10YR 5/3) coarse sand; single grain; loose; few fine roots; 2 percent fine gravel; strongly acid; abrupt smooth boundary.
- C2—15 to 22 inches; dark brown (10YR 3/3) loamy fine sand with lenses of coarse sand; single grain; loose; few fine roots; strongly acid; abrupt smooth boundary.
- C3—22 to 32 inches; pale brown (10YR 6/3) medium and coarse sand; single grain; loose; strongly acid; abrupt smooth boundary.
- C4—32 to 42 inches; dark grayish brown (10YR 4/2) fine and medium sand; single grain; loose; strongly acid; abrupt smooth boundary.
- C5—42 to 65 inches; dark grayish brown (10YR 4/2) sand; single grain; loose; 10 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 6 to 10 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid

Ap or A horizon:

Hue—10YR or 2.5Y

Value—3 or 4

Chroma—1 to 3

Content of rock fragments—0 to 10 percent

C horizons:

Hue—7.5YR to 5Y

Value—3 to 6

Chroma—1 to 4

Texture of the fine earth fraction—stratified loamy fine sand to coarse sand

Content of rock fragments—0 to 14 percent (above 40 inches), 0 to 34 percent (below 40 inches)

Sutton Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderately rapid in the surface layer, moderate or moderately rapid in the subsoil, and moderately rapid in the substratum

Landform: Hills

Parent material: Melt-out till derived from granite, gneiss and schist

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Canton (well drained, coarse-loamy over sandy or sandy-skeletal)

Charlton (well drained)
Leicester (poorly drained)

Associated similar soils:

Rainbow (dense substratum)
Wapping (finer texture in the subsoil)
Woodbridge (dense substratum)

Associated other soils:

Hollis (somewhat excessively drained and well drained, shallow)
Ridgebury (poorly drained, dense substratum)
Whitman (very poorly drained, dense substratum)

Taxonomic class: Coarse-loamy, mixed, active, mesic Aquic Dystrudepts

Typical Pedon

Sutton fine sandy loam, 2 to 15 percent slopes, extremely stony, located in the town of Prospect, 400 feet southeast along Merriman Lane from the intersection with Summit Road, and 70 feet north of Merriman Lane, on the Southington USGS topographic quadrangle, lat. 41 degrees 30 minutes 31 seconds N., long. 72 degrees 58 minutes 45 seconds W., NAD 27, in a wooded area:

Oe–0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials.

A–1 to 6 inches; very dark brown (10YR 2/2) fine sandy loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; very friable; common fine and medium roots; 5 percent gravel; strongly acid; clear wavy boundary.

Bw1–6 to 12 inches; brown (7.5YR 4/4) fine sandy loam; weak fine and medium subangular blocky structure; friable; common fine and medium roots; 10 percent gravel and cobbles; moderately acid; gradual wavy boundary.

Bw2–12 to 24 inches; yellowish brown (10YR 5/6) fine sandy loam; weak medium subangular blocky structure; friable; few medium roots; common fine and medium prominent light brownish gray (2.5Y 6/2) iron depletions and faint yellowish red (5YR 5/6) soft masses of iron accumulation; 10 percent gravel and cobbles; moderately acid; gradual wavy boundary.

Bw3–24 to 28 inches; yellowish brown (10YR 5/4) fine sandy loam; weak medium subangular blocky structure; friable; common medium distinct light brownish gray (2.5Y 6/2) iron depletions, reddish brown (5YR 4/4), and strong brown (7.5YR 5/6) soft masses of iron accumulation; 10 percent gravel and cobbles; moderately acid; gradual wavy boundary.

C1–28 to 36 inches; brown (10YR 5/3) gravelly fine sandy loam; weak thick platy structure; firm; common medium faint light brownish gray (2.5Y 6/2) iron depletions and prominent strong brown (7.5YR 5/6) soft masses of iron accumulation; 15 percent gravel and cobbles; moderately acid; gradual wavy boundary.

C2–36 to 65 inches; light olive brown (2.5Y 5/4) gravelly sandy loam; massive; friable; 25 percent gravel and cobbles; moderately acid.

Range in Characteristics

Solum thickness: 20 to 38 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

In some pedons the Oe horizon is absent.

A horizon:

Hue–10YR or 7.5YR

Value–2 to 4 (If Ap horizon is present instead of A, value is 3 or 4)

Chroma—1 to 3 (If Ap horizon is present instead of A, chroma is 2 through 4)

Content of rock fragments—5 to 14 percent

Upper Bw horizons:

Hue—7.5YR or 10YR

Value—4 to 6

Chroma—4 to 6

Texture of the fine earth fraction—fine sandy loam, loam or sandy loam

Content of rock fragments—5 to 34 percent

Lower Bw horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—4 to 6

Texture of the fine earth fraction—fine sandy loam, loam, or sandy loam

Content of rock fragments—5 to 34 percent

C horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam or sandy loam

Content of rock fragments—5 to 34 percent

Taconic Series

Depth class: Shallow

Drainage class: Somewhat excessively drained

Permeability: Moderate or moderately rapid

Landform: Bedrock controlled hills and ridges

Parent material: Melt-out till derived from phyllite and schist

Slope range: 3 to 70 percent

Associated similar soil:

Macomber (moderately deep, well drained)

Associated other soils:

Dummerston (*very deep*, well drained)

Fullam (*very deep*, moderately well drained)

Brayton (*very deep*, poorly drained)

Taxonomic class: Loamy-skeletal, mixed, active, frigid Lithic Dystrudepts

Typical Pedon

Taconic very gravelly loam, in an area of Taconic-Rock Outcrop complex, 45 to 70 percent slopes, located in the town of Salisbury, 500 feet northeast along the Appalachian Trail from the summit of Bear Mountain, on the Bash Bish Falls USGS topographic quadrangle, lat. 42 degrees 2 minutes 45 seconds N., long. 73 degrees 27 minutes 18 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch moderately decomposed plant materials derived from leaves and ferns

A—1 to 4 inches black (10YR 2/1) very gravelly loam, very dark grayish brown (10YR3/2) dry; friable; weak fine and medium granular structure; many very fine to medium roots; 45 percent gravel, 5 percent flagstones; very strongly acid; clear wavy boundary.

Bw—4 to 11 inches dark brown (7.5YR 3/3) very gravelly loam; friable; weak medium to coarse subangular blocky structure; many very fine to very coarse roots; 40 percent gravel, 10 percent stones; very strongly acid; abrupt irregular boundary.

2R—11 inches hard phyllite bedrock

Range in Characteristics

Solum thickness: 10 to 20 inches

Depth to bedrock: 10 to 20 inches

Reaction: Very strongly acid or strongly acid

A horizon:

Hue—7.5YR to 2.5Y

Value—2 to 5

Chroma—1 to 3

Content of rock fragments—35 to 50 percent

Bw horizons:

Hue—7.5YR to 5Y

Value—3 to 5

Chroma—2 to 6

Texture of the fine earth fraction—silt loam or loam

Content of rock fragments—35 to 59 percent

C horizon (where present):

Hue—2.5Y or 10YR

Value—4 or 5

Chroma—2 to 4

Texture of the fine earth fraction—silt loam or loam

Content of rock fragments—45 to 65 percent

Timakwa Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate to very rapid in the organic materials, and rapid or very rapid in the sandy substratum

Landform: Depressions

Parent Material: Organic materials over glaciofluvial deposits

Slope range: 0 to 2 percent

Associated similar soils:

Catden (organic materials greater than 51 inches deep)

Freetown (more acid, organic materials greater than 51 inches deep)

Natchaug (16 to 51 inches of organic materials, over loamy deposits)

Taxonomic class: Sandy or sandy-skeletal, mixed, euic, mesic Terric Haplosaprists

Typical Pedon

Timakwa muck, in an area of Timakwa and Natchaug soils, located in the town of Thompson, 750 feet NE on New Road from the intersection of Town Farm Road and New Road, 125 feet NW of New Road, on the Thompson USGS topographic quadrangle, lat. 41 degrees 59 minutes 57 seconds N., long. 71 degrees 48 minutes 30 seconds W., NAD 27, in a swamp:

Oa1—0 to 10 inches; black (10YR 2/1) broken face and rubbed muck; 20 percent fibers, 5 percent rubbed; massive; very friable, nonsticky and nonplastic; strongly acid; clear wavy boundary.

Oa2—10 to 21 inches; black (10YR 2/1) broken face and rubbed muck; 15 percent fibers, 2 percent rubbed; massive; very friable, nonsticky and nonplastic; 5 percent woody fragments; strongly acid; clear wavy boundary.

Oa3—21 to 24 inches; black (7.5YR 2.5/1) broken face and rubbed muck; 5 percent fibers, 0 percent rubbed; massive; very friable, nonsticky and nonplastic; strongly acid; clear wavy boundary.

- Oa4—24 to 37 inches; black (5YR 2.5/1) broken face and rubbed muck; 40 percent fibers, 10 percent rubbed; massive; very friable, nonsticky and nonplastic; 10 percent woody fragments; strongly acid; abrupt wavy boundary.
- 2Cg1—37 to 47 inches; dark gray (5Y 4/1) very gravelly loamy coarse sand; single grain; loose, nonsticky and nonplastic; 40 percent gravel; moderately acid; clear wavy boundary.
- 2Cg2—47 to 60 inches; gray (5Y 5/1) gravelly loamy very fine sand; massive; friable, nonsticky and nonplastic; 20 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: Organic layers 16 to 51 inches deep

Depth to bedrock: More than 80 inches

Reaction: Ultra acid to moderately acid (in 0.01M calcium chloride) in the organic layers and strongly acid to neutral in the sandy substratum

Woody fragments in organic soil material: 2 to 10 percent

Upper Oa horizons:

Hue—10YR to 5YR, or is neutral

Value—2 to 4

Chroma—0 to 6

Lower Oa horizons:

Hue—10YR to 5YR, or is neutral

Value—2 or 3

Chroma—0 to 4

2Cg horizons:

Hue—7.5YR to 5Y, or is neutral

Value—3 to 6

Chroma—0 to 8

Texture of the fine earth fraction—loamy very fine sand, loamy fine sand, loamy coarse sand, loamy sand, sand, or coarse sand

Content of rock fragments—0 to 40 percent

Tisbury Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Outwash plains and terraces

Parent material: Eolian deposits over glaciofluvial deposits derived from granite, gneiss, and schist

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Enfield (well drained)

Haven (well drained, coarse-loamy over sandy or sandy-skeletal)

Raypol (poorly drained, coarse-loamy over sandy or sandy-skeletal)

Associated similar soils:

Deerfield (sandy)

Ellington (redder, coarse-loamy over sandy or sandy-skeletal)

Ninigret (coarse-loamy over sandy or sandy-skeletal)

Sudbury (sandy)

Taxonomic class: Coarse-silty over sandy or sandy-skeletal, mixed, active, mesic Aquic Dystrudepts

Typical Pedon

Tisbury silt loam, in an area of Ninigret and Tisbury soils, 0 to 5 percent slopes, located in the town of Ledyard, 250 feet west along Route 214 from the intersection with Spicer Hill Road, 600 feet south of Route 214, and 50 feet west of Lee Brook, on the Uncasville USGS topographic quadrangle, lat. 41 degrees 26 minutes 15 seconds N., long. 72 degrees 00 minutes 06 seconds W., NAD 27, in a grassy field:

- Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) silt loam, light brownish gray (10 YR 6/2) dry; weak coarse granular structure; friable; many very fine and fine roots; few scattered pebbles; strongly acid; abrupt smooth boundary.
- Bw1—8 to 18 inches; yellowish brown (10YR 5/6) silt loam; weak medium and coarse subangular blocky structure; very friable; common very fine and fine roots; few scattered pebbles; strongly acid; clear wavy boundary.
- Bw2—18 to 26 inches; brownish yellow (10YR 6/6) silt loam; massive; very friable; few fine roots; common medium prominent grayish brown (2.5Y 5/2) iron depletions and faint strong brown (7.5YR 5/6) soft masses of iron accumulation; few scattered pebbles; strongly acid; clear wavy boundary.
- 2C—26 to 60 inches; grayish brown (10YR 5/2) very gravelly sand; single grain; loose; common medium prominent strong brown (7.5YR 5/6) soft masses of iron accumulation and common medium faint light brownish gray (10YR 6/2) iron depletions; 60 percent gravel; strongly acid.

Range in Characteristics

Solum thickness: 17 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—7.5YR or 10YR

Value—2 to 4 (dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 through 3)

Content of rock fragments—0 to 5 percent

Upper Bw horizon:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 5 percent

Lower Bw horizon:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—0 to 5 percent

2C horizon:

Hue—2.5YR to 2.5Y

Value—4 to 6

Chroma—0 to 6

Texture of the fine earth fraction— stratified sand and loamy sand

Content of rock fragments—0 to 59 percent

Udifuluents

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderately rapid in the surface layer and moderate to very rapid in the substratum

Landform: Flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soil in a drainage sequence:

Fluvaquents (somewhat poorly drained to very poorly drained)

Associated similar soils:

Occum (well drained, subsoil development)

Pootatuck (moderately well drained, subsoil development)

Rippowam (poorly drained)

Saco (very poorly drained)

Taxonomic class: Udifuluents

Sample Pedon

Udifuluents, in an area of Fluvaquents-Udifuluents complex, frequently flooded, located in the town of Cornwall, 1800 feet southwest of the intersection of Dawn Hill Road and River Road, 100 feet west of the Housatonic River, on the Ellsworth USGS topographic quadrangle, lat. 41 degrees 48 minutes 12 seconds N, long. 73 degrees 23 minutes 48 seconds W, NAD27, in an old field:

A—0 to 2 inches; very dark grayish brown (10YR 3/2) fine sandy loam, gray (5Y 5/1) dry; weak fine granular structure; very friable; many fine to medium roots; slightly alkaline; abrupt smooth boundary.

C—2 to 4 inches; light yellowish brown (2.5Y 6/3) loamy fine sand; massive; very friable; many fine to medium roots; slightly alkaline; abrupt smooth boundary.

Ap—4 to 12 inches; very dark grayish brown (10YR 3/2) fine sandy loam; weak coarse subangular blocky structure; very friable; many fine to medium roots; slightly alkaline; clear smooth boundary.

AC—12 to 18 inches; brown (10YR 4/3) fine sandy loam; weak medium subangular blocky structure; very friable; few fine to medium roots; 2 percent gravel; slightly alkaline; clear smooth boundary.

C1—18 to 35 inches; olive brown (2.5Y 4/4) loamy sand; massive; very friable; few fine to medium roots; 2 percent gravel; slightly alkaline; clear smooth boundary.

C2—35 to 38 inches; olive brown (2.5Y 4/3) very gravelly loamy sand; single grain; loose; 45 percent gravel; slightly alkaline; clear smooth boundary.

C3—38 to 60 inches; light yellowish brown (2.5Y 6/3) very gravelly coarse sand; single grain; loose; 55 percent gravel; slightly alkaline.

Range in Characteristics

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral

Content of rock fragments: 0 to 59 percent

A horizons:

Hue—5YR to 2.5Y

Value—3 to 5

Chroma—1 to 4

Texture of the fine earth fraction—loamy sand, loamy fine sand, loamy very fine sand, very fine sandy loam, fine sandy loam, sandy loam, loam, or silt loam

C horizons:

Hue—5YR to 2.5Y

Value—3 to 7

Chroma—2 to 6

Texture of the fine earth fraction—coarse sand, sand, fine sand, very fine sand, loamy sand, loamy fine sand, loamy very fine sand, very fine sandy loam, fine sandy loam, sandy loam, loam, or silt loam

Udipsamments

Depth class: Very deep*Drainage class:* Excessively drained*Permeability:* Rapid or very rapid*Landform:* Coastal dunes*Parent material:* Beach sands*Slope range:* 0 to 8 percent*Associated soils:*

Merrimac (somewhat excessively drained, more soil profile development)

Hinckley (sandy-skeletal, more soil profile development)

Windsor (more soil profile development)

Ipswich (very poorly drained organic soil)

Pawcatuck (very poorly drained organic materials over sandy or sandy skeletal)

Westbrook (very poorly drained organic materials over loamy)

Taxonomic class: Udipsamments

Sample Pedon

Udipsamments, in an area of Beaches-Udipsamments complex, coastal, located in the town of Madison in Hammonasset State Park , 1500 feet west of the traffic circle and 3000 feet west of Webster Point, on the Clinton USGS topographic quadrangle, lat. 41 degrees 15 minutes 43 seconds N., long. 72 degrees 33 minutes 39 seconds W., NAD 27, on Hammonasset Beach:

C1—0 to 38 inches; 85 percent light brownish gray (2.5Y 6/2), 10 percent yellow (2.5Y 7/6) and 5 percent black (10 YR 2/1) sand; single grain; loose; few fine and medium roots; moderately acid; clear smooth boundary.

C2—38 to 50 inches; 85 percent light brownish gray (2.5Y 6/2), 10 percent yellow (2.5Y 7/6) and 5 percent black (10 YR 2/1) coarse sand; single grain; loose; slightly acid; abrupt smooth boundary.

C3—50 to 65 inches; 50 percent light olive brown (2.5Y 5/4) and 50 percent olive yellow (2.5Y 6/6) sand; single grain; loose; neutral.

Range in Characteristics

Depth to bedrock: More than 80 inches*Reaction:* Moderately acid to neutral*Content of rock fragments:* 0 to 34 percent

C horizons:

Hue—10YR to 5Y

Value—2 to 7

Chroma—1 to 6

Texture of the fine earth fraction— fine sand, sand or coarse sand

Udorthents

Depth class: Very deep

Drainage class: Moderately well drained and well drained

Permeability: Very slow to very rapid

Landform: Variable

Parent material: Glaciofluvial deposits, till, or glaciolacustrine deposits

Slope range: 0 to 70 percent

Associated similar soil:

Udipsamments (excessively drained, sandy)

Taxonomic class: Udorthents

Sample Pedon

Udorthents, in an area of Udorthents-Urban land complex, located in the town of Hartford, 4400 feet north along Mark Twain Drive from the intersection with Connecticut Route 44, and 50 feet west of Mark Twain Drive, on the Hartford North USGS topographic quadrangle, lat. 41 degrees 47 minutes 32 seconds N., long. 72 degrees 42 minutes 35 seconds W., NAD 27, in a wooded area:

- A—0 to 5 inches; very dark grayish brown (10YR 3/2) loam; moderate medium to coarse subangular blocky structure parting to moderate medium granular structure; firm; common fine to very fine and few medium to coarse roots; 8 percent gravel; 2 percent concrete, asphalt, and brick fragments; neutral; clear wavy boundary.
- C1—5 to 21 inches; brown (10YR 4/3) gravelly loam; massive; firm; common very fine to fine roots; 10 percent gravel; 10 percent concrete, asphalt, shale, and brick fragments; neutral; clear wavy boundary.
- C2—21 to 80 inches; dark brown (10YR 3/3) very gravelly sandy loam; massive; firm; 15 percent gravel; 25 percent concrete, asphalt, and brick fragments; slightly alkaline.

Range in Characteristics

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly alkaline

Content of rock fragments: 0 to 65 percent

A horizon:

Hue—5YR to 5Y

Value—2 to 4

Chroma—1 to 4

Texture of the fine earth fraction—loamy coarse sand, loamy sand, loamy fine sand, loamy very fine sand, sandy loam, fine sandy loam, very fine sandy loam, loam, silt loam, or silty clay loam

C horizons:

Hue—5YR to 5Y

Value—3 to 7

Chroma—1 to 6

Texture of the fine earth fraction—coarse sand, sand, fine sand, very fine sand, loamy coarse sand, loamy sand, loamy fine sand, loamy very fine sand, very fine sandy loam, fine sandy loam, sandy loam, loam, silt loam, or silty clay loam

Walpole Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderately rapid in the surface layer and subsoil, and rapid or very rapid in the substratum

Landform: Depressions and drainageways on outwash plains and terraces

Parent material: Glaciofluvial deposits derived from granite, gneiss, and schist

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Hinckley (excessively drained, sandy-skeletal)

Merrimac (somewhat excessively drained)

Sudbury (moderately well drained)

Associated similar soils:

Raypol (coarse-loamy over sandy or sandy-skeletal)

Scarboro (very poorly drained)

Taxonomic class: Sandy, mixed, mesic Aeric Endoaquepts

Typical Pedon

Walpole sandy loam, located in the town of Killingly, 400 feet north along North Shore Drive from the intersection with Route 101, 500 feet east of North Shore Drive, and 700 feet south of the outlet of Alexander Lake, on the Danielson USGS topographic quadrangle, lat. 41 degrees 50 minutes 58 seconds N., long. 71 degrees 54 minutes 28 seconds W., NAD 27 in a wooded area:

Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials

A—1 to 7 inches; very dark brown (10YR 2/2) sandy loam, grayish brown (10 YR 5/2) dry; weak medium granular structure; very friable; many fine and medium roots; 8 percent gravel; very strongly acid; clear smooth boundary.

Bg—7 to 21 inches; dark grayish brown (2.5Y 4/2) sandy loam; massive; friable; common fine and medium roots in the upper part of the horizon and few fine roots in the lower part; common medium prominent strong brown (7.5YR 5/6) and yellowish brown (10YR 5/6) , and distinct yellowish brown (10YR 5/4) soft masses of iron accumulation and light brownish gray (10YR 6/2) iron depletions; 10 percent gravel; strongly acid; gradual smooth boundary.

Bw—21 to 25 inches; light olive brown (2.5Y 5/4) gravelly sandy loam; massive; friable; common medium distinct yellowish brown (10YR 5/6) soft masses of iron accumulation, light brownish gray (10YR 6/2) and dark grayish brown (2.5Y 4/2) iron depletions; 20 percent gravel; strongly acid; clear smooth boundary.

C1—25 to 41 inches; light yellowish brown (2.5Y 6/4) very gravelly loamy sand; single grain; loose; common medium prominent strong brown (7.5YR 5/6) and faint yellowish brown (10YR 5/4) soft masses of iron accumulation; 30 percent gravel and 5 percent cobbles; strongly acid; gradual smooth boundary.

C2—41 to 65 inches; light brownish gray (10YR 6/2) very gravelly sand; few brown (10YR 5/3) streaks; single grain; loose; 35 percent gravel and 5 percent cobbles, moderately acid.

Range in Characteristics

Solum thickness: 18 to 28 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to neutral

In some pedons the Oe horizon is absent.

A horizon:

Hue–10YR

Value–2 or 3 (If Ap horizon present instead of A, value 2 to 4)

Chroma–1 or 2 (If Ap horizon present instead of A, chroma 1 to 3)

Content of rock fragments–0 to 14 percent

Bg horizon:

Hue–10YR to 5Y

Value–4 to 6

Chroma–1 or 2

Texture of the fine earth fraction–sandy loam or fine sandy loam

Content of rock fragments–0 to 25 percent

Bw horizon:

Hue–10YR to 5Y

Value–4 to 6

Chroma–3 or 4

Texture of the fine earth fraction–sandy loam or fine sandy loam

Content of rock fragments–0 to 25 percent

C horizons:

Hue–10YR to 5Y

Value–4 to 6

Chroma–1 to 4

Texture of the fine earth fraction–stratified loamy fine sand to coarse sand

Content of rock fragments–0 to 50 percent

Wapping Series

Depth class: Very deep*Drainage class:* Moderately well drained*Permeability:* Moderate in the surface layer and subsoil, and moderately rapid in the upper substratum, and moderately rapid or rapid in the lower substratum*Landform:* Hills and till plains*Parent material:* Eolian deposits over melt-out till derived from sandstone, shale, basalt, and gneiss*Slope range:* 0 to 8 percent*Associated soil in a drainage sequence:*

Narragansett (well drained, coarse-loamy over sandy or sandy-skeletal)

Associated similar soils:

Ludlow (dense substratum, redder)

Rainbow (dense substratum)

Taxonomic class: Coarse-loamy, mixed, active, mesic Aquic Dystrudepts

Typical Pedon

Wapping very fine sandy loam, 0 to 3 percent slopes, located in the town of South Windsor, 4,400 feet north along Barber Hill Road from the intersection with Miller Road and 600 feet east of Barber Hill Road, on the Manchester USGS Topographic map, lat. 41 degrees 51 minutes 39 seconds N., long. 72 degrees 31 minutes 45 seconds W., NAD27, in a corn field:

Ap–0 to 11 inches; dark brown (7.5YR 3/3) very fine sandy loam, pale brown (10YR 6/3) dry; weak medium granular structure; very friable; common very fine and fine roots; 5 percent gravel; moderately acid; abrupt smooth boundary.

- Bw1—11 to 16 inches; brown (7.5YR 4/4) very fine sandy loam; moderate medium subangular blocky structure; very friable; few fine roots; 5 percent gravel; moderately acid; clear smooth boundary.
- Bw2—16 to 20 inches; yellowish brown (10YR 5/4) very fine sandy loam; moderate coarse subangular blocky structure; friable; few fine roots; common fine to coarse distinct light brownish gray (10 YR 6/2) iron depletions, and common fine distinct strong brown (7.5YR 5/6) and reddish brown (5 YR 4/4) soft masses of iron accumulation; 5 percent gravel; moderately acid; clear smooth boundary.
- 2C1—20 to 28 inches; reddish brown (5YR 4/4) gravelly sandy loam; massive; friable; 20 percent gravel; moderately acid; clear smooth boundary.
- 2C2—28 to 36 inches; reddish brown (2.5YR 4/4) gravelly loamy sand; massive; friable; 30 percent gravel; moderately acid. clear smooth boundary.
- 2C3—36 to 80 inches; reddish brown (2.5YR 4/4) gravelly loamy sand; massive; friable; 20 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: 20 to 38 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—7.5YR or 10YR

Value—3 or 4 (dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—0 to 14 percent

Bw horizons:

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—4 to 6

Texture of the fine earth fraction—very fine sandy loam, silt loam, or loam

Content of rock fragments—0 to 14 percent

2C1 horizon:

Hue—2.5YR to 5Y

Value—2 to 6

Chroma—2 to 6

Texture of the fine earth fraction— sandy loam or fine sandy loam

Content of rock fragments—15 to 34 percent

2C2 and 2C3 horizons:

Hue—2.5YR to 5Y

Value—2 to 6

Chroma—2 to 6

Texture of the fine earth fraction— sandy loam, fine sandy loam, or loamy sand

Content of rock fragments—15 to 50 percent

Watchaug Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, and moderate or moderately rapid in the substratum

Landform: Till plains and hills

Parent material: Melt-out till derived from sandstone, shale and basalt

Slope range: 0 to 8 percent

Associated soil in a drainage sequence:

Cheshire (well drained)

Associated similar soils:

Ludlow (dense substratum)

Rainbow (dense substratum, browner)

Wapping (yellowish in the subsoil)

Taxonomic class: Coarse-loamy, mixed, semiactive, mesic Aquic Dystrudepts

Typical Pedon

Watchaug fine sandy loam, 0 to 3 percent slopes, located in the town of Wallingford, 400 feet north of the intersection of Cook Hill Road and Schoolhouse Road, on the Wallingford USGS topographic quadrangle, lat. 41 degrees 27 minutes 22 seconds N., long. 72 degrees 51 minutes 54 seconds W., NAD 27, in a wooded area:

Ap—0 to 8 inches; dark reddish brown (5YR 3/3) fine sandy loam, light reddish brown (5YR 6/3) dry; weak medium and fine granular structure; friable; common fine and medium roots; 8 percent gravel; strongly acid; clear wavy boundary.

Bw1—8 to 18 inches; reddish brown (5YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; common fine and medium roots; 10 percent gravel; strongly acid; gradual wavy boundary.

Bw2—18 to 24 inches; yellowish red (5YR 5/6) fine sandy loam; weak medium subangular blocky structure; very friable; few fine and medium roots; common fine and medium faint strong brown (7.5YR 5/6) soft masses of iron accumulation and prominent pinkish gray (5YR 6/2) iron depletions; 10 percent gravel; strongly acid; gradual wavy boundary.

C—24 to 65 inches; reddish brown (5YR 4/3) gravelly sandy loam; massive; friable; few fine roots above 48 inches; streaks of pale red (2.5YR 6/2) and reddish brown (2.5YR 5/4); 25 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid in the surface layer and subsoil, and very strongly acid to slightly acid in the substratum

Ap horizon:

Hue—5YR to 10YR

Value—2 to 4 (dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 to 3)

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—2.5YR or 5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, silt loam, or loam

Content of rock fragments—5 to 34 percent

C horizon:

Hue—2.5YR or 5YR

Value—3 to 6

Chroma—3 to 6

Texture of the fine earth fraction—sandy loam or fine sandy loam

Content of rock fragments—5 to 34 percent

Westbrook Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate to rapid in the organic layers, and moderate to very slow in the substratum

Landform: Tidal marshes and salt marshes

Parent material: Grassy organic materials over loamy deposits

Slope range: 0 to 2 percent

Associated similar soils:

Ipswich (organic materials deeper than 51 inches)

Pawcatuck (organic materials over sandy deposits)

Taxonomic class: Loamy, mixed, euic, mesic Terric Sulfihemists

Typical Pedon

Westbrook mucky peat, located in the town of Westbrook, 250 feet south of US Route 1 and 500 feet west of the south end of Hammock Road and Hammock Dock, on the Essex USGS topographic quadrangle, lat. 41 degrees 16 minutes 30 seconds N., long. 72 degrees 28 minutes 18 seconds N., NAD 27, in a salt grass tidal marsh:

Oe1—0 to 10 inches; very dark gray (10YR 3/1) mucky peat; dark gray (10YR 4/1) dry; 65 percent fiber, 30 percent rubbed; dense mat of roots, stems and leaves; massive; many very fine, fine, and medium roots; fibers (herbaceous); thin lenses and coatings of silt; 45 percent organic matter; strongly saline; slightly acid; clear wavy boundary.

Oe2—10 to 40 inches; very dark gray (10YR 3/1) mucky peat, dark gray (10YR 4/1) dry; 50 percent fiber, 25 percent rubbed; massive; few very fine, fine, and medium roots; fibers (herbaceous); thin lenses and coatings of silt; 44 percent organic matter; strongly saline; moderately acid; gradual wavy boundary.

Oe3—40 to 48 inches; dark olive gray (5Y 3/2) mucky peat, dark gray (10YR 4/1) dry; 35 percent fiber, 25 percent rubbed; massive; fibers (herbaceous); 24 percent organic matter; strongly saline; neutral; clear wavy boundary.

Cg1—48 to 64 inches; very dark gray (5Y 3/1) silt loam, dark gray (10YR 4/1) dry; massive; 12 percent organic matter; strongly saline; neutral; diffuse wavy boundary.

Cg2—64 to 99 inches; dark gray (N 4/) silt loam, dark gray (10YR 4/1) dry; massive; 10 percent organic matter; few small shell fragments; strongly saline; slightly acid.

Range in Characteristics

Thickness of organic materials: 16 to 51 inches

Depth to bedrock: More than 80 inches

Reaction: Strongly acid to slightly alkaline

Salinity: Very slightly saline to strongly saline

Surface tier:

Hue—neutral or 5YR to 5Y

Value—2 to 5

Chroma—0 to 3

Content of rock fragments—none

Subsurface tier:

Hue—neutral or 5YR to 5Y

Value—2 to 5

Chroma—0 to 3

Texture of the organic fraction—mucky peat

Content of rock fragments—none

Bottom tier:

Hue—neutral or 5YR to 5Y
 Value—2 to 5
 Chroma—0 to 3
 Texture of the organic fraction—mucky peat
 Content of rock fragments—none

Cg horizons:

Hue—neutral or 10YR to 5BG
 Value—2 to 7
 Chroma—0 to 2
 Texture of the fine earth fraction—silt loam, silty clay loam or sandy loam
 Content of rock fragments—none

Westminster Series

Depth class: Shallow

Drainage class: Somewhat excessively drained

Permeability: Moderately rapid

Landform: Bedrock controlled hills and ridges

Parent material: Melt-out till derived from schist, gneiss, or granite

Slope range: 3 to 70 percent

Associated similar soil:

Millsite (moderately deep, well drained)

Associated other soils:

Ashfield (very deep, moderately well drained, dense substratum)

Bice (very deep, well drained)

Schroon (very deep, moderately well drained)

Loonmeadow (very deep, very poorly drained)

Taxonomic class: Loamy, mixed, active, frigid Lithic Dystrudepts

Typical Pedon

Westminster fine sandy loam, in an area of Westminster-Millsite-Rock Outcrop complex, 3 to 15 percent slopes, located in the town of Norfolk, 2000 feet southeast along Loon Meadow Drive from the intersection with Lovers Lane and 1000 feet south of Loon Meadow Drive, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees 00 minutes 31 seconds N., long. 73 degrees 10 minutes 51 seconds W., NAD 27, in a wooded area:

Oi—0 to 1 inch; slightly decomposed plant materials derived from leaf litter

Oe—1 to 2 inches; moderately decomposed plant materials derived from leaf litter

A—2 to 5 inches; dark brown (7.5YR 3/3) fine sandy loam, light yellowish brown (10 YR 6/4) dry; weak fine to medium granular structure; very friable; many very fine to coarse roots; 4 percent gravel, 3 percent stones, 3 percent cobbles; very strongly acid; abrupt wavy boundary.

Bw1—5 to 12 inches; dark yellowish brown (10Y 3/4) fine sandy loam; weak fine to coarse subangular blocky structure; very friable; common very fine to coarse roots; 5 percent gravel, 5 percent cobbles, 2 percent stones; very strongly acid; clear wavy boundary.

Bw2—12 to 16 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak fine to coarse subangular blocky structure; very friable; few very fine to medium roots; 5 percent gravel, 5 percent stones; strongly acid; irregular boundary.

2R—16 inches; green schist bedrock with bands of quartz

Range in Characteristics

Solum thickness: 10 to 20 inches

Depth to bedrock: 10 to 20 inches

Reaction: Extremely acid to strongly acid

A horizon:

Hue—5YR to 10YR

Value—2 or 3

Chroma—0 to 3

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—7.5YR or 10YR

Value—3 or 4

Chroma—2 to 4

Texture of the fine earth fraction—fine sandy loam or loam

Content of rock fragments—5 to 25 percent

2R horizon:

Schist bedrock, but in places is gneiss or granite

Wethersfield Series

Depth class: Very deep

Drainage class: Well drained

Permeability: Moderate in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Hills and drumlins

Parent material: Lodgement till derived from sandstone, shale or basalt

Slope range: 3 to 35 percent

Associated soils in a drainage sequence:

Ludlow (moderately well drained)

Wilbraham (poorly drained)

Menlo (very poorly drained)

Associated similar soils:

Broadbrook (brownier)

Cheshire (friable substratum)

Narragansett (coarse-silty over sandy or sandy-skeletal, brownier)

Yalesville (moderately deep over bedrock)

Taxonomic class: Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts

Typical Pedon

Wethersfield loam, 3 to 8 percent slopes, very stony, located in the town of Middlefield, 2000 feet northeast along Route 66 from the intersection with Route 147, 50 feet south of Route 66 at the south end of Mt. Higbee Reservoir, on the Middletown USGS topographic quadrangle, lat. 41 degrees 32 minutes 17 seconds N., long. 72 degrees 43 minutes 32 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials

A—1 to 3 inches; dark brown (7.5YR 3/2) loam, brown (7.5 YR 5/2); moderate medium granular structure; friable; many fine and medium roots; 10 percent gravel; strongly acid; clear wavy boundary.

Bw1—3 to 13 inches; reddish brown (5YR 4/4) loam; weak medium subangular blocky structure; friable; common fine and medium roots; 10 percent gravel; strongly acid; clear wavy boundary.

Bw2—13 to 27 inches; dark reddish brown (5YR 3/3) gravelly loam; weak medium subangular blocky structure; friable; few medium roots; 15 percent gravel and cobbles; strongly acid; clear wavy boundary.

Cd—27 to 65 inches; reddish brown (2.5YR 4/4) gravelly loam; weak thick platy structure; very firm, brittle; few silt films and black coatings on some plates; 20 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid to a depth of 60 inches, and strongly acid to slightly alkaline below 60 inches.

In some pedons, the Oe horizon may be absent.

Oe horizon (where present):

Hue—10YR

Value—2

Chroma—1

Content of rock fragments—none

A horizon:

Hue—5YR to 10YR

Value—2 or 3 (If Ap horizon is present instead of A, value is 3 or 4)

Chroma—1 to 3 (If Ap horizon is present instead of A, chroma is 2 to 4)

Content of rock fragments—5 to 14 percent

Bw horizons:

Hue—2.5YR or 5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—loam or fine sandy loam

Content of rock fragments—5 to 25 percent

Cd horizon:

Hue—10R to 5YR

Value—3 to 5

Chroma—2 to 6

Texture of the fine earth fraction— loam or fine sandy loam

Content of rock fragments—10 to 34 percent

Whitman Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Depressions and drainageways on hills and drumlins

Parent material: Lodgement till derived from granite, gneiss and schist

Slope range: 0 to 2 percent

Associated soils in a drainage sequence:

Montauk (well drained)

Paxton (well drained)

Woodbridge (moderately well drained)

Ridgebury taxadjunct (poorly drained)

Associated similar soil:

Leicester (poorly drained, friable substratum)

Taxonomic class: Loamy, mixed, active, nonacid, mesic shallow Typic Humaquepts

Typical Pedon

Whitman fine sandy loam, in an area of Ridgebury, Leicester, and Whitman soils, extremely stony, located in the town of Lebanon, 300 feet southwest along Browning Road from the intersection with Madley Road, and 200 feet west of Browning Road, on; the Fitchville USGS topographic quadrangle, lat. 41 degrees 37 minutes 15 seconds N., long. 72 degrees 12 minutes 39 seconds W., NAD 27, in a wooded area:

Oi–0 to 1 inch; slightly decomposed plant materials

A–1 to 9 inches; black (10YR 2/1) fine sandy loam, dark gray (10 YR 4/1) dry; weak medium granular structure; friable; common fine and medium roots; strongly acid; abrupt wavy boundary.

Bg–9 to 16 inches; dark grayish brown (10YR 4/2) fine sandy loam; weak medium subangular blocky structure; friable; few fine roots; few fine prominent yellowish brown (10YR 5/8) soft masses of iron accumulation; 5 percent rock fragments; moderately acid; clear wavy boundary.

Cdg1–16 to 22 inches; grayish brown (2.5Y 5/2) fine sandy loam; moderate medium platy structure; very firm, brittle; common medium prominent strong brown (7.5YR 5/8) soft masses of iron accumulation and few medium faint light brownish gray (2.5Y 6/2) iron depletions; 5 percent rock fragments; slightly acid; gradual wavy boundary.

Cdg2–22 to 60 inches; grayish brown (2.5Y 5/2) fine sandy loam; massive; firm, brittle; common medium prominent strong brown (7.5YR 5/8) soft masses of iron accumulation and few medium faint light brownish gray (2.5Y 6/2) iron depletions; 5 percent rock fragments; slightly acid.

Range in Characteristics

Solum thickness: 12 to 20 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to slightly acid (some horizon within a depth of 40 inches is moderately acid or slightly acid)

In some pedons, the Oi horizon may be absent.

A horizon:

Hue–10YR

Value–2 or 3

Chroma–1 or 2

Content of rock fragments–5 to 14 percent

Bg horizon:

Hue–neutral or 10YR to 5Y

Value–4 to 6

Chroma–1 or 2

Texture of the fine earth fraction–fine sandy loam or loam

Content of rock fragments–5 to 34 percent

Cdg horizons:

Hue–neutral or 10YR to 5Y

Value–4 to 6

Chroma–1 or 2 (chroma 3 below 30 inches)

Texture of the fine earth fraction–loam, fine sandy loam, or sandy loam

Content of rock fragments–5 to 34 percent

Wilbraham Series

Depth class: Very deep

Drainage class: Poorly drained

Permeability: Moderate in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Depressions and drainageways on hills and drumlins

Parent material: Lodgement till derived from sandstone, shale and/or basalt

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Wethersfield (well drained)

Ludlow (moderately well drained)

Menlo (very poorly drained)

Associated other soils:

Broadbrook (well drained)

Cheshire (well drained, friable substratum)

Rainbow (moderately well drained)

Watchaug (moderately well drained, friable substratum)

Taxonomic class: Coarse-loamy, mixed, active, mesic Aquic Dystrudepts

Typical Pedon

Wilbraham silt loam, in an area of Wilbraham and Menlo soils, extremely stony, located in the town of Middlefield, 1300 feet northeast along Laurel Brook Road from the intersection with Cherry Hill Road and 700 feet north of Laurel Brook Road, on the Middletown USGS topographic quadrangle, lat. 41 degrees 30 minutes 40 seconds N., long. 72 degrees 41 minutes 35 seconds W., NAD 27, in an unimproved pasture:

A—0 to 4 inches; very dark gray (10YR 3/1) silt loam, gray (10 YR 5/1) dry; weak medium granular structure; very friable; many fine roots; 5 percent gravel; strongly acid; abrupt wavy boundary.

Bw1—4 to 8 inches; dark reddish brown (5YR 3/3) silt loam; weak coarse subangular blocky structure; very friable; few fine roots; common medium prominent pinkish gray (7.5YR 6/2) iron depletions; 10 percent gravel; strongly acid; clear wavy boundary.

Bw2—8 to 20 inches; reddish brown (5YR 4/4) silt loam; weak coarse subangular blocky structure; friable; few fine roots; common distinct reddish gray (5YR 5/2) iron depletions; 13 percent gravel and cobbles; strongly acid; clear wavy boundary.

Cd—20 to 65 inches; dark reddish brown (5YR 3/3) gravelly loam; weak thick platy structure; very firm, brittle; silt films and black (10YR 2/1) coatings on some plates; many distinct brown (7.5YR 5/2) and faint dark brown (7.5YR 4/4) soft masses of iron accumulation; 25 percent gravel and cobbles; strongly acid.

Range in Characteristics

Solum thickness: 20 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

A horizon:

Hue—5YR to 10YR

Value—2 or 3 (If Ap horizon is present instead of A, value is 2 to 4)

Chroma—1 or 2 (If Ap horizon is present instead of A, chroma is 2 or 3)

Content of rock fragments—5 to 14 percent

Bw1 horizon:

Hue—5YR to 10YR

Value—3 to 6

Chroma—3 or 4

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—5 to 25 percent

Bw2 horizon:

Hue—2.5YR or 5YR

Value—3 to 5

Chroma—3 to 6

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—5 to 25 percent

Cd horizon:

Hue—10R to 5YR

Value—3 to 6

Chroma—2 to 6

Texture of the fine earth fraction—silt loam, loam, or very fine sandy loam

Content of rock fragments—10 to 34 percent

Windsor Series*Depth class:* Very deep*Drainage class:* Excessively drained*Permeability:* Rapid in the surface layer and upper subsoil, rapid or very rapid in the lower subsoil and substratum*Landform:* Outwash plains, terraces, and kames*Parent material:* Eolian deposits over glaciofluvial deposits derived from granite, gneiss, and schist*Slope range:* 0 to 15 percent*Associated soils in a drainage sequence:*

Deerfield (moderately well drained)

Scarboro (very poorly drained)

Associated similar soils:

Hinckley (sandy-skeletal)

Merrimac (somewhat excessively drained)

Taxonomic class: Mixed, mesic Typic Udipsamments**Typical Pedon**

Windsor loamy sand, 0 to 3 percent slopes, located in the town of South Windsor, 0.35 miles east of Chapel Road and U.S. Route 5, 50 feet south of Chapel Road, on the Manchester USGS topographic quadrangle, lat. 41 degrees 48 minutes 35 seconds N., long. 72 degrees 36 minutes 24 seconds W., NAD 27, in a wooded area:

Oe—0 to 1 inch; black (10YR 2/1) moderately decomposed plant materials; many very fine and fine roots; very strongly acid; abrupt smooth boundary.

A—1 to 3 inches; very dark grayish brown (10YR 3/2) loamy sand, pale brown (10 YR 6/3) dry; weak medium granular structure; very friable; many very fine and fine roots; strongly acid; abrupt wavy boundary.

Bw1—3 to 9 inches; strong brown (7.5YR 5/6) loamy sand; very weak fine granular structure; very friable; many fine and medium roots; strongly acid; gradual wavy boundary.

Bw2—9 to 21 inches; yellowish brown (10YR 5/6) loamy sand; very weak fine granular structure; very friable; common fine and medium roots; strongly acid; gradual wavy boundary.

Bw3—21 to 25 inches; light yellowish brown (10YR 6/4) sand; single grain; loose; few coarse roots; strongly acid; clear wavy boundary.

C—25 to 65 inches; pale brown (10YR 6/3) and light brownish gray (10YR 6/2) sand; single grain; loose; few coarse roots; strongly acid.

Range in Characteristics

Solum thickness: 18 to 36 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid in the surface layer and subsoil, and very strongly acid to slightly acid in the substratum

In some pedons, the Oe horizon may be absent.

A horizon:

Hue—7.5YR or 10YR

Value—2 or 3 (If Ap horizon present instead of A, value 3 or 4)

Chroma—1 to 3 (If Ap horizon present instead of A, chroma 2 to 4)

Content of rock fragments—0 to 10 percent

Upper Bw horizons:

Hue—7.5YR to 2.5Y

Value—4 to 6

Chroma—4 to 8

Texture of the fine earth fraction—loamy sand or loamy fine sand

Content of rock fragments—0 to 10 percent

Lower Bw horizons:

Hue—7.5YR to 5Y

Value—4 to 7

Chroma—3 to 6

Texture of the fine earth fraction—loamy sand, fine sand or sand

Content of rock fragments—0 to 10 percent

C horizon:

Hue—5YR to 5Y

Value—4 to 7

Chroma—1 to 6

Texture of the fine earth fraction— loamy sand, fine sand or sand

Content of rock fragments—0 to 14 percent

Winooski Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, and moderate or moderately rapid in the substratum

Landform: Flood plains

Parent material: Alluvium

Slope range: 0 to 3 percent

Associated soils in a drainage sequence:

Hadley (well drained)

Bash (somewhat poorly drained, coarse-loamy)

Lim (coarse-loamy, poorly drained)

Limerick (poorly drained)
Saco (very poorly drained)

Associated similar soil:

Pootatuck (coarse-loamy)

Taxonomic class: Coarse-silty, mixed, active, nonacid, mesic Fluvaquentic
Dystrudepts

Typical Pedon

Winooski silt loam, located in the town of South Windsor, 2,300 feet west along Newbury Road from the intersection with Main Street, and 100 feet south of Newbury Road, on the Hartford North USGS topographic quadrangle, lat. 41 degrees 49 minutes 54 seconds N., long. 72 degrees 37 minutes 34 seconds W., NAD 27, in a cropped field:

- Ap—0 to 12 inches; very dark grayish brown (2.5Y 3/2) silt loam; light brownish gray (2.5Y 6/2) dry; weak medium granular structure; very friable; few very fine and fine roots; moderately acid; clear smooth boundary.
- Bw1—12 to 18 inches; dark grayish brown (2.5Y 4/2) silt loam; massive friable; few fine roots; moderately acid; clear smooth boundary.
- Bw2—18 to 36 inches; dark grayish brown (2.5Y 4/2) silt loam; massive; friable; common medium distinct olive brown (2.5Y 4/4) and dark yellowish brown (10YR 4/4) soft masses of iron accumulation and distinct light olive gray (5Y 6/2) iron depletions; slightly acid; clear smooth boundary.
- C1—36 to 52 inches; olive gray (5Y 4/2) very fine sandy loam; massive; very friable; common medium prominent yellowish brown (10YR 5/6) and dark yellowish brown (10YR 4/4) soft masses of iron accumulation, and faint gray (5Y 5/1) iron depletions; slightly acid; clear smooth boundary.
- C2—52 to 65 inches; olive gray (5Y 4/2) silt loam; massive; friable; common medium prominent yellowish brown (10YR 5/6) and dark yellowish brown (10YR 4/4) soft masses of iron accumulation and faint gray (5Y 5/1) iron depletions; slightly acid.

Range in Characteristics

Solum thickness: 18 to 43 inches

Depth to bedrock: More than 80 inches

Reaction: Moderately acid to slightly acid in the surface layer and subsoil, moderately acid to neutral in the substratum

Ap or A horizon:

Hue—10YR to 5Y

Value—3 or 4

Chroma—2 or 3

Content of rock fragments—less than 1 percent

Bw horizons:

Hue—7.5YR to 5Y

Value—2 to 5

Chroma—2 to 4

Texture of the fine earth fraction—silt loam or very fine sandy loam

Content of rock fragments—less than 1 percent

C horizons:

Hue—10YR to 5Y

Value—3 to 6

Chroma—2 to 4

Texture of the fine earth fraction—silt loam, very fine sandy loam or loamy very fine sand

Content of rock fragments—less than 1 percent

Wonsqueak Series

Depth class: Very deep

Drainage class: Very poorly drained

Permeability: Moderately slow to moderately rapid in the organic material and moderate or moderately slow in the mineral substratum

Landform: Depressions

Parent material: Organic materials over drift

Slope range: 0 to 2 percent

Associated similar soil:

Bucksport (organic materials greater than 51 inches thick)

Associated other soils:

Brayton (poorly drained mineral soil, till from phyllite, schist, slate, and shale)

Loonmeadow (very poorly drained mineral soil, till from gneiss, schist, and dolomite)

Taxonomic class: Loamy, mixed, euic, frigid Terric Haplosaprists

Typical Pedon

Wonsqueak mucky peat, located in the town of Norfolk, 500 feet south along Barry Hill Road from the intersection with Wheeler Street, 50 feet west of Barry Hill Road, on the South Sandisfield USGS topographic quadrangle, lat. 42 degrees 01 minutes 44 seconds N., long. 73 degrees 10 minutes 58 seconds W., NAD 27, in a wooded swamp:

Oe—0 to 2 inches; moderately decomposed black (7.5YR 2.5/1) mucky peat; about 60 percent fiber, 35 percent rubbed; massive; many fine to very coarse roots; about 10 percent woody and 90 percent herbaceous fibers; moderately acid; clear wavy boundary.

Oa1—2 to 11 inches; highly decomposed very dark brown (7.5YR 2.5/2) muck; about 45 percent fiber, 15 percent rubbed; massive; many fine to very coarse roots; about 10 percent woody and 90 percent herbaceous fibers; moderately acid; clear smooth boundary.

Oa2—11 to 22 inches; highly decomposed black (10YR 2/1) muck; about 5 percent fiber, 2 percent rubbed; massive; few fine to medium roots; about 40 percent woody and 60 percent herbaceous fibers; moderately acid; clear broken boundary.

2Cg1—22 to 25 inches; black (N 2.5/1) mucky silt loam; massive; friable; slightly acid clear smooth boundary;

2Cg2—25 to 45 inches; dark gray (5Y 4/1) gravelly fine sandy loam; massive; friable; 15 percent gravel, 5 percent cobbles and 5 percent stones; slightly acid; clear smooth boundary

2Cg3—45 to 60 inches; dark grayish brown (2.5Y 4/2) fine sandy loam with lenses of loamy fine sand; massive; friable; 10 percent gravel, 2 percent stones, 2 percent cobbles; slightly acid.

Range in Characteristics

Thickness of organic layers: 16 to 51 inches

Depth to bedrock: More than 80 inches

Reaction: Extremely acid to slightly acid in the surface tier, very strongly acid to slightly acid in the subsurface and bottom tiers, and very strongly acid to neutral in the substratum

Woody fragments in organic layers: 0 to 20 percent, 3/4 to 1 inch in diameter

Surface tier:

Hue—neutral or 2.5YR to 10YR

Value—2 or 3
 Chroma—0 to 2

Subsurface and bottom tiers:
 Hue—neutral or 2.5YR to 2.5Y
 Value—2 to 4
 Chroma—0 to 2

2C horizons:
 Hue—neutral or 5YR to 5GY
 Value—3 to 6
 Chroma—0 to 4
 Texture—fine sandy loam, silt loam, or loam
 Rock fragments—0 to 20 percent, mostly gravel

Woodbridge Series

Depth class: Very deep

Drainage class: Moderately well drained

Permeability: Moderate in the surface layer and subsoil, and slow or very slow in the substratum

Landform: Hills and drumlins

Parent material: Lodgement till derived from granite, gneiss and schist

Slope range: 0 to 15 percent

Associated soils in a drainage sequence:

Paxton (well drained)
 Ridgebury taxadjunct (poorly drained)
 Whitman (very poorly drained)

Associated similar soil:

Sutton (friable substratum)

Taxonomic class: Coarse-loamy, mixed, active, mesic Aquic Dystrudepts

Typical Pedon

Woodbridge fine sandy loam, 3 to 8 percent slopes, located in the town of Mansfield, 0.75 miles south of the intersection of Connecticut Routes 275 and 195 along Rte. 195, and 0.25 miles east on the University of Connecticut Agronomy Farm, 800 feet north of the greenhouses, on the Spring Hill USGS topographic quadrangle, lat. 41 degrees 47 minutes 53 seconds N., long. 72 degrees 13 minutes 48 seconds W., NAD 27, in a field:

Ap—0 to 7 inches; very dark grayish brown (10YR 3/2) fine sandy loam, light brownish gray (10YR 6/2) dry; moderate medium granular structure; friable; many fine and medium roots; few very dark brown (10YR 2/2) earthworm casts; 5 percent gravel; moderately acid; abrupt wavy boundary.

Bw1—7 to 18 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; common fine roots; few very dark brown (10YR 2/2) earthworm casts; 10 percent gravel; moderately acid; gradual wavy boundary.

Bw2—18 to 26 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; common fine roots; few very dark brown (10YR 2/2) earthworm casts; few medium distinct strong brown (7.5YR 5/6) soft masses of iron accumulation and distinct light brownish gray (10YR 6/2) iron depletions; 10 percent gravel; moderately acid; gradual wavy boundary.

Bw3—26 to 30 inches; light olive brown (2.5Y 5/4) fine sandy loam; weak medium subangular blocky structures; friable; few fine roots; common medium prominent

strong brown (7.5YR 5/6) soft masses of iron accumulation and distinct light brownish gray (10YR 6/2) iron depletions; 10 percent gravel; moderately acid; clear wavy boundary.

Cd1—30 to 43 inches; light olive brown (2.5Y 5/4) gravelly fine sandy loam; weak thick platy structure; very firm, brittle; many medium prominent strong brown (7.5YR 5/8) soft masses of iron accumulation and distinct light brownish gray (10YR 6/2) iron depletions; 20 percent gravel; moderately acid; gradual wavy boundary.

Cd2—43 to 65 inches; light olive brown (2.5Y 5/4) gravelly fine sandy loam; weak thick platy structure; very firm, brittle; few fine distinct very dark brown (10YR 2/2) coatings on plates; common fine prominent brown (7.5YR 5/8) soft masses of iron accumulation; 25 percent gravel; moderately acid.

Range in Characteristics

Solum thickness: 20 to 40 inches

Depth to bedrock: More than 80 inches

Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—10YR

Value—3 or 4 (Dry value 6 or more) (If A horizon is present instead of Ap, value is 2 or 3)

Chroma—2 to 4 (If A horizon is present instead of Ap, chroma is 1 or 2)

Content of rock fragments—5 to 14 percent

Upper Bw horizons:

Hue—7.5YR to 2.5Y

Value—3 to 6

Chroma—3 to 8

Texture of the fine earth fraction—fine sandy loam, loam, or sandy loam

Content of rock fragments—5 to 34 percent

Lower Bw horizons:

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture of the fine earth fraction—fine sandy loam, loam, or sandy loam

Content of rock fragments—5 to 34 percent

Cd horizons:

Hue—10YR to 5Y

Value—4 to 6

Chroma—1 to 4

Texture of the fine earth fraction— fine sandy loam, sandy loam, or loam

Content of rock fragments—5 to 34 percent

Yalesville Series

Depth class: Moderately deep

Drainage class: Well drained

Permeability: Moderate or moderately rapid in the surface layer and subsoil, moderately rapid in the substratum

Landform: Bedrock-controlled hills and ridges

Parent material: Melt-out till derived from red sandstone, shale, and/or basalt

Slope range: 3 to 15 percent

Associated soils:

Cheshire (very deep, well drained)

Holyoke (shallow, well drained)
 Wethersfield (very deep, dense substratum)
 Ludlow (very deep, moderately well drained)
 Watchaug (very deep, moderately well drained)
 Wilbraham (very deep, poorly drained)

Taxonomic class: Coarse-loamy, mixed, active, mesic Typic Dystrudepts

Typical Pedon

Yalesville fine sandy loam, 3 to 8 percent slopes, located in the town of North Branford, 2000 feet north of the junction of Village Street and Clintonville Road, 250 feet west of Village Street, on the Wallingford USGS topographic quadrangle, lat. 41 degrees 23 minutes 57 seconds N., long. 72 degrees 48 minutes 26 seconds W., NAD 27, in a hay field:

- Ap—0 to 8 inches; dark brown (7.5YR 3/2) fine sandy loam, pinkish gray (7.5YR 6/2) dry; weak medium granular structure; friable; common very fine, fine, and medium roots; 5 percent gravel; moderately acid; abrupt wavy boundary.
- Bw1—8 to 14 inches; reddish brown (5YR 4/4) fine sandy loam; weak medium subangular blocky structure; friable; common fine and medium roots; few very dark grayish brown earthworm casts; 5 percent gravel; moderately acid; gradual wavy boundary.
- Bw2—14 to 25 inches; reddish brown (5YR 4/4) loam; weak medium subangular blocky structure; friable; few fine roots; 5 percent gravel; moderately acid; gradual wavy boundary.
- C—25 to 36 inches; reddish brown (2.5YR 4/4) gravelly sandy loam; massive; firm; 12 percent gravel and 3 percent cobbles; moderately acid; abrupt wavy boundary.
- 2R—36 inches; reddish brown (2.5YR 4/4) hard sandstone bedrock

Range in Characteristics

Solum thickness: 18 to 36 inches
Depth to bedrock: 20 to 40 inches
Reaction: Very strongly acid to moderately acid

Ap horizon:

Hue—5YR to 10YR
 Value—3 or 4 (Dry value 6 or more)
 Chroma—2 or 3
 Content of rock fragments—2 to 14 percent

Bw horizons:

Hue—10R, 2.5YR, or 5YR
 Value—3 to 5
 Chroma—4 to 6
 Texture of the fine earth fraction—fine sandy loam, sandy loam, or loam
 Content of rock fragments—2 to 34 percent

C horizon:

Hue—10R, 2.5YR, or 5YR
 Value—3 to 5
 Chroma—3 to 6
 Texture of the fine earth fraction—sandy loam, fine sandy loam, or loam
 Content of rock fragments—5 to 50 percent

2R horizon:

Reddish brown sandstone, shale, or basalt

Formation of the Soils

This first part of this section describes the factors of soil formation as they relate to the soils of Connecticut. The second part defines the processes of soil horizon formation.

Factors of Soil Formation

Soils are products of weathering and other physical and chemical processes that act on parent material. The properties of a soil at a given point on the earth depend on the combination of the following factors: the physical and chemical composition of the parent material, the topography, the climate, the plant and animal life, and time (Jenny, 1941). The relative influence of each of these factors differs from place to place, and each modifies the effect of the others. For example, the impact of climate over a given area is tempered by topography or parent material. The variability of Connecticut soils are primarily attributed to parent material, topography, and time. [Table 30](#) shows the relationship between the soil series in Connecticut and their parent material, position on the landscape, and drainage.

Parent Material

Parent material is the unconsolidated earthy material in which soils are formed. It influences the physical, chemical, and mineralogical composition of the soils. It also influences the rate at which soil forming processes will proceed.

Most of the soils in Connecticut formed in deposits left as a result of glaciation. Glacial till is the most extensive type of parent material. Less extensive parent materials are glaciofluvial deposits (outwash), glaciolacustrine deposits, alluvial deposits, and organic deposits.

Soils formed in glacial till have a wide range of characteristics as a result of the heterogeneous nature of the till, its rock and soil particles. Some soils such as Paxton, Broadbrook, Brayton, Shelburne, and Wethersfield, which are formed in very deep glacial till deposits, have a dense substratum. Other soils such as Charlton, Stockbridge, Schroon, and Dummerston, which also formed in very deep glacial till, do not have a dense layer. Narragansett is an example of a soil with a silty windblown layer above the glacial till. In some places, the glacial till is moderately deep or shallow over bedrock. Hollis, Farmington, and Taconic are examples of soils that are shallow over bedrock. Chatfield, Yalesville, and Millsite are soils that are moderately deep over bedrock. Some areas have bedrock exposed at the surface. Rock outcrop is mapped in these areas.

As the glacial ice melted, large quantities of meltwater transported and sorted soil and rock debris. This material is referred to as glacial outwash and was redeposited in layers of sand and gravel on outwash plains and terraces. Hinckley, Manchester, and Copake are examples of soils formed in these materials. A layer of silty windblown material is above the sand and gravel in Agawam and Tisbury soils, for example.

At one time, much of the Connecticut River Valley and some smaller valleys in other areas of the state contained glacial lakes where glacial meltwater was

impounded. Most of the stone-free sediment deposited in the still lake waters was silty or clayey. Belgrade, Brancroft, Scitico, and Berlin are examples of soils that formed in these fine textured deposits.

In more recent times, overflowing streams have deposited fresh, dark alluvial material on flood plains. This material tends to be variable in texture. Soils formed in this material show weak soil profile development. Suncook, Hadley, Pootatuck, Rumney, and Saco are examples of alluvial soils.

Soils formed in organic deposits are mainly in closed depressions throughout the state and in tidal marshes along Long Island Sound and associated estuaries of major rivers of the state. Catden, Bucksport, and Westbrook soils, for example, formed in well decomposed remains of trees or other plants.

Topography

Topography refers to the shape of the land surface, and also (for soils) to the effect of the shape of the land surface on the movement of water in soil and across the land. Components of topography include: slope configuration (e.g. convex, concave, flat), slope aspect (direction downhill slope faces, e.g. southwest), slope steepness, slope position (e.g., bottom, lower slope, mid-slope, shoulder, ridge top), and elevation (where elevation affects soil properties through its inter-relationship to climate). The components of topography may be viewed at multiple scales; for example, a large, generally concave slope may have a convex bulge. The appropriate scale for considering topography depends upon the soil forming process being considered.

Topography has a great influence on the formation of the soils. Soils that formed in convex positions, where little or no runoff accumulates, are generally well drained and do not contain iron depletions in the subsoil. Examples of soils in this category are Stockbridge and Charlton. In level or slightly depressional areas, the water table is usually closer to the surface for extended periods. This results in gray iron depletions close to the surface and often, accumulation of sediment at the surface.

Some soils are wet because they occupy a position where the water table is at or near the surface for long periods. This wetness is evident by a thick, dark surface layer and strong redoximorphic features or gray subsoil. Leicester and Scarboro soils are examples of soils with high water tables.

Climate

Climate, in particular temperature and precipitation, is one of the most influential of the soil forming factors. It determines to a large degree the kind of weathering processes that occur. It also affects the growth and kind of vegetation and the leaching and translocation of weathered materials.

Connecticut has a humid, temperate climate that promotes the development of moderately weathered, leached soils. The difference in elevation in the state, which ranges from sea level along Long Island Sound to over 2,000 feet in some areas of northern and northwestern Connecticut, results in two temperature regimes, mesic and frigid. Mesic soils have a mean annual soil temperature greater than 47 degrees F and frigid soils have a mean annual soil temperature less than 47 degrees F. Westminster, Ondawa, Ashfield, and Loonmeadow are examples of frigid soils. Cooler temperatures tend to slow down the weathering processes and shorten the growing season. More detailed and specific data on the climate of Connecticut are in the climate section under "General Nature of the Survey Area".

Plant and Animal Life

All living organisms, including plants, animals, bacteria, and fungi, influence soil formation. Vegetation is generally responsible for the amount of organic matter and nutrients in the soil and for the color and structure of the surface layer. Earthworms and burrowing animals help to keep the soil porous and more permeable for air and water. Their waste products cause aggregations of soil particles which improves soil structure. Bacteria and fungi decompose vegetation, which results in the release of nutrients.

As the last glacier began retreating northward, a long process of revegetation began in Connecticut. Initially the vegetation was tundra; by the time European settlement began, the forest was primarily an oak-chestnut cover type, with Northern Hardwoods found in the northern part of the state, especially at the higher elevations.

Because the rooting depth is shallow in many of the upland soils, trees are susceptible to windthrow, which has caused much mixing of the soil materials.

Human activity, through clearing trees, cultivating the land, artificial drainage, grading, and the introduction of new plants, has also influenced changes that occur in the soils. This has added nutrients by fertilization, has mixed some soil horizons by plowing or land development activities, and has accelerated erosion in many areas.

Time

The degree of profile development not only reflects the age of a soil, but it also reflects the influence of other factors. In geologic terms, the deposits in which Connecticut soils formed are relatively young, having being deposited when the last glacier receded about 10,000 to 15,000 years ago. The soils have not all reached the same stage of soil profile development because the other soil forming factors also influence the rate of soil profile development.

An immature soil is one that has not had enough time to develop distinct horizons. Lim soils and Fluvaquents, formed in recent alluvium, are examples. They are regularly flooded and more sediment is deposited. The time for soil development is constantly interrupted and thin or irregular soil profiles develop.

Processes of Soil Formation

This section contains a brief explanation of soil horizon nomenclature and a discussion of the processes involved in soil horizon development as they relate to soil formation.

The soil forming factors cause the formation of different layers, or soil horizons. These soil horizons can be viewed in a vertical cut of the soil, known as a soil profile. The soil profile extends from the surface downward into material that is little altered by the soil forming processes. Most soils contain three major horizons, called A, B, and C horizons.

Several processes cause the formation of soil horizons. They include the accumulation of organic matter, the leaching of soluble salts and minerals, the translocation of clay minerals, the reduction and transfer of iron, and the formation of dense and compact layers in the subsoil (Simonson, 1959).

The accumulation of organic matter takes place as plant residue decomposes. This process darkens the surface layer and helps to form the A horizon. It takes a long time to replace this organic matter once it has been lost. The organic matter content of the surface layers of soils in Connecticut averages about 4 percent.

For soils to develop distinct subsoil, some of the lime and other soluble salts must be leached before other soil processes such as translocation of clay minerals can

take place. Factors that affect leaching include the kinds of salts originally present, the rate and depth of percolation, and the texture of the soil.

One of the more important processes of soil horizon development in some of the soils is the translocation of silicate clay minerals. The amount of clay minerals in a soil is inherent in the parent material, but clay content varies from one soil horizon to another. Clay particles are transported (eluviation) downward from the A horizon and redeposited (illuviation) in the B horizon as clay films on ped faces, as linings along pores and root channels, and as coatings on some rock fragments. In some soils, an E horizon has formed by considerable eluviation of clay minerals to the B horizon. The Scitico soil is an example of a soil where the clay content is higher in the B horizon than in the A horizon because of translocation.

The reduction and transfer of iron compounds occur mainly in the wetter, more poorly drained soils. This process is known as gleying. In poorly drained and very poorly drained soils, such as Whitman, the grayish subsoil indicates the reduction of iron. In moderately well drained soils, such as Woodbridge, yellowish brown and reddish brown mottles indicate the segregation of iron compounds. A bright colored soil indicates a well drained soil where no reduction and transfer of iron have taken place. Canton soils are an example.

References

- Adams, F. 1984. Soil acidity and liming. American Society of Agronomy, Agronomy Monograph 12, 2nd edition.
- Allan, P.F., L.E. Garland, and R. Dugan. 1963. Rating northeastern soils for their suitability for wildlife habitat. In Transactions of the twenty-eighth North American wildlife and natural resources conference, pp. 247-261.
- Alerich, C.L. In press. Forest statistics for Connecticut—1985 and 1998. U. S. Department of Agriculture, Forest Service, Northeastern Research Station.
- American Association of State Highway and Transportation Officials (AASHTO). 1998. Standard specifications for transportation materials and methods of sampling and testing. 19th edition, 2 volumes.
- American Society for Testing and Materials (ASTM). 1998. Standard classification of soils for engineering purposes. ASTM Standard D 2487.
- Appelquist, M.B. 1959. Soil-site studies of southern hardwoods. In Southern forest soils—Eighth annual forestry symposium, pp. 49-63.
- Beck, Donald E. 1962. Yellow-poplar site index curves. U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station Research Note 180.
- Bell, Michael. 1985. The face of Connecticut: people, geology, and the land. Bulletin 110, State Geological and Natural History Survey of Connecticut, Connecticut Department of Environmental Protection.
- Birkeland, Peter W. 1974. Pedology, weathering, and geomorphological research.
- Birkeland, Peter W. 1984. Soils and geomorphology. 2nd edition.
- Black, C.A. 1968. Soil-plant relationships. 2nd edition.
- Broadfoot, Walter M. 1960. Field guide for evaluating cottonwood sites. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station Occasional Paper 178.
- Buol, S.W., F.D. Hole, and R.J. McCracken. 1980. Soil genesis and classification. 3rd edition.
- Carmean, Willard H. 1967. Soil survey refinements for predicting black oak site quality in southeastern Ohio. Soil Science Society of America Proceedings 31: 805-810.

- Coleman, Steven M. 1981. Rock-weathering rates as functions of time. *Quaternary Research* 15: 250-264.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Doolittle, James A. 1982. Characterizing soil map units with the ground-penetrating radar. *Soil Survey Horizons* 23(4): 3-10.
- Doolittle, James A. 1983. Investigating Histosols with the ground penetrating radar. *Soil Survey Horizons* 24(3): 23-28.
- Dowhan, Joseph J. and Robert J. Craig. 1976. Rare and endangered species of Connecticut and their habitat. Report of Investigations No. 6. State Geological and Natural History Survey of Connecticut, Connecticut Department of Environmental Protection.
- Eyre, F.H., editor. 1980. Forest cover types of the United States and Canada. Society of American Forestry.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. February 24, 1995. Hydric soils of the United States.
- Hurt, G.W., P.M. Whited, and R.F. Pringle, editors. 1996. Field indicators of hydric soils in the United States.
- Jenny, Hans. 1941. Factors of soil formation.
- Jenny, Hans. 1980. The soil resource—Origin and behavior. *Ecological Studies* 37.
- Johnson, R.W., R. Glaccum, and R. Wojtasinski. 1980. Application of ground penetrating radar to soil survey. *Soil and Crop Science Society of Florida Proceedings* 39: 68-72. (Reprinted in *Soil Survey Horizons* 23(3): 17-25)
- Keys, J.E. et al. 1995. Ecological units of the eastern United States. U.S. Department of Agriculture, Forest Service.
- Kilpatrick, Howard J. et al. 2001. Connecticut deer program summary 2001. Connecticut Department of Environmental Protection.
- Khasawneh, F.E., E.C. Sample, and E.J. Kamprath, editors. 1980. The role of phosphorus in agriculture. American Society of Agronomy.
- Michigan State University, Departments of Crop and Soil Sciences and Horticulture. 1985. Fertilizer recommendations for vegetables and field crops in Michigan. Extension Bulletin E-550.
- Mokma, D.L. 1978. Soil management units and land use planning. Michigan State University, Extension Bulletin E-1262.
- Munson, Robert D., editor. 1985. Potassium in agriculture. American Society of Agronomy.

- National Research Council. 1995. Wetlands: Characteristics and boundaries.
- Ohio Cooperative Extension Service. 1985. Ohio agronomy guide. Ohio State University, Bulletin 472, Agdex 100.
- Portland Cement Association. 1973. PCA soil primer.
- Ruhe, Robert V. 1956. Geomorphic surfaces and the nature of soils. *Soil Science* 82: 441-455.
- Ruhe, Robert V. 1969. Quaternary landscapes in Iowa.
- Simonson, Roy W. 1959. Outline of a generalized theory of soil genesis. *Soil Science Society of America Proceedings* 23: 152-156.
- Stevenson, F.J. 1982. Humus chemistry: Genesis, composition, reactions.
- Stevenson, F.J., editor. 1982. Nitrogen in agricultural soils. American Society of Agronomy, Agronomy Monograph 22.
- Storie, R.E. 1933. An index for rating the agricultural value of soils. University of California Agricultural Experiment Station Bulletin 556.
- Tennessee Valley Authority. Site curves for eastern redcedar. (Unpublished, processed curves based on 271 observations from plots throughout the Tennessee Valley)
- Thornbury, William D. 1969. Principles of geomorphology. 2nd edition.
- Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.
- United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.
- United States Department of Agriculture, Forest Service. 1976. Volume, yield, and stand tables for second-growth southern pines. Forest Service Miscellaneous Publication 50.
- United States Department of Agriculture, Natural Resources Conservation Service. National engineering handbook.
- United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual.
- United States Department of Agriculture, Natural Resources Conservation Service. 1996. National soil survey handbook, title 430-VI. Soil Survey Staff.
- United States Department of Agriculture, Natural Resources Conservation Service. 1996. Soil survey laboratory methods manual. Soil Survey Investigations Report 42.

- United States Department of Agriculture, Natural Resources Conservation Service. 1998. Keys to soil taxonomy. 8th edition. Soil Survey Staff.
- United States Department of Agriculture, Natural Resources Conservation Service. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Soil Survey Staff. U.S. Department of Agriculture Handbook 436.
- United States Department of Agriculture, Natural Resources Conservation Service and Statistical Laboratory, Iowa State University. 2000. Summary report: 1997 national resources inventory (revised December 2000).
- United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210.
- United States Department of Agriculture, Soil Conservation Service. 1981. Land resource regions and major land resource areas of the United States. U.S. Department of Agriculture Handbook 296.
- United States Department of Agriculture, Soil Conservation Service. 1985. Site index and yield of second growth baldcypress. Soil Conservation Service Technical Note 5.
- United States Department of Agriculture, Soil Conservation Service. 1987. Basic statistics, 1982 national resources inventory. Statistical Bulletin 756.
- United States Department of Agriculture, Soil Conservation Service. 1993. Soil survey manual. Soil Survey Staff, U.S. Department of Agriculture Handbook 18.
- Walsh, L.M., and J.D. Beaton, editors. 1973. Soil testing and plant analysis. Soil Science Society of America.

Glossary

ABC soil. A soil having an A, a B, and a C horizon.

AC soil. A soil having only an A and a C horizon. Commonly, such soil formed in recent alluvium or on steep, rocky slopes.

Aeration, soil. The exchange of air in soil with air from the atmosphere. The air in a well aerated soil is similar to that in the atmosphere; the air in a poorly aerated soil is considerably higher in carbon dioxide and lower in oxygen.

Aggregate, soil. Many fine particles held in a single mass or cluster. Natural soil aggregates, such as granules, blocks, or prisms, are called peds. Clods are aggregates produced by tillage or logging.

Alluvial fan. The fanlike deposit of a stream where it issues from a gorge upon a plain or of a tributary stream near or at its junction with its main stream.

Alluvium. Material, such as sand, silt, or clay, deposited on land by streams.

Alpha, alpha-dipyridyl. A dye that when dissolved in 1N ammonium acetate is used to detect the presence of reduced iron (Fe II) in the soil. A positive reaction indicates a type of redoximorphic feature.

Anaerobic. The absence of molecular oxygen.

Animal unit month (AUM). The amount of forage required by one mature cow of approximately 1,000 pounds weight, with or without a calf, for 1 month.

Aquic conditions. Current soil wetness characterized by saturation, reduction, and redoximorphic features.

Area reclaim (in tables). An area difficult to reclaim after the removal of soil for construction and other uses. Revegetation and erosion control are extremely difficult.

Argillic horizon. A subsoil horizon characterized by an accumulation of illuvial clay.

Aspect. The direction in which a slope faces.

Association, soil. A group of soils or miscellaneous areas geographically associated in a characteristic repeating pattern and defined and delineated as a single map unit.

Available water capacity (available moisture capacity). The capacity of soils to hold water available for use by most plants. It is commonly defined as the difference between the amount of soil water at field moisture capacity and the amount at wilting point. It is commonly expressed as inches of water per inch of soil. The capacity, in inches, in a 60-inch profile or to a limiting layer is expressed as:

Very low	0 to 2.4
Low	2.4 to 3.2
Moderate	3.2 to 5.2
High	more than 5.2
Very high	more than 10

Backslope. The position that forms the steepest and generally linear, middle portion of a hillslope. In profile, backslopes are commonly bounded by a convex shoulder above and a concave footslope below.

- Basal area.** The area of a cross section of a tree, generally referring to the section at breast height and measured outside the bark. It is a measure of stand density, commonly expressed in square feet.
- Base saturation.** The degree to which material having cation-exchange properties is saturated with exchangeable bases (sum of Ca, Mg, Na, and K), expressed as a percentage of the total cation-exchange capacity.
- Base slope.** A geomorphic component of hills consisting of the concave to linear (perpendicular to the contour) slope that, regardless of the lateral shape, forms an apron or wedge at the bottom of a hillside dominated by colluvium and slope-wash sediments (for example, slope alluvium).
- Bedding planes.** Fine strata, less than 5 millimeters thick, in unconsolidated alluvial, eolian, lacustrine, or marine sediment.
- Bedrock.** The solid rock that underlies the soil and other unconsolidated material or that is exposed at the surface.
- Bedrock-controlled topography.** A landscape where the configuration and relief of the landforms are determined or strongly influenced by the underlying bedrock.
- Bench terrace.** A raised, level or nearly level strip of earth constructed on or nearly on a contour, supported by a barrier of rocks or similar material, and designed to make the soil suitable for tillage and to prevent accelerated erosion.
- Bisequum.** Two sequences of soil horizons, each of which consists of an illuvial horizon and the overlying eluvial horizons.
- Blowout.** A shallow depression from which all or most of the soil material has been removed by the wind. A blowout has a flat or irregular floor formed by a resistant layer or by an accumulation of pebbles or cobbles. In some blowouts the water table is exposed.
- Bottom land.** The normal flood plain of a stream, subject to flooding.
- Boulders.** Rock fragments larger than 2 feet (60 centimeters) in diameter.
- Breaks.** The steep and very steep broken land at the border of an upland summit that is dissected by ravines.
- Breast height.** An average height of 4.5 feet above the ground surface; the point on a tree where diameter measurements are ordinarily taken.
- Brush management.** Use of mechanical, chemical, or biological methods to make conditions favorable for reseeding or to reduce or eliminate competition from woody vegetation and thus allow understory grasses and forbs to recover. Brush management increases forage production and thus reduces the hazard of erosion. It can improve the habitat for some species of wildlife.
- Buffer strip.** Streamside vegetation consisting of trees, shrubs, and grasses for intercepting pollutants from a farm field. See conservation buffers.
- Cable yarding.** A method of moving felled trees to a nearby central area for transport to a processing facility. Most cable yarding systems involve use of a drum, a pole, and wire cables in an arrangement similar to that of a rod and reel used for fishing. To reduce friction and soil disturbance, felled trees generally are reeled in while one end is lifted or the entire log is suspended.
- Calcareous soil.** A soil containing enough calcium carbonate (commonly combined with magnesium carbonate) to effervesce visibly when treated with cold, dilute hydrochloric acid.
- Canopy.** The leafy crown of trees or shrubs. (See Crown.)
- Capillary water.** Water held as a film around soil particles and in tiny spaces between particles. Surface tension is the adhesive force that holds capillary water in the soil.
- Catena.** A sequence, or "chain," of soils on a landscape that formed in similar kinds of parent material but have different characteristics as a result of differences in relief and drainage.

- Cation.** An ion carrying a positive charge of electricity. The common soil cations are calcium, potassium, magnesium, sodium, and hydrogen.
- Cation-exchange capacity.** The total amount of exchangeable cations that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. The term, as applied to soils, is synonymous with base-exchange capacity but is more precise in meaning.
- Catsteps.** Very small, irregular terraces on steep hillsides, especially in pasture, formed by the trampling of cattle or the slippage of saturated soil.
- Channer.** Flat fragments of rock that are longer than they are thick, ranging in size from 2 millimeters to 150 millimeters.
- Channery soil material.** Soil material that has, by volume, 15 to 35 percent thin, flat fragments of sandstone, shale, slate, limestone, or schist as much as 6 inches (15 centimeters) along the longest axis. A single piece is called a channer.
- Chemical treatment.** Control of unwanted vegetation through the use of chemicals.
- Chiseling.** Tillage with an implement having one or more soil-penetrating points that shatter or loosen hard, compacted layers to a depth below normal plow depth.
- Clay.** As a soil separate, the mineral soil particles less than 0.002 millimeter in diameter. As a soil textural class, soil material that is 40 percent or more clay, less than 45 percent sand, and less than 40 percent silt.
- Clay depletions.** Low-chroma zones having a low content of iron, manganese, and clay because of the chemical reduction of iron and manganese and the removal of iron, manganese, and clay. A type of redoximorphic depletion.
- Clay film.** A thin coating of oriented clay on the surface of a soil aggregate or lining pores or root channels. Synonyms: clay coating, clay skin.
- Clayey.** Have 35 percent or more (by weight) clay and are in a shallow family or the layer is an element in a strongly contrasting particle-size class.
- Climax plant community.** The array of plant species that would be present upon a moist, fertile, adequately drained site if the site remained free of vegetation disturbance (such as wind, fire, insects, disease, human-wrought changes) long enough for the shade tolerant plant species best adapted to the climate to be both present and reproducing themselves.
- Coarse textured soil.** Sand or loamy sand.
- Coarse-loamy.** Have, in the fine-earth and gravel fraction, 15 percent or more (by weight) particles that are fine sand or coarser and in the fine-earth fraction, less than 18 percent (by weight) clay.
- Coarse-silty.** Have, in the fine-earth and gravel fraction, less than 15 percent (by weight) fine sand or coarser and, in the fine-earth fraction, less than 18 percent (by weight) clay.
- Cobble (or cobblestone).** A rounded or partly rounded fragment of rock 3 to 10 inches (7.6 to 25 centimeters) in diameter.
- Cobbly soil material.** Material that has 15 to 35 percent, by volume, rounded or partially rounded rock fragments 3 to 10 inches (7.6 to 25 centimeters) in diameter. Very cobbly soil material has 35 to 60 percent of these rock fragments, and extremely cobbly soil material has more than 60 percent.
- COLE (coefficient of linear extensibility).** See Linear extensibility.
- Complex slope.** Irregular or variable slope. Planning or establishing terraces, diversions, and other water-control structures on a complex slope is difficult.
- Complex, soil.** A map unit of two or more kinds of soil or miscellaneous areas in such an intricate pattern or so small in area that it is not practical to map them separately at the selected scale of mapping. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas.
- Component, soil.** The collection of polypedons or parts of polypedons within a map unit that are members of the taxon (or a kind of miscellaneous unit) for which the map unit is named. Simple or complex names for the component soils are formed

from a class name (taxon name) from some categorical level of the U.S. system of soil taxonomy, with or without an additional phase identification for utilitarian features.

Concretions. Cemented bodies with crude internal symmetry organized around a point, a line, or a plane. They typically take the form of concentric layers visible to the naked eye. Calcium carbonate, iron oxide, and manganese oxide are common compounds making up concretions. If formed in place, concretions of iron oxide or manganese oxide are generally considered a type of redoximorphic concentration.

Conglomerate. A coarse grained, clastic rock composed of rounded or subangular rock fragments more than 2 millimeters in diameter. It commonly has a matrix of sand and finer textured material. Conglomerate is the consolidated equivalent of gravel.

Conservation buffers. Conservation buffers are small areas or strips of land in permanent vegetation, designed to slow water runoff, provide shelter and stabilize riparian areas. Strategically placed buffer strips in the agricultural landscape can effectively mitigate the movement of sediment, nutrients, and pesticides within farm fields and from farm fields. Buffers include: contour buffer strips, field borders, filter strips, grassed waterways, living snow fences, riparian buffers, shelterbelts/windbreaks, (grass, shrubs, and trees), and wetlands.

Conservation cropping system. Growing crops in combination with needed cultural and management practices. In a good conservation cropping system, the soil-improving crops and practices more than offset the effects of the soil-depleting crops and practices. Cropping systems are needed on all tilled soils. Soil-improving practices in a conservation cropping system include the use of rotations that contain grasses and legumes and the return of crop residue to the soil. Other practices include the use of green manure crops of grasses and legumes, proper tillage, adequate fertilization, and weed and pest control.

Conservation tillage. Any tillage and planting system that covers 30 percent or more of the soil surface with crop residue, after planting, to reduce soil erosion by water.

Consistence, soil. Refers to the degree of cohesion and adhesion of soil material and its resistance to deformation when ruptured. Consistence includes resistance of soil material to rupture and to penetration; plasticity, toughness, and stickiness of puddled soil material; and the manner in which the soil material behaves when subject to compression. Terms describing consistence are defined in the "Soil Survey Manual."

Consociation, soil. A kind of map unit comprised of delineations, each of which shows the size, shape, and location of a landscape unit composed of one kind of component soil, or one kind of miscellaneous area, plus allowable inclusions in either case. See also component soil, soil complex, soil association, undifferentiated group, miscellaneous areas.

Contour farming. Farming with row patterns nearly level around the hill, not up and down the slope.

Contour stripcropping. Growing crops in strips that follow the contour. Strips of grass or close-growing crops are alternated with strips of clean-tilled crops or summer fallow.

Control section. The part of the soil on which classification is based. The thickness varies among different kinds of soil, but for many it is that part of the soil profile between depths of 10 inches and 40 or 80 inches.

Corrosion. Soil-induced electrochemical or chemical action that dissolves or weakens concrete or uncoated steel.

Cover crop. A close-growing crop grown primarily to improve and protect the soil between periods of regular crop production, or a crop grown between trees and vines in orchards and vineyards.

- Cropping system.** Growing crops according to a planned system of rotation and management practices.
- Crop residue management.** Returning crop residue to the soil, which helps to maintain soil structure, organic matter content, and fertility and helps to control erosion.
- Cross-slope farming.** Deliberately conducting farming operations on sloping farmland in such a way that tillage is across the general slope.
- Crown.** The upper part of a tree or shrub, including the living branches and their foliage.
- Cutbanks cave** (in tables). The walls of excavations tend to cave in or slough.
- Deferred grazing.** Postponing grazing or resting grazing land for a prescribed period.
- Delta.** A body of alluvium having a surface that is nearly flat and fan shaped; deposited at or near the mouth of a river or stream where it enters a body of relatively quiet water, generally a sea or lake.
- Dense layer (in tables).** A very firm, massive layer that has a bulk density of more than 1.8 grams per cubic centimeter. Such a layer affects the ease of digging and can affect filling and compacting.
- Dense till.** Earthy, compact material deposited by a glacier. The dense till in Connecticut has a bulk density of at least 1.65 g cm⁻³. See lodgement till.
- Densic material.** Relatively unaltered materials that have a noncemented rupture resistance class. The bulk density is such that roots cannot enter, except in cracks. Densic material is mostly earthy materials, such as till.
- Depth, soil.** Generally, the thickness of the soil over bedrock. Very deep soils are more than 60 inches deep over bedrock; deep soils, 40 to 60 inches; moderately deep, 20 to 40 inches; shallow, 10 to 20 inches; and very shallow, less than 10 inches.
- Depth to rock** (in tables). Bedrock is too near the surface for the specified use.
- Dip slope.** A slope of the land surface, roughly determined by and approximately conforming to the dip of the underlying bedrock.
- Diversion (or diversion terrace).** A ridge of earth, generally a terrace, built to protect downslope areas by diverting runoff from its natural course.
- Drainage class** (natural). Refers to the frequency and duration of wet periods under conditions similar to those under which the soil formed. Alterations of the water regime by human activities, either through drainage or irrigation, are not a consideration unless they have significantly changed the morphology of the soil. Seven classes of natural soil drainage are recognized—*excessively drained, somewhat excessively drained, well drained, moderately well drained, somewhat poorly drained, poorly drained, and very poorly drained*. These classes are defined in the “Soil Survey Manual.”
- Drainage, surface.** Runoff, or surface flow of water, from an area.
- Drumlin.** A low, smooth, elongated oval hill, mound, or ridge of compact glacial till. The longer axis is parallel to the path of the glacier and commonly has a blunt nose pointing in the direction from which the ice approached.
- Duff.** A generally firm organic layer on the surface of mineral soils. It consists of fallen plant material that is in the process of decomposition and includes everything from the litter on the surface to underlying pure humus.
- Dysic.** Histosols that have a pH value, on undried samples, of less than 4.5 (in 0.01 M CaCl₂) in one or more layers of organic soil materials within the control section.
- Ecological site.** An area where climate, soil, and relief are sufficiently uniform to produce a distinct natural plant community. An ecological site is the product of all the environmental factors responsible for its development. It is typified by an association of species that differ from those on other ecological sites in kind and/or proportion of species or in total production.

- Eluviation.** The movement of material in true solution or colloidal suspension from one place to another within the soil. Soil horizons that have lost material through eluviation are eluvial; those that have received material are illuvial.
- Endosaturation.** A type of saturation of the soil in which all horizons between the upper boundary of saturation and a depth of 2 meters are saturated.
- Eolian soil material.** Earthy parent material accumulated through wind action; commonly refers to sandy material in dunes or to loess in blankets on the surface.
- Ephemeral stream.** A stream, or reach of a stream, that flows only in direct response to precipitation. It receives no long-continued supply from melting snow or other source, and its channel is above the water table at all times.
- Episaturation.** A type of saturation indicating a perched water table in a soil in which saturated layers are underlain by one or more unsaturated layers within 2 meters of the surface.
- Erosion.** The wearing away of the land surface by water, wind, ice, or other geologic agents and by such processes as gravitational creep.
Erosion (geologic). Erosion caused by geologic processes acting over long geologic periods and resulting in the wearing away of mountains and the building up of such landscape features as flood plains and coastal plains. Synonym: natural erosion.
Erosion (accelerated). Erosion much more rapid than geologic erosion, mainly as a result of human or animal activities or of a catastrophe in nature, such as a fire, that exposes the surface.
- Erosion pavement.** A layer of gravel or stones that remains on the surface after fine particles are removed by sheet or rill erosion.
- Escarpment.** A relatively continuous and steep slope or cliff breaking the general continuity of more gently sloping land surfaces and resulting from erosion or faulting. Synonym: scarp.
- Esker.** A narrow, winding ridge of stratified gravelly and sandy drift deposited by a stream flowing in a tunnel beneath a glacier.
- Euic.** Histosols that have a pH value, on undried samples, of 4.5 or more (in 0.01 M CaCl₂) in one or more layers of organic soil materials within the control section.
- Excess fines (in tables).** Excess silt and clay in the soil. The soil does not provide a source of gravel or sand for construction purposes.
- Excess salts (in tables).** Excess water-soluble salts in the soil that restrict the growth of most plants.
- Excess sulfur (in tables).** Excessive amount of sulfur in the soil. The sulfur causes extreme acidity if the soil is drained, and the growth of most plants is restricted.
- Extrusive rock.** Igneous rock derived from deep-seated molten matter (magma) emplaced on the earth's surface.
- Fallow.** Cropland left idle in order to restore productivity through accumulation of moisture. Summer fallow is common in regions of limited rainfall where cereal grain is grown. The soil is tilled for at least one growing season for weed control and decomposition of plant residue.
- Fast intake (in tables).** The rapid movement of water into the soil.
- Fertility, soil.** The quality that enables a soil to provide plant nutrients, in adequate amounts and in proper balance, for the growth of specified plants when light, moisture, temperature, tilth, and other growth factors are favorable.
- Fibric soil material (peat).** The least decomposed of all organic soil material. Peat contains a large amount of well preserved fiber that is readily identifiable according to botanical origin. Peat has the lowest bulk density and the highest water content at saturation of all organic soil material.
- Field moisture capacity.** The moisture content of a soil, expressed as a percentage of the oven-dry weight, after the gravitational, or free, water has drained away; the

field moisture content 2 or 3 days after a soaking rain; also called *normal field capacity*, *normal moisture capacity*, or *capillary capacity*.

- Fill slope.** A sloping surface consisting of excavated soil material from a road cut. It commonly is on the downhill side of the road.
- Fine.** Have 35 percent or more (by weight) clay and less than 60 percent (by weighted average) in the fine-earth fraction.
- Fine textured soil.** Sandy clay, silty clay, or clay.
- Fine-loamy.** Have, in the fine-earth and gravel fraction, 15 percent or more (by weight) fine sand or coarser, and 18 to 35 percent (by weight) clay.
- Fine-silty.** Have, in the fine-earth and gravel fraction, less than 15 percent (by weight) fine sand or coarser and in the fine-earth fraction 18 to 35 percent clay.
- Firebreak.** Area cleared of flammable material to stop or help control creeping or running fires. It also serves as a line from which to work and to facilitate the movement of firefighters and equipment. Designated roads also serve as firebreaks.
- Flaggy soil material.** Material that has, by volume, 15 to 35 percent flagstones. Very flaggy soil material has 35 to 60 percent flagstones, and extremely flaggy soil material has more than 60 percent flagstones.
- Flagstone.** A thin fragment of sandstone, limestone, slate, shale, or (rarely) schist 6 to 15 inches (15 to 38 centimeters) long.
- Flood plain.** A nearly level alluvial plain that borders a stream and is subject to flooding unless protected artificially.
- Flooding.** Accumulation of large amounts of runoff on the landscape as a result of rainfall in excess of the soils ability to drain water from the landscape before extensive inundation and ponding occurs. Flooding is described by frequency, duration, and month. Flooding is a result of overflow from a stream or other flowing water. See ponding.
- Fluvial.** Of or pertaining to rivers; produced by river action, as a fluvial plain.
- Foothill.** A steeply sloping upland that has relief of as much as 1,000 feet (300 meters) and fringes a mountain range or high-plateau escarpment.
- Footslope.** The position that forms the inner, gently inclined surface at the base of a hillslope. In profile, footslopes are commonly concave. A footslope is a transition zone between upslope sites of erosion and transport (shoulders and backslopes) and downslope sites of deposition (toeslopes).
- Forb.** Any herbaceous plant not a grass or a sedge.
- Forest cover.** All trees and other woody plants (underbrush) covering the ground in a forest.
- Forest type.** A stand of trees similar in composition and development because of given physical and biological factors by which it may be differentiated from other stands.
- Fragipan.** A loamy, brittle subsurface horizon low in porosity and content of organic matter and low or moderate in clay but high in silt or very fine sand. A fragipan appears cemented and restricts roots. When dry, it is hard or very hard and has a higher bulk density than the horizon or horizons above. When moist, it tends to rupture suddenly under pressure rather than to deform slowly.
- Frost action** (in tables). Freezing and thawing of soil moisture. Frost action can damage roads, buildings and other structures, and plant roots.
- Genesis, soil.** The mode of origin of the soil. Refers especially to the processes or soil-forming factors responsible for the formation of the solum, or true soil, from the unconsolidated parent material.
- Glacial drift.** Pulverized and other rock material transported by glacial ice and then deposited. Also, the sorted and unsorted material deposited by streams flowing from glaciers.

- Glacial outwash.** Gravel, sand, and silt, commonly stratified, deposited by glacial meltwater.
- Glacial till (till).** Unsorted, nonstratified glacial drift consisting of clay, silt, sand, and boulders transported and deposited by glacial ice.
- Glaciofluvial deposits.** Material moved by glaciers and subsequently sorted and deposited by streams flowing from the melting ice. The deposits are stratified and occur as kames, eskers, deltas, and outwash plains.
- Glaciolacustrine deposits.** Material ranging from fine clay to sand derived from glaciers and deposited in glacial lakes mainly by glacial meltwater. Many deposits are interbedded or laminated.
- Gleyed soil.** Soil that formed under poor drainage, resulting in the reduction of iron and other elements in the profile and in gray colors.
- Grassed waterway.** A natural or constructed waterway, typically broad and shallow, seeded to grass as protection against erosion. Conducts surface water away from cropland.
- Gravel.** Rounded or angular fragments of rock as much as 3 inches (2 millimeters to 7.6 centimeters) in diameter. An individual piece is a pebble.
- Gravelly soil material.** Material that has 15 to 35 percent, by volume, rounded or angular rock fragments, not prominently flattened, as much as 3 inches (7.6 centimeters) in diameter.
- Green manure crop (agronomy).** A soil-improving crop grown to be plowed under in an early stage of maturity or soon after maturity.
- Ground water.** Water filling all the unblocked pores of the material below the water table.
- Gully.** A miniature valley with steep sides cut by running water and through which water ordinarily runs only after rainfall. The distinction between a gully and a rill is one of depth. A gully generally is an obstacle to farm machinery and is too deep to be obliterated by ordinary tillage; a rill is of lesser depth and can be smoothed over by ordinary tillage.
- Hard bedrock.** Bedrock that cannot be excavated except by blasting or by the use of special equipment that is not commonly used in construction.
- Hardpan.** A hardened or cemented soil horizon, or layer. The soil material is sandy, loamy, or clayey and is cemented by iron oxide, silica, calcium carbonate, or other substance.
- Head slope.** A geomorphic component of hills consisting of a laterally concave area of a hillside, especially at the head of a drainageway. The overland waterflow is converging.
- Headcut.** Small abrupt elevation drops (1-5 centimeters) on the floor of rills or irrigation furrows that result in accelerated erosion as they undercut the rill floor and migrate upstream.
- Hemic soil material (mucky peat).** Organic soil material intermediate in degree of decomposition between the less decomposed fibric material and the more decomposed sapric material.
- High-residue crops.** Such crops as small grain and corn used for grain. If properly managed, residue from these crops can be used to control erosion until the next crop in the rotation is established. These crops return large amounts of organic matter to the soil.
- Hill.** A natural elevation of the land surface, rising as much as 1,000 feet above surrounding lowlands, commonly of limited summit area and having a well defined outline; hillsides generally have slopes of more than 15 percent. The distinction between a hill and a mountain is arbitrary and is dependent on local usage.
- Histic epipedon.** A surface organic soil layer that is characterized by saturation and reduction for some time during normal years. The histic epipedon contains more organic carbon than the mollic epipedon.

Horizon, soil. A layer of soil, approximately parallel to the surface, having distinct characteristics produced by soil-forming processes. In the identification of soil horizons, an uppercase letter represents the major horizons. Numbers or lowercase letters that follow represent subdivisions of the major horizons. An explanation of the subdivisions is given in the "Soil Survey Manual." The major horizons of mineral soil are as follows:

O horizon.—An organic layer of fresh and decaying plant residue.

A horizon.—The mineral horizon at or near the surface in which an accumulation of humified organic matter is mixed with the mineral material. Also, a plowed surface horizon, most of which was originally part of a B horizon.

E horizon.—The mineral horizon in which the main feature is loss of silicate clay, iron, aluminum, or some combination of these.

B horizon.—The mineral horizon below an A horizon. The B horizon is in part a layer of transition from the overlying A to the underlying C horizon. The B horizon also has distinctive characteristics, such as (1) accumulation of clay, sesquioxides, humus, or a combination of these; (2) prismatic or blocky structure; (3) redder or browner colors than those in the A horizon; or (4) a combination of these.

C horizon.—The mineral horizon or layer, excluding indurated bedrock, that is little affected by soil-forming processes and does not have the properties typical of the overlying soil material. The material of a C horizon may be either like or unlike that in which the solum formed. If the material is known to differ from that in the solum, an Arabic numeral, commonly a 2, precedes the letter C.

Cr horizon.—Soft, consolidated bedrock beneath the soil.

R layer.—Consolidated bedrock beneath the soil. The bedrock commonly underlies a C horizon, but it can be directly below an A or a B horizon.

Humus. The well decomposed, more or less stable part of the organic matter in mineral soils.

Hydric soil. A soil that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part. The concept of hydric soils includes soils developed under sufficiently wet conditions to support the growth and regeneration of hydrophytic vegetation. Soils that are sufficiently wet because of artificial measures are included in the concept of hydric soils. Also, soils in which the hydrology has been artificially modified are hydric if the soil, in an unaltered state, was hydric. Some series designated as hydric have phases that are not hydric depending on water table, flooding, and ponding characteristics.

Hydrologic soil groups. Refers to soils grouped according to their runoff potential. The soil properties that influence this potential are those that affect the minimum rate of water infiltration on a bare soil during periods after prolonged wetting when the soil is not frozen. These properties are depth to a seasonal high water table, the infiltration rate and permeability after prolonged wetting, and depth to a very slowly permeable layer. The slope and the kind of plant cover are not considered but are separate factors in predicting runoff.

Hydrophytic vegetation. Plant life growing in water or on a substrate that is at least periodically deficient in oxygen as a result of excessive water content.

Igneous rock. Rock formed by solidification from a molten or partially molten state. Major varieties include plutonic and volcanic rock. Examples are andesite, basalt, and granite.

Illuviation. The movement of soil material from one horizon to another in the soil profile. Generally, material is removed from an upper horizon and deposited in a lower horizon.

Impervious soil. A soil through which water, air, or roots penetrate slowly or not at all. No soil is absolutely impervious to air and water all the time.

Inclusion. See minor component.

Infiltration. The downward entry of water into the immediate surface of soil or other material, as contrasted with percolation, which is movement of water through soil layers or material.

Infiltration capacity. The maximum rate at which water can infiltrate into a soil under a given set of conditions.

Infiltration rate. The rate at which water penetrates the surface of the soil at any given instant, usually expressed in inches per hour. The rate can be limited by the infiltration capacity of the soil or the rate at which water is applied at the surface.

Intake rate. The average rate of water entering the soil under irrigation. Most soils have a fast initial rate; the rate decreases with application time. Therefore, intake rate for design purposes is not a constant but is a variable depending on the net irrigation application. The rate of water intake, in inches per hour, is expressed as follows:

Less than 0.2	very low
0.2 to 0.4	low
0.4 to 0.75	moderately low
0.75 to 1.25	moderate
1.25 to 1.75	moderately high
1.75 to 2.5	high
More than 2.5	very high

Interfluvium. An elevated area between two drainageways that sheds water to those drainageways.

Intermittent stream. A stream, or reach of a stream, that flows for prolonged periods only when it receives ground-water discharge or long, continued contributions from melting snow or other surface and shallow subsurface sources.

Invasive species. A species that is 1) non-native to the ecosystem under consideration and 2) whose introduction causes or is likely to cause economic or environmental harm or harm to human health.

Iron depletions. Low-chroma zones having a low content of iron and manganese oxide because of chemical reduction and removal, but having a clay content similar to that of the adjacent matrix. A type of redoximorphic depletion.

Irrigation. Application of water to soils to assist in production of crops. Methods of irrigation are:

Basin.—Water is applied rapidly to nearly level plains surrounded by levees or dikes.

Controlled flooding.—Water is released at intervals from closely spaced field ditches and distributed uniformly over the field.

Corrugation.—Water is applied to small, closely spaced furrows or ditches in fields of close-growing crops or in orchards so that it flows in only one direction.

Drip (or trickle).—Water is applied slowly and under low pressure to the surface of the soil or into the soil through such applicators as emitters, porous tubing, or perforated pipe.

Furrow.—Water is applied in small ditches made by cultivation implements. Furrows are used for tree and row crops.

Sprinkler.—Water is sprayed over the soil surface through pipes or nozzles from a pressure system.

Subirrigation.—Water is applied in open ditches or tile lines until the water table is raised enough to wet the soil.

Wild flooding.—Water, released at high points, is allowed to flow onto an area without controlled distribution.

Kame. An irregular, short ridge or hill of stratified glacial drift.

Knoll. A small, low, rounded hill rising above adjacent landforms.

- K_{sat} .** Saturated hydraulic conductivity (K_{sat}) quantifies the soil's ability to transmit water under saturated conditions. It is used to compare water movement in different soils, layers, or materials. (See Permeability.)
- Lacustrine deposit.** Material deposited in lake water and exposed when the water level is lowered or the elevation of the land is raised.
- Landslide.** The rapid downhill movement of a mass of soil and loose rock, generally when wet or saturated. The speed and distance of movement, as well as the amount of soil and rock material, vary greatly.
- Large stones** (in tables). Rock fragments 3 inches (7.6 centimeters) or more across. Large stones adversely affect the specified use of the soil.
- Leaching.** The removal of soluble material from soil or other material by percolating water.
- Linear extensibility.** Refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. Linear extensibility is used to determine the shrink-swell potential of soils. It is an expression of the volume change between the water content of the clod at $1/3$ - or $1/10$ -bar tension (33kPa or 10kPa tension) and oven dryness. Volume change is influenced by the amount and type of clay minerals in the soil. The volume change is the percent change for the whole soil. If it is expressed as a fraction, the resulting value is COLE, coefficient of linear extensibility.
- Liquid limit.** The moisture content at which the soil passes from a plastic to a liquid state.
- Loam.** Soil material that is 7 to 27 percent clay particles, 28 to 50 percent silt particles, and less than 52 percent sand particles.
- Loamy.** Have a texture of loamy very fine sand, very fine sand, or finer, including less than 35 percent (by weight) clay in the fine-earth fraction.
- Loamy-skeletal.** Soil material that has 35 percent or more (by volume) rock fragments, texture finer than loamy sand, and less than 35 percent (by weight) clay.
- Lodgement till.** Compact glacial till deposited beneath the ice.
- Loess.** Fine grained material, dominantly of silt-sized particles, deposited by wind.
- Loose till.** Earthy, friable material deposited by a glacier. Loose till in Connecticut has a bulk density less than 1.65 g cm⁻³. See melt-out till.
- Low-residue crops.** Such crops as corn used for silage, peas, beans, and potatoes. Residue from these crops is not adequate to control erosion until the next crop in the rotation is established. These crops return little organic matter to the soil.
- Low strength.** The soil is not strong enough to support loads.
- Major component.** The dominant soil(s) in a named mapping unit. Individually, major components account for no less than 15 percent of the composition of soils in a mapping unit.
- Marl.** An earthy, unconsolidated deposit consisting chiefly of calcium carbonate mixed with clay in approximately equal amounts.
- Masses.** Concentrations of substances in the soil matrix that do not have a clearly defined boundary with the surrounding soil material and cannot be removed as a discrete unit. Common compounds making up masses are calcium carbonate, gypsum or other soluble salts, iron oxide, and manganese oxide. Masses consisting of iron oxide or manganese oxide generally are considered a type of redoximorphic concentration.
- Mechanical treatment.** Use of mechanical equipment for seeding, brush management, and other management practices.
- Medium textured soil.** Very fine sandy loam, loam, silt loam, or silt.
- Meltout till.** Loose, permeable till deposited during the final downwasting of glacial ice. Lenses of crudely sorted sand and gravel are common. See loose till.

- Metamorphic rock.** Rock of any origin altered in mineralogical composition, chemical composition, or structure by heat, pressure, and movement. Nearly all such rocks are crystalline.
- Mineral soil.** Soil that is mainly mineral material and low in organic material. Its bulk density is more than that of organic soil.
- Minimum tillage.** Only the tillage essential to crop production and prevention of soil damage.
- Minor component.** The unnamed and lesser soils in a mapping unit, sometimes called inclusions. Individually, minor components account for less than 15 percent of the composition of soils in a mapping unit.
- Miscellaneous area.** An area that has little or no natural soil and supports little or no vegetation without major reclamation. Examples of miscellaneous areas are beaches, dumps, urban land, and rock outcrop.
- Moderately coarse textured soil.** Coarse sandy loam, sandy loam, or fine sandy loam.
- Moderately fine textured soil.** Clay loam, sandy clay loam, or silty clay loam.
- Mollic epipedon.** A thick, dark, humus-rich surface horizon (or horizons) that has high base saturation and pedogenic soil structure. It may include the upper part of the subsoil.
- Moraine.** An accumulation of earth, stones, and other debris deposited by a glacier. Some types are terminal, lateral, medial, and ground.
- Morphology, soil.** The physical makeup of the soil, including the texture, structure, porosity, consistence, color, and other physical, mineral, and biological properties of the various horizons, and the thickness and arrangement of those horizons in the soil profile.
- Mottling, soil.** Irregular spots of different colors that vary in number and size. Descriptive terms are as follows: abundance—*few*, *common*, and *many*; size—*fine*, *medium*, and *coarse*; and contrast—*faint*, *distinct*, and *prominent*. The size measurements are of the diameter along the greatest dimension. *Fine* indicates less than 5 millimeters (about 0.2 inch); *medium*, from 5 to 15 millimeters (about 0.2 to 0.6 inch); and *coarse*, more than 15 millimeters (about 0.6 inch).
- Muck.** Dark, finely divided, well decomposed organic soil material. (See Sapric soil material.)
- Mudstone.** Sedimentary rock formed by induration of silt and clay in approximately equal amounts.
- Munsell notation.** A designation of color by degrees of three simple variables—hue, value, and chroma. For example, a notation of 10YR 6/4 is a color with hue of 10YR, value of 6, and chroma of 4.
- Neutral soil.** A soil having a pH value of 6.6 to 7.3. (See Reaction, soil.)
- Nodules.** Cemented bodies lacking visible internal structure. Calcium carbonate, iron oxide, and manganese oxide are common compounds making up nodules. If formed in place, nodules of iron oxide or manganese oxide are considered types of redoximorphic concentrations.
- Nose slope.** A geomorphic component of hills consisting of the projecting end (laterally convex area) of a hillside. The overland waterflow is predominantly divergent.
- Nutrient, plant.** Any element taken in by a plant essential to its growth. Plant nutrients are mainly nitrogen, phosphorus, potassium, calcium, magnesium, sulfur, iron, manganese, copper, boron, and zinc obtained from the soil and carbon, hydrogen, and oxygen obtained from the air and water.
- Ochric epipedon.** A surface horizon of mineral soil that is too light in color, too high in chroma, too low in organic carbon, or too thin to be a plaggen, mollic, umbric, anthropic, or histic epipedon, or is both hard and massive when dry. Typically, ochric epipedons have color value of 4 or more.

Organic matter. Plant and animal residue in the soil in various stages of decomposition. The content of organic matter in the surface layer is described as follows:

Very low	less than 0.5 percent
Low	0.5 to 1.0 percent
Moderately low	1.0 to 2.0 percent
Moderate	2.0 to 4.0 percent
High	4.0 to 8.0 percent
Very high	more than 8.0 percent

Outwash plain. A landform of mainly sandy or coarse textured material of glaciofluvial origin. An outwash plain is commonly smooth; where pitted, it generally is low in relief.

Pan. A compact, dense layer in a soil that impedes the movement of water and the growth of roots. For example, *hardpan*, *fragipan*, *claypan*, *plowpan*, and *traffic pan*.

Parent material. The unconsolidated organic and mineral material in which soil forms.

Peat. Unconsolidated material, largely undecomposed organic matter, which has accumulated under excess moisture. (See Fibric soil material.)

Ped. An individual natural soil aggregate, such as a granule, a prism, or a block.

Pedon. The smallest volume that can be called "a soil." A pedon is three dimensional and large enough to permit study of all horizons. Its area ranges from about 10 to 100 square feet (1 square meter to 10 square meters), depending on the variability of the soil.

Percolation. The movement of water through the soil.

Percs slowly (in tables). The slow movement of water through the soil adversely affects the specified use.

Permeability. The quality of the soil that enables water or air to move downward through the profile. The rate at which a saturated soil transmits water is accepted as a measure of this quality. In soil physics, the rate is referred to as "saturated hydraulic conductivity," which is defined in the "Soil Survey Manual." In line with conventional usage in the engineering profession and with traditional usage in published soil surveys, this rate of flow continues to be expressed as "permeability." Permeability rates are listed in the tables for both mineral and organic soil horizons. The permeability ranges listed in the detailed soil map unit descriptions and soil series descriptions refer only to the permeability of the mineral soil horizons. The permeability ranges are measured in inches per hour and are as follows:

Extremely slow	0.0 to 0.01 inch
Very slow	0.01 to 0.06 inch
Slow	0.06 to 0.2 inch
Moderately slow	0.2 to 0.6 inch
Moderate	0.6 inch to 2.0 inches
Moderately rapid	2.0 to 6.0 inches
Rapid	6.0 to 20 inches
Very rapid	more than 20 inches

Phase, soil. A subdivision of a soil series based on features that affect its use and management, such as slope, stoniness, and flooding.

pH value. A numerical designation of acidity and alkalinity in soil. (See Reaction, soil.)

Piping (in tables). Formation of subsurface tunnels or pipelike cavities by water moving through the soil.

- Plasticity index.** The numerical difference between the liquid limit and the plastic limit; the range of moisture content within which the soil remains plastic.
- Plastic limit.** The moisture content at which a soil changes from semisolid to plastic.
- Plowpan.** A compacted layer formed in the soil directly below the plowed layer.
- Ponding.** Standing water on soils in closed depressions. Unless the soils are artificially drained, the water can be removed only by percolation or evapotranspiration. Ponding is described by depth, duration, frequency class, and month.
- Poor filter** (in tables). Because of rapid or very rapid permeability, the soil may not adequately filter effluent from a waste disposal system.
- Poorly graded.** Refers to a coarse grained soil or soil material consisting mainly of particles of nearly the same size. Because there is little difference in size of the particles, density can be increased only slightly by compaction.
- Potential native plant community.** See Climax plant community.
- Potential rooting depth (effective rooting depth).** Depth to which roots could penetrate if the content of moisture in the soil were adequate. The soil has no properties restricting the penetration of roots to this depth.
- Prescribed burning.** Deliberately burning an area for specific management purposes, under the appropriate conditions of weather and soil moisture and at the proper time of day.
- Productivity, soil.** The capability of a soil for producing a specified plant or sequence of plants under specific management.
- Profile, soil.** A vertical section of the soil extending through all its horizons and into the parent material.
- Proper grazing use.** Grazing at an intensity that maintains enough cover to protect the soil and maintain or improve the quantity and quality of the desirable vegetation. This practice increases the vigor and reproduction capacity of the key plants and promotes the accumulation of litter and mulch necessary to conserve soil and water.
- Reaction, soil.** A measure of acidity or alkalinity of a soil, expressed in pH values. A soil that tests to pH 7.0 is described as precisely neutral in reaction because it is neither acid nor alkaline. The reaction ranges listed in the detailed soil map unit descriptions and soil series descriptions refer does not include the reaction of Oi or Oe horizons. The degrees of acidity or alkalinity, expressed as pH values, are:
- | | |
|------------------------------|----------------|
| Ultra acid | less than 3.5 |
| Extremely acid | 3.5 to 4.4 |
| Very strongly acid | 4.5 to 5.0 |
| Strongly acid | 5.1 to 5.5 |
| Moderately acid | 5.6 to 6.0 |
| Slightly acid | 6.1 to 6.5 |
| Neutral | 6.6 to 7.3 |
| Slightly alkaline | 7.4 to 7.8 |
| Moderately alkaline | 7.9 to 8.4 |
| Strongly alkaline | 8.5 to 9.0 |
| Very strongly alkaline | 9.1 and higher |
- Red beds.** Sedimentary strata that are mainly red and are made up largely of sandstone and shale.
- Redoximorphic concentrations.** Nodules, concretions, soft masses, pore linings, and other features resulting from the accumulation of iron or manganese oxide. An indication of chemical reduction and oxidation resulting from saturation.
- Redoximorphic depletions.** Low-chroma zones from which iron and manganese oxide or a combination of iron and manganese oxide and clay has been removed. These zones are indications of the chemical reduction of iron resulting from saturation.

- Redoximorphic features.** Redoximorphic concentrations, redoximorphic depletions, reduced matrices, a positive reaction to alpha,alpha-dipyridyl, and other features indicating the chemical reduction and oxidation of iron and manganese compounds resulting from saturation.
- Reduced matrix.** A soil matrix that has low chroma in situ because of chemically reduced iron (Fe II). The chemical reduction results from nearly continuous wetness. The matrix undergoes a change in hue or chroma within 30 minutes after exposure to air as the iron is oxidized (Fe III). A type of redoximorphic feature.
- Regolith.** The unconsolidated mantle of weathered rock and soil material on the earth's surface; the loose earth material above the solid rock.
- Relief.** The elevations or inequalities of a land surface, considered collectively.
- Rill.** A steep-sided channel resulting from accelerated erosion. A rill generally is a few inches deep and not wide enough to be an obstacle to farm machinery.
- Rill erosion.** The removal of soil by concentrated water running through little streamlets, or headcuts. Detachment in a rill occurs if the sediment in the flow is below the amount the water can transport and if the flow exceeds the soil's resistance to detachment. As detachment continues or flow increases, rills will become wider and deeper.
- Riparian buffer.** Strips of grass, shrubs, and/or trees along the banks of rivers and streams that filter polluted runoff and provide a transition zone between water and human land use.
- Road cut.** A sloping surface produced by mechanical means during road construction. It is commonly on the uphill side of the road.
- Rock fragments.** Rock or mineral fragments having a diameter of 2 millimeters or more; for example, pebbles, cobbles, stones, and boulders.
- Root zone.** The part of the soil that can be penetrated by plant roots.
- Rooting depth** (in tables). Shallow root zone. The soil is shallow over a layer that greatly restricts roots.
- Runoff.** The precipitation discharged into stream channels from an area. The water that flows off the surface of the land without sinking into the soil is called surface runoff. Water that enters the soil before reaching surface streams is called ground-water runoff or seepage flow from ground water.
- Salic horizon.** A mineral soil horizon of enrichment with secondary salts more soluble in cold water than gypsum. A salic horizon is 15 centimeters or more in thickness, contains at least 20 grams per kilogram salt, and the product of the thickness in centimeters and amount of salt by weight is more than 600 grams per kilogram.
- Saline soil.** A soil containing soluble salts in an amount that impairs growth of plants. A saline soil does not contain excess exchangeable sodium.
- Salty water** (in tables). Water that is too salty for consumption by livestock.
- Sand.** As a soil separate, individual rock or mineral fragments from 0.05 millimeter to 2.0 millimeters in diameter. Most sand grains consist of quartz. As a soil textural class, a soil that is 85 percent or more sand and not more than 10 percent clay.
- Sandstone.** Sedimentary rock containing dominantly sand-sized particles.
- Sandy.** Have a texture of sand or loamy sand, including less than 50 percent (by weight) very fine sand in the fine-earth.
- Sandy loam.** Soil material that contains 20% clay or less and the percentage of silt plus twice the percentage of clay exceeds 30, and 52% or more sand; OR less than 7% clay, less than 50% silt and between 43% and 52% sand. Sandy loam texture includes coarse sandy loam, fine sandy loam, and very fine sandy loam.
- Sandy-skeletal.** Soil material that has 35 percent or more (by volume) rock fragments and a fine-earth fraction with a texture of sand or loamy sand, including less than 50 percent (by weight) very fine sand.

- Sapric soil material (muck).** The most highly decomposed of all organic soil material. Muck has the least amount of plant fiber, the highest bulk density, and the lowest water content at saturation of all organic soil material.
- Saprolite.** Unconsolidated residual material underlying the soil and grading to hard bedrock below.
- Saturation.** Wetness characterized by zero or positive pressure of the soil water. Under conditions of saturation, the water will flow from the soil matrix into an unlined auger hole.
- Saturated conditions.** The condition in which all pores (voids) between soil particles are filled with water.
- Scarification.** The act of abrading, scratching, loosening, crushing, or modifying the surface to increase water absorption or to provide a more tillable soil.
- Sedimentary rock.** Rock made up of particles deposited from suspension in water. The chief kinds of sedimentary rock are conglomerate, formed from gravel; sandstone, formed from sand; shale, formed from clay; and limestone, formed from soft masses of calcium carbonate. There are many intermediate types. Some wind-deposited sand is consolidated into sandstone.
- Seepage** (in tables). The movement of water through the soil. Seepage adversely affects the specified use.
- Sequum.** A sequence consisting of an illuvial horizon and the overlying eluvial horizon. (See Eluviation.)
- Series, soil.** A group of soils that have profiles that are almost alike, except for differences in texture of the surface layer. All the soils of a series have horizons that are similar in composition, thickness, and arrangement.
- Shale.** Sedimentary rock formed by the hardening of a clay deposit.
- Sheet erosion.** The removal of a fairly uniform layer of soil material from the land surface by the action of rainfall and surface runoff.
- Shoulder.** The position that forms the uppermost inclined surface near the top of a hillslope. It is a transition from backslope to summit. The surface is dominantly convex in profile and erosional in origin.
- Shrink-swell** (in tables). The shrinking of soil when dry and the swelling when wet. Shrinking and swelling can damage roads, dams, building foundations, and other structures. It can also damage plant roots.
- Side slope.** A geomorphic component of hills consisting of a laterally planar area of a hillside. The overland waterflow is predominantly parallel.
- Silica.** A combination of silicon and oxygen. The mineral form is called quartz.
- Silica-sesquioxide ratio.** The ratio of the number of molecules of silica to the number of molecules of alumina and iron oxide. The more highly weathered soils or their clay fractions in warm-temperate, humid regions, and especially those in the tropics, generally have a low ratio.
- Silt.** As a soil separate, individual mineral particles that range in diameter from the upper limit of clay (0.002 millimeter) to the lower limit of very fine sand (0.05 millimeter). As a soil textural class, soil that is 80 percent or more silt and less than 12 percent clay.
- Silt loam.** Soil material that contains 50 percent or more silt and 7 percent to 27 percent clay; or 50 percent to 80 percent silt and less than 12 percent clay.
- Silty clay.** Soil material that contains 40 percent or more clay and 40 percent or more silt.
- Silty clay loam.** Soil material that contains 27 percent to 40 percent clay and less than 20 percent sand.
- Siltstone.** Sedimentary rock made up of dominantly silt-sized particles.
- Similar soils.** Soils that share limits of diagnostic criteria, behave and perform in a similar manner, and have similar conservation needs or management requirements for the major land uses in the survey area.

- Site index.** A designation of the quality of a forest site based on the height of the dominant stand at an arbitrarily chosen age. For example, if the average height attained by dominant and codominant trees in a fully stocked stand at the age of 50 years is 75 feet, the site index is 75.
- Slickensides.** Polished and grooved surfaces produced by one mass sliding past another. In soils, slickensides may occur at the bases of slip surfaces on the steeper slopes; on faces of blocks, prisms, and columns; and in swelling clayey soils, where there is marked change in moisture content.
- Slippage** (in tables). Soil mass susceptible to movement downslope when loaded, excavated, or wet.
- Slope.** The inclination of the land surface from the horizontal. Percentage of slope is the vertical distance divided by horizontal distance, then multiplied by 100. Thus, a slope of 20 percent is a drop of 20 feet in 100 feet of horizontal distance.
- Slope** (in tables). Slope is great enough that special practices are required to ensure satisfactory performance of the soil for a specific use.
- Slow intake** (in tables). The slow movement of water into the soil.
- Slow refill** (in tables). The slow filling of ponds, resulting from restricted permeability in the soil.
- Small stones** (in tables). Rock fragments less than 3 inches (7.6 centimeters) in diameter. Small stones adversely affect the specified use of the soil.
- Soft bedrock.** Bedrock that can be excavated with trenching machines, backhoes, small rippers, and other equipment commonly used in construction.
- Soil.** A natural, three-dimensional body at the earth's surface. It is capable of supporting plants and has properties resulting from the integrated effect of climate and living matter acting on earthy parent material, as conditioned by relief over periods of time.
- Soil quality.** The capacity of a specific kind of soil to function within natural or managed ecosystem boundaries, to sustain plant and animal productivity, maintain or enhance water and air quality, and support human health and habitation.
- Soil separates.** Mineral particles less than 2 millimeters in equivalent diameter and ranging between specified size limits. The names and sizes, in millimeters, of separates recognized in the United States are as follows:
- | | |
|------------------------|-----------------|
| Very coarse sand | 2.0 to 1.0 |
| Coarse sand | 1.0 to 0.5 |
| Medium sand | 0.5 to 0.25 |
| Fine sand | 0.25 to 0.10 |
| Very fine sand | 0.10 to 0.05 |
| Silt | 0.05 to 0.002 |
| Clay | less than 0.002 |
- Solum.** The upper part of a soil profile, above the C horizon, in which the processes of soil formation are active. The solum in soil consists of the A, E, and B horizons. Generally, the characteristics of the material in these horizons are unlike those of the material below the solum. The living roots and plant and animal activities are largely confined to the solum.
- Stone line.** A concentration of coarse fragments in a soil. Generally, it is indicative of an old weathered surface. In a cross section, the line may be one fragment or more thick. It generally overlies material that weathered in place and is overlain by recent sediment of variable thickness.
- Stones.** Rock fragments 10 to 24 inches (25 to 60 centimeters) in diameter if rounded or 15 to 24 inches (38 to 60 centimeters) in length if flat.
- Stony.** Refers to a soil containing stones in numbers that interfere with or prevent tillage.

- Stripcropping.** Growing crops in a systematic arrangement of strips or bands that provide vegetative barriers to wind erosion and water erosion.
- Structure, soil.** The arrangement of primary soil particles into compound particles or aggregates. The principal forms of soil structure are—*platy* (laminated), *prismatic* (vertical axis of aggregates longer than horizontal), *columnar* (prisms with rounded tops), *blocky* (angular or subangular), and *granular*. *Structureless* soils are either *single grained* (each grain by itself, as in dune sand) or *massive* (the particles adhering without any regular cleavage, as in many hardpans).
- Stubble mulch.** Stubble or other crop residue left on the soil or partly worked into the soil. It protects the soil from wind erosion and water erosion after harvest, during preparation of a seedbed for the next crop, and during the early growing period of the new crop.
- Subsoil.** Technically, the B horizon; roughly, the part of the solum below plow depth.
- Subsoiling.** Tilling a soil below normal plow depth, ordinarily to shatter a hardpan or claypan.
- Substratum.** The part of the soil below the solum.
- Subsurface layer.** Any surface soil horizon (A, E, AB, or EB) below the surface layer.
- Summit.** The topographically highest position of a hillslope. It has a nearly level (planar or only slightly convex) surface.
- Surface layer.** The soil ordinarily moved in tillage, or its equivalent in uncultivated soil, ranging in depth from 4 to 10 inches (10 to 25 centimeters). Frequently designated as the “plow layer,” or the “Ap horizon.”
- Surface soil.** The A, E, AB, and EB horizons, considered collectively. It includes all subdivisions of these horizons.
- Talus.** Fragments of rock and other soil material accumulated by gravity at the foot of cliffs or steep slopes.
- Taxadjuncts.** Soils that cannot be classified in a series recognized in the classification system. Such soils are named for a series they strongly resemble and are designated as taxadjuncts to that series because they differ in ways too small to be of consequence in interpreting their use and behavior. Soils are recognized as taxadjuncts only when one or more of their characteristics are slightly outside the range defined for the family of the series for which the soils are named.
- Terminal moraine.** A belt of thick glacial drift that generally marks the termination of important glacial advances.
- Terrace.** An embankment, or ridge, constructed across sloping soils on the contour or at a slight angle to the contour. The terrace intercepts surface runoff so that water soaks into the soil or flows slowly to a prepared outlet. A terrace in a field generally is built so that the field can be farmed. A terrace intended mainly for drainage has a deep channel that is maintained in permanent sod.
- Terrace (geologic).** An old alluvial plain, ordinarily flat or undulating, bordering a river, a lake, or the sea.
- Texture, soil.** The relative proportions of sand, silt, and clay particles in a mass of soil. The basic textural classes, in order of increasing proportion of fine particles, are *sand*, *loamy sand*, *sandy loam*, *loam*, *silt loam*, *silt*, *sandy clay loam*, *clay loam*, *silty clay loam*, *sandy clay*, *silty clay*, and *clay*. The sand, loamy sand, and sandy loam classes may be further divided by specifying “coarse,” “fine,” or “very fine.”
- Thin layer** (in tables). Otherwise suitable soil material that is too thin for the specified use.
- Till plain.** An extensive area of nearly level to undulating soils underlain by glacial till.
- Tilth, soil.** The physical condition of the soil as related to tillage, seedbed preparation, seedling emergence, and root penetration.
- Toeslope.** The position that forms the gently inclined surface at the base of a hillslope. Toeslopes in profile are commonly gentle and linear and are

constructional surfaces forming the lower part of a hillslope continuum that grades to valley or closed-depression floors.

Topsoil. The upper part of the soil, which is the most favorable material for plant growth. It is ordinarily rich in organic matter and is used to topdress roadbanks, lawns, and land affected by mining.

Trace elements. Chemical elements, for example, zinc, cobalt, manganese, copper, and iron, in soils in extremely small amounts. They are essential to plant growth.

Umbric epipedon. A surface layer of mineral soil that has the same requirements as the mollic epipedon with respect to color, thickness, organic carbon content, consistence, structure, and phosphorus content, but that has a base saturation less than 50% when measured at pH 7.

Undifferentiated group, soil. A kind of map unit used in soil surveys comprised of two or more taxa components that are not consistently associated geographically and do not always occur together in the same map delineation. These taxa are included as the same named map unit because use and management are the same or very similar for common uses.

Upland. Land at a higher elevation, in general, than the alluvial plain or stream terrace; land above the lowlands along streams.

Valley fill. In glaciated regions, material deposited in stream valleys by glacial meltwater. In nonglaciated regions, alluvium deposited by heavily loaded streams.

Variiegation. Refers to patterns of contrasting colors assumed to be inherited from the parent material rather than to be the result of poor drainage.

Varve. A sedimentary layer or a lamina or sequence of laminae deposited in a body of still water within a year. Specifically, a thin pair of graded glaciolacustrine layers seasonally deposited, usually by meltwater streams, in a glacial lake or other body of still water in front of a glacier.

Water bars. Smooth, shallow ditches or depressional areas that are excavated at an angle across a sloping road. They are used to reduce the downward velocity of water and divert it off and away from the road surface. Water bars can easily be driven over if constructed properly.

Water quality. Used to describe the chemical, physical, and biological characteristics of water, usually in respect to its suitability for a particular use.

Weathering. All physical and chemical changes produced in rocks or other deposits at or near the earth's surface by atmospheric agents. These changes result in disintegration and decomposition of the material.

Well graded. Refers to soil material consisting of coarse grained particles that are well distributed over a wide range in size or diameter. Such soil normally can be easily increased in density and bearing properties by compaction. Contrasts with poorly graded soil.

Wetlands, Connecticut. Land, including submerged land, which consists of any of the soil types designated as poorly drained, very poorly drained, alluvial, and floodplain by the National Cooperative Soil Survey, as may be amended from time to time, of the Natural Resources Conservation Service of the United States Department of Agriculture.

Wetlands, federal. Land that has 1) a predominance of hydric soils; and 2) is inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and under normal circumstances does support, a prevalence of hydrophytic vegetation typically adapted for life in saturated soil conditions.

Wilting point (or permanent wilting point). The moisture content of soil, on an oven-dry basis, at which a plant (specifically a sunflower) wilts so much that it does not recover when placed in a humid, dark chamber.

Windthrow. The uprooting and tipping over of trees by the wind.

Tables

Table 1.--Temperature and Precipitation
(Recorded in the period 1961-90 at BRIDGEPORT WSO ARPT, CT0806)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
°F	°F	°F	°F	°F	Units	In	In	In	Days	In	
January-----	35.9	21.9	28.9	57	-1	9	3.24	1.45	4.77	6	8.4
February-----	37.6	23.2	30.4	58	2	11	3.01	1.75	4.14	6	8.0
March-----	46.2	30.9	38.6	69	11	68	3.75	2.03	5.26	6	4.0
April-----	56.7	39.8	48.3	81	25	256	3.75	1.99	5.31	6	0.4
May-----	66.8	49.9	58.3	86	36	569	3.93	1.93	5.66	7	0.0
June-----	76.0	59.1	67.6	92	46	828	3.46	1.36	5.23	5	0.0
July-----	81.6	65.6	73.6	95	53	1041	3.78	1.61	5.63	5	0.0
August-----	80.7	65.0	72.8	92	50	1018	3.25	1.72	4.59	5	0.0
September---	74.0	57.5	65.7	89	40	772	3.07	1.56	4.38	5	0.0
October-----	63.7	46.9	55.3	79	29	476	3.11	1.68	4.38	5	0.0
November----	52.8	38.1	45.5	71	21	194	3.81	1.89	5.49	6	0.6
December----	41.0	27.5	34.2	61	6	37	3.50	1.80	4.98	6	4.9
Yearly:											
Average---	59.4	43.8	51.6	---	---	---	---	---	---	---	---
Extreme---	100	-7	---	95	-2	---	---	---	---	---	---
Total-----	---	---	---	---	---	5279	41.67	33.28	49.63	68	26.3

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 1.—Temperature and Precipitation—Continued
 (Recorded in the period 1961-90 at COCKAPONSET RANGER ST, CT1488)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>	
January-----	37.6	15.8	26.7	57	-13	2	4.28	1.85	6.36	6	9.8
February----	39.4	18.1	28.7	60	-4	8	3.73	2.24	5.06	6	11.0
March-----	49.4	25.0	37.2	79	5	86	4.28	2.63	5.76	6	5.6
April-----	57.6	34.5	46.1	87	20	197	4.61	2.46	6.50	6	1.3
May-----	69.7	45.7	57.7	94	28	549	4.64	2.93	6.19	7	0.0
June-----	78.3	53.5	65.9	94	35	776	3.71	1.69	5.44	6	0.0
July-----	81.9	59.7	70.8	95	41	954	3.89	1.80	5.68	5	0.0
August-----	80.8	57.2	69.0	95	37	899	3.68	1.96	5.20	6	0.0
September---	72.6	46.7	59.6	88	29	553	4.01	2.32	5.51	6	0.0
October-----	62.5	34.8	48.6	82	18	279	4.17	1.99	6.05	5	0.1
November----	52.2	27.7	39.9	74	3	98	4.78	2.67	6.64	7	1.1
December----	38.0	18.3	28.2	62	-13	16	4.55	2.26	6.53	7	6.7
Yearly:											
Average---	60.0	36.4	48.2	---	---	---	---	---	---	---	---
Extreme---	98	-18	---	101	-15	---	---	---	---	---	---
Total-----	---	---	---	---	---	4419	50.32	42.61	54.81	73	35.7

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 1.--Temperature and Precipitation--Continued
 (Recorded in the period 1961-90 at FALLS VILLAGE, CT2658)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>	
January-----	33.6	11.8	22.7	59	-19	3	2.95	1.48	4.22	6	10.4
February-----	36.4	14.3	25.3	60	-15	8	2.73	1.51	3.80	6	9.5
March-----	46.3	24.0	35.2	76	0	50	3.07	1.74	4.25	6	6.4
April-----	59.1	33.1	46.1	85	15	211	3.64	2.14	4.99	7	1.7
May-----	71.7	43.8	57.8	90	25	548	4.00	2.37	5.46	8	0.1
June-----	79.3	52.6	65.9	93	34	778	4.26	2.12	6.12	7	0.0
July-----	83.8	57.5	70.6	96	41	950	3.66	1.91	5.19	6	0.0
August-----	81.4	56.1	68.8	93	37	892	4.38	2.65	5.94	7	0.0
September---	73.6	48.6	61.1	90	29	632	3.54	1.95	4.95	6	0.0
October-----	62.7	37.1	49.9	81	18	320	3.28	1.94	4.47	6	0.1
November----	50.1	29.4	39.8	73	10	98	3.68	2.32	4.90	7	1.2
December----	37.6	18.4	28.0	62	-10	14	3.42	1.90	4.76	6	10.3
Yearly:											
Average---	59.6	35.6	47.6	---	---	---	---	---	---	---	---
Extreme---	98	-30	---	96	-20	---	---	---	---	---	---
Total-----	---	---	---	---	---	4504	42.60	36.52	48.46	78	39.7

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 1.--Temperature and Precipitation--Continued
 (Recorded in the period 1961-90 at GROTON, CT3207)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>
January-----	36.9	18.5	27.7	58	-7	10	4.16	2.08	5.96	7	7.7
February-----	38.5	20.3	29.4	58	-4	12	3.83	2.27	5.23	6	6.8
March-----	46.6	28.5	37.5	68	9	57	4.34	2.73	5.79	7	3.2
April-----	56.4	37.0	46.7	79	21	212	4.28	2.47	5.89	6	0.3
May-----	66.0	46.6	56.3	86	33	505	3.96	2.25	5.48	7	0.0
June-----	75.0	55.7	65.4	91	41	761	3.47	1.49	5.15	6	0.0
July-----	80.7	62.1	71.4	94	50	973	3.30	1.55	4.81	5	0.0
August-----	79.9	61.3	70.6	92	46	950	3.71	1.96	5.24	5	0.0
September---	73.1	53.6	63.3	89	36	700	3.61	1.91	5.10	5	0.0
October-----	63.0	43.0	53.0	79	25	406	3.89	2.19	5.39	6	0.0
November----	52.9	34.8	43.9	70	16	162	4.94	2.61	6.98	7	0.6
December----	41.7	24.3	33.0	63	0	35	4.66	2.53	6.53	7	4.4
Yearly:											
Average---	59.2	40.5	49.8	---	---	---	---	---	---	---	---
Extreme---	99	-14	---	95	-9	---	---	---	---	---	---
Total-----	---	---	---	---	---	4783	48.12	41.31	54.21	74	23.0

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 1.--Temperature and Precipitation--Continued
 (Recorded in the period 1961-90 at HARTFORD BRAINARD FIELD, CT3451)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>	
January-----	34.7	16.2	25.4	60	-8	7	3.25	1.29	4.90	5	6.4
February-----	37.5	18.7	28.1	61	-4	12	2.98	1.58	4.22	5	7.8
March-----	46.9	27.6	37.2	74	8	66	3.35	1.95	4.59	6	5.8
April-----	58.6	36.9	47.8	85	21	247	3.88	2.24	5.34	6	0.8
May-----	70.0	47.2	58.6	91	31	576	4.01	2.15	5.64	7	0.0
June-----	78.4	56.5	67.5	95	42	824	3.48	2.01	4.78	6	0.0
July-----	83.5	62.2	72.9	97	48	1015	3.57	1.99	4.96	6	0.0
August-----	81.5	60.3	70.9	95	44	959	3.60	1.50	5.38	5	0.0
September---	74.4	51.6	63.0	92	33	689	3.63	1.97	5.09	5	0.0
October-----	63.3	40.5	51.9	83	23	374	3.56	1.94	5.00	5	0.0
November----	51.6	32.9	42.2	74	16	135	3.64	1.95	5.12	6	0.6
December----	38.9	21.9	30.4	64	-1	22	3.71	1.91	5.28	6	6.8
Yearly:											
Average---	59.9	39.4	49.7	---	---	---	---	---	---	---	---
Extreme---	102	-17	---	98	-9	---	---	---	---	---	---
Total-----	---	---	---	---	---	4926	42.65	34.87	48.00	68	28.4

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 1.--Temperature and Precipitation--Continued
 (Recorded in the period 1961-90 at MOUNT CARMEL, CT5077)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>	
January-----	36.3	18.2	27.3	59	-9	10	3.84	1.65	5.70	6	9.3
February----	38.8	20.2	29.5	60	-6	17	3.40	2.00	4.65	6	9.2
March-----	48.0	28.5	38.3	72	6	76	4.19	2.37	5.81	6	4.9
April-----	59.3	37.2	48.2	83	20	258	4.56	2.55	6.33	6	0.5
May-----	69.8	46.7	58.2	89	30	566	4.57	2.34	6.53	7	0.0
June-----	78.5	55.9	67.2	93	39	816	4.25	1.70	6.41	6	0.0
July-----	83.3	61.8	72.5	95	46	1009	4.17	1.76	6.22	6	0.0
August-----	81.7	60.7	71.2	93	43	967	3.92	1.84	5.71	6	0.0
September---	74.6	53.0	63.8	91	34	714	4.32	2.14	6.21	5	0.0
October-----	64.3	42.2	53.3	82	22	414	3.91	1.97	5.60	5	0.1
November----	52.7	34.2	43.5	73	14	159	4.39	2.29	6.23	7	1.2
December----	40.3	23.4	31.8	64	-2	30	4.11	2.15	5.82	6	6.4
Yearly:											
Average---	60.6	40.2	50.4	---	---	---	---	---	---	---	---
Extreme---	99	-17	---	96	-11	---	---	---	---	---	---
Total-----	---	---	---	---	---	5035	49.62	39.48	57.84	72	31.6

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 1.—Temperature and Precipitation—Continued
 (Recorded in the period 1961-90 at NORFOLK 2 SW, CT5445)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>
January-----	27.4	10.8	19.1	53	-14	2	3.74	2.04	5.24	7	21.7
February----	29.5	11.9	20.7	55	-12	4	3.88	2.33	5.27	7	22.6
March-----	39.6	21.5	30.5	70	-2	27	4.16	2.56	5.60	7	17.5
April-----	52.4	32.2	42.3	81	14	140	4.45	2.69	6.03	8	7.1
May-----	64.8	43.3	54.1	85	27	440	4.59	2.60	6.36	8	0.9
June-----	72.8	52.4	62.6	88	36	678	4.60	2.37	6.55	7	0.0
July-----	77.5	57.5	67.5	89	44	854	4.19	2.52	5.70	7	0.0
August-----	75.4	55.9	65.7	87	40	796	4.55	2.88	6.06	7	0.0
September---	67.6	48.3	58.0	84	31	540	4.13	2.28	5.77	6	0.0
October-----	56.4	37.7	47.0	76	21	245	3.95	2.43	5.31	6	0.9
November---	44.4	29.3	36.8	67	10	67	4.56	3.04	5.94	8	6.9
December---	32.0	17.2	24.6	58	-8	9	4.44	2.66	6.03	8	20.9
Yearly:											
Average---	53.3	34.8	44.1	---	---	---	---	---	---	---	---
Extreme---	92	-20	---	90	-16	---	---	---	---	---	---
Total-----	---	---	---	---	---	3802	51.24	43.84	58.11	86	98.6

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees. F).

Table 1.--Temperature and Precipitation--Continued
 (Recorded in the period 1961-90 at SHEPAUG DAM, CT7373)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>	
January-----	33.1	13.6	23.3	57	-16	4	3.52	1.66	5.12	6	14.7
February-----	35.7	15.5	25.6	59	-12	8	3.36	1.77	4.76	6	13.8
March-----	45.1	24.9	35.0	74	0	49	3.90	2.22	5.39	7	9.4
April-----	57.5	34.6	46.0	84	16	208	4.24	2.55	5.75	8	2.3
May-----	68.2	45.0	56.6	88	29	515	4.57	2.56	6.35	9	0.1
June-----	75.8	53.9	64.9	91	37	744	4.08	2.10	5.81	7	0.0
July-----	80.3	59.2	69.7	93	45	921	4.21	2.48	5.76	7	0.0
August-----	78.7	57.8	68.3	91	41	876	4.25	2.44	5.86	6	0.0
September---	71.5	50.6	61.1	88	32	629	4.06	2.07	5.79	6	0.0
October-----	61.8	40.0	50.9	81	22	345	3.95	2.11	5.57	6	0.4
November----	50.1	31.8	40.9	71	13	116	4.34	2.73	5.79	7	3.0
December----	37.6	20.2	28.9	62	-6	17	3.94	2.15	5.52	7	10.8
Yearly:											
Average---	57.9	37.3	47.6	---	---	---	---	---	---	---	---
Extreme---	95	-27	---	93	-17	---	---	---	---	---	---
Total-----	---	---	---	---	---	4433	48.41	39.62	55.96	82	54.4

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 1.--Temperature and Precipitation--Continued
(Recorded in the period 1961-90 at STORRS, CT8138)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>	
January-----	32.9	16.5	24.7	58	-8	8	3.93	1.65	5.86	6	10.1
February-----	35.4	18.7	27.1	60	-4	11	3.45	2.07	4.68	6	10.7
March-----	44.8	27.4	36.1	73	6	59	3.50	2.14	4.73	6	6.5
April-----	56.5	36.6	46.5	82	20	218	4.24	2.39	5.88	7	0.8
May-----	67.6	46.1	56.8	88	31	521	4.20	2.57	5.66	7	0.1
June-----	75.4	54.8	65.1	90	40	751	3.85	1.89	5.54	7	0.0
July-----	79.8	60.5	70.1	93	47	933	4.14	2.36	5.72	6	0.0
August-----	78.2	59.1	68.7	90	43	887	3.96	2.09	5.59	6	0.0
September---	71.2	51.4	61.3	87	34	638	3.94	2.15	5.51	5	0.0
October-----	61.4	41.3	51.4	80	24	360	4.14	2.40	5.68	5	0.2
November----	49.8	33.4	41.6	72	14	128	4.47	2.65	6.10	7	1.8
December----	37.3	21.9	29.6	63	-2	23	4.21	2.21	5.97	6	7.4
Yearly:											
Average---	57.5	39.0	48.2	---	---	---	---	---	---	---	---
Extreme---	98	-13	---	94	-10	---	---	---	---	---	---
Total-----	---	---	---	---	---	4538	48.02	39.30	53.09	74	37.7

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees. F)

Table 1.--Temperature and Precipitation--Continued
 (Recorded in the period 1961-90 at WEST THOMPSON LAKE, CT9388)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>°F</u>	<u>Units</u>	<u>In</u>	<u>In</u>	<u>In</u>	<u>Days</u>	<u>In</u>	
January-----	34.5	11.9	23.2	61	-14	7	3.93	1.74	5.81	6	10.3
February-----	36.9	14.2	25.6	62	-13	9	3.22	1.65	4.59	6	9.5
March-----	46.2	24.4	35.3	76	1	54	4.06	2.55	5.43	7	5.1
April-----	57.6	34.1	45.9	85	16	202	4.32	2.50	5.95	7	1.1
May-----	68.9	43.9	56.4	90	27	509	4.39	3.08	5.60	7	0.0
June-----	76.9	53.2	65.0	93	36	744	4.13	2.07	5.93	7	0.0
July-----	82.6	58.9	70.7	96	43	939	4.18	2.42	5.74	6	0.0
August-----	80.7	57.5	69.1	94	39	890	4.16	2.04	6.00	6	0.0
September---	73.0	48.3	60.6	90	31	610	3.79	1.92	5.42	5	0.0
October-----	62.5	36.7	49.6	82	18	303	4.28	2.51	5.86	5	0.1
November----	51.1	29.7	40.4	74	10	111	4.70	2.75	6.43	7	2.0
December----	38.8	18.4	28.6	64	-8	19	4.37	2.27	6.21	7	6.4
Yearly:											
Average---	59.1	35.9	47.5	---	---	---	---	---	---	---	---
Extreme---	100	-23	---	97	-18	---	---	---	---	---	---
Total-----	---	---	---	---	---	4397	49.54	43.79	55.12	76	34.4

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (40 degrees F).

Table 2.—Freeze Dates in Spring and Fall

(Recorded in the period 1961-90 at BRIDGEPORT WSO ARPT, CT0806)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 2	April 8	April 24
2 year in 10 later than--	March 29	April 5	April 20
5 year in 10 later than--	March 21	March 29	April 12
First freezing temperature in fall:			
1 yr in 10 earlier than--	November 16	November 2	October 17
2 yr in 10 earlier than--	November 21	November 7	October 23
5 yr in 10 earlier than--	December 1	November 18	November 2

Table 2.--Freeze Dates in Spring and Fall--Continued
 (Recorded in the period 1961-90 at COCKAPONSET RANGER ST, CT1488)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 23	May 16	May 18
2 year in 10 later than--	April 19	May 11	May 13
5 year in 10 later than--	April 12	May 1	May 5
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 4	October 3	September 14
2 yr in 10 earlier than--	October 10	October 6	September 17
5 yr in 10 earlier than--	October 21	October 11	September 21

Table 2.—Freeze Dates in Spring and Fall—Continued
 (Recorded in the period 1961-90 at FALLS VILLAGE, CT2658)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	May 3	May 14	June 2
2 year in 10 later than--	April 28	May 11	May 28
5 year in 10 later than--	April 18	May 4	May 18
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 6	September 23	September 12
2 yr in 10 earlier than--	October 11	September 28	September 17
5 yr in 10 earlier than--	October 20	October 9	September 26

Table 2.--Freeze Dates in Spring and Fall--Continued
 (Recorded in the period 1961-90 at GROTON, CT3207)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 8	April 21	May 4
2 year in 10 later than--	April 4	April 16	April 30
5 year in 10 later than--	March 28	April 8	April 22
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 28	October 15	October 4
2 yr in 10 earlier than--	November 4	October 21	October 9
5 yr in 10 earlier than--	November 16	November 1	October 18

Table 2.—Freeze Dates in Spring and Fall—Continued
 (Recorded in the period 1961-90 at HARTFORD BRAINARD FIELD, CT3451)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 11	April 25	May 11
2 year in 10 later than--	April 7	April 20	May 6
5 year in 10 later than--	March 29	April 12	April 27
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 17	October 9	September 26
2 yr in 10 earlier than--	October 25	October 14	October 1
5 yr in 10 earlier than--	November 7	October 24	October 11

Table 2.—Freeze Dates in Spring and Fall—Continued
 (Recorded in the period 1961-90 at MOUNT CARMEL, CT5077)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 12	April 27	May 15
2 year in 10 later than--	April 7	April 23	May 10
5 year in 10 later than--	March 29	April 15	May 1
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 17	October 10	September 27
2 yr in 10 earlier than--	October 25	October 15	October 1
5 yr in 10 earlier than--	November 7	October 25	October 10

Table 2.—Freeze Dates in Spring and Fall—Continued
 (Recorded in the period 1961-90 at NORFOLK 2 SW, CT5445)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 22	May 9	May 28
2 year in 10 later than--	April 19	May 5	May 23
5 year in 10 later than--	April 13	April 27	May 13
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 14	October 4	September 19
2 yr in 10 earlier than--	October 19	October 8	September 24
5 yr in 10 earlier than--	October 30	October 17	October 3

Table 2.--Freeze Dates in Spring and Fall--Continued
 (Recorded in the period 1961-90 at SHEPAUG DAM, CT7373)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 19	April 29	May 17
2 year in 10 later than--	April 15	April 26	May 14
5 year in 10 later than--	April 6	April 20	May 7
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 18	October 4	September 22
2 yr in 10 earlier than--	October 23	October 9	September 27
5 yr in 10 earlier than--	November 3	October 20	October 7

Table 2.—Freeze Dates in Spring and Fall—Continued
 (Recorded in the period 1961-90 at STORRS, CT8138)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 12	April 24	May 13
2 year in 10 later than--	April 8	April 21	May 8
5 year in 10 later than--	April 1	April 14	April 29
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 23	October 9	September 28
2 yr in 10 earlier than--	October 29	October 15	October 3
5 yr in 10 earlier than--	November 9	October 25	October 14

Table 2.--Freeze Dates in Spring and Fall--Continued
 (Recorded in the period 1961-90 at WEST THOMPSON LAKE, CT9388)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	April 27	May 8	May 28
2 year in 10 later than--	April 21	May 5	May 23
5 year in 10 later than--	April 10	April 28	May 13
First freezing temperature in fall:			
1 yr in 10 earlier than--	October 11	October 4	September 21
2 yr in 10 earlier than--	October 16	October 8	September 25
5 yr in 10 earlier than--	October 27	October 14	October 3

Table 3.—Growing Season

(Recorded for the period 1961-90 at BRIDGEPORT WSO ARPT, CT0806)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	234	212	182
8 years in 10	241	219	190
5 years in 10	253	233	203
2 years in 10	266	246	217
1 year in 10	273	253	224

Table 3.—Growing Season

(Recorded for the period 1961-90 at COCKAPONSET RANGER ST, CT1488)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	176	143	131
8 years in 10	183	151	134
5 years in 10	194	166	141
2 years in 10	206	180	147
1 year in 10	212	188	150

Table 3.—Growing Season—Continued

(Recorded for the period 1961-90 at FALLS VILLAGE, CT2658)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	164	138	109
8 years in 10	171	144	117
5 years in 10	184	157	130
2 years in 10	197	170	144
1 year in 10	204	176	151

Table 3.—Growing Season

(Recorded for the period 1961-90 at GROTON, CT3207)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	208	185	158
8 years in 10	216	192	165
5 years in 10	232	204	178
2 years in 10	247	216	191
1 year in 10	255	223	198

Table 3.—Growing Season—Continued

(Recorded for the period 1961-90 at HARTFORD BRAINARD FIELD, CT3451)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	193	171	143
8 years in 10	203	179	151
5 years in 10	221	194	164
2 years in 10	240	209	178
1 year in 10	250	217	185

Table 3.—Growing Season

(Recorded for the period 1961-90 at MOUNT CARMEL, CT5077)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	193	171	142
8 years in 10	202	178	149
5 years in 10	220	192	162
2 years in 10	239	205	176
1 year in 10	248	212	182

Table 3.—Growing Season—Continued

(Recorded for the period 1961-90 at NORFOLK 2 SW, CT5445)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	180	154	123
8 years in 10	187	161	130
5 years in 10	199	173	142
2 years in 10	212	185	155
1 year in 10	218	191	161

Table 3.—Growing Season

(Recorded for the period 1961-90 at SHEPAUG DAM, CT7373)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	190	161	137
8 years in 10	197	169	142
5 years in 10	210	183	153
2 years in 10	223	197	163
1 year in 10	230	204	169

Table 3.—Growing Season—Continued
 (Recorded for the period 1961-90 at STORRS, CT8138)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	204	175	146
8 years in 10	211	181	153
5 years in 10	224	193	166
2 years in 10	237	205	179
1 year in 10	243	212	186

Table 3.—Growing Season
 (Recorded for the period 1961-90 at WEST THOMPSON LAKE, CT9388)

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	178	151	126
8 years in 10	185	157	131
5 years in 10	199	168	142
2 years in 10	212	179	152
1 year in 10	219	185	158

Table 4.—Acreage and Proportionate Extent of the Soils

Map symbol	Soil name	Acres	Percent
2	Ridgebury fine sandy loam-----	8,683	0.3
3	Ridgebury, Leicester, and Whitman soils, extremely stony-----	154,932	4.8
4	Leicester fine sandy loam-----	3,274	0.1
5	Wilbraham silt loam-----	4,814	0.1
6	Wilbraham and Menlo soils, extremely stony-----	13,950	0.4
7	Mudgepond silt loam-----	1,050	*
8	Mudgepond and Alden soils, extremely stony-----	1,803	*
9	Scitico, Shaker, and Maybid soils-----	10,560	0.3
10	Raynham silt loam-----	1,923	*
12	Raypol silt loam-----	14,946	0.5
13	Walpole sandy loam-----	13,482	0.4
14	Fredon silt loam-----	866	*
15	Scarboro muck-----	16,168	0.5
16	Halsey silt loam-----	1,548	*
17	Timakwa and Natchaug soils-----	30,322	0.9
18	Catden and Freetown soils-----	40,679	1.3
20A	Ellington silt loam, 0 to 5 percent slopes-----	3,823	0.1
21A	Ninigret and Tisbury soils, 0 to 5 percent slopes-----	25,048	0.8
22A	Hero gravelly loam, 0 to 3 percent slopes-----	873	*
22B	Hero gravelly loam, 3 to 8 percent slopes-----	487	*
23A	Sudbury sandy loam, 0 to 5 percent slopes-----	11,590	0.4
24A	Deerfield loamy fine sand, 0 to 3 percent slopes-----	1,029	*
25A	Brancroft silt loam, 0 to 3 percent slopes-----	899	*
25B	Brancroft silt loam, 3 to 8 percent slopes-----	2,941	*
25C	Brancroft silt loam, 8 to 15 percent slopes-----	399	*
26A	Berlin silt loam, 0 to 3 percent slopes-----	735	*
26B	Berlin silt loam, 3 to 8 percent slopes-----	408	*
27A	Belgrade silt loam, 0 to 5 percent slopes-----	1,179	*
28A	Elmridge fine sandy loam, 0 to 3 percent slopes-----	3,800	0.1
28B	Elmridge fine sandy loam, 3 to 8 percent slopes-----	2,968	*
29A	Agawam fine sandy loam, 0 to 3 percent slopes-----	12,368	0.4
29B	Agawam fine sandy loam, 3 to 8 percent slopes-----	21,231	0.7
29C	Agawam fine sandy loam, 8 to 15 percent slopes-----	1,990	*
30A	Branford silt loam, 0 to 3 percent slopes-----	3,460	0.1
30B	Branford silt loam, 3 to 8 percent slopes-----	6,546	0.2
30C	Branford silt loam, 8 to 15 percent slopes-----	983	*
31A	Copake fine sandy loam, 0 to 3 percent slopes-----	774	*
31B	Copake fine sandy loam, 3 to 8 percent slopes-----	2,472	*
31C	Copake gravelly loam, 8 to 15 percent slopes-----	822	*
32A	Haven and Enfield soils, 0 to 3 percent slopes-----	11,987	0.4
32B	Haven and Enfield soils, 3 to 8 percent slopes-----	13,934	0.4
32C	Haven and Enfield soils, 8 to 15 percent slopes-----	1,095	*
33A	Hartford sandy loam, 0 to 3 percent slopes-----	3,464	0.1
33B	Hartford sandy loam, 3 to 8 percent slopes-----	3,855	0.1
34A	Merrimac sandy loam, 0 to 3 percent slopes-----	15,827	0.5
34B	Merrimac sandy loam, 3 to 8 percent slopes-----	19,101	0.6
34C	Merrimac sandy loam, 8 to 15 percent slopes-----	4,369	0.1
35A	Penwood loamy sand, 0 to 3 percent slopes-----	2,539	*
35B	Penwood loamy sand, 3 to 8 percent slopes-----	2,141	*
36A	Windsor loamy sand, 0 to 3 percent slopes-----	10,277	0.3
36B	Windsor loamy sand, 3 to 8 percent slopes-----	10,845	0.3
36C	Windsor loamy sand, 8 to 15 percent slopes-----	1,546	*
37A	Manchester gravelly sandy loam, 0 to 3 percent slopes-----	1,721	*
37C	Manchester gravelly sandy loam, 3 to 15 percent slopes-----	19,084	0.6
37E	Manchester gravelly sandy loam, 15 to 45 percent slopes-----	8,690	0.3
38A	Hinckley gravelly sandy loam, 0 to 3 percent slopes-----	5,607	0.2
38C	Hinckley gravelly sandy loam, 3 to 15 percent slopes-----	72,730	2.3
38E	Hinckley gravelly sandy loam, 15 to 45 percent slopes-----	33,382	1.0
39A	Groton gravelly sandy loam, 0 to 3 percent slopes-----	204	*
39C	Groton gravelly sandy loam, 3 to 15 percent slopes-----	1,214	*
39E	Groton gravelly sandy loam, 15 to 45 percent slopes-----	957	*
40A	Ludlow silt loam, 0 to 3 percent slopes-----	1,959	*

See footnote at end of table.

Table 4.--Acreage and Proportionate Extent of the Soils--Continued

Map symbol	Soil name	Acres	Percent
40B	Ludlow silt loam, 3 to 8 percent slopes-----	6,865	0.2
41B	Ludlow silt loam, 2 to 8 percent slopes, very stony-----	2,108	*
42C	Ludlow silt loam, 2 to 15 percent slopes, extremely stony-----	4,194	0.1
43A	Rainbow silt loam, 0 to 3 percent slopes-----	1,089	*
43B	Rainbow silt loam, 3 to 8 percent slopes-----	1,525	*
44B	Rainbow silt loam, 2 to 8 percent slopes, very stony-----	1,318	*
45A	Woodbridge fine sandy loam, 0 to 3 percent slopes-----	13,430	0.4
45B	Woodbridge fine sandy loam, 3 to 8 percent slopes-----	48,606	1.5
45C	Woodbridge fine sandy loam, 8 to 15 percent slopes-----	7,688	0.2
46B	Woodbridge fine sandy loam, 2 to 8 percent slopes, very stony-----	52,163	1.6
46C	Woodbridge fine sandy loam, 8 to 15 percent slopes, very stony-----	6,134	0.2
47C	Woodbridge fine sandy loam, 2 to 15 percent slopes, extremely stony-----	59,508	1.9
48B	Georgia and Amenia silt loams, 2 to 8 percent slopes-----	4,809	0.1
48C	Georgia and Amenia silt loams, 8 to 15 percent slopes-----	1,172	*
49B	Georgia and Amenia silt loams, 3 to 8 percent slopes, very stony-----	668	*
49C	Georgia and Amenia silt loams, 8 to 15 percent slopes, very stony-----	1,273	*
50A	Sutton fine sandy loam, 0 to 3 percent slopes-----	1,916	*
50B	Sutton fine sandy loam, 3 to 8 percent slopes-----	11,127	0.3
51B	Sutton fine sandy loam, 2 to 8 percent slopes, very stony-----	21,172	0.7
52C	Sutton fine sandy loam, 2 to 15 percent slopes, extremely stony-----	31,190	1.0
53A	Wapping very fine sandy loam, 0 to 3 percent slopes-----	669	*
53B	Wapping very fine sandy loam, 3 to 8 percent slopes-----	698	*
54B	Wapping very fine sandy loam, 2 to 8 percent slopes, very stony-----	655	*
55A	Watchaug fine sandy loam, 0 to 3 percent slopes-----	636	*
55B	Watchaug fine sandy loam, 3 to 8 percent slopes-----	1,895	*
56B	Watchaug fine sandy loam, 2 to 8 percent slopes, very stony-----	454	*
57B	Gloucester gravelly sandy loam, 3 to 8 percent slopes-----	2,260	*
57C	Gloucester gravelly sandy loam, 8 to 15 percent slopes-----	2,186	*
57D	Gloucester gravelly sandy loam, 15 to 25 percent slopes-----	456	*
58B	Gloucester gravelly sandy loam, 3 to 8 percent slopes, very stony-----	2,666	*
58C	Gloucester gravelly sandy loam, 8 to 15 percent slopes, very stony-----	3,282	0.1
59C	Gloucester gravelly sandy loam, 3 to 15 percent slopes, extremely stony--	4,767	0.1
59D	Gloucester gravelly sandy loam, 15 to 35 percent slopes, extremely stony--	3,057	*
60B	Canton and Charlton soils, 3 to 8 percent slopes-----	54,509	1.7
60C	Canton and Charlton soils, 8 to 15 percent slopes-----	24,403	0.8
60D	Canton and Charlton soils, 15 to 25 percent slopes-----	10,397	0.3
61B	Canton and Charlton soils, 3 to 8 percent slopes, very stony-----	64,355	2.0
61C	Canton and Charlton soils, 8 to 15 percent slopes, very stony-----	53,878	1.7
62C	Canton and Charlton soils, 3 to 15 percent slopes, extremely stony-----	92,372	2.9
62D	Canton and Charlton soils, 15 to 35 percent slopes, extremely stony-----	68,412	2.1
63B	Cheshire fine sandy loam, 3 to 8 percent slopes-----	11,276	0.4
63C	Cheshire fine sandy loam, 8 to 15 percent slopes-----	3,942	0.1
63D	Cheshire fine sandy loam, 15 to 25 percent slopes-----	1,446	*
64B	Cheshire fine sandy loam, 3 to 8 percent slopes, very stony-----	3,707	0.1
64C	Cheshire fine sandy loam, 8 to 15 percent slopes, very stony-----	1,621	*
65C	Cheshire fine sandy loam, 3 to 15 percent slopes, extremely stony-----	2,472	*
65D	Cheshire fine sandy loam, 15 to 35 percent slopes, extremely stony-----	467	*
66B	Narragansett silt loam, 2 to 8 percent slopes-----	6,004	0.2
66C	Narragansett silt loam, 8 to 15 percent slopes-----	998	*
67B	Narragansett silt loam, 3 to 8 percent slopes, very stony-----	2,756	*
67C	Narragansett silt loam, 8 to 15 percent slopes, very stony-----	615	*
68C	Narragansett silt loam, 3 to 15 percent slopes, extremely stony-----	1,990	*
68D	Narragansett silt loam, 15 to 25 percent slopes, extremely stony-----	1,998	*
69B	Yalesville fine sandy loam, 3 to 8 percent slopes-----	4,528	0.1
69C	Yalesville fine sandy loam, 8 to 15 percent slopes-----	3,530	0.1
70C	Branford-Holyoke complex, 3 to 15 percent slopes, very rocky-----	524	*
71C	Brookfield-Brimfield-Rock outcrop complex, 3 to 15 percent slopes-----	4,886	0.2
71E	Brookfield-Brimfield-Rock outcrop complex, 15 to 45 percent slopes-----	2,698	*
73C	Charlton-Chatfield complex, 3 to 15 percent slopes, very rocky-----	269,531	8.4
73E	Charlton-Chatfield complex, 15 to 45 percent slopes, very rocky-----	134,058	4.2
74C	Narragansett-Hollis complex, 3 to 15 percent slopes, very rocky-----	2,763	*
75C	Hollis-Chatfield-Rock outcrop complex, 3 to 15 percent slopes-----	75,037	2.3
75E	Hollis-Chatfield-Rock outcrop complex, 15 to 45 percent slopes-----	110,455	3.4

See footnote at end of table.

Table 4.--Acreage and Proportionate Extent of the Soils--Continued

Map symbol	Soil name	Acres	Percent
76E	Rock outcrop-Hollis complex, 3 to 45 percent slopes-----	10,453	0.3
76F	Rock outcrop-Hollis complex, 45 to 60 percent slopes-----	7,527	0.2
77C	Cheshire-Holyoke complex, 3 to 15 percent slopes, very rocky-----	12,446	0.4
77D	Cheshire-Holyoke complex, 15 to 35 percent slopes, very rocky-----	9,361	0.3
78C	Holyoke-Rock outcrop complex, 3 to 15 percent slopes-----	9,355	0.3
78E	Holyoke-Rock outcrop complex, 15 to 45 percent slopes-----	14,185	0.4
79E	Rock outcrop-Holyoke complex, 3 to 45 percent slopes-----	6,331	0.2
80B	Bernardston silt loam, 3 to 8 percent slopes-----	451	*
80C	Bernardston silt loam, 8 to 15 percent slopes-----	560	*
81C	Bernardston silt loam, 3 to 15 percent slopes, extremely stony-----	456	*
81D	Bernardston silt loam, 15 to 25 percent slopes, extremely stony-----	515	*
82B	Broadbrook silt loam, 3 to 8 percent slopes-----	5,717	0.2
82C	Broadbrook silt loam, 8 to 15 percent slopes-----	1,394	*
82D	Broadbrook silt loam, 15 to 25 percent slopes-----	671	*
83B	Broadbrook silt loam, 3 to 8 percent slopes, very stony-----	1,321	*
83C	Broadbrook silt loam, 8 to 15 percent slopes, very stony-----	647	*
84B	Paxton and Montauk fine sandy loams, 3 to 8 percent slopes-----	72,531	2.3
84C	Paxton and Montauk fine sandy loams, 8 to 15 percent slopes-----	31,139	1.0
84D	Paxton and Montauk fine sandy loams, 15 to 25 percent slopes-----	11,895	0.4
85B	Paxton and Montauk fine sandy loams, 3 to 8 percent slopes, very stony---	41,472	1.3
85C	Paxton and Montauk fine sandy loams, 8 to 15 percent slopes, very stony--	25,132	0.8
86C	Paxton and Montauk fine sandy loams, 3 to 15 percent slopes, extremely stony-----	27,302	0.9
86D	Paxton and Montauk fine sandy loams, 15 to 35 percent slopes, extremely stony-----	25,318	0.8
87B	Wethersfield loam, 3 to 8 percent slopes-----	19,779	0.6
87C	Wethersfield loam, 8 to 15 percent slopes-----	8,915	0.3
87D	Wethersfield loam, 15 to 25 percent slopes-----	5,380	0.2
88B	Wethersfield loam, 3 to 8 percent slopes, very stony-----	3,563	0.1
88C	Wethersfield loam, 8 to 15 percent slopes, very stony-----	2,264	*
89C	Wethersfield loam, 3 to 15 percent slopes, extremely stony-----	4,124	0.1
89D	Wethersfield loam, 15 to 35 percent slopes, extremely stony-----	3,899	0.1
90B	Stockbridge loam, 3 to 8 percent slopes-----	6,151	0.2
90C	Stockbridge loam, 8 to 15 percent slopes-----	4,900	0.2
90D	Stockbridge loam, 15 to 25 percent slopes-----	1,778	*
91B	Stockbridge loam, 3 to 8 percent slopes, very stony-----	171	*
91C	Stockbridge loam, 8 to 15 percent slopes, very stony-----	1,387	*
91D	Stockbridge loam, 15 to 35 percent slopes, very stony-----	1,689	*
92B	Nellis fine sandy loam, 3 to 8 percent slopes-----	1,566	*
92C	Nellis fine sandy loam, 8 to 15 percent slopes-----	584	*
92D	Nellis fine sandy loam, 15 to 25 percent slopes-----	68	*
93C	Nellis fine sandy loam, 3 to 15 percent slopes, very stony-----	492	*
94C	Farmington-Nellis complex, 3 to 15 percent slopes, very rocky-----	4,746	0.1
94E	Farmington-Nellis complex, 15 to 35 percent slopes, very rocky-----	2,302	*
95C	Farmington-Rock outcrop complex, 3 to 15 percent slopes-----	666	*
95E	Farmington-Rock outcrop complex, 15 to 45 percent slopes-----	802	*
96	Ipswich mucky peat-----	500	*
97	Pawcatuck mucky peat-----	840	*
98	Westbrook mucky peat-----	7,121	0.2
99	Westbrook mucky peat, low salt-----	4,108	0.1
100	Suncook loamy fine sand-----	4,050	0.1
101	Occum fine sandy loam-----	4,887	0.2
102	Pootatuck fine sandy loam-----	9,310	0.3
103	Rippowam fine sandy loam-----	21,028	0.7
104	Bash silt loam-----	3,708	0.1
105	Hadley silt loam-----	4,000	0.1
106	Winooski silt loam-----	3,669	0.1
107	Limerick and Lim soils-----	7,651	0.2
108	Saco silt loam-----	21,212	0.7
109	Fluvaquents-Udifluvents complex, frequently flooded-----	8,667	0.3
221A	Ninigret-Urban land complex, 0 to 5 percent slopes-----	2,288	*
224A	Deerfield-Urban land complex, 0 to 3 percent slopes-----	301	*
225B	Brancroft-Urban land complex, 0 to 8 percent slopes-----	1,362	*

See footnote at end of table.

Table 4.--Acreage and Proportionate Extent of the Soils--Continued

Map symbol	Soil name	Acres	Percent
226B	Berlin-Urban land complex, 0 to 8 percent slopes-----	441	*
228B	Elmridge-Urban land complex, 0 to 8 percent slopes-----	1,541	*
229B	Agawam-Urban land complex, 0 to 8 percent slopes-----	8,926	0.3
229C	Agawam-Urban land complex, 8 to 15 percent slopes-----	150	*
230B	Branford-Urban land complex, 0 to 8 percent slopes-----	2,403	*
230C	Branford-Urban land complex, 8 to 15 percent slopes-----	236	*
232B	Haven-Urban land complex, 0 to 8 percent slopes-----	2,967	*
234B	Merrimac-Urban land complex, 0 to 8 percent slopes-----	2,617	*
235B	Penwood-Urban land complex, 0 to 8 percent slopes-----	6,100	0.2
236B	Windsor-Urban land complex, 0 to 8 percent slopes-----	3,403	0.1
237A	Manchester-Urban land complex, 0 to 3 percent slopes-----	1,566	*
237C	Manchester-Urban land complex, 3 to 15 percent slopes-----	4,343	0.1
238A	Hinckley-Urban land complex, 0 to 3 percent slopes-----	966	*
238C	Hinckley-Urban land complex, 3 to 15 percent slopes-----	3,754	0.1
240B	Ludlow-Urban land complex, 0 to 8 percent slopes-----	2,080	*
243B	Rainbow-Urban land complex, 0 to 8 percent slopes-----	252	*
245B	Woodbridge-Urban land complex, 0 to 8 percent slopes-----	3,121	*
245C	Woodbridge-Urban land complex, 8 to 15 percent slopes-----	431	*
248B	Georgia-Urban land complex, 2 to 8 percent slopes-----	11	*
250B	Sutton-Urban land complex, 0 to 8 percent slopes-----	2,658	*
253B	Wapping-Urban land complex, 0 to 8 percent slopes-----	18	*
255B	Watchaug-Urban land complex, 0 to 8 percent slopes-----	262	*
260B	Charlton-Urban land complex, 3 to 8 percent slopes-----	8,905	0.3
260C	Charlton-Urban land complex, 8 to 15 percent slopes-----	3,884	0.1
260D	Charlton-Urban land complex, 15 to 25 percent slopes-----	816	*
263B	Cheshire-Urban land complex, 3 to 8 percent slopes-----	3,184	*
263C	Cheshire-Urban land complex, 8 to 15 percent slopes-----	2,274	*
266B	Narragansett-Urban land complex, 3 to 8 percent slopes-----	13	*
269B	Yalesville-Urban land complex, 3 to 8 percent slopes-----	686	*
269C	Yalesville-Urban land complex, 8 to 15 percent slopes-----	519	*
273C	Urban land-Charlton-Chatfield complex, rocky, 3 to 15 percent slopes-----	9,664	0.3
273E	Urban land-Charlton-Chatfield complex, rocky, 15 to 45 percent slopes-----	1,125	*
275C	Urban land-Chatfield complex, rocky, 3 to 15 percent slopes-----	1,286	*
275E	Urban land-Chatfield-Rock outcrop complex, 15 to 45 percent slopes-----	600	*
282B	Broadbrook-Urban land complex, 3 to 8 percent slopes-----	804	*
284B	Paxton-Urban land complex, 3 to 8 percent slopes-----	6,466	0.2
284C	Paxton-Urban land complex, 8 to 15 percent slopes-----	3,373	0.1
284D	Paxton-Urban land complex, 15 to 25 percent slopes-----	814	*
287B	Wethersfield-Urban land complex, 3 to 8 percent slopes-----	7,083	0.2
287C	Wethersfield-Urban land complex, 8 to 15 percent slopes-----	1,985	*
287D	Wethersfield-Urban land complex, 15 to 25 percent slopes-----	369	*
290B	Stockbridge-Urban land complex, 3 to 8 percent slopes-----	140	*
290C	Stockbridge-Urban land complex, 8 to 15 percent slopes-----	69	*
290D	Stockbridge-Urban land complex, 15 to 25 percent slopes-----	2	*
301	Beaches-Udipsamments complex, coastal-----	1,672	*
302	Dumps-----	2,274	*
303	Pits, quarries-----	2,099	*
304	Udorthents, loamy, very steep-----	2,216	*
305	Udorthents-Pits complex, gravelly-----	11,843	0.4
306	Udorthents-Urban land complex-----	120,353	3.7
307	Urban land-----	44,596	1.4
308	Udorthents, smoothed-----	20,687	0.6
309	Udorthents, flood control-----	753	*
310	Udorthents, periodically flooded-----	98	*
401C	Macomber-Taconic complex, 3 to 15 percent slopes, very rocky-----	2,042	*
402D	Taconic-Macomber-Rock outcrop complex, 15 to 25 percent slopes-----	428	*
403C	Taconic-Rock outcrop complex, 3 to 15 percent slopes-----	145	*
403E	Taconic-Rock outcrop complex, 15 to 45 percent slopes-----	787	*
403F	Taconic-Rock outcrop complex, 45 to 70 percent slopes-----	948	*
405C	Dummerston gravelly loam, 3 to 15 percent slopes, very stony-----	367	*
405E	Dummerston gravelly loam, 15 to 45 percent slopes, very stony-----	356	*
407C	Lanesboro loam, 3 to 15 percent slopes, very stony-----	318	*
407E	Lanesboro loam, 15 to 45 percent slopes, very stony-----	157	*

See footnote at end of table.

Table 4.--Acreage and Proportionate Extent of the Soils--Continued

Map symbol	Soil name	Acres	Percent
408C	Fullam silt loam, 3 to 15 percent slopes, very stony-----	953	*
409B	Brayton mucky silt loam, 0 to 8 percent slopes, very stony-----	371	*
412B	Bice fine sandy loam, 3 to 8 percent slopes-----	514	*
412C	Bice fine sandy loam, 8 to 15 percent slopes-----	260	*
412D	Bice fine sandy loam, 15 to 25 percent slopes-----	117	*
413C	Bice-Millsite complex, 3 to 15 percent slopes, very rocky-----	9,270	0.3
413E	Bice-Millsite complex, 15 to 45 percent slopes, very rocky-----	6,001	0.2
414	Fredon silt loam, cold-----	44	*
415C	Westminster-Millsite-Rock outcrop complex, 3 to 15 percent slopes-----	3,104	*
415E	Westminster-Millsite-Rock outcrop complex, 15 to 45 percent slopes-----	5,623	0.2
416E	Rock outcrop-Westminster complex, 8 to 45 percent slopes-----	844	*
416F	Rock outcrop-Westminster complex, 45 to 70 percent slopes-----	460	*
417B	Bice fine sandy loam, 3 to 8 percent slopes, very stony-----	1,391	*
417C	Bice fine sandy loam, 8 to 15 percent slopes, very stony-----	4,399	0.1
417D	Bice fine sandy loam, 15 to 25 percent slopes, very stony-----	3,736	0.1
418C	Schroon fine sandy loam, 2 to 15 percent slopes, very stony-----	2,237	*
420A	Schroon fine sandy loam, 0 to 3 percent slopes-----	11	*
420B	Schroon fine sandy loam, 3 to 8 percent slopes-----	112	*
421A	Ninigret fine sandy loam, cold, 0 to 3 percent slopes-----	100	*
423A	Sudbury sandy loam, cold, 0 to 3 percent slopes-----	69	*
424B	Shelburne fine sandy loam, 3 to 8 percent slopes-----	992	*
424C	Shelburne fine sandy loam, 8 to 15 percent slopes-----	419	*
424D	Shelburne fine sandy loam, 15 to 25 percent slopes-----	74	*
425B	Shelburne fine sandy loam, 3 to 8 percent slopes, very stony-----	1,283	*
425C	Shelburne fine sandy loam, 8 to 15 percent slopes, very stony-----	3,284	0.1
426D	Shelburne fine sandy loam, 15 to 35 percent slopes, extremely stony-----	1,666	*
427B	Ashfield fine sandy loam, 2 to 8 percent slopes, very stony-----	1,424	*
427C	Ashfield fine sandy loam, 8 to 15 percent slopes, very stony-----	3,641	0.1
428A	Ashfield fine sandy loam, 0 to 3 percent slopes-----	106	*
428B	Ashfield fine sandy loam, 3 to 8 percent slopes-----	454	*
428C	Ashfield fine sandy loam, 8 to 15 percent slopes-----	104	*
429A	Agawam fine sandy loam, cold, 0 to 3 percent slopes-----	7	*
429B	Agawam fine sandy loam, cold, 3 to 8 percent slopes-----	63	*
429C	Agawam fine sandy loam, cold, 8 to 15 percent slopes-----	26	*
433	Moosilauke sandy loam-----	107	*
434A	Merrimac sandy loam, cold, 0 to 3 percent slopes-----	18	*
434B	Merrimac sandy loam, cold, 3 to 8 percent slopes-----	138	*
434C	Merrimac sandy loam, cold, 8 to 15 percent slopes-----	37	*
435	Scarboro muck, cold-----	96	*
436	Halsey silt loam, cold-----	138	*
437	Wonsqueak mucky peat-----	1,132	*
438	Bucksport muck-----	1,768	*
440A	Boscawen gravelly sandy loam, 0 to 3 percent slopes-----	54	*
440C	Boscawen gravelly sandy loam, 3 to 15 percent slopes-----	578	*
440E	Boscawen gravelly sandy loam, 15 to 45 percent slopes-----	363	*
442	Brayton loam-----	147	*
443	Brayton-Loonmeadow complex, extremely stony-----	3,877	0.1
448B	Hogansburg loam, 3 to 8 percent slopes-----	114	*
449B	Hogansburg loam, 3 to 8 percent slopes, very stony-----	61	*
449C	Hogansburg loam, 8 to 15 percent slopes, very stony-----	176	*
450B	Pyrities loam, 3 to 8 percent slopes-----	211	*
450C	Pyrities loam, 8 to 15 percent slopes-----	85	*
450D	Pyrities loam, 15 to 25 percent slopes-----	8	*
451B	Pyrities loam, 3 to 8 percent slopes, very stony-----	17	*
451C	Pyrities loam, 8 to 15 percent slopes, very stony-----	86	*
451D	Pyrities loam, 15 to 25 percent slopes, very stony-----	6	*
457	Mudgepond silt loam, cold-----	16	*
458	Mudgepond and Alden soils, extremely stony, cold-----	131	*
501	Ondawa fine sandy loam-----	14	*
503	Rumney fine sandy loam-----	131	*
508	Medomak silt loam-----	210	*
W	Water-----	348,652	10.9
	Total-----	3,211,700	100.0

Table 5.—Prime and other Important Farmland

(Only the soils considered prime or important farmland are listed. Urban or built-up areas of the soils listed are not considered prime or important farmland. If a soil is prime or important farmland only under certain conditions, the conditions are specified in parentheses after the soil name.)

Map symbol	Map unit name	Farmland Classification
20A	Ellington silt loam, 0 to 5 percent slopes	All areas are prime farmland
21A	Ninigret and Tisbury soils, 0 to 5 percent slopes	All areas are prime farmland
22A	Hero gravelly loam, 0 to 3 percent slopes	All areas are prime farmland
22B	Hero gravelly loam, 3 to 8 percent slopes	All areas are prime farmland
23A	Sudbury sandy loam, 0 to 5 percent slopes	All areas are prime farmland
26A	Berlin silt loam, 0 to 3 percent slopes	All areas are prime farmland
27A	Belgrade silt loam, 0 to 5 percent slopes	All areas are prime farmland
28A	Elmridge fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
28B	Elmridge fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
29A	Agawam fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
29B	Agawam fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
30A	Branford silt loam, 0 to 3 percent slopes	All areas are prime farmland
30B	Branford silt loam, 3 to 8 percent slopes	All areas are prime farmland
31A	Copake fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
31B	Copake fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
32A	Haven and Enfield soils, 0 to 3 percent slopes	All areas are prime farmland
32B	Haven and Enfield soils, 3 to 8 percent slopes	All areas are prime farmland
33A	Hartford sandy loam, 0 to 3 percent slopes	All areas are prime farmland
33B	Hartford sandy loam, 3 to 8 percent slopes	All areas are prime farmland
34A	Merrimac sandy loam, 0 to 3 percent slopes	All areas are prime farmland
34B	Merrimac sandy loam, 3 to 8 percent slopes	All areas are prime farmland
40A	Ludlow silt loam, 0 to 3 percent slopes	All areas are prime farmland
40B	Ludlow silt loam, 3 to 8 percent slopes	All areas are prime farmland
43A	Rainbow silt loam, 0 to 3 percent slopes	All areas are prime farmland
43B	Rainbow silt loam, 3 to 8 percent slopes	All areas are prime farmland
45A	Woodbridge fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
45B	Woodbridge fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
48B	Georgia and Amenia silt loams, 2 to 8 percent slopes	All areas are prime farmland
50A	Sutton fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
50B	Sutton fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
53A	Wapping very fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
53B	Wapping very fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
55A	Watchaug fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
55B	Watchaug fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
57B	Gloucester gravelly sandy loam, 3 to 8 percent slopes	All areas are prime farmland
60B	Canton and Charlton soils, 3 to 8 percent slopes	All areas are prime farmland
63B	Cheshire fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
66B	Narragansett silt loam, 2 to 8 percent slopes	All areas are prime farmland
69B	Yalesville fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
80B	Bernardston silt loam, 3 to 8 percent slopes	All areas are prime farmland
82B	Broadbrook silt loam, 3 to 8 percent slopes	All areas are prime farmland
84B	Paxton and Montauk fine sandy loams, 3 to 8 percent slopes	All areas are prime farmland
87B	Wethersfield loam, 3 to 8 percent slopes	All areas are prime farmland
90B	Stockbridge loam, 3 to 8 percent slopes	All areas are prime farmland
92B	Nellis fine sandy loam, 3 to 8 percent slopes	All areas are prime farmland
101	Occum fine sandy loam	All areas are prime farmland
102	Pootatuck fine sandy loam	All areas are prime farmland
105	Hadley silt loam	All areas are prime farmland
106	Winooski silt loam	All areas are prime farmland
420A	Schroon fine sandy loam, 0 to 3 percent slopes	All areas are prime farmland
421A	Ninigret fine sandy loam, cold, 0 to 3 percent slopes	All areas are prime farmland
423A	Sudbury sandy loam, cold, 0 to 3 percent slopes	All areas are prime farmland
429A	Agawam fine sandy loam, cold, 0 to 3 percent slopes	All areas are prime farmland
429B	Agawam fine sandy loam, cold, 3 to 8 percent slopes	All areas are prime farmland
434A	Merrimac sandy loam, cold, 0 to 3 percent slopes	All areas are prime farmland
434B	Merrimac sandy loam, cold, 3 to 8 percent slopes	All areas are prime farmland
448B	Hogansburg loam, 3 to 8 percent slopes	All areas are prime farmland
450B	Pyrities loam, 3 to 8 percent slopes	All areas are prime farmland
501	Ondawa fine sandy loam	All areas are prime farmland
2	Ridgebury fine sandy loam	Farmland of statewide importance

Table 5.—Prime and other Important Farmland—Continued

Map symbol	Map unit name	Farmland Classification
4	Leicester fine sandy loam	Farmland of statewide importance
5	Wilbraham silt loam	Farmland of statewide importance
7	Mudgepond silt loam	Farmland of statewide importance
9	Scitico, Shaker, and Maybid soils	Farmland of statewide importance
10	Raynham silt loam	Farmland of statewide importance
12	Raypol silt loam	Farmland of statewide importance
13	Walpole sandy loam	Farmland of statewide importance
14	Fredon silt loam	Farmland of statewide importance
24A	Deerfield loamy fine sand, 0 to 3 percent slopes	Farmland of statewide importance
25A	Brancroft silt loam, 0 to 3 percent slopes	Farmland of statewide importance
25B	Brancroft silt loam, 3 to 8 percent slopes	Farmland of statewide importance
25C	Brancroft silt loam, 8 to 15 percent slopes	Farmland of statewide importance
26B	Berlin silt loam, 3 to 8 percent slopes	Farmland of statewide importance
29C	Agawam fine sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
30C	Branford silt loam, 8 to 15 percent slopes	Farmland of statewide importance
31C	Copake gravelly loam, 8 to 15 percent slopes	Farmland of statewide importance
32C	Haven and Enfield soils, 8 to 15 percent slopes	Farmland of statewide importance
34C	Merrimac sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
35A	Penwood loamy sand, 0 to 3 percent slopes	Farmland of statewide importance
35B	Penwood loamy sand, 3 to 8 percent slopes	Farmland of statewide importance
36A	Windsor loamy sand, 0 to 3 percent slopes	Farmland of statewide importance
36B	Windsor loamy sand, 3 to 8 percent slopes	Farmland of statewide importance
36C	Windsor loamy sand, 8 to 15 percent slopes	Farmland of statewide importance
37A	Manchester gravelly sandy loam, 0 to 3 percent slopes	Farmland of statewide importance
37C	Manchester gravelly sandy loam, 3 to 15 percent slopes	Farmland of statewide importance
38A	Hinckley gravelly sandy loam, 0 to 3 percent slopes	Farmland of statewide importance
38C	Hinckley gravelly sandy loam, 3 to 15 percent slopes	Farmland of statewide importance
39A	Groton gravelly sandy loam, 0 to 3 percent slopes	Farmland of statewide importance
39C	Groton gravelly sandy loam, 3 to 15 percent slopes	Farmland of statewide importance
45C	Woodbridge fine sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
48C	Georgia and Amenia silt loams, 8 to 15 percent slopes	Farmland of statewide importance
57C	Gloucester gravelly sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
60C	Canton and Charlton soils, 8 to 15 percent slopes	Farmland of statewide importance
63C	Cheshire fine sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
66C	Narragansett silt loam, 8 to 15 percent slopes	Farmland of statewide importance
69C	Yalesville fine sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
80C	Bernardston silt loam, 8 to 15 percent slopes	Farmland of statewide importance
82C	Broadbrook silt loam, 8 to 15 percent slopes	Farmland of statewide importance
84C	Paxton and Montauk fine sandy loams, 8 to 15 percent slopes	Farmland of statewide importance
87C	Wethersfield loam, 8 to 15 percent slopes	Farmland of statewide importance
90C	Stockbridge loam, 8 to 15 percent slopes	Farmland of statewide importance
92C	Nellis fine sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
100	Suncook loamy fine sand	Farmland of statewide importance
103	Rippowam fine sandy loam	Farmland of statewide importance
104	Bash silt loam	Farmland of statewide importance
107	Limerick and Lim soils	Farmland of statewide importance
412B	Bice fine sandy loam, 3 to 8 percent slopes	Farmland of statewide importance
412C	Bice fine sandy loam, 8 to 15 percent slopes	Farmland of statewide importance
414	Fredon silt loam, cold	Farmland of statewide importance
420B	Schroon fine sandy loam, 3 to 8 percent slopes	Farmland of statewide importance
429C	Agawam fine sandy loam, cold, 8 to 15 percent slopes	Farmland of statewide importance
433	Moosilauke sandy loam	Farmland of statewide importance
434C	Merrimac sandy loam, cold, 8 to 15 percent slopes	Farmland of statewide importance
440A	Boscawen gravelly sandy loam, 0 to 3 percent slopes	Farmland of statewide importance
440C	Boscawen gravelly sandy loam, 3 to 15 percent slopes	Farmland of statewide importance
450C	Pyrities loam, 8 to 15 percent slopes	Farmland of statewide importance
457	Mudgepond silt loam, cold	Farmland of statewide importance
503	Rumney fine sandy loam	Farmland of statewide importance

Table 6.—Non-Irrigated Yields by Map Unit Component

(Yields are those that can be expected under a high level of management. They are for nonirrigated areas. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil.)

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
2: Ridgebury-----	4w	---	---	2.00	1.50	---
3: Ridgebury-----	7s	---	---	---	0.50	---
Leicester-----	7s	---	---	---	0.50	---
Whitman-----	7s	---	---	---	1.00	---
4: Leicester-----	4w	---	---	---	3.50	---
5: Wilbraham-----	4w	---	---	---	1.50	---
6: Wilbraham-----	7s	---	---	---	---	---
Menlo-----	7s	---	---	---	---	---
7: Mudgepond-----	4w	---	34.00	---	4.50	---
8: Mudgepond-----	7s	---	---	---	1.50	---
Alden-----	7s	---	---	---	1.50	---
9: Scitico-----	4w	---	35.00	---	5.50	---
Shaker-----	4w	---	32.00	---	4.00	---
Maybid-----	6w	---	---	---	6.00	---
10: Raynham-----	4w	---	33.00	---	6.00	---
12: Raypol-----	4w	---	27.00	---	2.50	---
13: Walpole-----	4w	---	25.00	---	2.00	---
14: Fredon-----	4w	---	32.00	---	4.00	---
15: Scarboro-----	5w	---	---	---	---	---
16: Halsey-----	5w	---	---	---	1.00	---
17: Timakwa-----	5w	---	---	---	4.50	---
Natchaug-----	5w	---	---	---	5.00	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
18: Catden-----	5w	---	---	---	5.00	---
Freetown-----	5w	---	---	---	5.00	---
20A: Ellington-----	2w	161.00	31.00	3.50	---	---
21A: Ninigret-----	2w	158.00	30.00	3.50	---	1,850.00
Tisbury-----	2w	167.00	33.00	4.50	---	2,000.00
22A: Hero-----	2w	155.00	29.00	3.50	3.50	---
22B: Hero-----	2w	153.00	29.00	4.00	3.50	---
23A: Sudbury-----	2w	141.00	26.00	2.50	---	---
24A: Deerfield-----	2w	141.00	26.00	2.50	---	1,550.00
25A: Brancroft-----	2w	180.00	35.00	5.00	5.50	---
25B: Brancroft-----	2e	177.00	35.00	4.50	5.00	---
25C: Brancroft-----	3e	---	33.00	4.00	4.50	---
26A: Berlin-----	2w	180.00	35.00	5.00	5.50	---
26B: Berlin-----	2e	178.00	35.00	4.50	5.50	---
27A: Belgrade-----	2w	180.00	35.00	5.00	6.00	---
28A: Elmridge-----	2w	176.00	35.00	4.50	5.00	2,100.00
28B: Elmridge-----	2w	172.00	34.00	4.50	5.00	2,100.00
29A: Agawam-----	1	156.00	30.00	---	---	1,800.00
29B: Agawam-----	2e	153.00	29.00	---	---	1,750.00
29C: Agawam-----	3e	141.00	28.00	---	---	---
30A: Branford-----	1	161.00	31.00	---	---	---
30B: Branford-----	2e	158.00	30.00	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
30C: Branford-----	3e	153.00	29.00	---	---	---
31A: Copake-----	1	164.00	32.00	4.00	4.00	---
31B: Copake-----	2e	161.00	31.00	3.50	4.00	---
31C: Copake-----	3e	155.00	29.00	3.50	3.50	---
32A: Haven-----	1	156.00	30.00	---	---	1,850.00
Enfield-----	1	169.00	33.00	---	---	2,050.00
32B: Haven-----	2e	155.00	29.00	---	---	1,800.00
Enfield-----	2e	166.00	32.00	---	---	2,000.00
32C: Haven-----	3e	150.00	28.00	---	---	---
Enfield-----	3e	161.00	31.00	---	---	---
33A: Hartford-----	1	145.00	27.00	---	---	1,600.00
33B: Hartford-----	2e	144.00	26.00	---	---	1,600.00
34A: Merrimac-----	1	142.00	26.00	---	2.50	1,550.00
34B: Merrimac-----	2e	141.00	26.00	---	2.50	1,550.00
34C: Merrimac-----	3e	138.00	25.00	---	2.00	---
35A: Penwood-----	2s	139.00	25.00	---	---	1,500.00
35B: Penwood-----	2s	138.00	25.00	---	---	1,450.00
36A: Windsor-----	2s	134.00	24.00	---	2.00	1,400.00
36B: Windsor-----	2s	133.00	24.00	---	---	1,400.00
36C: Windsor-----	3e	131.00	23.00	---	1.50	---
37A: Manchester-----	3s	135.00	24.00	---	2.00	1,450.00
37C: Manchester-----	4e	133.00	24.00	---	1.50	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
37E: Manchester-----	7e	---	---	---	---	---
38A: Hinckley-----	3s	135.00	24.00	---	2.00	---
38C: Hinckley-----	4e	133.00	24.00	---	1.50	---
38E: Hinckley-----	6e	---	---	---	0.50	---
39A: Groton-----	2s	137.00	25.00	2.50	2.00	---
39C: Groton-----	3e	134.00	24.00	2.00	1.50	---
39E: Groton-----	6e	---	---	---	0.50	---
40A: Ludlow-----	2w	145.00	27.00	3.00	2.50	---
40B: Ludlow-----	2e	144.00	26.00	2.50	2.50	---
41B: Ludlow-----	6s	---	---	---	1.00	---
42C: Ludlow-----	7s	---	---	---	1.00	---
43A: Rainbow-----	2w	146.00	27.00	3.00	2.50	1,650.00
43B: Rainbow-----	2e	144.00	27.00	2.50	2.50	---
44B: Rainbow-----	6s	---	---	---	1.00	---
45A: Woodbridge-----	2w	147.00	27.00	3.00	3.00	---
45B: Woodbridge-----	2w	145.00	27.00	3.00	2.50	---
45C: Woodbridge-----	3e	142.00	26.00	2.50	2.50	---
46B: Woodbridge-----	6s	---	---	---	1.00	---
46C: Woodbridge-----	6s	---	---	---	1.00	---
47C: Woodbridge-----	7s	---	---	---	1.00	---
48B: Georgia-----	2e	176.00	35.00	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
48B: Amenia-----	2e	175.00	35.00	---	---	---
48C: Georgia-----	3e	168.00	30.00	---	---	---
Amenia-----	3e	167.00	29.00	---	---	---
49B: Georgia-----	6s	---	---	---	1.50	---
Amenia-----	6s	---	---	---	1.50	---
49C: Georgia-----	6s	---	---	---	1.50	---
Amenia-----	6s	---	---	---	1.50	---
50A: Sutton-----	2w	---	---	---	---	---
50B: Sutton-----	2w	---	---	---	---	---
51B: Sutton-----	6s	---	---	---	---	---
52C: Sutton-----	7s	---	---	---	---	---
53A: Wapping-----	2w	168.00	31.00	4.00	4.50	---
53B: Wapping-----	2e	166.00	31.00	4.00	3.50	---
54B: Wapping-----	6s	---	---	---	1.50	---
55A: Watchaug-----	2w	165.00	31.00	4.00	4.00	---
55B: Watchaug-----	2w	162.00	31.00	3.50	4.00	---
56B: Watchaug-----	6s	---	---	---	1.50	---
57B: Gloucester-----	2s	138.00	25.00	2.50	2.00	---
57C: Gloucester-----	3e	135.00	24.00	2.00	2.00	---
57D: Gloucester-----	4e	---	---	2.00	1.50	---
58B: Gloucester-----	6s	---	---	---	0.50	---
58C: Gloucester-----	6s	---	---	---	0.50	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
59C: Gloucester-----	7s	---	---	---	0.50	---
59D: Gloucester-----	7s	---	---	---	---	---
60B: Canton-----	2e	144.00	27.00	3.00	---	1,600.00
Charlton-----	2e	152.00	29.00	3.50	---	1,750.00
60C: Canton-----	3e	141.00	26.00	2.50	2.50	---
Charlton-----	3e	141.00	28.00	3.00	3.00	---
60D: Canton-----	4e	133.00	24.00	2.00	1.50	---
Charlton-----	4e	139.00	25.00	2.50	2.00	---
61B: Canton-----	6s	---	---	---	1.00	---
Charlton-----	6s	---	---	---	1.00	---
61C: Canton-----	6s	---	---	---	1.00	---
Charlton-----	6s	---	---	---	1.00	---
62C: Canton-----	7s	---	---	---	---	---
Charlton-----	7s	---	---	---	---	---
62D: Canton-----	7s	---	---	---	---	---
Charlton-----	7s	---	---	---	---	---
63B: Cheshire-----	2e	160.00	31.00	3.50	---	1,900.00
63C: Cheshire-----	3e	155.00	29.00	3.50	3.50	---
63D: Cheshire-----	4e	143.00	26.00	2.50	2.50	---
64B: Cheshire-----	6s	---	---	---	1.00	---
64C: Cheshire-----	6s	---	---	---	1.00	---
65C: Cheshire-----	7s	---	---	---	1.00	---
65D: Cheshire-----	7s	---	---	---	1.00	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
66B: Narragansett-----	2e	150.00	28.00	3.50	3.00	---
66C: Narragansett-----	3e	146.00	27.00	3.00	2.50	---
67B: Narragansett-----	6s	---	---	---	1.00	---
67C: Narragansett-----	6s	---	---	---	1.00	---
68C: Narragansett-----	7s	---	---	---	1.00	---
68D: Narragansett-----	7s	---	---	---	---	---
69B: Yalesville-----	2e	150.00	28.00	3.00	3.00	---
69C: Yalesville-----	3e	145.00	27.00	3.00	2.50	---
70C: Branford-----	6s	---	---	---	3.50	---
Holyoke-----	6s	---	---	---	0.50	---
71C: Brookfield-----	6s	---	---	---	1.00	---
Brimfield-----	6s	---	---	---	---	---
Rock outcrop-----	---	---	---	---	---	---
71E: Brookfield-----	7s	---	---	---	---	---
Brimfield-----	7s	---	---	---	---	---
Rock outcrop-----	---	---	---	---	---	---
73C: Charlton-----	6s	---	---	---	1.00	---
Chatfield-----	6s	---	---	---	0.50	---
73E: Charlton-----	7s	---	---	---	0.50	---
Chatfield-----	7s	---	---	---	0.50	---
74C: Narragansett-----	6s	---	---	---	3.00	---
Hollis-----	6s	---	---	---	0.50	---
75C: Hollis-----	6s	---	---	---	0.50	---
Chatfield-----	6s	---	---	---	1.00	---
Rock outcrop-----	8	---	---	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
75E:						
Hollis-----	7s	---	---	---	0.50	---
Chatfield-----	7s	---	---	---	0.50	---
Rock outcrop-----	8	---	---	---	---	---
76E:						
Rock outcrop-----	8	---	---	---	---	---
Hollis-----	7s	---	---	---	---	---
76F:						
Rock outcrop-----	8	---	---	---	---	---
Hollis-----	7s	---	---	---	---	---
77C:						
Cheshire-----	6s	---	---	---	1.00	---
Holyoke-----	6s	---	---	---	0.50	---
77D:						
Cheshire-----	7s	---	---	---	1.00	---
Holyoke-----	7s	---	---	---	0.50	---
78C:						
Holyoke-----	6s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
78E:						
Holyoke-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
79E:						
Rock outcrop-----	8	---	---	---	---	---
Holyoke-----	7s	---	---	---	---	---
80B:						
Bernardston-----	2e	---	---	2.50	2.50	---
80C:						
Bernardston-----	3e	---	---	2.00	2.00	---
81C:						
Bernardston-----	7s	---	---	---	---	---
81D:						
Bernardston-----	7s	---	---	---	---	---
82B:						
Broadbrook-----	2e	143.00	26.00	---	---	---
82C:						
Broadbrook-----	3e	140.00	26.00	---	---	---
82D:						
Broadbrook-----	4e	---	---	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
83B: Broadbrook-----	6s	---	---	---	1.00	---
83C: Broadbrook-----	6s	---	---	---	1.00	---
84B: Paxton-----	2e	139.00	25.00	2.50	2.00	---
Montauk-----	2e	138.00	25.00	2.50	2.00	---
84C: Paxton-----	3e	136.00	24.00	2.00	2.00	---
Montauk-----	3e	135.00	24.00	2.00	2.00	---
84D: Paxton-----	4e	130.00	23.00	2.00	1.50	---
Montauk-----	4e	129.00	23.00	2.00	1.50	---
85B: Paxton-----	6s	---	---	---	0.50	---
Montauk-----	6s	---	---	---	0.50	---
85C: Paxton-----	6s	---	---	---	0.50	---
Montauk-----	6s	---	---	---	0.50	---
86C: Paxton-----	7s	---	---	---	0.50	---
Montauk-----	7s	---	---	---	0.50	---
86D: Paxton-----	7s	---	---	---	0.50	---
Montauk-----	7s	---	---	---	0.50	---
87B: Wethersfield-----	2e	142.00	26.00	2.50	2.50	---
87C: Wethersfield-----	3e	139.00	25.00	2.50	2.00	---
87D: Wethersfield-----	4e	132.00	23.00	2.00	1.50	---
88B: Wethersfield-----	6s	---	---	1.50	0.50	---
88C: Wethersfield-----	6s	---	---	1.50	0.50	---
89C: Wethersfield-----	7s	---	---	1.50	0.50	---
89D: Wethersfield-----	7s	---	---	1.50	0.50	---
90B: Stockbridge-----	2e	176.00	35.00	4.50	5.00	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
90C: Stockbridge-----	3e	169.00	33.00	4.00	4.50	---
90D: Stockbridge-----	4e	---	---	4.00	3.50	---
91B: Stockbridge-----	6s	---	---	---	1.50	---
91C: Stockbridge-----	6s	---	---	---	1.50	---
91D: Stockbridge-----	7s	---	---	---	1.50	---
92B: Nellis-----	2e	161.00	31.00	3.50	4.00	---
92C: Nellis-----	3e	156.00	30.00	3.50	3.50	---
92D: Nellis-----	4e	143.00	26.00	2.50	2.50	---
93C: Nellis-----	6s	---	---	---	1.00	---
94C: Farmington-----	6s	---	---	---	0.50	---
Nellis-----	6s	---	---	---	3.50	---
94E: Farmington-----	7s	---	---	---	0.50	---
Nellis-----	7s	---	---	---	1.50	---
95C: Farmington-----	6s	---	---	---	1.00	---
Rock outcrop-----	8	---	---	---	---	---
95E: Farmington-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
96: Ipswich-----	8	---	---	---	---	---
97: Pawcatuck-----	8	---	---	---	---	---
98: Westbrook-----	8	---	---	---	---	---
99: Westbrook, low salt----	8	---	---	---	---	---
100: Suncook-----	2s	139.00	25.00	2.50	1.50	---
101: Occum-----	1	150.00	30.00	3.50	3.50	1,850.00

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
102: Pootatuck-----	2w	148.00	28.00	3.00	3.00	---
103: Rippowam-----	---	---	---	---	---	---
104: Bash-----	4w	159.00	31.00	3.50	3.50	---
105: Hadley-----	1	180.00	35.00	5.00	6.00	2,100.00
106: Winooski-----	2w	179.00	35.00	4.50	5.50	---
107: Limerick-----	---	---	---	---	---	---
Lim-----	4w	155.00	29.00	3.50	3.50	---
108: Saco-----	6w	---	---	---	---	---
109: Fluvaquents, Frequently Flooded-----	6w	---	---	---	3.50	---
Udifluvents, Frequently Flooded-----	6w	---	---	---	2.00	---
221A: Ninigret-----	2w	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
224A: Deerfield-----	2w	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
225B: Brancroft-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
226B: Berlin-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
228B: Elmridge-----	2w	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
229B: Agawam-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
229C: Agawam-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---

Table 6.-Non-Irrigated Yields by Map Unit Component-Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
230B: Branford-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
230C: Branford-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
232B: Haven-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
234B: Merrimac-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
235B: Penwood-----	2s	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
236B: Windsor-----	2s	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
237A: Manchester-----	2s	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
237C: Manchester-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
238A: Hinckley-----	3s	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
238C: Hinckley-----	4e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
240B: Ludlow-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
243B: Rainbow-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
245B: Woodbridge-----	2w	---	---	---	---	---
Urban land-----	8	---	---	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
245C:						
Woodbridge-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
248B:						
Georgia-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
250B:						
Sutton-----	2w	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
253B:						
Wapping-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
255B:						
Watchaug-----	2w	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
260B:						
Charlton-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
260C:						
Charlton-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
260D:						
Charlton-----	4e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
263B:						
Cheshire-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
263C:						
Cheshire-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
266B:						
Narragansett-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
269B:						
Yalesville-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
269C:						
Yalesville-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---

Table 6.-Non-Irrigated Yields by Map Unit Component-Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
273C:						
Urban land-----	8	---	---	---	---	---
Charlton-----	3e	---	---	---	---	---
Chatfield-----	3e	---	---	---	---	---
273E:						
Urban land-----	8	---	---	---	---	---
Charlton-----	6e	---	---	---	---	---
Chatfield-----	6e	---	---	---	---	---
275C:						
Urban land-----	8	---	---	---	---	---
Chatfield-----	6s	---	---	---	---	---
275E:						
Urban land-----	8	---	---	---	---	---
Chatfield-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
282B:						
Broadbrook-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
284B:						
Paxton-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
284C:						
Paxton-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
284D:						
Paxton-----	4e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
287B:						
Wethersfield-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
287C:						
Wethersfield-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
287D:						
Wethersfield-----	4e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
290B:						
Stockbridge-----	2e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
290C:						
Stockbridge-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
290D:						
Stockbridge-----	4e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
301:						
Beaches-----	8	---	---	---	---	---
Udipsamments-----	3s	---	---	---	---	---
302:						
Dumps-----	8	---	---	---	---	---
303:						
Pits, quarries-----	8	---	---	---	---	---
304:						
Udorthents-----	7e	---	---	---	---	---
305:						
Udorthents-----	4e	---	---	---	---	---
Pits-----	8	---	---	---	---	---
306:						
Udorthents-----	3e	---	---	---	---	---
Urban land-----	8	---	---	---	---	---
307:						
Urban land-----	8	---	---	---	---	---
308:						
Udorthents-----	4e	---	---	---	---	---
309:						
Udorthents-----	4e	---	---	---	---	---
310:						
Udorthents, Periodically Flooded-----	4e	---	---	---	---	---
401C:						
Macomber-----	6s	---	---	---	---	---
Taconic-----	6s	---	---	---	---	---
402D:						
Macomber-----	7s	---	---	---	---	---
Taconic-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
403C:						
Taconic-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
403E:						
Taconic-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
403F:						
Taconic-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
405C:						
Dummerston-----	6s	---	---	---	---	---
405E:						
Dummerston-----	7s	---	---	---	---	---
407C:						
Lanesboro-----	6s	---	---	---	---	---
407E:						
Lanesboro-----	7s	---	---	---	---	---
408C:						
Fullam-----	6s	---	---	---	---	---
409B:						
Brayton-----	6s	---	---	---	---	---
412B:						
Bice-----	2e	---	---	---	3.00	---
412C:						
Bice-----	3e	---	---	---	2.50	---
412D:						
Bice-----	4e	---	---	---	2.00	---
413C:						
Bice-----	6s	---	---	---	2.00	---
Millsite-----	6s	---	---	---	2.00	---
413E:						
Bice-----	7s	---	---	---	---	---
Millsite-----	7s	---	---	---	---	---
414:						
Fredon, cold-----	4w	155.00	30.00	---	4.00	---
415C:						
Millsite-----	7s	---	---	---	---	---
Westminster-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---
415E:						
Millsite-----	7s	---	---	---	---	---
Westminster-----	7s	---	---	---	---	---
Rock outcrop-----	8	---	---	---	---	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
416E: Rock outcrop-----	8	---	---	---	---	---
Westminster-----	7s	---	---	---	---	---
416F: Rock outcrop-----	8	---	---	---	---	---
Westminster-----	7s	---	---	---	---	---
417B: Bice-----	6s	---	---	---	2.50	---
417C: Bice-----	6s	---	---	---	2.00	---
417D: Bice-----	6s	---	---	---	2.00	---
418C: Schroon-----	6s	---	---	---	2.00	---
420A: Schroon-----	2w	---	---	---	3.00	---
420B: Schroon-----	2w	---	---	---	3.00	---
421A: Ninigret, cold-----	2w	151.00	29.00	4.00	3.50	---
423A: Sudbury, cold-----	2w	134.00	24.00	3.00	2.50	---
424B: Shelburne-----	2s	---	---	2.50	1.50	---
424C: Shelburne-----	3e	---	---	2.50	1.50	---
424D: Shelburne-----	4e	---	---	2.00	1.00	---
425B: Shelburne-----	6s	---	---	---	1.00	---
425C: Shelburne-----	6s	---	---	---	1.00	---
426D: Shelburne-----	7s	---	---	---	---	---
427B: Ashfield-----	6s	---	---	---	1.00	---
427C: Ashfield-----	6s	---	---	---	1.00	---
428A: Ashfield-----	1	---	---	2.50	2.00	---
428B: Ashfield-----	2e	---	---	2.50	1.50	---
428C: Ashfield-----	3e	---	---	2.50	1.50	---

Table 6.—Non-Irrigated Yields by Map Unit Component—Continued

Map symbol and soil name	Land capability	Corn	Corn silage	Grass-legume hay	Pasture	Tobacco
		Bu	Tons	Tons	AUM	Lbs
429A: Agawam, cold-----	1	148.00	28.00	4.00	3.50	---
429B: Agawam, cold-----	2e	145.00	27.00	3.50	3.00	---
429C: Agawam, cold-----	3e	141.00	26.00	3.50	3.00	---
433: Moosilauke-----	4w	---	---	---	---	---
434A: Merrimac, cold-----	1	135.00	24.00	---	2.50	---
434B: Merrimac, cold-----	2e	135.00	24.00	---	2.50	---
434C: Merrimac, cold-----	3e	131.00	23.00	---	2.00	---
435: Scarboro-----	5w	---	---	---	---	---
436: Halsey-----	5w	---	---	---	---	---
437: Wonsqueak-----	5w	---	---	---	---	---
438: Bucksport-----	5w	---	---	---	---	---
440A: Boscawen-----	3s	126.00	22.00	---	1.50	---
440C: Boscawen-----	4e	124.00	21.00	---	1.50	---
440E: Boscawen-----	6e	---	---	---	---	---
442: Brayton-----	4w	---	---	---	1.00	---
443: Brayton-----	7s	---	---	---	---	---
Loonmeadow-----	7s	---	---	---	---	---
448B: Hogansburg-----	2e	160.00	32.00	4.50	5.00	---
449B: Hogansburg-----	6s	---	---	---	4.50	---
449C: Hogansburg-----	6s	---	---	---	4.00	---
450B: Pyrities-----	2e	167.00	33.00	5.00	5.00	---
450C: Pyrities-----	3e	162.00	32.00	4.50	5.00	---

Table 7.—Acreage by Capability Class and Subclass

Capability class	Capability subclass	Acreage
Unclassified	---	370,616
1	---	46,174
2	e	238,649
2	w	123,027
2	s	31,134
3	e	156,420
3	s	6,877
4	e	127,589
4	w	52,345
5	w	75,142
6	e	28,278
6	w	25,921
6	s	545,521
7	e	8,946
7	s	638,984
8	---	183,220

Table 8.—Forestland Productivity

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
2: Ridgebury-----	eastern white pine--	63	114	green ash, red maple, tuliptree
	elm-----	---	---	
	northern red oak----	57	43	
	red maple-----	---	---	
	sugar maple-----	52	29	
3: Ridgebury-----	eastern white pine--	63	114	green ash, red maple, tuliptree
	elm-----	---	---	
	northern red oak----	57	43	
	red maple-----	---	---	
	sugar maple-----	52	29	
Leicester-----	eastern white pine--	69	129	green ash, red maple, tuliptree
	northern red oak----	56	43	
	red maple-----	70	43	
	yellow birch-----	---	---	
Whitman-----	eastern white pine--	56	100	---
	red maple-----	55	29	
	swamp tupelo-----	---	---	
4: Leicester-----	eastern white pine--	69	129	green ash, red maple, tuliptree
	northern red oak----	56	43	
	red maple-----	70	43	
	yellow birch-----	---	---	
5: Wilbraham-----	eastern white pine--	65	114	green ash, red maple, tuliptree
	northern red oak----	63	43	
	red maple-----	70	43	
	sugar maple-----	55	29	
	white oak-----	---	---	
6: Wilbraham-----	eastern white pine--	65	114	green ash, red maple, tuliptree
	northern red oak----	63	43	
	red maple-----	70	43	
	sugar maple-----	55	29	
Menlo-----	eastern white pine--	55	86	---
	elm-----	---	---	
	red maple-----	55	29	
7: Mudgepond-----	eastern white pine--	75	143	green ash, red maple, tuliptree
	green ash-----	---	---	
	northern red oak----	70	57	
	red maple-----	75	43	
8: Mudgepond-----	eastern white pine--	75	143	green ash, red maple, tuliptree
	green ash-----	---	---	
	northern red oak----	70	57	
	red maple-----	75	43	
Alden-----	red maple-----	50	29	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
9:				
Scitico-----	eastern white pine--	57	100	green ash, red maple, tuliptree
	elm-----	---	---	
	pin oak-----	---	---	
	red maple-----	55	29	
Shaker-----	eastern white pine--	57	100	green ash, red maple, tuliptree
	elm-----	---	---	
	pin oak-----	---	---	
	red maple-----	55	29	
Maybid-----	red maple-----	55	29	---
10:				
Raynham-----	eastern hemlock-----	---	0	green ash, red maple, tuliptree
	eastern white pine--	65	114	
	elm-----	---	0	
	red maple-----	65	43	
	sugar maple-----	---	0	
	tamarack-----	---	0	
12:				
Raypol-----	eastern white pine--	68	114	green ash, red maple, tuliptree
	elm-----	---	---	
	red maple-----	75	43	
13:				
Walpole-----	eastern hemlock-----	54	114	green ash, red maple, tuliptree
	eastern white pine--	68	114	
	red maple-----	75	43	
	white ash-----	61	43	
14:				
Fredon-----	eastern white pine--	70	129	green ash, red maple, tuliptree
	northern red oak----	60	43	
	red maple-----	70	43	
	tuliptree-----	80	72	
15:				
Scarboro-----	Atlantic white cedar	45	0	---
	eastern white pine--	55	86	
	red maple-----	55	29	
16:				
Halsey-----	red maple-----	55	29	---
	river birch-----	---	0	
	swamp white oak----	---	0	
17:				
Timakwa-----	Atlantic white cedar	---	0	---
	eastern hemlock-----	---	---	
	eastern white pine--	---	---	
	red maple-----	55	29	
	silver maple-----	80	29	
	tamarack-----	61	57	
	white ash-----	---	0	
Natchaug-----	Atlantic white cedar	---	0	---
	eastern hemlock-----	---	---	
	eastern white pine--	---	---	
	red maple-----	55	29	
	silver maple-----	80	29	
	tamarack-----	61	57	
	white ash-----	---	0	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
18: Catden-----	eastern hemlock----- eastern white pine-- red maple-----	--- --- 56	--- --- 29	---
Freetown-----	eastern hemlock----- eastern white pine-- red maple-----	--- --- 56	--- --- 29	---
20A: Ellington-----	eastern white pine-- northern red oak---- red maple----- white oak-----	75 70 --- ---	143 57 0 ---	eastern white pine, northern red oak, white oak
21A: Ninigret-----	eastern white pine-- northern red oak---- red maple----- sugar maple----- white oak-----	75 65 60 55 ---	143 43 43 29 ---	eastern white pine, northern red oak, white oak
Tisbury-----	eastern white pine-- northern red oak---- tuliptree----- white oak-----	75 70 --- ---	143 57 --- ---	eastern white pine, northern red oak, tuliptree, white oak
22A: Hero-----	eastern hemlock----- eastern white pine-- northern red oak---- white oak-----	--- 70 65 ---	--- 129 43 ---	eastern hemlock, eastern white pine, northern red oak, white oak
22B: Hero-----	eastern hemlock----- eastern white pine-- northern red oak---- white oak-----	--- 70 65 ---	--- 129 43 ---	eastern hemlock, eastern white pine, northern red oak, white oak
23A: Sudbury-----	eastern white pine-- northern red oak---- white oak-----	60 45 ---	100 29 ---	eastern white pine, northern red oak, white oak
24A: Deerfield-----	eastern white pine-- northern red oak---- pitch pine----- white oak-----	65 55 --- ---	114 43 --- ---	eastern white pine
25A: Brancroft-----	eastern hemlock----- eastern white pine-- northern red oak---- red maple----- white ash----- white oak-----	--- 65 55 --- --- ---	0 114 43 0 0 ---	eastern hemlock, eastern white pine

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
25B: Brancroft-----	eastern hemlock----	---	0	eastern hemlock, eastern white pine
	eastern white pine--	65	114	
	northern red oak----	55	43	
	red maple-----	---	0	
	white ash-----	---	0	
	white oak-----	---	---	
25C: Brancroft-----	eastern hemlock----	---	0	eastern hemlock, eastern white pine
	eastern white pine--	65	114	
	northern red oak----	55	43	
	red maple-----	---	0	
	white ash-----	---	0	
	white oak-----	---	---	
26A: Berlin-----	eastern white pine--	65	114	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	red maple-----	---	---	
	white oak-----	---	---	
26B: Berlin-----	eastern white pine--	65	114	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	red maple-----	---	---	
	white oak-----	---	---	
27A: Belgrade-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak----	62	43	
	white oak-----	---	---	
28A: Elmridge-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	red maple-----	---	0	
	shagbark hickory----	60	43	
	white oak-----	---	---	
28B: Elmridge-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	red maple-----	---	0	
	shagbark hickory----	60	43	
	white oak-----	---	---	
29A: Agawam-----	eastern white pine--	70	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	
29B: Agawam-----	eastern white pine--	70	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	
29C: Agawam-----	eastern white pine--	70	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
30A: Branford-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	shagbark hickory----	---	0	
	white oak-----	---	---	
30B: Branford-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	shagbark hickory----	---	0	
	white oak-----	---	---	
30C: Branford-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	shagbark hickory----	---	0	
	white oak-----	---	---	
31A: Copake-----	eastern white pine--	65	114	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	55	29	
	white oak-----	---	---	
31B: Copake-----	eastern white pine--	65	114	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	55	29	
	white oak-----	---	---	
31C: Copake-----	eastern white pine--	65	114	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	55	29	
	white oak-----	---	---	
32A: Haven-----	eastern white pine--	75	143	eastern white pine, sugar maple, white oak
	northern red oak----	55	43	
	sugar maple-----	65	43	
	white oak-----	---	---	
Enfield-----	eastern white pine--	74	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	43	
	shagbark hickory----	---	0	
	white oak-----	---	---	
32B: Haven-----	eastern white pine--	75	143	eastern white pine, sugar maple, white oak
	northern red oak----	55	43	
	sugar maple-----	65	43	
	white oak-----	---	---	
Enfield-----	eastern white pine--	74	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	43	
	shagbark hickory----	---	0	
	white oak-----	---	---	
32C: Haven-----	eastern white pine--	75	143	eastern white pine, sugar maple, white oak
	northern red oak----	55	43	
	sugar maple-----	65	43	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
32C: Enfield-----	eastern white pine--	74	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	43	
	shagbark hickory----	---	0	
	white oak-----	---	---	
33A: Hartford-----	eastern white pine--	65	114	eastern white pine, sugar maple, white oak
	northern red oak----	59	43	
	sugar maple-----	60	43	
	white oak-----	---	---	
33B: Hartford-----	eastern white pine--	65	114	eastern white pine, sugar maple, white oak
	northern red oak----	59	43	
	sugar maple-----	60	43	
	white oak-----	---	---	
34A: Merrimac-----	eastern white pine--	64	114	eastern white pine, white oak
	northern red oak----	51	29	
	sugar maple-----	58	43	
	white oak-----	---	---	
34B: Merrimac-----	eastern white pine--	64	114	eastern white pine, white oak
	northern red oak----	51	29	
	sugar maple-----	58	43	
	white oak-----	---	---	
34C: Merrimac-----	eastern white pine--	64	114	eastern white pine, white oak
	northern red oak----	51	29	
	sugar maple-----	58	43	
	white oak-----	---	---	
35A: Penwood-----	eastern white pine--	57	100	eastern white pine
	northern red oak----	50	29	
	pitch pine-----	50	0	
35B: Penwood-----	eastern white pine--	57	100	eastern white pine
	northern red oak----	50	29	
	pitch pine-----	50	0	
36A: Windsor-----	eastern white pine--	57	100	eastern white pine
	northern red oak----	52	29	
	pitch pine-----	---	---	
	sugar maple-----	55	29	
36B: Windsor-----	eastern white pine--	57	100	eastern white pine
	northern red oak----	52	29	
	pitch pine-----	---	---	
	sugar maple-----	55	29	
36C: Windsor-----	eastern white pine--	57	100	eastern white pine
	northern red oak----	52	29	
	pitch pine-----	---	---	
	sugar maple-----	55	29	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
37A: Manchester-----	eastern white pine--	55	86	eastern white pine
	northern red oak----	50	29	
	pitch pine-----	---	---	
	white oak-----	---	---	
37C: Manchester-----	eastern white pine--	55	86	eastern white pine
	northern red oak----	50	29	
	pitch pine-----	---	---	
	white oak-----	---	---	
37E: Manchester-----	eastern white pine--	55	86	eastern white pine
	northern red oak----	50	29	
	pitch pine-----	---	---	
	white oak-----	---	---	
38A: Hinckley-----	eastern white pine--	60	100	eastern white pine
	northern red oak----	49	29	
	sugar maple-----	57	29	
	white oak-----	---	---	
38C: Hinckley-----	eastern white pine--	60	100	eastern white pine
	northern red oak----	49	29	
	sugar maple-----	57	29	
	white oak-----	---	---	
38E: Hinckley-----	eastern white pine--	60	100	eastern white pine
	northern red oak----	49	29	
	sugar maple-----	57	29	
	white oak-----	---	---	
39A: Groton-----	eastern white pine--	60	100	eastern white pine
	northern red oak----	55	43	
	sugar maple-----	55	43	
39C: Groton-----	eastern white pine--	60	100	eastern white pine
	northern red oak----	55	43	
	sugar maple-----	55	43	
39E: Groton-----	eastern white pine--	60	100	eastern white pine
	northern red oak----	55	43	
	sugar maple-----	55	43	
40A: Ludlow-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	red maple-----	---	0	
40B: Ludlow-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	red maple-----	---	0	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
41B: Ludlow-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	red maple-----	---	0	
42C: Ludlow-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	red maple-----	---	0	
43A: Rainbow-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	54	29	
43B: Rainbow-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	54	29	
44B: Rainbow-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	54	29	
45A: Woodbridge-----	eastern white pine--	67	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
45B: Woodbridge-----	eastern white pine--	67	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
45C: Woodbridge-----	eastern white pine--	67	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
46B: Woodbridge-----	eastern white pine--	67	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
46C: Woodbridge-----	eastern white pine--	67	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
47C:				
Woodbridge-----	eastern white pine--	67	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
48B:				
Georgia-----	basswood-----	65	43	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	quaking aspen-----	---	0	
	red maple-----	70	43	
	sugar maple-----	60	43	
	white ash-----	66	43	
	yellow birch-----	60	43	
Amenia-----	American basswood---	---	0	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
48C:				
Georgia-----	basswood-----	65	43	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	quaking aspen-----	---	0	
	red maple-----	70	43	
	sugar maple-----	60	43	
	white ash-----	66	43	
	yellow birch-----	60	43	
Amenia-----	American basswood---	---	0	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
49B:				
Georgia-----	basswood-----	65	43	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
Amenia-----	American basswood---	---	0	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
49C:				
Georgia-----	basswood-----	65	43	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
Amenia-----	American basswood---	---	0	eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
50A: Sutton-----	black cherry-----	72	43	eastern white pine, northern red oak, white oak
	eastern white pine--	62	114	
	northern red oak----	62	43	
	sugar maple-----	54	29	
	white oak-----	---	---	
50B: Sutton-----	black cherry-----	72	43	eastern white pine, northern red oak, white oak
	eastern white pine--	62	114	
	northern red oak----	62	43	
	sugar maple-----	54	29	
	white oak-----	---	---	
51B: Sutton-----	black cherry-----	72	43	eastern white pine, northern red oak, white oak
	eastern white pine--	62	114	
	northern red oak----	62	43	
	sugar maple-----	54	29	
	white oak-----	---	---	
52C: Sutton-----	black cherry-----	72	43	eastern white pine, northern red oak, white oak
	eastern white pine--	62	114	
	northern red oak----	62	43	
	sugar maple-----	54	29	
	white oak-----	---	---	
53A: Wapping-----	black cherry-----	72	43	eastern white pine, northern red oak
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	54	29	
53B: Wapping-----	black cherry-----	72	43	eastern white pine, northern red oak
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	54	29	
54B: Wapping-----	black cherry-----	72	43	eastern white pine, northern red oak
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	54	29	
55A: Watchaug-----	eastern white pine--	65	114	eastern white pine, northern red oak
	northern red oak----	65	43	
	red maple-----	68	43	
55B: Watchaug-----	eastern white pine--	65	114	eastern white pine, northern red oak
	northern red oak----	65	43	
	red maple-----	68	43	
56B: Watchaug-----	eastern white pine--	65	114	eastern white pine, northern red oak
	northern red oak----	65	43	
	red maple-----	68	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
57B: Gloucester-----	eastern white pine--	61	100	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	53	29	
	white oak-----	---	---	
57C: Gloucester-----	eastern white pine--	61	100	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	53	29	
	white oak-----	---	---	
57D: Gloucester-----	eastern white pine--	61	100	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	53	29	
	white oak-----	---	---	
58B: Gloucester-----	eastern white pine--	61	100	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	53	29	
	white oak-----	---	---	
58C: Gloucester-----	eastern white pine--	61	100	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	53	29	
	white oak-----	---	---	
59C: Gloucester-----	eastern white pine--	61	100	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	53	29	
	white oak-----	---	---	
59D: Gloucester-----	eastern white pine--	61	100	eastern white pine, northern red oak, white oak
	northern red oak----	60	43	
	sugar maple-----	53	29	
	white oak-----	---	---	
60B: Canton-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, white oak
	eastern white pine--	58	100	
	northern red oak----	52	29	
	white oak-----	---	---	
Charlton-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
60C: Canton-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, white oak
	eastern white pine--	58	100	
	northern red oak----	52	29	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
60C:				
Charlton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
60D:				
Canton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, white oak
	eastern white pine--	58	100	
	northern red oak----	52	29	
	white oak-----	---	---	
Charlton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
61B:				
Canton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, white oak
	eastern white pine--	58	100	
	northern red oak----	52	29	
	white oak-----	---	---	
Charlton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
61C:				
Canton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, white oak
	eastern white pine--	58	100	
	northern red oak----	52	29	
	white oak-----	---	---	
Charlton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
62C:				
Canton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, white oak
	eastern white pine--	58	100	
	northern red oak----	52	29	
	white oak-----	---	---	
Charlton-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
62D:				
Canton-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	58	100	eastern white
	northern red oak----	52	29	pine, white oak
	white oak-----	---	---	
Charlton-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	65	43	pine, northern red
	red maple-----	55	29	oak, white oak
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
63B:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	60	43	pine, northern red
	shagbark hickory----	---	0	oak
	sugar maple-----	---	0	
63C:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	60	43	pine, northern red
	shagbark hickory----	---	0	oak
	sugar maple-----	---	0	
63D:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	60	43	pine, northern red
	shagbark hickory----	---	0	oak
	sugar maple-----	---	0	
64B:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	60	43	pine, northern red
	shagbark hickory----	---	0	oak
	sugar maple-----	---	0	
64C:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	60	43	pine, northern red
	shagbark hickory----	---	0	oak
	sugar maple-----	---	0	
65C:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	60	43	pine, northern red
	shagbark hickory----	---	0	oak
	sugar maple-----	---	0	
65D:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock,
	eastern white pine--	65	114	eastern white
	northern red oak----	60	43	pine, northern red
	shagbark hickory----	---	0	oak
	sugar maple-----	---	0	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
66B: Narragansett-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
66C: Narragansett-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
67B: Narragansett-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
67C: Narragansett-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
68C: Narragansett-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
68D: Narragansett-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
69B: Yalesville-----	eastern white pine--	65	114	eastern white pine, northern red oak
	northern red oak----	60	43	
	sugar maple-----	---	0	
69C: Yalesville-----	eastern white pine--	65	114	eastern white pine, northern red oak
	northern red oak----	60	43	
	sugar maple-----	---	0	
70C: Branford-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	shagbark hickory----	---	0	
	white oak-----	---	---	
Holyoke-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	
71C: Brookfield-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	sugar maple-----	55	29	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
71C:				
Brimfield-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	
Rock outcrop-----	---	---	---	---
71E:				
Brookfield-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	sugar maple-----	55	29	
	white oak-----	---	---	
Brimfield-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	
Rock outcrop-----	---	---	---	---
73C:				
Charlton-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
Chatfield-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
	white oak-----	---	---	
73E:				
Charlton-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
Chatfield-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
	white oak-----	---	---	
74C:				
Narragansett-----	eastern hemlock-----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
Hollis-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern hemlock-----	---	---	
	eastern white pine--	55	86	
	northern red oak----	47	29	
	sugar maple-----	56	29	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
75C:				
Hollis-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern hemlock----	---	---	
	eastern white pine--	55	86	
	northern red oak----	47	29	
	sugar maple-----	56	29	
Chatfield-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
	white oak-----	---	---	
Rock outcrop-----	---	---	---	---
75E:				
Hollis-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern hemlock----	---	---	
	eastern white pine--	55	86	
	northern red oak----	47	29	
	sugar maple-----	56	29	
Chatfield-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
	white oak-----	---	---	
Rock outcrop-----	---	---	---	---
76E:				
Rock outcrop-----	---	---	---	---
Hollis-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern hemlock----	---	---	
	eastern white pine--	55	86	
	northern red oak----	47	29	
	sugar maple-----	56	29	
76F:				
Rock outcrop-----	---	---	---	---
Hollis-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern hemlock----	---	---	
	eastern white pine--	55	86	
	northern red oak----	47	29	
	sugar maple-----	56	29	
77C:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	65	114	
	northern red oak----	60	43	
	shagbark hickory----	---	0	
	sugar maple-----	---	0	
Holyoke-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
77D:				
Cheshire-----	eastern hemlock----	---	---	eastern hemlock, eastern white pine, northern red oak
	eastern white pine--	65	114	
	northern red oak----	60	43	
	shagbark hickory----	---	0	
	sugar maple-----	---	0	
Holyoke-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	
78C:				
Holyoke-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	
Rock outcrop-----	---	---	---	---
78E:				
Holyoke-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	
Rock outcrop-----	---	---	---	---
79E:				
Rock outcrop-----	---	---	---	---
Holyoke-----	chestnut oak-----	---	---	chestnut oak, eastern white pine
	eastern white pine--	55	86	
	northern red oak----	47	29	
80B:				
Bernardston-----	eastern hemlock----	65	0	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern white pine--	65	114	
	northern red oak----	55	43	
	sugar maple-----	65	43	
80C:				
Bernardston-----	eastern hemlock----	65	0	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern white pine--	65	114	
	northern red oak----	55	43	
	sugar maple-----	65	43	
81C:				
Bernardston-----	eastern hemlock----	65	0	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern white pine--	65	114	
	northern red oak----	55	43	
	sugar maple-----	65	43	
81D:				
Bernardston-----	eastern hemlock----	65	0	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern white pine--	65	114	
	northern red oak----	55	43	
	sugar maple-----	65	43	
82B:				
Broadbrook-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	tuliptree-----	93	100	
	white ash-----	76	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
82C:				
Broadbrook-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	tuliptree-----	93	100	
	white ash-----	76	43	
82D:				
Broadbrook-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	tuliptree-----	93	100	
	white ash-----	76	43	
83B:				
Broadbrook-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	tuliptree-----	93	100	
	white ash-----	76	43	
83C:				
Broadbrook-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	tuliptree-----	93	100	
	white ash-----	76	43	
84B:				
Paxton-----	eastern white pine--	66	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Montauk-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
84C:				
Paxton-----	eastern white pine--	66	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Montauk-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
84D:				
Paxton-----	eastern white pine--	66	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Montauk-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
85B:				
Paxton-----	eastern white pine--	66	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
85B:				
Montauk-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
85C:				
Paxton-----	eastern white pine--	66	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Montauk-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
86C:				
Paxton-----	eastern white pine--	66	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Montauk-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
86D:				
Paxton-----	eastern white pine--	66	114	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Montauk-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
87B:				
Wethersfield-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
87C:				
Wethersfield-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
87D:				
Wethersfield-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
88B:				
Wethersfield-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
88C: Wethersfield-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
89C: Wethersfield-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
89D: Wethersfield-----	eastern white pine--	75	143	ash, northern red oak, sugar maple, tuliptree, white oak
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
90B: Stockbridge-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
90C: Stockbridge-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
90D: Stockbridge-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
91B: Stockbridge-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
91C: Stockbridge-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
91D: Stockbridge-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
92B: Nellis-----	American basswood---	80	57	American basswood, eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	85	143	
	northern red oak----	80	57	
	sugar maple-----	70	43	
	white ash-----	85	57	
92C: Nellis-----	American basswood---	80	57	American basswood, eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	85	143	
	northern red oak----	80	57	
	sugar maple-----	70	43	
	white ash-----	85	57	
92D: Nellis-----	American basswood---	80	57	American basswood, eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	85	143	
	northern red oak----	80	57	
	sugar maple-----	70	43	
	white ash-----	85	57	
93C: Nellis-----	American basswood---	80	57	American basswood, eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	85	143	
	northern red oak----	80	57	
	sugar maple-----	70	43	
	white ash-----	85	57	
94C: Farmington-----	eastern redcedar----	---	---	eastern redcedar, eastern white pine
	eastern white pine--	---	---	
	sugar maple-----	---	---	
Nellis-----	American basswood---	80	57	American basswood, eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	85	143	
	northern red oak----	80	57	
	sugar maple-----	70	43	
	white ash-----	85	57	
94E: Farmington-----	eastern redcedar----	---	---	eastern redcedar, eastern white pine
	eastern white pine--	---	---	
	sugar maple-----	---	---	
Nellis-----	American basswood---	80	57	American basswood, eastern white pine, northern red oak, sugar maple, white ash
	eastern white pine--	85	143	
	northern red oak----	80	57	
	sugar maple-----	70	43	
	white ash-----	85	57	
95C: Farmington-----	eastern redcedar----	---	---	eastern redcedar, eastern white pine
	eastern white pine--	---	---	
	sugar maple-----	---	---	
Rock outcrop-----	---	---	---	---
95E: Farmington-----	eastern redcedar----	---	---	eastern redcedar, eastern white pine
	eastern white pine--	---	---	
	sugar maple-----	---	---	
Rock outcrop-----	---	---	---	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
96: Ipswich-----	---	---	---	---
97: Pawcatuck-----	---	---	---	---
98: Westbrook-----	---	---	---	---
99: Westbrook, low salt----	---	---	---	---
100: Suncook-----	American sycamore--- eastern white pine-- pin oak----- red maple----- silver maple-----	--- 55 --- 50 ---	--- 86 --- 29 ---	---
101: Occum-----	American sycamore--- eastern white pine-- red maple----- silver maple-----	--- 70 --- ---	--- 129 --- ---	---
102: Pootatuck-----	American sycamore--- eastern white pine-- pin oak----- red maple----- yellow birch-----	--- 75 --- 60 60	--- 143 --- 43 43	---
103: Rippowam-----	eastern white pine-- pin oak----- red maple-----	65 --- 75	114 --- 43	---
104: Bash-----	red maple----- white ash-----	70 70	43 43	---
105: Hadley-----	eastern white pine-- red maple----- silver maple-----	70 --- ---	129 --- ---	---
106: Winooski-----	eastern white pine-- pin oak----- silver maple-----	75 --- ---	143 --- ---	---
107: Limerick-----	eastern white pine-- red maple----- silver maple-----	65 40 ---	114 29 ---	---
Lim-----	American elm----- eastern white pine-- green ash----- red maple----- yellow birch-----	--- 60 --- 60 ---	0 100 0 43 0	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
108:				
Saco-----	eastern white pine--	50	86	---
	red maple-----	50	29	
109:				
Fluvaquents, Frequently Flooded-----	eastern white pine--	50	86	---
	red maple-----	55	29	
Udifluents, Frequently Flooded-----	American sycamore---	---	0	---
	eastern white pine--	50	86	
	red maple-----	55	29	
	silver maple-----	55	14	
221A:				
Ninigret-----	eastern white pine--	75	143	---
	northern red oak----	65	43	
	red maple-----	60	43	
	sugar maple-----	55	29	
	white oak-----	---	---	
Urban land-----	---	---	---	---
224A:				
Deerfield-----	eastern white pine--	65	114	---
	northern red oak----	55	43	
	pitch pine-----	---	---	
	white oak-----	---	---	
Urban land-----	---	---	---	---
225B:				
Brancroft-----	eastern hemlock----	---	0	---
	eastern white pine--	65	114	
	northern red oak----	55	43	
	red maple-----	---	0	
	white ash-----	---	0	
	white oak-----	---	---	
Urban land-----	---	---	---	---
226B:				
Berlin-----	eastern white pine--	65	114	---
	northern red oak----	60	43	
	red maple-----	---	---	
	white oak-----	---	---	
Urban land-----	---	---	---	---
228B:				
Elmridge-----	eastern white pine--	75	143	---
	northern red oak----	70	57	
	red maple-----	---	0	
	shagbark hickory----	60	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
229B:				
Agawam-----	eastern white pine--	70	129	---
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	
Urban land-----	---	---	---	---
229C:				
Agawam-----	eastern white pine--	70	129	---
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	
Urban land-----	---	---	---	---
230B:				
Branford-----	eastern white pine--	75	143	---
	northern red oak----	70	57	
	shagbark hickory----	---	0	
	white oak-----	---	---	
Urban land-----	---	---	---	---
230C:				
Branford-----	eastern white pine--	75	143	---
	northern red oak----	70	57	
	shagbark hickory----	---	0	
	white oak-----	---	---	
Urban land-----	---	---	---	---
232B:				
Haven-----	eastern white pine--	75	143	---
	northern red oak----	55	43	
	sugar maple-----	65	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---
234B:				
Merrimac-----	eastern white pine--	64	114	---
	northern red oak----	51	29	
	sugar maple-----	58	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---
235B:				
Penwood-----	eastern white pine--	57	100	---
	northern red oak----	50	29	
	pitch pine-----	50	0	
Urban land-----	---	---	---	---
236B:				
Windsor-----	eastern white pine--	57	100	---
	northern red oak----	52	29	
	pitch pine-----	---	---	
	sugar maple-----	55	29	
Urban land-----	---	---	---	---

Table 8.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
237A:				
Manchester-----	eastern white pine--	55	86	---
	northern red oak----	50	29	
	pitch pine-----	---	---	
	white oak-----	---	---	
Urban land-----	---	---	---	---
237C:				
Manchester-----	eastern white pine--	55	86	---
	northern red oak----	50	29	
	pitch pine-----	---	---	
	white oak-----	---	---	
Urban land-----	---	---	---	---
238A:				
Hinckley-----	eastern white pine--	60	100	---
	northern red oak----	49	29	
	sugar maple-----	57	29	
	white oak-----	---	---	
Urban land-----	---	---	---	---
238C:				
Hinckley-----	eastern white pine--	60	100	---
	northern red oak----	49	29	
	sugar maple-----	57	29	
	white oak-----	---	---	
Urban land-----	---	---	---	---
240B:				
Ludlow-----	eastern white pine--	75	143	---
	northern red oak----	70	57	
	red maple-----	---	0	
Urban land-----	---	---	---	---
243B:				
Rainbow-----	eastern white pine--	75	143	---
	northern red oak----	70	57	
	sugar maple-----	54	29	
Urban land-----	---	---	---	---
245B:				
Woodbridge-----	eastern white pine--	67	114	---
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---
245C:				
Woodbridge-----	eastern white pine--	67	114	---
	northern red oak----	72	57	
	sugar maple-----	65	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
248B:				
Georgia-----	basswood-----	65	43	---
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
Urban land-----	---	---	---	---
250B:				
Sutton-----	black cherry-----	72	43	---
	eastern white pine--	62	114	
	northern red oak----	62	43	
	sugar maple-----	54	29	
	white oak-----	---	---	
Urban land-----	---	---	---	---
253B:				
Wapping-----	black cherry-----	72	43	---
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	54	29	
Urban land-----	---	---	---	---
255B:				
Watchaug-----	eastern white pine--	65	114	---
	northern red oak----	65	43	
	red maple-----	68	43	
Urban land-----	---	---	---	---
260B:				
Charlton-----	eastern hemlock-----	---	---	---
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
Urban land-----	---	---	---	---
260C:				
Charlton-----	eastern hemlock-----	---	---	---
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
Urban land-----	---	---	---	---
260D:				
Charlton-----	eastern hemlock-----	---	---	---
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
Urban land-----	---	---	---	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
263B:				
Cheshire-----	eastern hemlock----	---	---	---
	eastern white pine--	65	114	
	northern red oak----	60	43	
	shagbark hickory----	---	0	
	sugar maple-----	---	0	
Urban land-----	---	---	---	---
263C:				
Cheshire-----	eastern hemlock----	---	---	---
	eastern white pine--	65	114	
	northern red oak----	60	43	
	shagbark hickory----	---	0	
	sugar maple-----	---	0	
Urban land-----	---	---	---	---
266B:				
Narragansett-----	eastern hemlock----	---	---	---
	eastern white pine--	68	114	
	northern red oak----	60	43	
	sugar maple-----	55	29	
Urban land-----	---	---	---	---
269B:				
Yalesville-----	eastern white pine--	65	114	---
	northern red oak----	60	43	
	sugar maple-----	---	0	
Urban land-----	---	---	---	---
269C:				
Yalesville-----	eastern white pine--	65	114	---
	northern red oak----	60	43	
	sugar maple-----	---	0	
Urban land-----	---	---	---	---
273C:				
Urban land-----	---	---	---	---
Charlton-----	eastern hemlock----	---	---	---
	eastern white pine--	65	114	
	northern red oak----	65	43	
	red maple-----	55	29	
	shagbark hickory----	---	0	
	sugar maple-----	55	29	
	white oak-----	---	---	
Chatfield-----	eastern hemlock----	---	---	---
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
273C:				
Chatfield-----	eastern hemlock----	---	---	---
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
	white oak-----	---	---	
275E:				
Urban land-----	---	---	---	---
Chatfield-----	eastern hemlock----	---	---	---
	northern red oak----	70	57	
	sugar maple-----	65	43	
	white ash-----	75	43	
	white oak-----	---	---	
Rock outcrop-----	---	---	---	---
282B:				
Broadbrook-----	eastern white pine--	75	143	---
	northern red oak----	70	57	
	tuliptree-----	93	100	
	white ash-----	76	43	
Urban land-----	---	---	---	---
284B:				
Paxton-----	eastern white pine--	66	114	---
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---
284C:				
Paxton-----	eastern white pine--	66	114	---
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---
284D:				
Paxton-----	eastern white pine--	66	114	---
	northern red oak----	65	43	
	sugar maple-----	75	43	
	white oak-----	---	---	
Urban land-----	---	---	---	---
287B:				
Wethersfield-----	eastern white pine--	75	143	---
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
Urban land-----	---	---	---	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
287C:				
Wethersfield-----	eastern white pine--	75	143	---
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
Urban land-----	---	---	---	---
287D:				
Wethersfield-----	eastern white pine--	75	143	---
	northern red oak----	74	57	
	sugar maple-----	63	43	
	tuliptree-----	87	86	
Urban land-----	---	---	---	---
290B:				
Stockbridge-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
Urban land-----	---	---	---	---
290C:				
Stockbridge-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
Urban land-----	---	---	---	---
290D:				
Stockbridge-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
Urban land-----	---	---	---	---
301:				
Beaches-----	---	---	---	---
Udipsamments-----	American holly-----	---	0	---
	eastern redcedar----	---	0	
	pitch pine-----	---	0	
302:				
Dumps-----	---	---	---	---
303:				
Pits, quarries-----	---	---	---	---
304:				
Udorthents-----	American beech-----	---	---	---
	eastern hemlock-----	---	---	
	eastern white pine--	65	114	
	northern red oak----	55	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
305:				
Udorthents-----	---	---	---	---
Pits-----	---	---	---	---
306:				
Udorthents-----	---	---	---	---
Urban land-----	---	---	---	---
307:				
Urban land-----	---	---	---	---
308:				
Udorthents-----	---	---	---	---
309:				
Udorthents-----	---	---	---	---
310:				
Udorthents, Periodically Flooded-----	---	---	---	---
401C:				
Macomber-----	American beech-----	---	0	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	northern red oak----	70	57	
	paper birch-----	60	57	
	sugar maple-----	65	43	
	white oak-----	70	57	
Taconic-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	jack pine-----	---	---	
	northern red oak----	50	29	
	paper birch-----	53	43	
	sugar maple-----	50	29	
402D:				
Macomber-----	American beech-----	---	0	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	northern red oak----	70	57	
	paper birch-----	60	57	
	sugar maple-----	65	43	
	white oak-----	70	57	
Taconic-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	jack pine-----	---	---	
	northern red oak----	50	29	
	paper birch-----	53	43	
	sugar maple-----	50	29	
Rock outcrop-----	---	---	---	---
403C:				
Taconic-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	jack pine-----	---	---	
	northern red oak----	50	29	
	paper birch-----	53	43	
	sugar maple-----	50	29	
Rock outcrop-----	---	---	---	---

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
403E:				
Taconic-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	jack pine-----	---	---	
	northern red oak----	50	29	
	paper birch-----	53	43	
	sugar maple-----	50	29	
Rock outcrop-----	---	---	---	---
403F:				
Taconic-----	American beech-----	---	0	---
	eastern hemlock-----	---	0	
	jack pine-----	---	---	
	northern red oak----	50	29	
	paper birch-----	53	43	
	sugar maple-----	50	29	
Rock outcrop-----	---	---	---	---
405C:				
Dummerston-----	American beech-----	---	0	eastern white pine,
	eastern hemlock-----	---	0	northern red oak,
	eastern white pine--	---	0	sugar maple
	northern red oak----	---	0	
	paper birch-----	---	0	
	red maple-----	60	43	
	sugar maple-----	61	43	
	white ash-----	---	0	
	white oak-----	---	0	
405E:				
Dummerston-----	American beech-----	---	0	eastern white pine,
	eastern hemlock-----	---	0	northern red oak,
	eastern white pine--	---	0	sugar maple
	northern red oak----	---	0	
	paper birch-----	---	0	
	red maple-----	60	43	
	sugar maple-----	61	43	
	white ash-----	---	0	
	white oak-----	---	0	
407C:				
Lanesboro-----	American beech-----	---	---	ash, northern red
	eastern hemlock-----	65	0	oak, sugar maple,
	eastern white pine--	65	114	tuliptree, white
	northern red oak----	55	43	oak
	sugar maple-----	65	43	
407E:				
Lanesboro-----	American beech-----	---	---	ash, northern red
	eastern hemlock-----	65	0	oak, sugar maple,
	eastern white pine--	65	114	tuliptree, white
	northern red oak----	55	43	oak
	sugar maple-----	65	43	

Table 8.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
408C: Fullam-----	American beech-----	---	0	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	---	0	
	eastern white pine--	---	0	
	northern red oak----	65	43	
	paper birch-----	---	0	
	sugar maple-----	60	43	
	white ash-----	---	0	
	yellow birch-----	---	0	
409B: Brayton-----	eastern hemlock-----	---	---	green ash, red maple, tuliptree
	eastern white pine--	67	114	
	red maple-----	65	43	
	red spruce-----	50	114	
	tamarack-----	60	57	
412B: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak----	65	43	
412C: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak----	65	43	
412D: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak----	65	43	
413C: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak----	65	43	
Millsite-----	American beech-----	---	---	American beech, eastern hemlock, eastern white pine, northern red oak
	eastern hemlock-----	---	---	
	northern red oak----	60	43	
	sugar maple-----	73	43	
	white ash-----	75	43	
413E: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak----	65	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
413E: Millsite-----	American beech-----	---	---	American beech, eastern hemlock, eastern white pine, northern red oak
	eastern hemlock-----	---	---	
	northern red oak----	60	43	
	sugar maple-----	73	43	
	white ash-----	75	43	
414: Fredon, cold-----	eastern white pine--	70	129	green ash, red maple, tuliptree
	northern red oak----	60	43	
	red maple-----	70	43	
	tuliptree-----	80	72	
415C: Millsite-----	American beech-----	---	---	American beech, eastern hemlock, eastern white pine, northern red oak
	eastern hemlock-----	---	---	
	northern red oak----	60	43	
	sugar maple-----	73	43	
	white ash-----	75	43	
Westminster-----	American beech-----	---	0	eastern hemlock, eastern white pine
	eastern hemlock-----	---	0	
	eastern white pine--	56	100	
	northern red oak----	54	43	
	paper birch-----	---	0	
	sugar maple-----	---	0	
Rock outcrop-----	---	---	---	---
415E: Millsite-----	American beech-----	---	---	American beech, eastern hemlock, eastern white pine, northern red oak
	eastern hemlock-----	---	---	
	northern red oak----	60	43	
	sugar maple-----	73	43	
	white ash-----	75	43	
Westminster-----	American beech-----	---	0	eastern hemlock, eastern white pine
	eastern hemlock-----	---	0	
	eastern white pine--	56	100	
	northern red oak----	54	43	
	paper birch-----	---	0	
	sugar maple-----	---	0	
Rock outcrop-----	---	---	---	---
416E: Rock outcrop-----	---	---	---	---
Westminster-----	American beech-----	---	0	eastern hemlock, eastern white pine
	eastern hemlock-----	---	0	
	eastern white pine--	56	100	
	northern red oak----	54	43	
	paper birch-----	---	0	
	sugar maple-----	---	0	
416F: Rock outcrop-----	---	---	---	---
Westminster-----	American beech-----	---	0	eastern hemlock, eastern white pine
	eastern hemlock-----	---	0	
	eastern white pine--	56	100	
	northern red oak----	54	43	
	paper birch-----	---	0	
	sugar maple-----	---	0	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
417B: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak---	65	43	
417C: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak---	65	43	
417D: Bice-----	American beech-----	---	---	eastern hemlock, eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	---	
	paper birch-----	---	---	
	sugar maple-----	---	---	
	eastern white pine--	65	114	
	northern red oak---	65	43	
418C: Schroon-----	eastern hemlock-----	---	---	black cherry, eastern hemlock, eastern white pine, northern red oak
	northern red oak---	---	---	
	sugar maple-----	55	29	
	eastern white pine--	55	86	
	black cherry-----	72	43	
420A: Schroon-----	eastern hemlock-----	---	---	black cherry, eastern hemlock, eastern white pine, northern red oak
	northern red oak---	---	---	
	sugar maple-----	55	29	
	eastern white pine--	55	86	
	black cherry-----	72	43	
420B: Schroon-----	eastern hemlock-----	---	---	black cherry, eastern hemlock, eastern white pine, northern red oak
	northern red oak---	---	---	
	sugar maple-----	55	29	
	eastern white pine--	55	86	
	black cherry-----	72	43	
421A: Ninigret, cold-----	eastern white pine--	75	143	eastern white pine, northern red oak, white oak
	northern red oak---	65	43	
	red maple-----	60	43	
	sugar maple-----	55	29	
	white oak-----	---	---	
423A: Sudbury, cold-----	eastern white pine--	60	100	eastern white pine, northern red oak, white oak
	northern red oak---	45	29	
	white oak-----	---	---	
424B: Shelburne-----	American beech-----	---	---	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	---	---	
	eastern white pine--	60	100	
	northern red oak---	60	43	
	sugar maple-----	70	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
424C: Shelburne-----	American beech-----	---	---	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	---	---	
	eastern white pine--	60	100	
	northern red oak----	60	43	
	sugar maple-----	70	43	
424D: Shelburne-----	American beech-----	---	---	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	---	---	
	eastern white pine--	60	100	
	northern red oak----	60	43	
	sugar maple-----	70	43	
425B: Shelburne-----	American beech-----	---	---	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	---	---	
	eastern white pine--	60	100	
	northern red oak----	60	43	
	sugar maple-----	70	43	
425C: Shelburne-----	American beech-----	---	---	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	---	---	
	eastern white pine--	60	100	
	northern red oak----	60	43	
	sugar maple-----	70	43	
426D: Shelburne-----	American beech-----	---	---	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	---	---	
	eastern white pine--	60	100	
	northern red oak----	60	43	
	sugar maple-----	70	43	
427B: Ashfield-----	American beech-----	66	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	60	0	
	eastern white pine--	71	129	
	northern red oak----	---	---	
	paper birch-----	59	57	
	sugar maple-----	57	29	
	white ash-----	61	43	
427C: Ashfield-----	American beech-----	66	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	60	0	
	eastern white pine--	71	129	
	northern red oak----	---	---	
	paper birch-----	59	57	
	sugar maple-----	57	29	
	white ash-----	61	43	
428A: Ashfield-----	American beech-----	66	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	60	0	
	eastern white pine--	71	129	
	northern red oak----	---	---	
	paper birch-----	59	57	
	sugar maple-----	57	29	
	white ash-----	61	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
428B: Ashfield-----	American beech-----	66	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	60	0	
	eastern white pine--	71	129	
	northern red oak----	---	---	
	paper birch-----	59	57	
	sugar maple-----	57	29	
	white ash-----	61	43	
428C: Ashfield-----	American beech-----	66	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern hemlock-----	60	0	
	eastern white pine--	71	129	
	northern red oak----	---	---	
	paper birch-----	59	57	
	sugar maple-----	57	29	
	white ash-----	61	43	
429A: Agawam, cold-----	eastern white pine--	70	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	
429B: Agawam, cold-----	eastern white pine--	70	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	
429C: Agawam, cold-----	eastern white pine--	70	129	eastern white pine, northern red oak, white oak
	northern red oak----	65	129	
	sugar maple-----	---	0	
	white oak-----	---	---	
433: Moosilauke-----	eastern hemlock-----	54	114	green ash, red maple, tuliptree
	eastern white pine--	68	114	
	red maple-----	75	43	
	white ash-----	61	43	
434A: Merrimac, cold-----	eastern white pine--	64	114	eastern white pine, white oak
	northern red oak----	51	29	
	sugar maple-----	58	43	
	white oak-----	---	---	
434B: Merrimac, cold-----	eastern white pine--	64	114	eastern white pine, white oak
	northern red oak----	51	29	
	sugar maple-----	58	43	
	white oak-----	---	---	
434C: Merrimac, cold-----	eastern white pine--	64	114	eastern white pine, white oak
	northern red oak----	51	29	
	sugar maple-----	58	43	
	white oak-----	---	---	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
435: Scarboro-----	Atlantic white cedar	45	0	---
	eastern white pine--	55	86	
	red maple-----	55	29	
436: Halsey-----	red maple-----	55	29	---
	river birch-----	---	0	
	swamp white oak----	---	0	
437: Wonsqueak-----	black spruce-----	20	29	---
	eastern hemlock----	---	0	
	red maple-----	---	---	
	red spruce-----	---	0	
	tamarack-----	---	---	
438: Bucksport-----	black spruce-----	25	29	---
	eastern hemlock----	---	---	
	red maple-----	---	0	
	red spruce-----	---	---	
	tamarack-----	---	0	
	yellow birch-----	---	---	
440A: Boscawen-----	American beech-----	---	---	eastern hemlock, eastern white pine
	eastern hemlock----	---	---	
	eastern white pine--	60	100	
	northern red oak----	49	29	
	sweet birch-----	---	---	
440C: Boscawen-----	American beech-----	---	---	eastern hemlock, eastern white pine
	eastern hemlock----	---	---	
	eastern white pine--	60	100	
	northern red oak----	49	29	
	sweet birch-----	---	---	
440E: Boscawen-----	American beech-----	---	---	eastern hemlock, eastern white pine
	eastern hemlock----	---	---	
	eastern white pine--	60	100	
	northern red oak----	49	29	
	sweet birch-----	---	---	
442: Brayton-----	eastern hemlock----	---	---	green ash, red maple, tuliptree
	eastern white pine--	67	114	
	red maple-----	65	43	
	red spruce-----	50	114	
	tamarack-----	60	57	
443: Brayton-----	eastern hemlock----	---	---	green ash, red maple, tuliptree
	eastern white pine--	67	114	
	red maple-----	65	43	
	red spruce-----	50	114	
	tamarack-----	60	57	
Loonmeadow-----	eastern hemlock----	---	---	---
	red maple-----	---	---	
	yellow birch-----	---	---	
	eastern white pine--	65	114	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
448B: Hogansburg-----	basswood-----	65	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern white pine--	75	143	
	northern red oak----	70	57	
	quaking aspen-----	---	0	
	red maple-----	70	43	
	sugar maple-----	60	43	
	white ash-----	66	43	
	yellow birch-----	60	43	
449B: Hogansburg-----	basswood-----	65	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
449C: Hogansburg-----	basswood-----	65	43	ash, northern red oak, sugar maple, tuliptree, white oak
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
450B: Pyrities-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
450C: Pyrities-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
450D: Pyrities-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
451B: Pyrities-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
451C: Pyrities-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	
451D: Pyrities-----	American beech-----	---	0	eastern white pine, northern red oak, sugar maple
	eastern hemlock-----	---	0	
	eastern white pine--	75	143	
	northern red oak----	70	57	
	sugar maple-----	60	43	

Table 8.—Forestland Productivity—Continued

Map symbol and soil name	Potential productivity			Trees to manage
	Common trees	Site index	Volume of wood fiber cu ft/ac	
457: Mudgepond-----	eastern white pine--	75	143	green ash, red maple, tuliptree
	green ash-----	---	---	
	northern red oak----	70	57	
	red maple-----	75	43	
458: Mudgepond-----	eastern white pine--	75	143	green ash, red maple, tuliptree
	green ash-----	---	---	
	northern red oak----	70	57	
	red maple-----	75	43	
Alden-----	red maple-----	50	29	---
501: Ondawa-----	eastern hemlock----	---	---	---
	eastern white pine--	70	129	
	northern red oak----	65	43	
	sugar maple-----	60	43	
	yellow birch-----	---	---	
503: Rumney-----	eastern white pine--	65	114	---
	red maple-----	75	43	
508: Medomak-----	eastern white pine--	50	86	---
	red maple-----	50	29	

Table 9.--Forestland Management

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Slight		Poorly suited Wetness	1.00	Severe Wetness Strength Slope	0.77 0.38 0.10
3: Ridgebury-----	40	Moderate Stoniness	0.50	Poorly suited Wetness Rock fragments	1.00 0.50	Severe Wetness Strength Slope	0.77 0.38 0.10
Leicester-----	35	Moderate Stoniness	0.50	Poorly suited Wetness Rock fragments	1.00 0.50	Moderate Wetness Strength Slope	0.58 0.38 0.10
Whitman-----	15	Severe Wetness Stoniness	1.00 0.50	Poorly suited Ponding Wetness Rock fragments	1.00 1.00 0.50	Severe Wetness Strength Slope	1.00 0.38 0.05
4: Leicester-----	80	Slight		Poorly suited Wetness	1.00	Moderate Wetness Strength Slope	0.58 0.38 0.10
5: Wilbraham-----	80	Slight		Poorly suited Wetness Low strength	1.00 0.50	Severe Strength Wetness Slope	0.75 0.58 0.10
6: Wilbraham-----	60	Moderate Stoniness	0.50	Poorly suited Wetness Rock fragments Low strength	1.00 0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.10
Menlo-----	25	Severe Wetness Stoniness	1.00 0.50	Poorly suited Ponding Wetness Rock fragments	1.00 1.00 0.50	Severe Wetness Strength Slope	1.00 0.75 0.05
7: Mudgepond-----	85	Severe Wetness Low strength	1.00 0.50	Poorly suited Wetness Low strength	1.00 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8: Mudgepond-----	45	Severe Wetness Stoniness Low strength	1.00 0.50 0.50	Poorly suited Wetness Rock fragments Low strength	1.00 0.50 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
Alden-----	35	Moderate Stoniness Low strength	0.50 0.50	Poorly suited Ponding Wetness Rock fragments Low strength	1.00 1.00 0.50 0.50	Severe Wetness Strength Slope	0.75 0.75 0.10
9: Scitico-----	40	Moderate Low strength	0.50	Poorly suited Wetness Low strength	1.00 0.50	Severe Wetness Strength Slope	0.75 0.75 0.10
Shaker-----	30	Slight		Poorly suited Wetness	1.00	Severe Wetness Strength Slope	0.75 0.38 0.10
Maybid-----	15	Severe Wetness Low strength	1.00 0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
10: Raynham-----	80	Moderate Low strength	0.50	Poorly suited Wetness Low strength	1.00 0.50	Severe Strength Wetness Slope	0.75 0.58 0.10
12: Raypol-----	80	Moderate Low strength	0.50	Poorly suited Wetness Low strength	1.00 0.50	Severe Strength Wetness Slope	0.75 0.58 0.10
13: Walpole-----	80	Moderate Sandiness	0.50	Poorly suited Wetness Sandiness	1.00 0.50	Severe Wetness Strength Slope	0.66 0.38 0.10
14: Fredon-----	85	Severe Wetness Low strength Sandiness	1.00 0.50 0.50	Poorly suited Wetness Sandiness Low strength	1.00 0.50 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
15: Scarboro-----	80	Severe Wetness	1.00	Poorly suited Ponding Low strength	1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
16: Halsey-----	80	Severe Wetness Low strength Sandiness	1.00 0.50 0.50	Poorly suited Ponding Wetness Sandiness Low strength	1.00 1.00 0.50 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
17: Timakwa-----	45	Severe Wetness Sandiness	1.00 0.50	Poorly suited Ponding Low strength Wetness Sandiness	1.00 1.00 1.00 0.50	Severe Wetness Strength Slope	1.00 0.75 0.05
Natchaug-----	40	Severe Wetness	1.00	Poorly suited Ponding Wetness	1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05
18: Catden-----	40	Severe Wetness	1.00	Poorly suited Ponding Low strength Wetness	1.00 1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05
Freetown-----	40	Severe Wetness	1.00	Poorly suited Ponding Low strength Wetness	1.00 1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05
20A: Ellington-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.50 0.15
21A: Ninigret-----	60	Slight		Well suited		Severe Wetness Strength Slope	0.67 0.38 0.15
Tisbury-----	25	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.67 0.10
22A: Hero-----	85	Slight		Well suited		Moderate Wetness Strength Slope	0.50 0.38 0.10
22B: Hero-----	85	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
23A: Sudbury-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.42 0.38 0.15
24A: Deerfield-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.42 0.38 0.10
25A: Brancroft-----	80	Moderate Low strength	0.50	Moderately suited Low strength Wetness	0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.10
25B: Brancroft-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.30
25C: Brancroft-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.58
26A: Berlin-----	80	Moderate Low strength	0.50	Moderately suited Low strength Wetness	0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.10
26B: Berlin-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.30
27A: Belgrade-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.50 0.15
28A: Elmridge-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.58 0.38 0.10
28B: Elmridge-----	80	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.58 0.38 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
29A: Agawam-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.10
29B: Agawam-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness Slope	0.50 0.50	Slight Strength Slope	0.38 0.30
29C: Agawam-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.60 0.38
30A: Branford-----	80	Moderate Low strength Sandiness	0.50 0.50	Moderately suited Sandiness Low strength	0.50 0.50	Severe Strength Slope	0.75 0.10
30B: Branford-----	80	Moderate Low strength Sandiness	0.50 0.50	Moderately suited Sandiness Low strength	0.50 0.50	Severe Strength Slope	0.75 0.23
30C: Branford-----	80	Moderate Low strength Sandiness	0.50 0.50	Moderately suited Slope Sandiness Low strength	0.50 0.50 0.50	Severe Strength Slope	0.75 0.60
31A: Copake-----	85	Slight		Well suited		Slight Strength Slope	0.38 0.10
31B: Copake-----	85	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
31C: Copake-----	85	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
32A: Haven-----	60	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Slope	0.75 0.10
Enfield-----	25	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Slope	0.75 0.10
32B: Haven-----	60	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
32B: Enfield-----	25	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
32C: Haven-----	60	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
Enfield-----	25	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
33A: Hartford-----	80	Slight		Well suited		Slight Strength Slope	0.38 0.10
33B: Hartford-----	80	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
34A: Merrimac-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.10
34B: Merrimac-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness Slope	0.50 0.50	Slight Strength Slope	0.38 0.30
34C: Merrimac-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.60 0.38
35A: Penwood-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.10
35B: Penwood-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness Slope	0.50 0.50	Slight Strength Slope	0.38 0.30
36A: Windsor-----	80	Slight		Well suited		Slight Strength Slope	0.38 0.10
36B: Windsor-----	80	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
36C: Windsor-----	80	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
37A: Manchester-----	80	Slight		Well suited		Slight Slope Strength	0.38 0.10
37C: Manchester-----	80	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
37E: Manchester-----	80	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
38A: Hinckley-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Slope Strength	0.38 0.10
38C: Hinckley-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.45 0.38
38E: Hinckley-----	80	Moderate Slope Sandiness	0.50 0.50	Poorly suited Slope Sandiness	1.00 0.50	Moderate Slope Strength	1.00 0.38
39A: Groton-----	85	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Slope Strength	0.38 0.10
39C: Groton-----	85	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.45 0.38
39E: Groton-----	85	Moderate Slope Sandiness	0.50 0.50	Poorly suited Slope Sandiness	1.00 0.50	Moderate Slope Strength	1.00 0.38
40A: Ludlow-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Slope Strength Wetness	0.75 0.50 0.10
40B: Ludlow-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Slope Strength Wetness	0.75 0.50 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
41B: Ludlow-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.50 0.25
42C: Ludlow-----	80	Moderate Stoniness Low strength	0.50 0.50	Moderately suited Slope Rock fragments Low strength	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.50 0.45
43A: Rainbow-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.58 0.10
43B: Rainbow-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.30
44B: Rainbow-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.30
45A: Woodbridge-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.58 0.38 0.10
45B: Woodbridge-----	80	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.58 0.38 0.30
45C: Woodbridge-----	80	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.58 0.38
46B: Woodbridge-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.58 0.38 0.25
46C: Woodbridge-----	80	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.58 0.38

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
47C: Woodbridge-----	80	Moderate Stoniness	0.50	Moderately suited Slope Rock fragments	0.50 0.50	Moderate Wetness Slope Strength	0.58 0.45 0.38
48B: Georgia-----	50	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.58 0.25
Amenia-----	35	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.58 0.25
48C: Georgia-----	50	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.58
Amenia-----	35	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.58
49B: Georgia-----	50	Moderate Low strength Stoniness	0.50 0.50	Moderately suited Low strength Rock fragments Slope	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.30
Amenia-----	35	Moderate Low strength Stoniness	0.50 0.50	Moderately suited Low strength Rock fragments Slope	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.30
49C: Georgia-----	50	Moderate Stoniness Low strength	0.50 0.50	Moderately suited Slope Low strength Rock fragments	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.58
Amenia-----	35	Moderate Stoniness Low strength	0.50 0.50	Moderately suited Slope Low strength Rock fragments	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.58
50A: Sutton-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.50 0.38 0.10
50B: Sutton-----	80	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
51B: Sutton-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.50 0.38 0.25
52C: Sutton-----	80	Moderate Stoniness	0.50	Moderately suited Slope Rock fragments	0.50 0.50	Moderate Wetness Slope Strength	0.50 0.45 0.38
53A: Wapping-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.50 0.10
53B: Wapping-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Wetness Slope	0.75 0.50 0.30
54B: Wapping-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.50 0.25
55A: Watchaug-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.50 0.38 0.10
55B: Watchaug-----	80	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30
56B: Watchaug-----	80	Slight		Well suited		Moderate Wetness Strength Slope	0.50 0.38 0.25
57B: Gloucester-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness Slope	0.50 0.50	Slight Strength Slope	0.38 0.30
57C: Gloucester-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.60 0.38

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
57D: Gloucester-----	80	Moderate Slope Sandiness	0.50 0.50	Poorly suited Slope Sandiness	1.00 0.50	Moderate Slope Strength	1.00 0.38
58B: Gloucester-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness Slope	0.50 0.50	Slight Strength Slope	0.38 0.30
58C: Gloucester-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.60 0.38
59C: Gloucester-----	80	Moderate Stoniness Sandiness	0.50 0.50	Moderately suited Slope Rock fragments Sandiness	0.50 0.50 0.50	Slight Slope Strength	0.45 0.38
59D: Gloucester-----	80	Moderate Slope Stoniness Sandiness	0.50 0.50 0.50	Poorly suited Slope Rock fragments Sandiness	1.00 0.50 0.50	Moderate Slope Strength	1.00 0.38
60B: Canton-----	45	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
Charlton-----	35	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
60C: Canton-----	45	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
Charlton-----	35	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
60D: Canton-----	45	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Charlton-----	35	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
61B: Canton-----	45	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
Charlton-----	35	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
61C: Canton-----	45	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
Charlton-----	35	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
62C: Canton-----	45	Moderate Stoniness	0.50	Moderately suited Slope Rock fragments	0.50 0.50	Slight Slope Strength	0.45 0.38
Charlton-----	35	Moderate Stoniness	0.50	Moderately suited Slope Rock fragments	0.50 0.50	Slight Slope Strength	0.45 0.38
62D: Canton-----	45	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments	1.00 0.50	Moderate Slope Strength	1.00 0.38
Charlton-----	35	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments	1.00 0.50	Moderate Slope Strength	1.00 0.38
63B: Cheshire-----	80	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
63C: Cheshire-----	80	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
63D: Cheshire-----	80	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
64B: Cheshire-----	80	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
64C: Cheshire-----	80	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
65C: Cheshire-----	80	Moderate Stoniness	0.50	Moderately suited Slope Rock fragments	0.50 0.50	Slight Slope Strength	0.45 0.38
65D: Cheshire-----	80	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments	1.00 0.50	Moderate Slope Strength	1.00 0.38

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
66B: Narragansett-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Slope	0.75 0.25
66C: Narragansett-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
67B: Narragansett-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
67C: Narragansett-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
68C: Narragansett-----	80	Moderate Stoniness Low strength	0.50 0.50	Moderately suited Slope Rock fragments Low strength	0.50 0.50 0.50	Severe Strength Slope	0.75 0.45
68D: Narragansett-----	80	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments Low strength	1.00 0.50 0.50	Severe Slope Strength	1.00 0.75
69B: Yalesville-----	75	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
69C: Yalesville-----	75	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
70C: Branford-----	50	Moderate Low strength Sandiness	0.50 0.50	Moderately suited Slope Sandiness Low strength	0.50 0.50 0.50	Severe Strength Slope	0.75 0.45
Holyoke-----	30	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Severe Strength Slope	0.75 0.45
71C: Brookfield-----	45	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
71C: Brimfield-----	30	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
Rock outcrop-----	15	Not rated		Not rated		Not rated	
71E: Brookfield-----	45	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Brimfield-----	30	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Rock outcrop-----	15	Not rated		Not rated		Not rated	
73C: Charlton-----	45	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
Chatfield-----	30	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
73E: Charlton-----	45	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Chatfield-----	30	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
74C: Narragansett-----	55	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.45
Hollis-----	20	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
75C: Hollis-----	35	Severe Restrictive layer Stoniness	1.00 0.50	Moderately suited Slope Rock fragments	0.50 0.50	Slight Slope Strength	0.45 0.38
Chatfield-----	30	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
Rock outcrop-----	15	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
75E: Hollis-----	35	Severe Restrictive layer	1.00	Poorly suited Slope	1.00	Moderate Slope	1.00
		Slope	0.50	Rock fragments	0.50	Strength	0.38
		Stoniness	0.50				
Chatfield-----	30	Severe Restrictive layer	1.00	Poorly suited Slope	1.00	Moderate Slope	1.00
		Slope	0.50			Strength	0.38
Rock outcrop-----	15	Not rated		Not rated		Not rated	
76E: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Severe Restrictive layer	1.00	Poorly suited Slope	1.00	Moderate Slope	1.00
		Slope	0.50	Rock fragments	0.50	Strength	0.38
		Stoniness	0.50				
76F: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Severe Slope	1.00	Poorly suited Slope	1.00	Moderate Slope	1.00
		Stoniness	0.50	Rock fragments	0.50	Strength	0.38
76F: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Severe Slope	1.00	Poorly suited Slope	1.00	Moderate Slope	1.00
		Stoniness	0.50	Rock fragments	0.50	Strength	0.38
77C: Cheshire-----	45	Slight		Moderately suited Slope	0.50	Slight Slope	0.45
						Strength	0.38
Holyoke-----	35	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Severe Strength	0.75
						Slope	0.45
77D: Cheshire-----	45	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope	1.00
						Strength	0.38
Holyoke-----	35	Severe Restrictive layer	1.00	Poorly suited Slope	1.00	Severe Slope	1.00
		Slope	0.50			Strength	0.75
78C: Holyoke-----	50	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Severe Strength	0.75
						Slope	0.45
Rock outcrop-----	25	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
78E: Holyoke-----	50	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Severe Slope Strength	1.00 0.75
Rock outcrop-----	25	Not rated		Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Holyoke-----	25	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Severe Slope Strength	1.00 0.75
80B: Bernardston-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.30 0.25
80C: Bernardston-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.25
81C: Bernardston-----	80	Moderate Stoniness Low strength	0.50 0.50	Moderately suited Slope Rock fragments Low strength Wetness	0.50 0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.45 0.25
81D: Bernardston-----	80	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments Low strength Wetness	1.00 0.50 0.50 0.50	Severe Slope Strength Wetness	1.00 0.75 0.25
82B: Broadbrook-----	80	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Slope Wetness	0.75 0.25 0.17
82C: Broadbrook-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.17

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
82D: Broadbrook-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	1.00 0.75 0.17
83B: Broadbrook-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope Wetness	0.75 0.30 0.17
83C: Broadbrook-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.17
84B: Paxton-----	55	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30
Montauk-----	30	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30
84C: Paxton-----	55	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.50 0.38
Montauk-----	30	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.50 0.38
84D: Paxton-----	55	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Wetness Strength	1.00 0.50 0.38
Montauk-----	30	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Wetness Strength	1.00 0.50 0.38
85B: Paxton-----	55	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30
Montauk-----	30	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
85C: Paxton-----	55	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.50 0.38
Montauk-----	30	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.50 0.38
86C: Paxton-----	55	Moderate Stoniness	0.50	Moderately suited Slope Rock fragments	0.50 0.50	Moderate Wetness Slope Strength	0.50 0.45 0.38
Montauk-----	30	Moderate Stoniness	0.50	Moderately suited Slope Rock fragments	0.50 0.50	Moderate Wetness Slope Strength	0.50 0.45 0.38
86D: Paxton-----	55	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments	1.00 0.50	Moderate Slope Wetness Strength	1.00 0.50 0.38
Montauk-----	30	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments	1.00 0.50	Moderate Slope Wetness Strength	1.00 0.50 0.38
87B: Wethersfield-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope Wetness	0.75 0.30 0.25
87C: Wethersfield-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.25
87D: Wethersfield-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	1.00 0.75 0.25
88B: Wethersfield-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.30 0.25
88C: Wethersfield-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.25

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
89C: Wethersfield-----	80	Moderate Stoniness Low strength	0.50 0.50	Moderately suited Slope Rock fragments Low strength Wetness	0.50 0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.45 0.25
89D: Wethersfield-----	80	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments Low strength Wetness	1.00 0.50 0.50 0.50	Severe Slope Strength Wetness	1.00 0.75 0.25
90B: Stockbridge-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
90C: Stockbridge-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
90D: Stockbridge-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
91B: Stockbridge-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
91C: Stockbridge-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
91D: Stockbridge-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
92B: Nellis-----	85	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
92C: Nellis-----	85	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
92D: Nellis-----	85	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
93C: Nellis-----	85	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
94C: Farmington-----	40	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
Nellis-----	35	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
94E: Farmington-----	40	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Nellis-----	35	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
95C: Farmington-----	60	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
Rock outcrop-----	20	Not rated		Not rated		Not rated	
95E: Farmington-----	60	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Rock outcrop-----	20	Not rated		Not rated		Not rated	
96: Ipswich-----	85	Severe Flooding Wetness	1.00 1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05
97: Pawcatuck-----	85	Severe Flooding Wetness	1.00 1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
98: Westbrook-----	80	Severe Flooding Wetness	1.00 1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05
99: Westbrook, low salt-	80	Severe Flooding Wetness	1.00 1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05
100: Suncook-----	80	Slight		Well suited		Slight Strength Wetness Slope	0.38 0.33 0.10
101: Occum-----	80	Moderate Flooding Sandiness	0.50 0.50	Moderately suited Flooding Sandiness	0.50 0.50	Moderate Wetness Strength Slope	0.50 0.38 0.10
102: Pootatuck-----	80	Severe Flooding Sandiness	1.00 0.50	Poorly suited Flooding	1.00	Moderate Wetness Strength Slope	0.50 0.38 0.10
103: Rippowam-----	80	Severe Flooding Sandiness	1.00 0.50	Poorly suited Flooding Wetness	1.00 1.00	Severe Wetness Strength Slope	0.83 0.38 0.10
104: Bash-----	80	Severe Flooding Wetness Low strength	1.00 0.50 0.50	Poorly suited Flooding Low strength	1.00 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
105: Hadley-----	80	Moderate Flooding Low strength	0.50 0.50	Moderately suited Flooding Low strength	0.50 0.50	Severe Strength Wetness Slope	0.75 0.50 0.10
106: Winooski-----	80	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Low strength	1.00 0.50	Severe Strength Wetness Slope	0.75 0.50 0.10
107: Limerick-----	50	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50	Severe Wetness Strength Slope	0.83 0.75 0.10

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
107: Lim-----	30	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50	Severe Wetness Strength Slope	0.83 0.75 0.10
108: Saco-----	80	Severe Flooding Wetness Low strength Sandiness	1.00 1.00 0.50 0.50	Poorly suited Ponding Flooding Wetness Sandiness Low strength	1.00 1.00 1.00 0.50 0.50	Severe Wetness Strength Slope	1.00 0.75 0.05
109: Fluvaquents, Frequently Flooded-	50	Severe Flooding Wetness Sandiness	1.00 1.00 0.50	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50	Severe Wetness Strength Slope	1.00 0.75 0.05
Udifulvents, Frequently Flooded-	35	Severe Flooding	1.00	Poorly suited Flooding	1.00	Severe Wetness Strength Slope	1.00 0.38 0.05
221A: Ninigret-----	40	Slight		Well suited		Severe Wetness Strength Slope	0.67 0.38 0.15
Urban land-----	35	Not rated		Not rated		Not rated	
224A: Deerfield-----	40	Slight		Well suited		Moderate Wetness Strength Slope	0.42 0.38 0.10
Urban land-----	35	Not rated		Not rated		Not rated	
225B: Brancroft-----	40	Moderate Low strength	0.50	Moderately suited Low strength Wetness	0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
226B: Berlin-----	40	Moderate Low strength	0.50	Moderately suited Low strength Wetness	0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.20
Urban land-----	35	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
228B: Elmridge-----	40	Slight		Well suited		Moderate Wetness Strength Slope	0.58 0.38 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
229B: Agawam-----	40	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
229C: Agawam-----	40	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.60 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
230B: Branford-----	40	Moderate Low strength Sandiness	0.50 0.50	Moderately suited Sandiness Low strength	0.50 0.50	Severe Strength Slope	0.75 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
230C: Branford-----	40	Moderate Low strength Sandiness	0.50 0.50	Moderately suited Slope Sandiness Low strength	0.50 0.50 0.50	Severe Strength Slope	0.75 0.60
Urban land-----	35	Not rated		Not rated		Not rated	
232B: Haven-----	40	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Slope	0.75 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
234B: Merrimac-----	40	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.20
Urban land-----	35	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
235B: Penwood-----	40	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
236B: Windsor-----	40	Slight		Well suited		Slight Strength Slope	0.38 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
237A: Manchester-----	40	Slight		Well suited		Slight Strength Slope	0.38 0.10
Urban land-----	35	Not rated		Not rated		Not rated	
237C: Manchester-----	40	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
238A: Hinckley-----	40	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.10
Urban land-----	35	Not rated		Not rated		Not rated	
238C: Hinckley-----	40	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.45 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
240B: Ludlow-----	40	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.50 0.20
Urban land-----	35	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
243B: Rainbow-----	40	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.58 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
245B: Woodbridge-----	40	Slight		Well suited		Moderate Wetness Strength Slope	0.58 0.38 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
245C: Woodbridge-----	40	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.58 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
248B: Georgia-----	40	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.58 0.25
Urban land-----	35	Not rated		Not rated		Not rated	
250B: Sutton-----	40	Slight		Well suited		Moderate Wetness Strength Slope	0.50 0.38 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
253B: Wapping-----	40	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.50 0.20
Urban land-----	35	Not rated		Not rated		Not rated	
255B: Watchaug-----	40	Slight		Well suited		Moderate Wetness Strength Slope	0.50 0.38 0.20

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
255B: Urban land-----	35	Not rated		Not rated		Not rated	
260B: Charlton-----	40	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
Urban land-----	35	Not rated		Not rated		Not rated	
260C: Charlton-----	40	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
260D: Charlton-----	40	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
263B: Cheshire-----	40	Slight		Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
Urban land-----	35	Not rated		Not rated		Not rated	
263C: Cheshire-----	40	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
266B: Narragansett-----	40	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
Urban land-----	35	Not rated		Not rated		Not rated	
269B: Yalesville-----	40	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Strength Slope	0.38 0.30
Urban land-----	35	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
269C: Yalesville-----	40	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Slight		Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
Chatfield-----	15	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
273E: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Chatfield-----	15	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
275C: Urban land-----	45	Not rated		Not rated		Not rated	
Chatfield-----	30	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.45 0.38
275E: Urban land-----	35	Not rated		Not rated		Not rated	
Chatfield-----	25	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Rock outcrop-----	15	Not rated		Not rated		Not rated	
282B: Broadbrook-----	40	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope Wetness	0.75 0.30 0.17
Urban land-----	35	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
284B: Paxton-----	40	Slight		Moderately suited Slope	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30
Urban land-----	35	Not rated		Not rated		Not rated	
284C: Paxton-----	40	Slight		Moderately suited Slope	0.50	Moderate Slope Wetness Strength	0.60 0.50 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
284D: Paxton-----	40	Moderate Slope	0.50	Poorly suited Slope	1.00	Moderate Slope Wetness Strength	1.00 0.50 0.38
Urban land-----	35	Not rated		Not rated		Not rated	
287B: Wethersfield-----	40	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope Wetness	0.75 0.30 0.25
Urban land-----	35	Not rated		Not rated		Not rated	
287C: Wethersfield-----	40	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.25
Urban land-----	35	Not rated		Not rated		Not rated	
287D: Wethersfield-----	40	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	1.00 0.75 0.25
Urban land-----	35	Not rated		Not rated		Not rated	
290B: Stockbridge-----	40	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
Urban land-----	35	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
290C: Stockbridge-----	40	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
Urban land-----	35	Not rated		Not rated		Not rated	
290D: Stockbridge-----	40	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
Urban land-----	35	Not rated		Not rated		Not rated	
301: Beaches-----	50	Not rated		Not rated		Not rated	
Udipsamments-----	35	Severe Flooding Sandiness	1.00 0.50	Poorly suited Flooding Sandiness Slope	1.00 0.50 0.50	Severe Wetness Slope Strength	1.00 0.40 0.38
302: Dumps-----	95	Not rated		Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated		Not rated	
304: Udorthents-----	90	Severe Slope	1.00	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	1.00 0.75 0.50
305: Udorthents-----	65	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	0.90 0.75 0.50
Pits-----	25	Not rated		Not rated		Not rated	
306: Udorthents-----	50	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	0.90 0.75 0.50
Urban land-----	35	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
307: Urban land-----	80	Not rated		Not rated		Not rated	
308: Udorthents-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	0.90 0.75 0.50
309: Udorthents-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength Wetness	0.90 0.75 0.50
310: Udorthents, Periodically Flooded-----	85	Severe Flooding Slope	1.00 0.50	Poorly suited Flooding Slope Low strength	1.00 1.00 0.50	Severe Slope Strength Wetness	0.90 0.75 0.50
401C: Macomber-----	55	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.60 0.08
Taconic-----	30	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Slight Slope Strength	0.40 0.08
402D: Macomber-----	50	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.08
Taconic-----	25	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.08
Rock outcrop-----	15	Not rated		Not rated		Not rated	
403C: Taconic-----	70	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Slight Slope Strength	0.40 0.08
Rock outcrop-----	25	Not rated		Not rated		Not rated	
403E: Taconic-----	70	Severe Slope	1.00	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.08

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
403E: Rock outcrop-----	20	Not rated		Not rated		Not rated	
403F: Taconic-----	70	Severe Slope	1.00	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.08
Rock outcrop-----	20	Not rated		Not rated		Not rated	
405C: Dummerston-----	85	Severe Stoniness	1.00	Moderately suited Slope	0.50	Severe Strength Slope	0.75 0.40
405E: Dummerston-----	85	Moderate Slope	0.50	Poorly suited Slope	1.00	Severe Slope Strength	1.00 0.75
407C: Lanesboro-----	85	Slight		Poorly suited Low strength Slope Wetness	1.00 0.50 0.50	Severe Strength Slope Wetness	0.75 0.50 0.25
407E: Lanesboro-----	85	Moderate Slope	0.50	Poorly suited Slope Low strength Wetness	1.00 1.00 0.50	Severe Slope Strength Wetness	1.00 0.75 0.25
408C: Fullam-----	85	Moderate Low strength	0.50	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.50 0.50
409B: Brayton-----	85	Slight		Poorly suited Low strength Wetness	1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.25
412B: Bice-----	85	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
412C: Bice-----	85	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.50

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
412D: Bice-----	85	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
413C: Bice-----	45	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.40
Millsite-----	40	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.40 0.38
413E: Bice-----	45	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
Millsite-----	40	Moderate Slope Restrictive layer	0.50 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
414: Fredon, cold-----	85	Severe Wetness Low strength Sandiness	1.00 0.50 0.50	Poorly suited Wetness Sandiness Low strength	1.00 0.50 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
415C: Millsite-----	40	Moderate Restrictive layer	0.50	Moderately suited Slope	0.50	Slight Slope Strength	0.40 0.38
Westminster-----	40	Severe Restrictive layer	1.00	Moderately suited Slope	0.50	Slight Slope Strength	0.40 0.38
Rock outcrop-----	15	Not rated		Not rated		Not rated	
415E: Millsite-----	40	Moderate Slope Restrictive layer	0.50 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Westminster-----	40	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
Rock outcrop-----	15	Not rated		Not rated		Not rated	
416E: Rock outcrop-----	70	Not rated		Not rated		Not rated	

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
416E: Westminster-----	20	Severe Restrictive layer Slope	1.00 0.50	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
416F: Rock outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Severe Slope	1.00	Poorly suited Slope	1.00	Moderate Slope Strength	1.00 0.38
417B: Bice-----	85	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
417C: Bice-----	85	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.50
417D: Bice-----	85	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
418C: Schroon-----	85	Slight		Moderately suited Slope Wetness	0.50 0.50	Moderate Wetness Slope Strength	0.50 0.45 0.38
420A: Schroon-----	85	Slight		Moderately suited Wetness	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.05
420B: Schroon-----	85	Slight		Moderately suited Wetness	0.50	Moderate Wetness Strength Slope	0.50 0.38 0.25
421A: Ninigret, cold-----	85	Slight		Well suited		Severe Wetness Strength Slope	0.67 0.38 0.10
423A: Sudbury, cold-----	85	Slight		Well suited		Moderate Wetness Strength Slope	0.42 0.38 0.10

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
424B: Shelburne-----	85	Slight		Moderately suited Slope Wetness	0.50 0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30
424C: Shelburne-----	85	Slight		Moderately suited Slope Wetness	0.50 0.50	Moderate Slope Wetness Strength	0.60 0.50 0.38
424D: Shelburne-----	85	Moderate Slope	0.50	Poorly suited Slope Wetness	1.00 0.50	Moderate Slope Wetness Strength	1.00 0.50 0.38
425B: Shelburne-----	85	Slight		Moderately suited Slope Wetness	0.50 0.50	Moderate Wetness Strength Slope	0.50 0.38 0.30
425C: Shelburne-----	85	Slight		Moderately suited Slope Wetness	0.50 0.50	Moderate Slope Wetness Strength	0.60 0.50 0.38
426D: Shelburne-----	85	Moderate Slope Stoniness	0.50 0.50	Poorly suited Slope Rock fragments Wetness	1.00 0.50 0.50	Moderate Slope Wetness Strength	1.00 0.50 0.38
427B: Ashfield-----	85	Slight		Moderately suited Wetness Low strength Slope	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.42 0.30
427C: Ashfield-----	85	Slight		Moderately suited Slope Wetness Low strength	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.42
428A: Ashfield-----	85	Slight		Moderately suited Wetness Low strength	0.50 0.50	Severe Strength Wetness Slope	0.75 0.42 0.10
428B: Ashfield-----	85	Slight		Moderately suited Wetness Low strength Slope	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.42 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
428C: Ashfield-----	85	Slight		Moderately suited Slope Wetness Low strength	 0.50 0.50 0.50	Severe Strength Slope Wetness	 0.75 0.60 0.42
429A: Agawam, cold-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness	 0.50	Slight Strength Slope	 0.38 0.10
429B: Agawam, cold-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness Slope	 0.50 0.50	Slight Strength Slope	 0.38 0.30
429C: Agawam, cold-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	 0.50 0.50	Slight Slope Strength	 0.60 0.38
433: Moosilauke-----	80	Slight		Poorly suited Wetness Sandiness Low strength	 1.00 0.50 0.50	Severe Wetness Strength Slope	 0.92 0.75 0.10
434A: Merrimac, cold-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness	 0.50	Slight Strength Slope	 0.38 0.10
434B: Merrimac, cold-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness Slope	 0.50 0.50	Slight Strength Slope	 0.38 0.30
434C: Merrimac, cold-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	 0.50 0.50	Slight Slope Strength	 0.60 0.38
435: Scarboro-----	80	Severe Wetness	1.00	Poorly suited Ponding Low strength	 1.00 1.00	Severe Wetness Strength Slope	 1.00 0.75 0.05
436: Halsey-----	80	Severe Wetness Low strength Sandiness	1.00 0.50 0.50	Poorly suited Ponding Wetness Sandiness Low strength	 1.00 1.00 0.50 0.50	Severe Wetness Strength Slope	 1.00 0.75 0.10
437: Wonsqueak-----	85	Severe Wetness	1.00	Poorly suited Ponding Low strength Wetness	 1.00 1.00 1.00	Severe Wetness Strength Slope	 1.00 0.75 0.05

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
438: Bucksport-----	85	Severe Wetness	1.00	Poorly suited Ponding Low strength Wetness	1.00 1.00 1.00	Severe Wetness Strength Slope	1.00 0.75 0.05
440A: Boscawen-----	80	Moderate Sandiness	0.50	Moderately suited Sandiness	0.50	Slight Strength Slope	0.38 0.10
440C: Boscawen-----	80	Moderate Sandiness	0.50	Moderately suited Slope Sandiness	0.50 0.50	Slight Slope Strength	0.45 0.38
440E: Boscawen-----	85	Moderate Slope Sandiness	0.50 0.50	Poorly suited Slope Sandiness	1.00 0.50	Moderate Slope Strength	1.00 0.38
442: Brayton-----	85	Slight		Poorly suited Wetness	1.00	Severe Wetness Strength Slope	0.75 0.38 0.25
443: Brayton-----	50	Moderate Stoniness	0.50	Poorly suited Wetness Rock fragments	1.00 0.50	Severe Wetness Strength Slope	0.75 0.38 0.25
Loonmeadow-----	35	Severe Wetness Stoniness	1.00 0.50	Poorly suited Ponding Wetness Rock fragments	1.00 1.00 0.50	Severe Wetness Strength Slope	1.00 0.38 0.05
448B: Hogansburg-----	85	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Strength Wetness Slope	0.75 0.58 0.25
449B: Hogansburg-----	85	Moderate Low strength Stoniness	0.50 0.50	Moderately suited Low strength Rock fragments Slope	0.50 0.50 0.50	Severe Strength Wetness Slope	0.75 0.58 0.30
449C: Hogansburg-----	85	Moderate Stoniness Low strength	0.50 0.50	Moderately suited Slope Low strength Rock fragments	0.50 0.50 0.50	Severe Strength Slope Wetness	0.75 0.60 0.58
450B: Pyrities-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
450C: Pyrities-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
450D: Pyrities-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
451B: Pyrities-----	80	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Strength Slope	0.75 0.30
451C: Pyrities-----	80	Moderate Low strength	0.50	Moderately suited Slope Low strength	0.50 0.50	Severe Strength Slope	0.75 0.60
451D: Pyrities-----	80	Moderate Slope	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Slope Strength	1.00 0.75
457: Mudgepond-----	80	Severe Wetness Low strength	1.00 0.50	Poorly suited Wetness Low strength	1.00 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
458: Mudgepond-----	55	Severe Wetness Stoniness Low strength	1.00 0.50 0.50	Poorly suited Wetness Rock fragments Low strength	1.00 0.50 0.50	Severe Wetness Strength Slope	1.00 0.75 0.10
Alden-----	35	Moderate Stoniness Low strength	0.50 0.50	Poorly suited Ponding Wetness Rock fragments Low strength	1.00 1.00 0.50 0.50	Severe Wetness Strength Slope	0.75 0.75 0.10
501: Ondawa-----	85	Moderate Flooding Low strength	0.50 0.50	Moderately suited Flooding Low strength	0.50 0.50	Severe Strength Wetness Slope	0.75 0.50 0.10
503: Rumney-----	80	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50	Severe Wetness Strength Slope	0.83 0.75 0.10

Table 9.—Forestland Management—Continued

Map symbol and soil name	Pct. of map unit	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
508: Medomak-----	85	Severe Flooding Wetness Low strength Sandiness	 1.00 1.00 0.50 0.50	Poorly suited Ponding Flooding Wetness Sandiness Low strength	 1.00 1.00 1.00 0.50 0.50	Severe Wetness Strength Slope	 1.00 0.75 0.05

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Slight		Slight		Poorly suited Wetness	1.00
3: Ridgebury-----	40	Slight		Slight		Poorly suited Wetness Rock fragments	1.00 0.50
Leicester-----	35	Slight		Slight		Poorly suited Wetness Rock fragments	1.00 0.50
Whitman-----	15	Slight		Slight		Poorly suited Ponding Wetness Rock fragments	1.00 1.00 0.50
4: Leicester-----	80	Slight		Slight		Poorly suited Wetness	1.00
5: Wilbraham-----	80	Slight		Slight		Poorly suited Wetness Low strength	1.00 0.50
6: Wilbraham-----	60	Slight		Slight		Poorly suited Wetness Rock fragments Low strength	1.00 0.50 0.50
Menlo-----	25	Slight		Slight		Poorly suited Ponding Wetness Rock fragments	1.00 1.00 0.50
7: Mudgepond-----	85	Slight		Slight		Poorly suited Wetness Low strength	1.00 0.50
8: Mudgepond-----	45	Slight		Slight		Poorly suited Wetness Rock fragments Low strength	1.00 0.50 0.50
Alden-----	35	Slight		Slight		Poorly suited Ponding Wetness Rock fragments Low strength	1.00 1.00 0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
9: Scitico-----	40	Slight		Slight		Poorly suited Wetness Low strength	1.00 0.50
Shaker-----	30	Slight		Slight		Poorly suited Wetness	1.00
Maybid-----	15	Slight		Slight		Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50
10: Raynham-----	80	Slight		Slight		Poorly suited Wetness Low strength	1.00 0.50
12: Raypol-----	80	Slight		Slight		Poorly suited Wetness Low strength	1.00 0.50
13: Walpole-----	80	Slight		Slight		Poorly suited Wetness Sandiness	1.00 0.50
14: Fredon-----	85	Slight		Slight		Poorly suited Wetness Sandiness Low strength	1.00 0.50 0.50
15: Scarboro-----	80	Slight		Slight		Poorly suited Ponding Low strength	1.00 1.00
16: Halsey-----	80	Slight		Slight		Poorly suited Ponding Wetness Sandiness Low strength	1.00 1.00 0.50 0.50
17: Timakwa-----	45	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Low strength Wetness Sandiness	1.00 1.00 1.00 0.50
Natchaug-----	40	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Wetness	1.00 1.00

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
18: Catden-----	40	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Low strength Wetness	1.00 1.00 1.00
Freetown-----	40	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Low strength Wetness	1.00 1.00 1.00
20A: Ellington-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
21A: Ninigret-----	60	Slight		Moderate Slope/erodibility	0.50	Well suited	
Tisbury-----	25	Slight		Slight		Moderately suited Low strength	0.50
22A: Hero-----	85	Slight		Slight		Well suited	
22B: Hero-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
23A: Sudbury-----	80	Slight		Moderate Slope/erodibility	0.50	Well suited	
24A: Deerfield-----	80	Slight		Slight		Well suited	
25A: Brancroft-----	80	Slight		Slight		Moderately suited Low strength Wetness	0.50 0.50
25B: Brancroft-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50
25C: Brancroft-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50
26A: Berlin-----	80	Slight		Slight		Moderately suited Low strength Wetness	0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
26B: Berlin-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50
27A: Belgrade-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
28A: Elmridge-----	80	Slight		Slight		Well suited	
28B: Elmridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
29A: Agawam-----	80	Slight		Slight		Moderately suited Sandiness	0.50
29B: Agawam-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Slope	0.50 0.50
29C: Agawam-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness	0.50 0.50
30A: Branford-----	80	Slight		Slight		Moderately suited Sandiness Low strength	0.50 0.50
30B: Branford-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Low strength	0.50 0.50
30C: Branford-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness Low strength	0.50 0.50 0.50
31A: Copake-----	85	Slight		Slight		Well suited	
31B: Copake-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
31C: Copake-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
32A: Haven-----	60	Slight		Slight		Moderately suited Low strength	0.50
Enfield-----	25	Slight		Slight		Moderately suited Low strength	0.50
32B: Haven-----	60	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
Enfield-----	25	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
32C: Haven-----	60	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
Enfield-----	25	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
33A: Hartford-----	80	Slight		Slight		Well suited	
33B: Hartford-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
34A: Merrimac-----	80	Slight		Slight		Moderately suited Sandiness	0.50
34B: Merrimac-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Slope	0.50 0.50
34C: Merrimac-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness	0.50 0.50
35A: Penwood-----	80	Slight		Slight		Moderately suited Sandiness	0.50
35B: Penwood-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Slope	0.50 0.50
36A: Windsor-----	80	Slight		Slight		Well suited	
36B: Windsor-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
36C: Windsor-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
37A: Manchester-----	80	Slight		Slight		Well suited	
37C: Manchester-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
37E: Manchester-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
38A: Hinckley-----	80	Slight		Slight		Moderately suited Sandiness	0.50
38C: Hinckley-----	80	Slight		Slight		Moderately suited Slope Sandiness	0.50 0.50
38E: Hinckley-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Sandiness	1.00 0.50
39A: Groton-----	85	Slight		Slight		Moderately suited Sandiness	0.50
39C: Groton-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Sandiness	0.50 0.50
39E: Groton-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Sandiness	1.00 0.50
40A: Ludlow-----	80	Slight		Slight		Moderately suited Low strength	0.50
40B: Ludlow-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
41B: Ludlow-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
42C: Ludlow-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments Low strength	0.50 0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
43A: Rainbow-----	80	Slight		Slight		Moderately suited Low strength	0.50
43B: Rainbow-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
44B: Rainbow-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
45A: Woodbridge-----	80	Slight		Slight		Well suited	
45B: Woodbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
45C: Woodbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
46B: Woodbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Well suited	
46C: Woodbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
47C: Woodbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
48B: Georgia-----	50	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
Amenia-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
48C: Georgia-----	50	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
Amenia-----	35	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
49B: Georgia-----	50	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Rock fragments Slope	0.50 0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
49B: Amenia-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Rock fragments Slope	0.50 0.50 0.50
49C: Georgia-----	50	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength Rock fragments	0.50 0.50 0.50
Amenia-----	35	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength Rock fragments	0.50 0.50 0.50
50A: Sutton-----	80	Slight		Slight		Well suited	
50B: Sutton-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
51B: Sutton-----	80	Slight		Moderate Slope/erodibility	0.50	Well suited	
52C: Sutton-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
53A: Wapping-----	80	Slight		Slight		Moderately suited Low strength	0.50
53B: Wapping-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
54B: Wapping-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
55A: Watchaug-----	80	Slight		Slight		Well suited	
55B: Watchaug-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
56B: Watchaug-----	80	Slight		Moderate Slope/erodibility	0.50	Well suited	
57B: Gloucester-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Slope	0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
57C: Gloucester-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Sandiness	0.50 0.50
57D: Gloucester-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Sandiness	1.00 0.50
58B: Gloucester-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Slope	0.50 0.50
58C: Gloucester-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Sandiness	0.50 0.50
59C: Gloucester-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments Sandiness	0.50 0.50 0.50
59D: Gloucester-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments Sandiness	1.00 0.50 0.50
60B: Canton-----	45	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Charlton-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
60C: Canton-----	45	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope	0.50
Charlton-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
60D: Canton-----	45	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Charlton-----	35	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
61B: Canton-----	45	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Charlton-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
61C: Canton-----	45	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope	0.50
Charlton-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
62C: Canton-----	45	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Rock fragments	0.50 0.50
Charlton-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
62D: Canton-----	45	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
Charlton-----	35	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
63B: Cheshire-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
63C: Cheshire-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
63D: Cheshire-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
64B: Cheshire-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
64C: Cheshire-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
65C: Cheshire-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
65D: Cheshire-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
66B: Narragansett-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
66C: Narragansett-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
67B: Narragansett-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
67C: Narragansett-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
68C: Narragansett-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Rock fragments Low strength	0.50 0.50 0.50
68D: Narragansett-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments Low strength	1.00 0.50 0.50
69B: Yalesville-----	75	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
69C: Yalesville-----	75	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope	0.50
70C: Branford-----	50	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness Low strength	0.50 0.50 0.50
Holyoke-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
71C: Brookfield-----	45	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Brimfield-----	30	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope	0.50
Rock outcrop-----	15	Not rated		Not rated		Not rated	
71E: Brookfield-----	45	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
71E: Brimfield-----	30	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	
73C: Charlton-----	45	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Chatfield-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
73E: Charlton-----	45	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Chatfield-----	30	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
74C: Narragansett-----	55	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
Hollis-----	20	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
75C: Hollis-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
Chatfield-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Rock outcrop-----	15	Not rated		Not rated		Not rated	
75E: Hollis-----	35	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
Chatfield-----	30	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	

7Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
76E: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
76F: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very severe Slope/erodibility	0.95	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
77C: Cheshire-----	45	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Holyoke-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
77D: Cheshire-----	45	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Holyoke-----	35	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
78C: Holyoke-----	50	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Rock outcrop-----	25	Not rated		Not rated		Not rated	
78E: Holyoke-----	50	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	25	Not rated		Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Holyoke-----	25	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
80B: Bernardston-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50
80C: Bernardston-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50
81C: Bernardston-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Rock fragments Low strength Wetness	0.50 0.50 0.50 0.50
81D: Bernardston-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments Low strength Wetness	1.00 0.50 0.50 0.50
82B: Broadbrook-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
82C: Broadbrook-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
82D: Broadbrook-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
83B: Broadbrook-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
83C: Broadbrook-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
84B: Paxton-----	55	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Montauk-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
84C: Paxton-----	55	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Montauk-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
84D: Paxton-----	55	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Montauk-----	30	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
85B: Paxton-----	55	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Montauk-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
85C: Paxton-----	55	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Montauk-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
86C: Paxton-----	55	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
Montauk-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Rock fragments	0.50 0.50
86D: Paxton-----	55	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
Montauk-----	30	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments	1.00 0.50
87B: Wethersfield-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
87C: Wethersfield-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
87D: Wethersfield-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
88B: Wethersfield-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope Wetness	0.50 0.50 0.50
88C: Wethersfield-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50
89C: Wethersfield-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Rock fragments Low strength Wetness	0.50 0.50 0.50 0.50
89D: Wethersfield-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments Low strength Wetness	1.00 0.50 0.50 0.50
90B: Stockbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
90C: Stockbridge-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
90D: Stockbridge-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
91B: Stockbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
91C: Stockbridge-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Low strength	0.50 0.50
91D: Stockbridge-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
92B: Nellis-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
92C: Nellis-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
92D: Nellis-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
93C: Nellis-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
94C: Farmington-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Nellis-----	35	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
94E: Farmington-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Nellis-----	35	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
95C: Farmington-----	60	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Rock outcrop-----	20	Not rated		Not rated		Not rated	
95E: Farmington-----	60	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	20	Not rated		Not rated		Not rated	
96: Ipswich-----	85	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00
97: Pawcatuck-----	85	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
98: Westbrook-----	80	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00 1.00
99: Westbrook, low salt-	80	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Flooding Low strength Wetness	1.00 1.00 1.00 1.00
100: Suncook-----	80	Slight		Slight		Well suited	
101: Occum-----	80	Slight		Slight		Moderately suited Flooding Sandiness	0.50 0.50
102: Pootatuck-----	80	Slight		Slight		Poorly suited Flooding	1.00
103: Rippowam-----	80	Slight		Slight		Poorly suited Flooding Wetness	1.00 1.00
104: Bash-----	80	Slight		Slight		Poorly suited Flooding Low strength	1.00 0.50
105: Hadley-----	80	Slight		Slight		Moderately suited Flooding Low strength	0.50 0.50
106: Winooski-----	80	Slight		Slight		Poorly suited Flooding Low strength	1.00 0.50
107: Limerick-----	50	Slight		Slight		Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50
Lim-----	30	Slight		Slight		Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
108: Saco-----	80	Slight		Slight		Poorly suited Ponding Flooding Wetness Sandiness Low strength	1.00 1.00 1.00 0.50 0.50
109: Fluvaquents, Frequently Flooded-	50	Slight		Slight		Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50
Udifluents, Frequently Flooded-	35	Slight		Slight		Poorly suited Flooding	1.00
221A: Ninigret-----	40	Slight		Moderate Slope/erodibility	0.50	Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	
224A: Deerfield-----	40	Slight		Slight		Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	
225B: Brancroft-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Wetness	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
226B: Berlin-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Wetness	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
228B: Elmridge-----	40	Slight		Moderate Slope/erodibility	0.50	Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	
229B: Agawam-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness	0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
229B: Urban land-----	35	Not rated		Not rated		Not rated	
229C: Agawam-----	40	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
230B: Branford-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Low strength	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
230C: Branford-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness Low strength	0.50 0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
232B: Haven-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
234B: Merrimac-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
235B: Penwood-----	40	Slight		Slight		Moderately suited Sandiness	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
236B: Windsor-----	40	Slight		Slight		Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	
237A: Manchester-----	40	Slight		Slight		Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
237C: Manchester-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
238A: Hinckley-----	40	Slight		Slight		Moderately suited Sandiness	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
238C: Hinckley-----	40	Slight		Slight		Moderately suited Slope Sandiness	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
240B: Ludlow-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
243B: Rainbow-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
245B: Woodbridge-----	40	Slight		Slight		Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	
245C: Woodbridge-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
248B: Georgia-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
Urban land-----	35	Not rated		Not rated		Not rated	

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
250B: Sutton-----	40	Slight		Slight		Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	
253B: Wapping-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
255B: Watchaug-----	40	Slight		Slight		Well suited	
Urban land-----	35	Not rated		Not rated		Not rated	
260B: Charlton-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
260C: Charlton-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
260D: Charlton-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
263B: Cheshire-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
263C: Cheshire-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
266B: Narragansett-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
269B: Yalesville-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
269C: Yalesville-----	40	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Chatfield-----	15	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
273E: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Chatfield-----	15	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
275C: Urban land-----	45	Not rated		Not rated		Not rated	
Chatfield-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
275E: Urban land-----	35	Not rated		Not rated		Not rated	

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
275E: Chatfield-----	25	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	
282B: Broadbrook-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
284B: Paxton-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
284C: Paxton-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
284D: Paxton-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
287B: Wethersfield-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
287C: Wethersfield-----	40	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
287D: Wethersfield-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
290B: Stockbridge-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
290C: Stockbridge-----	40	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength	0.50 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
290D: Stockbridge-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
301: Beaches-----	50	Not rated		Not rated		Not rated	
Udipsamments-----	35	Slight		Moderate Slope/erodibility	0.50	Poorly suited Flooding Sandiness Slope	1.00 0.50 0.50
302: Dumps-----	95	Not rated		Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated		Not rated	
304: Udorthents-----	90	Severe Slope/erodibility	0.75	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
305: Udorthents-----	65	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
Pits-----	25	Not rated		Not rated		Not rated	
306: Udorthents-----	50	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
Urban land-----	35	Not rated		Not rated		Not rated	
307: Urban land-----	80	Not rated		Not rated		Not rated	
308: Udorthents-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
309: Udorthents-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
310: Udorthents, Periodically Flooded-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Flooding Slope Low strength	1.00 1.00 0.50
401C: Macomber-----	55	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Taconic-----	30	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
402D: Macomber-----	50	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Taconic-----	25	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
403C: Taconic-----	70	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Rock outcrop-----	25	Not rated		Not rated		Not rated	
403E: Taconic-----	70	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	20	Not rated		Not rated		Not rated	
403F: Taconic-----	70	Very severe Slope/erodibility	0.95	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	20	Not rated		Not rated		Not rated	
405C: Dummerston-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
405E: Dummerston-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
407C: Lanesboro-----	85	Slight		Moderate Slope/erodibility	0.50	Poorly suited Low strength Slope Wetness	1.00 0.50 0.50
407E: Lanesboro-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength Wetness	1.00 1.00 0.50
408C: Fullam-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Low strength Wetness	0.50 0.50 0.50
409B: Brayton-----	85	Slight		Moderate Slope/erodibility	0.50	Poorly suited Low strength Wetness	1.00 1.00

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
412B: Bice-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
412C: Bice-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Low strength	0.50 0.50
412D: Bice-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
413C: Bice-----	45	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Low strength	0.50 0.50
Millsite-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
413E: Bice-----	45	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
Millsite-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
414: Fredon, cold-----	85	Slight		Slight		Poorly suited Wetness Sandiness Low strength	1.00 0.50 0.50
415C: Millsite-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Westminster-----	40	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope	0.50
Rock outcrop-----	15	Not rated		Not rated		Not rated	
415E: Millsite-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Westminster-----	40	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
416E: Rock outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
416F: Rock outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very severe Slope/erodibility	0.95	Severe Slope/erodibility	0.95	Poorly suited Slope	1.00
417B: Bice-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
417C: Bice-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Low strength	0.50 0.50
417D: Bice-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
418C: Schroon-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Wetness	0.50 0.50
420A: Schroon-----	85	Slight		Slight		Moderately suited Wetness	0.50
420B: Schroon-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Wetness	0.50
421A: Ninigret, cold-----	85	Slight		Slight		Well suited	
423A: Sudbury, cold-----	85	Slight		Slight		Well suited	
424B: Shelburne-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Wetness	0.50 0.50
424C: Shelburne-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Wetness	0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
424D: Shelburne-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Wetness	1.00 0.50
425B: Shelburne-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Wetness	0.50 0.50
425C: Shelburne-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Wetness	0.50 0.50
426D: Shelburne-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Rock fragments Wetness	1.00 0.50 0.50
427B: Ashfield-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Wetness Low strength Slope	0.50 0.50 0.50
427C: Ashfield-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Wetness Low strength	0.50 0.50 0.50
428A: Ashfield-----	85	Slight		Slight		Moderately suited Wetness Low strength	0.50 0.50
428B: Ashfield-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Wetness Low strength Slope	0.50 0.50 0.50
428C: Ashfield-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Wetness Low strength	0.50 0.50 0.50
429A: Agawam, cold-----	80	Slight		Slight		Moderately suited Sandiness	0.50
429B: Agawam, cold-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Slope	0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
429C: Agawam, cold-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness	0.50 0.50
433: Moosilauke-----	80	Slight		Slight		Poorly suited Wetness Sandiness Low strength	1.00 0.50 0.50
434A: Merrimac, cold-----	80	Slight		Slight		Moderately suited Sandiness	0.50
434B: Merrimac, cold-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Sandiness Slope	0.50 0.50
434C: Merrimac, cold-----	80	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Sandiness	0.50 0.50
435: Scarboro-----	80	Slight		Slight		Poorly suited Ponding Low strength	1.00 1.00
436: Halsey-----	80	Slight		Slight		Poorly suited Ponding Wetness Sandiness Low strength	1.00 1.00 0.50 0.50
437: Wonsqueak-----	85	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Low strength Wetness	1.00 1.00 1.00
438: Bucksport-----	85	Very Severe Organic matter content high	1.00	Very Severe Organic matter content high	1.00	Poorly suited Ponding Low strength Wetness	1.00 1.00 1.00
440A: Boscawen-----	80	Slight		Slight		Moderately suited Sandiness	0.50
440C: Boscawen-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Sandiness	0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
440E: Boscawen-----	85	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Sandiness	1.00 0.50
442: Brayton-----	85	Slight		Moderate Slope/erodibility	0.50	Poorly suited Wetness	1.00
443: Brayton-----	50	Slight		Moderate Slope/erodibility	0.50	Poorly suited Wetness Rock fragments	1.00 0.50
Loonmeadow-----	35	Slight		Slight		Poorly suited Ponding Wetness Rock fragments	1.00 1.00 0.50
448B: Hogansburg-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength	0.50
449B: Hogansburg-----	85	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Rock fragments Slope	0.50 0.50 0.50
449C: Hogansburg-----	85	Slight		Severe Slope/erodibility	0.95	Moderately suited Slope Low strength Rock fragments	0.50 0.50 0.50
450B: Pyrities-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
450C: Pyrities-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Low strength	0.50 0.50
450D: Pyrities-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
451B: Pyrities-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Low strength Slope	0.50 0.50
451C: Pyrities-----	80	Slight		Moderate Slope/erodibility	0.50	Moderately suited Slope Low strength	0.50 0.50

Table 10.—Hazard of Erosion and Suitability for Roads on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
451D: Pyrities-----	80	Moderate Slope/erodibility	0.50	Severe Slope/erodibility	0.95	Poorly suited Slope Low strength	1.00 0.50
457: Mudgepond-----	80	Slight		Slight		Poorly suited Wetness Low strength	1.00 0.50
458: Mudgepond-----	55	Slight		Slight		Poorly suited Wetness Rock fragments Low strength	1.00 0.50 0.50
Alden-----	35	Slight		Slight		Poorly suited Ponding Wetness Rock fragments Low strength	1.00 1.00 0.50 0.50
501: Ondawa-----	85	Slight		Slight		Moderately suited Flooding Low strength	0.50 0.50
503: Rumney-----	80	Slight		Slight		Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50
508: Medomak-----	85	Slight		Slight		Poorly suited Ponding Flooding Wetness Sandiness Low strength	1.00 1.00 1.00 0.50 0.50

Table 11.—Damage by Fire and Seedling Mortality on Forestland

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
3: Ridgebury-----	40	Low Texture/rock fragments	0.10	High Wetness	1.00
Leicester-----	35	Low Texture/rock fragments	0.10	High Wetness	1.00
Whitman-----	15	Low Texture/rock fragments	0.10	High Wetness	1.00
4: Leicester-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
5: Wilbraham-----	80	Moderate Texture/surface depth/rock fragments	0.50	High Wetness	1.00
6: Wilbraham-----	60	Moderate Texture/surface depth/rock fragments	0.50	High Wetness	1.00
Menlo-----	25	Low Texture/rock fragments	0.10	High Wetness	1.00
7: Mudgepond-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
8: Mudgepond-----	45	Low Texture/rock fragments	0.10	High Wetness	1.00
Alden-----	35	Moderate Texture/surface depth/rock fragments	0.50	High Wetness	1.00

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
9: Scitico-----	40	Low Texture/rock fragments	0.10	High Wetness	1.00
Shaker-----	30	Moderate Texture/surface depth/rock fragments	0.50	High Wetness	1.00
Maybid-----	15	Low Texture/rock fragments	0.10	High Wetness	1.00
10: Raynham-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
12: Raypol-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
13: Walpole-----	80	Moderate Texture/rock fragments	0.50	High Wetness	1.00
14: Fredon-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
15: Scarboro-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
16: Halsey-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
17: Timakwa-----	45	Low		High Wetness	1.00
Natchaug-----	40	Low		High Wetness	1.00
18: Catden-----	40	Low		High Wetness	1.00
Freetown-----	40	Low		High Wetness Soil reaction	1.00 0.50

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
20A: Ellington-----	80	Low Texture/rock fragments	0.10	Low	
21A: Ninigret-----	60	Low Texture/rock fragments	0.10	Low	
Tisbury-----	25	Low Texture/rock fragments	0.10	Low	
22A: Hero-----	85	Low Texture/rock fragments	0.10	Low	
22B: Hero-----	85	Low Texture/rock fragments	0.10	Low	
23A: Sudbury-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
24A: Deerfield-----	80	Moderate Texture/rock fragments	0.50	Low	
25A: Brancroft-----	80	Low Texture/rock fragments	0.10	Low	
25B: Brancroft-----	80	Low Texture/rock fragments	0.10	Low	
25C: Brancroft-----	80	Low Texture/rock fragments	0.10	Low	
26A: Berlin-----	80	Low Texture/rock fragments	0.10	Low	
26B: Berlin-----	80	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
27A: Belgrade-----	80	Low Texture/rock fragments	0.10	Low	
28A: Elmridge-----	80	Low Texture/rock fragments	0.10	Low	
28B: Elmridge-----	80	Low Texture/rock fragments	0.10	Low	
29A: Agawam-----	80	Low Texture/rock fragments	0.10	Low	
29B: Agawam-----	80	Low Texture/rock fragments	0.10	Low	
29C: Agawam-----	80	Low Texture/rock fragments	0.10	Low	
30A: Branford-----	80	Low Texture/rock fragments	0.10	Low	
30B: Branford-----	80	Low Texture/rock fragments	0.10	Low	
30C: Branford-----	80	Low Texture/rock fragments	0.10	Low	
31A: Copake-----	85	Low Texture/rock fragments	0.10	Low	
31B: Copake-----	85	Low Texture/rock fragments	0.10	Low	
31C: Copake-----	85	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
32A: Haven-----	60	Low Texture/rock fragments	0.10	Low	
Enfield-----	25	Low Texture/rock fragments	0.10	Low	
32B: Haven-----	60	Low Texture/rock fragments	0.10	Low	
Enfield-----	25	Low Texture/rock fragments	0.10	Low	
32C: Haven-----	60	Low Texture/rock fragments	0.10	Low	
Enfield-----	25	Low Texture/rock fragments	0.10	Low	
33A: Hartford-----	80	Low Texture/rock fragments	0.10	Low	
33B: Hartford-----	80	Low Texture/rock fragments	0.10	Low	
34A: Merrimac-----	80	Low Texture/rock fragments	0.10	Low	
34B: Merrimac-----	80	Low Texture/rock fragments	0.10	Low	
34C: Merrimac-----	80	Low Texture/rock fragments	0.10	Low	
35A: Penwood-----	80	Moderate Texture/rock fragments	0.50	Low	
35B: Penwood-----	80	Moderate Texture/rock fragments	0.50	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
36A: Windsor-----	80	Moderate Texture/surface depth/rock fragments	0.50	Low	
36B: Windsor-----	80	Moderate Texture/surface depth/rock fragments	0.50	Low	
36C: Windsor-----	80	Moderate Texture/surface depth/rock fragments	0.50	Low	
37A: Manchester-----	80	Low Texture/rock fragments	0.10	Low	
37C: Manchester-----	80	Low Texture/rock fragments	0.10	Low	
37E: Manchester-----	80	Low Texture/rock fragments	0.10	Low	
38A: Hinckley-----	80	Low Texture/rock fragments	0.10	Low	
38C: Hinckley-----	80	Low Texture/rock fragments	0.10	Low	
38E: Hinckley-----	80	Low Texture/slope/roc k fragments	0.10	Low	
39A: Groton-----	85	Low Texture/rock fragments	0.10	Low	
39C: Groton-----	85	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
39E: Groton-----	85	Low Texture/slope/rock fragments	0.10	Low	
40A: Ludlow-----	80	Low Texture/rock fragments	0.10	Low	
40B: Ludlow-----	80	Low Texture/rock fragments	0.10	Low	
41B: Ludlow-----	80	Low Texture/rock fragments	0.10	Low	
42C: Ludlow-----	80	Low Texture/rock fragments	0.10	Low	
43A: Rainbow-----	80	Low Texture/rock fragments	0.10	Low	
43B: Rainbow-----	80	Low Texture/rock fragments	0.10	Low	
44B: Rainbow-----	80	Low Texture/rock fragments	0.10	Low	
45A: Woodbridge-----	80	Low Texture/rock fragments	0.10	Low	
45B: Woodbridge-----	80	Low Texture/rock fragments	0.10	Low	
45C: Woodbridge-----	80	Low Texture/rock fragments	0.10	Low	
46B: Woodbridge-----	80	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
46C: Woodbridge-----	80	Low Texture/rock fragments	0.10	Low	
47C: Woodbridge-----	80	Low Texture/rock fragments	0.10	Low	
48B: Georgia-----	50	Low Texture/rock fragments	0.10	Low	
Amenia-----	35	Low Texture/rock fragments	0.10	Low	
48C: Georgia-----	50	Low Texture/rock fragments	0.10	Low	
Amenia-----	35	Low Texture/rock fragments	0.10	Low	
49B: Georgia-----	50	Low Texture/rock fragments	0.10	Low	
Amenia-----	35	Low Texture/rock fragments	0.10	Low	
49C: Georgia-----	50	Low Texture/rock fragments	0.10	Low	
Amenia-----	35	Low Texture/rock fragments	0.10	Low	
50A: Sutton-----	80	Low Texture/rock fragments	0.10	Low	
50B: Sutton-----	80	Low Texture/rock fragments	0.10	Low	
51B: Sutton-----	80	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
52C: Sutton-----	80	Low Texture/rock fragments	0.10	Low	
53A: Wapping-----	80	Low Texture/rock fragments	0.10	Low	
53B: Wapping-----	80	Low Texture/rock fragments	0.10	Low	
54B: Wapping-----	80	Low Texture/rock fragments	0.10	Low	
55A: Watchaug-----	80	Low Texture/rock fragments	0.10	Low	
55B: Watchaug-----	80	Low Texture/rock fragments	0.10	Low	
56B: Watchaug-----	80	Low Texture/rock fragments	0.10	Low	
57B: Gloucester-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
57C: Gloucester-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
57D: Gloucester-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
58B: Gloucester-----	80	Low Texture/surface depth/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
58C: Gloucester-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
59C: Gloucester-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
59D: Gloucester-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
60B: Canton-----	45	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50
Charlton-----	35	Low Texture/surface depth/rock fragments	0.10	Low	
60C: Canton-----	45	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50
Charlton-----	35	Low Texture/surface depth/rock fragments	0.10	Low	
60D: Canton-----	45	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50
Charlton-----	35	Low Texture/surface depth/rock fragments	0.10	Low	
61B: Canton-----	45	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
61B: Charlton-----	35	Low Texture/surface depth/rock fragments	0.10	Low	
61C: Canton-----	45	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50
Charlton-----	35	Low Texture/surface depth/rock fragments	0.10	Low	
62C: Canton-----	45	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50
Charlton-----	35	Low Texture/surface depth/rock fragments	0.10	Low	
62D: Canton-----	45	Low Texture/surface depth/rock fragments	0.10	Moderate Soil reaction	0.50
Charlton-----	35	Low Texture/surface depth/rock fragments	0.10	Low	
63B: Cheshire-----	80	Low Texture/rock fragments	0.10	Low	
63C: Cheshire-----	80	Low Texture/rock fragments	0.10	Low	
63D: Cheshire-----	80	Low Texture/rock fragments	0.10	Low	
64B: Cheshire-----	80	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
64C: Cheshire-----	80	Low Texture/rock fragments	0.10	Low	
65C: Cheshire-----	80	Low Texture/rock fragments	0.10	Low	
65D: Cheshire-----	80	Low Texture/rock fragments	0.10	Low	
66B: Narragansett-----	80	Low Texture/rock fragments	0.10	Low	
66C: Narragansett-----	80	Low Texture/rock fragments	0.10	Low	
67B: Narragansett-----	80	Low Texture/rock fragments	0.10	Low	
67C: Narragansett-----	80	Low Texture/rock fragments	0.10	Low	
68C: Narragansett-----	80	Low Texture/rock fragments	0.10	Low	
68D: Narragansett-----	80	Low Texture/rock fragments	0.10	Low	
69B: Yalesville-----	75	Low Texture/rock fragments	0.10	Low	
69C: Yalesville-----	75	Low Texture/rock fragments	0.10	Low	
70C: Branford-----	50	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
70C: Holyoke-----	30	Moderate Texture/surface depth/rock fragments	0.50	Moderate Soil reaction	0.50
71C: Brookfield-----	45	Low Texture/surface depth/rock fragments	0.10	Low	
Brimfield-----	30	Low Texture/surface depth/rock fragments	0.10	Low	
Rock outcrop-----	15	Not rated		Not rated	
71E: Brookfield-----	45	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Brimfield-----	30	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Rock outcrop-----	15	Not rated		Not rated	
73C: Charlton-----	45	Low Texture/surface depth/rock fragments	0.10	Low	
Chatfield-----	30	Low Texture/rock fragments	0.10	Low	
73E: Charlton-----	45	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Chatfield-----	30	Low Texture/slope/roc k fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
74C: Narragansett-----	55	Low Texture/rock fragments	0.10	Low	
Hollis-----	20	Low Texture/rock fragments	0.10	Low	
75C: Hollis-----	35	Low Texture/rock fragments	0.10	Low	
Chatfield-----	30	Low Texture/rock fragments	0.10	Low	
Rock outcrop-----	15	Not rated		Not rated	
75E: Hollis-----	35	Low Texture/slope/roc k fragments	0.10	Low	
Chatfield-----	30	Low Texture/slope/roc k fragments	0.10	Low	
Rock outcrop-----	15	Not rated		Not rated	
76E: Rock outcrop-----	55	Not rated		Not rated	
Hollis-----	25	Low Texture/rock fragments	0.10	Low	
76F: Rock outcrop-----	55	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
76F: Hollis-----	25	Low Texture/slope/rock fragments	0.10	Low	
77C: Cheshire-----	45	Low Texture/rock fragments	0.10	Low	
Holyoke-----	35	Moderate Texture/surface depth/rock fragments	0.50	Moderate Soil reaction	0.50
77D: Cheshire-----	45	Low Texture/rock fragments	0.10	Low	
Holyoke-----	35	Moderate Texture/surface depth/rock fragments	0.50	Moderate Soil reaction	0.50
				Available water	0.50
78C: Holyoke-----	50	Moderate Texture/surface depth/rock fragments	0.50	Moderate Soil reaction	0.50
Rock outcrop-----	25	Not rated		Not rated	
78E: Holyoke-----	50	Moderate Texture/slope/surface depth/rock fragments	0.50	Moderate Soil reaction	0.50
				Available water	0.50
Rock outcrop-----	25	Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
79E: Holyoke-----	25	Moderate Texture/surface depth/rock fragments	0.50	Moderate Soil reaction	0.50
				Available water	0.50
80B: Bernardston-----	80	Low Texture/rock fragments	0.10	Low	
80C: Bernardston-----	80	Low Texture/rock fragments	0.10	Low	
81C: Bernardston-----	80	Low Texture/rock fragments	0.10	Low	
81D: Bernardston-----	80	Low Texture/rock fragments	0.10	Moderate Available water	0.50
82B: Broadbrook-----	80	Low Texture/rock fragments	0.10	Low	
82C: Broadbrook-----	80	Low Texture/rock fragments	0.10	Low	
82D: Broadbrook-----	80	Low Texture/rock fragments	0.10	Low	
83B: Broadbrook-----	80	Low Texture/rock fragments	0.10	Low	
83C: Broadbrook-----	80	Low Texture/rock fragments	0.10	Low	
84B: Paxton-----	55	Low Texture/rock fragments	0.10	Low	
Montauk-----	30	Moderate Texture/surface depth/rock fragments	0.50	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
84C: Paxton-----	55	Low Texture/rock fragments	0.10	Low	
Montauk-----	30	Moderate Texture/surface depth/rock fragments	0.50	Low	
84D: Paxton-----	55	Low Texture/rock fragments	0.10	Moderate Available water	0.50
Montauk-----	30	Moderate Texture/surface depth/rock fragments	0.50	Moderate Available water	0.50
85B: Paxton-----	55	Low Texture/rock fragments	0.10	Low	
Montauk-----	30	Moderate Texture/surface depth/rock fragments	0.50	Low	
85C: Paxton-----	55	Low Texture/rock fragments	0.10	Low	
Montauk-----	30	Moderate Texture/surface depth/rock fragments	0.50	Low	
86C: Paxton-----	55	Low Texture/rock fragments	0.10	Low	
Montauk-----	30	Moderate Texture/surface depth/rock fragments	0.50	Low	
86D: Paxton-----	55	Low Texture/rock fragments	0.10	Moderate Available water	0.50
Montauk-----	30	Moderate Texture/surface depth/rock fragments	0.50	Moderate Available water	0.50

Table 11.--Damage by Fire and Seedling Mortality on Forestland--Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
87B: Wethersfield-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
87C: Wethersfield-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
87D: Wethersfield-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
88B: Wethersfield-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
88C: Wethersfield-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
89C: Wethersfield-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
89D: Wethersfield-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
90B: Stockbridge-----	80	Low Texture/rock fragments	0.10	Low	
90C: Stockbridge-----	80	Low Texture/rock fragments	0.10	Low	
90D: Stockbridge-----	80	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
91B: Stockbridge-----	80	Low Texture/rock fragments	0.10	Low	
91C: Stockbridge-----	80	Low Texture/rock fragments	0.10	Low	
91D: Stockbridge-----	80	Low Texture/rock fragments	0.10	Low	
92B: Nellis-----	85	Low Texture/rock fragments	0.10	Low	
92C: Nellis-----	85	Low Texture/rock fragments	0.10	Low	
92D: Nellis-----	85	Low Texture/rock fragments	0.10	Low	
93C: Nellis-----	85	Low Texture/rock fragments	0.10	Low	
94C: Farmington-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Nellis-----	35	Low Texture/rock fragments	0.10	Low	
94E: Farmington-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Nellis-----	35	Low Texture/rock fragments	0.10	Low	
95C: Farmington-----	60	Low Texture/surface depth/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
95C: Rock outcrop-----	20	Not rated		Not rated	
95E: Farmington-----	60	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Rock outcrop-----	20	Not rated		Not rated	
96: Ipswich-----	85	Low		High Wetness	1.00
97: Pawcatuck-----	85	Low		High Wetness	1.00
98: Westbrook-----	80	Low		High Wetness	1.00
99: Westbrook, low salt-	80	Low		High Wetness	1.00
100: Suncook-----	80	Moderate Texture/rock fragments	0.50	Low	
101: Occum-----	80	Low Texture/rock fragments	0.10	Low	
102: Pootatuck-----	80	Low Texture/surface depth/rock fragments	0.10	Low	
103: Rippowam-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
104: Bash-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
105: Hadley-----	80	Low Texture/rock fragments	0.10	Low	
106: Winooski-----	80	Low Texture/rock fragments	0.10	Low	
107: Limerick-----	50	Low Texture/rock fragments	0.10	High Wetness	1.00
Lim-----	30	Low Texture/rock fragments	0.10	High Wetness	1.00
108: Saco-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
109: Fluvaquents, Frequently Flooded-	50	Low Texture/surface depth/rock fragments	0.10	High Wetness	1.00
Udifluents, Frequently Flooded-	35	Low Texture/surface depth/rock fragments	0.10	Low	
221A: Ninigret-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
224A: Deerfield-----	40	Moderate Texture/rock fragments	0.50	Low	
Urban land-----	35	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
225B: Brancroft-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
226B: Berlin-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
228B: Elmridge-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
229B: Agawam-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
229C: Agawam-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
230B: Branford-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
230C: Branford-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
232B: Haven-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
234B: Merrimac-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
235B: Penwood-----	40	Moderate Texture/rock fragments	0.50	Low	
Urban land-----	35	Not rated		Not rated	
236B: Windsor-----	40	Moderate Texture/surface depth/rock fragments	0.50	Low	
Urban land-----	35	Not rated		Not rated	
237A: Manchester-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
237C: Manchester-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
238A: Hinckley-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
238C: Hinckley-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
240B: Ludlow-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
243B: Rainbow-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
245B: Woodbridge-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
245C: Woodbridge-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
248B: Georgia-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
250B: Sutton-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
253B: Wapping-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
255B: Watchaug-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
260B: Charlton-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
260C: Charlton-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
260D: Charlton-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
263B: Cheshire-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
263C: Cheshire-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
266B: Narragansett-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
269B: Yalesville-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
269C: Yalesville-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Low Texture/surface depth/rock fragments	0.10	Low	
Chatfield-----	15	Low Texture/rock fragments	0.10	Low	
273E: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Moderate Texture/slope/surface depth/rock fragments	0.50	Low	
Chatfield-----	15	Low Texture/slope/rock fragments	0.10	Low	
275C: Urban land-----	45	Not rated		Not rated	
Chatfield-----	30	Low Texture/rock fragments	0.10	Low	
275E: Urban land-----	35	Not rated		Not rated	
Chatfield-----	25	Low Texture/slope/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
275E: Rock outcrop-----	15	Not rated		Not rated	
282B: Broadbrook-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
284B: Paxton-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
284C: Paxton-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
284D: Paxton-----	40	Low Texture/rock fragments	0.10	Moderate Available water	0.50
Urban land-----	35	Not rated		Not rated	
287B: Wethersfield-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
287C: Wethersfield-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
287D: Wethersfield-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
290B: Stockbridge-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
290C: Stockbridge-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
290D: Stockbridge-----	40	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
301: Beaches-----	50	Not rated		Not rated	
Udipsamments-----	35	High Texture/rock fragments	1.00	Moderate Available water	0.50
302: Dumps-----	95	Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated	
304: Udorthents-----	90	Low Texture/slope/roc k fragments	0.10	Low	
305: Udorthents-----	65	Low Texture/rock fragments	0.10	Low	
Pits-----	25	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
306: Udorthents-----	50	Low Texture/rock fragments	0.10	Low	
Urban land-----	35	Not rated		Not rated	
307: Urban land-----	80	Not rated		Not rated	
308: Udorthents-----	80	Low Texture/rock fragments	0.10	Low	
309: Udorthents-----	80	Low Texture/rock fragments	0.10	Low	
310: Udorthents, Periodically Flooded-----	85	Low Texture/rock fragments	0.10	Low	
401C: Macomber-----	55	Moderate Texture/surface depth/rock fragments	0.50	Low	
Taconic-----	30	Moderate Texture/surface depth/rock fragments	0.50	Low	
402D: Macomber-----	50	Moderate Texture/surface depth/rock fragments	0.50	Low	
Taconic-----	25	Moderate Texture/surface depth/rock fragments	0.50	Low	
Rock outcrop-----	15	Not rated		Not rated	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
403C: Taconic-----	70	Moderate Texture/surface depth/rock fragments	0.50	Low	
Rock outcrop-----	25	Not rated		Not rated	
403E: Taconic-----	70	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Rock outcrop-----	20	Not rated		Not rated	
403F: Taconic-----	70	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Rock outcrop-----	20	Not rated		Not rated	
405C: Dummerston-----	85	Moderate Texture/surface depth/rock fragments	0.50	Low	
405E: Dummerston-----	85	Moderate Texture/surface depth/rock fragments	0.50	Moderate Available water	0.50
407C: Lanesboro-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
407E: Lanesboro-----	85	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
408C: Fullam-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
409B: Brayton-----	85	Low Texture/surface depth/rock fragments	0.10	High Wetness	1.00
412B: Bice-----	85	Low Texture/rock fragments	0.10	Low	
412C: Bice-----	85	Low Texture/rock fragments	0.10	Low	
412D: Bice-----	85	Low Texture/rock fragments	0.10	Moderate Available water	0.50
413C: Bice-----	45	Low Texture/rock fragments	0.10	Low	
Millsite-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
413E: Bice-----	45	Low Texture/rock fragments	0.10	Low	
Millsite-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
414: Fredon, cold-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
415C: Millsite-----	40	Low Texture/surface depth/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
415C: Westminster-----	40	Low Texture/surface depth/rock fragments	0.10	Low	
Rock outcrop-----	15	Not rated		Not rated	
415E: Millsite-----	40	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Westminster-----	40	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
Rock outcrop-----	15	Not rated		Not rated	
416E: Rock outcrop-----	70	Not rated		Not rated	
Westminster-----	20	Low Texture/surface depth/rock fragments	0.10	Low	
416F: Rock outcrop-----	70	Not rated		Not rated	
Westminster-----	20	Moderate Texture/slope/sur face depth/rock fragments	0.50	Low	
417B: Bice-----	85	Low Texture/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
417C: Bice-----	85	Low Texture/rock fragments	0.10	Low	
417D: Bice-----	85	Low Texture/rock fragments	0.10	Low	
418C: Schroon-----	85	Low Texture/rock fragments	0.10	Low	
420A: Schroon-----	85	Low Texture/rock fragments	0.10	Low	
420B: Schroon-----	85	Low Texture/rock fragments	0.10	Low	
421A: Ninigret, cold-----	85	Low Texture/rock fragments	0.10	Low	
423A: Sudbury, cold-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
424B: Shelburne-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
424C: Shelburne-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
424D: Shelburne-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
425B: Shelburne-----	85	Low Texture/surface depth/rock fragments	0.10	Low	

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
425C: Shelburne-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
426D: Shelburne-----	85	Low Texture/surface depth/rock fragments	0.10	Low	
427B: Ashfield-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
				Soil reaction	0.50
427C: Ashfield-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
				Soil reaction	0.50
428A: Ashfield-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
				Soil reaction	0.50
428B: Ashfield-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
				Soil reaction	0.50
428C: Ashfield-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
				Soil reaction	0.50
429A: Agawam, cold-----	80	Low Texture/rock fragments	0.10	Low	
429B: Agawam, cold-----	80	Low Texture/rock fragments	0.10	Low	
429C: Agawam, cold-----	80	Low Texture/rock fragments	0.10	Low	
433: Moosilauke-----	80	Moderate Texture/rock fragments	0.50	High Wetness	1.00

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
434A: Merrimac, cold-----	80	Low Texture/rock fragments	0.10	Low	
434B: Merrimac, cold-----	80	Low Texture/rock fragments	0.10	Low	
434C: Merrimac, cold-----	80	Low Texture/rock fragments	0.10	Low	
435: Scarboro-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
436: Halsey-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
437: Wonsqueak-----	85	Low		High Wetness	1.00
438: Bucksport-----	85	Low		High Wetness	1.00
440A: Boscawen-----	80	Moderate Texture/surface depth/rock fragments	0.50	Moderate Available water	0.50
440C: Boscawen-----	80	Moderate Texture/surface depth/rock fragments	0.50	Moderate Available water	0.50
440E: Boscawen-----	85	Moderate Texture/slope/sur face depth/rock fragments	0.50	High Available water	1.00
442: Brayton-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00
443: Brayton-----	50	Low Texture/rock fragments	0.10	High Wetness	1.00

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
443: Loonmeadow-----	35	Low Texture/rock fragments	0.10	High Wetness	1.00
448B: Hogansburg-----	85	Low Texture/rock fragments	0.10	Low	
449B: Hogansburg-----	85	Low Texture/rock fragments	0.10	Low	
449C: Hogansburg-----	85	Low Texture/rock fragments	0.10	Low	
450B: Pyrities-----	80	Low Texture/rock fragments	0.10	Low	
450C: Pyrities-----	80	Low Texture/rock fragments	0.10	Low	
450D: Pyrities-----	80	Low Texture/rock fragments	0.10	Low	
451B: Pyrities-----	80	Low Texture/rock fragments	0.10	Low	
451C: Pyrities-----	80	Low Texture/rock fragments	0.10	Low	
451D: Pyrities-----	80	Low Texture/rock fragments	0.10	Low	
457: Mudgepond-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
458: Mudgepond-----	55	Low Texture/rock fragments	0.10	High Wetness	1.00

Table 11.—Damage by Fire and Seedling Mortality on Forestland—Continued

Map symbol and soil name	Pct. of map unit	Potential for damage to soil by fire		Potential for seedling mortality	
		Rating class and limiting features	Value	Rating class and limiting features	Value
458: Alden-----	35	Low Texture/surface depth/rock fragments	0.10	High Wetness	1.00
501: Ondawa-----	85	Low Texture/rock fragments	0.10	Low	
503: Rumney-----	80	Low Texture/rock fragments	0.10	High Wetness	1.00
508: Medomak-----	85	Low Texture/rock fragments	0.10	High Wetness	1.00

Table 12.—Conservation and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
2: Ridgebury-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
3: Ridgebury-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
Leicester-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
Whitman-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
4: Leicester-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
5: Wilbraham-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
6: Wilbraham-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
Menlo-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
7: Mudgepond-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
8: Mudgepond-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
Alden-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
9: Scitico-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
Shaker-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
Maybid-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
10: Raynham-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
12: Raypol-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
13: Walpole-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
14: Fredon-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch

Table 12.-Conservation and Environmental Plantings-Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
15: Scarboro-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple
16: Halsey-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
17: Timakwa-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple
Natchaug-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple
18: Catden-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple
Freetown-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple
20A: Ellington-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
21A: Ninigret-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Tisbury-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
22A: Hero-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine
22B: Hero-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine
23A: Sudbury-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
24A: Deerfield-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
25A: Brancroft-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
25B: Brancroft-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
25C: Brancroft-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
26A: Berlin-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
26B: Berlin-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
27A: Belgrade-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
28A: Elmridge-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
28B: Elmridge-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
29A: Agawam-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
29B: Agawam-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
29C: Agawam-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
30A: Branford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
30B: Branford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
30C: Branford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
31A: Copake-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine
31B: Copake-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
31C: Copake-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine
32A: Haven-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Enfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
32B: Haven-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Enfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
32C: Haven-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Enfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
33A: Hartford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
33B: Hartford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
34A: Merrimac-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
34B: Merrimac-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
34C: Merrimac-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
35A: Penwood-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
35B: Penwood-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
36A: Windsor-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
36B: Windsor-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
36C: Windsor-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
37A: Manchester-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
37C: Manchester-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
37E: Manchester-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
38A: Hinckley-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
38C: Hinckley-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
38E: Hinckley-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
39A: Groton-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine
39C: Groton-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine
39E: Groton-----	downy arrowwood; yew	shagbark hickory	mockernut hickory; sugar maple	chinkapin oak; eastern redcedar	eastern white pine

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
40A: Ludlow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
40B: Ludlow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
41B: Ludlow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
42C: Ludlow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
43A: Rainbow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
43B: Rainbow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
44B: Rainbow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
45A: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
45B: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
45C: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
46B: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
46C: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
47C: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
48B: Georgia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
48B: Amenia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
48C: Georgia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
Amenia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
49B: Georgia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
Amenia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
49C: Georgia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
Amenia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
50A: Sutton-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
50B: Sutton-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
51B: Sutton-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
52C: Sutton-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
53A: Wapping-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
53B: Wapping-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
54B: Wapping-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
55A: Watchaug-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
55B: Watchaug-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
56B: Watchaug-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
57B: Gloucester-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
57C: Gloucester-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
57D: Gloucester-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
58B: Gloucester-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
58C: Gloucester-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
59C: Gloucester-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
59D: Gloucester-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
60B: Canton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
60B: Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
60C: Canton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
60D: Canton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
61B: Canton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
61C: Canton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
62C: Canton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
62D: Canton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
63B: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
63C: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
63D: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
64B: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
64C: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
65C: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
65D: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
66B: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
66C: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
67B: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
67C: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
68C: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
68D: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
69B: Yalesville-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
69C: Yalesville-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
70C: Branford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Holyoke-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
71C: Brookfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Brimfield-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
71E: Brookfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Brimfield-----	beach plum; kinnikinnick; sweet fern	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
71E: Rock outcrop-----	---	---	---	---	---
73C: Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
73E: Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
74C: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
Hollis-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
75C: Hollis-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
75C: Rock outcrop-----	---	---	---	---	---
75E: Hollis-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Rock outcrop-----	---	---	---	---	---
76E: Rock outcrop-----	---	---	---	---	---
Hollis-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
76F: Rock outcrop-----	---	---	---	---	---
Hollis-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
77C: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
Holyoke-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
77D: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
Holyoke-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
78C: Holyoke-----	beach plum; kinnikinnick; sweet fern	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
78C: Rock outcrop-----	---	---	---	---	---
78E: Holyoke-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
79E: Rock outcrop-----	---	---	---	---	---
Holyoke-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
80B: Bernardston-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
80C: Bernardston-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
81C: Bernardston-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
81D: Bernardston-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
82B: Broadbrook-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
82C: Broadbrook-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
82D: Broadbrook-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
83B: Broadbrook-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
83C: Broadbrook-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
84B: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Montauk-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
84C: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Montauk-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
84D: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Montauk-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
85B: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Montauk-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
85C: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Montauk-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
86C: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
86C: Montauk-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
86D: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Montauk-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
87B: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
87C: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
87D: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
88B: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
88C: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
89C: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
89D: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
90B: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
90C: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
90D: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
91B: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
91C: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
91D: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
92B: Nellis-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
92C: Nellis-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
92D: Nellis-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
93C: Nellis-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
94C: Farmington-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Nellis-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
94E: Farmington-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Nellis-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
95C: Farmington-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
95E: Farmington-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
96: Ipswich-----	---	---	---	---	---
97: Pawcatuck-----	---	---	---	---	---
98: Westbrook-----	---	---	---	---	---
99: Westbrook, low salt----	---	---	---	---	---
100: Suncook-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
101: Occum-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
102: Pootatuck-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
103: Rippowam-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
104: Bash-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
105: Hadley-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
106: Winooski-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
107: Limerick-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
Lim-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
108: Saco-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
109: Fluvaquents, Frequently Flooded-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
Udifluents, Frequently Flooded-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
221A: Ninigret-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
221A: Urban land-----	---	---	---	---	---
224A: Deerfield-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
Urban land-----	---	---	---	---	---
225B: Brancroft-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
Urban land-----	---	---	---	---	---
226B: Berlin-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
Urban land-----	---	---	---	---	---
228B: Elmridge-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
Urban land-----	---	---	---	---	---
229B: Agawam-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
229C: Agawam-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
230B: Branford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
230C: Branford-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
232B: Haven-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
234B: Merrimac-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
235B: Penwood-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
235B: Urban land-----	---	---	---	---	---
236B: Windsor-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
Urban land-----	---	---	---	---	---
237A: Manchester-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
Urban land-----	---	---	---	---	---
237C: Manchester-----	huckleberry; sweet fern	beach plum; northern bayberry	pitch pine	---	---
Urban land-----	---	---	---	---	---
238A: Hinckley-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
Urban land-----	---	---	---	---	---
238C: Hinckley-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
Urban land-----	---	---	---	---	---
240B: Ludlow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
Urban land-----	---	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
243B: Rainbow-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
Urban land-----	---	---	---	---	---
245B: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
Urban land-----	---	---	---	---	---
245C: Woodbridge-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
Urban land-----	---	---	---	---	---
248B: Georgia-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
Urban land-----	---	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
250B: Sutton-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
Urban land-----	---	---	---	---	---
253B: Wapping-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
Urban land-----	---	---	---	---	---
255B: Watchaug-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
Urban land-----	---	---	---	---	---
260B: Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
260C: Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
260D: Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
260D: Urban land-----	---	---	---	---	---
263B: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
Urban land-----	---	---	---	---	---
263C: Cheshire-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
Urban land-----	---	---	---	---	---
266B: Narragansett-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
Urban land-----	---	---	---	---	---
269B: Yalesville-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
269C: Yalesville-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
273C: Urban land-----	---	---	---	---	---
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
273C: Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
273E: Urban land-----	---	---	---	---	---
Charlton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
275C: Urban land-----	---	---	---	---	---
Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
275E: Urban land-----	---	---	---	---	---
Chatfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Rock outcrop-----	---	---	---	---	---
282B: Broadbrook-----	silky dogwood; swamp rose	hobblebush	American holly; American hornbeam	swamp white oak; yellow birch	American elm; tuliptree
Urban land-----	---	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
284B: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
284C: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
284D: Paxton-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
287B: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
287C: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
287D: Wethersfield-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Urban land-----	---	---	---	---	---
290B: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
Urban land-----	---	---	---	---	---
290C: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
Urban land-----	---	---	---	---	---
290D: Stockbridge-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
Urban land-----	---	---	---	---	---
301: Beaches-----	---	---	---	---	---
Udipsamments-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
302: Dumps-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
303: Pits, quarries-----	---	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
304: Udorthents-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
305: Udorthents-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Pits-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
306: Udorthents-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Urban land-----	---	---	---	---	---
307: Urban land-----	---	---	---	---	---
308: Udorthents-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
309: Udorthents-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
310: Udorthents, Periodically Flooded-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
401C: Macomber-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
401C: Taconic-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
402D: Macomber-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Taconic-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
403C: Taconic-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
403E: Taconic-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
403F: Taconic-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
405C: Dummerston-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
405E: Dummerston-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
407C: Lanesboro-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
407E: Lanesboro-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
408C: Fullam-----	elderberry	Canadian serviceberry	butternut; common pricklyash	bur oak; common hackberry	American basswood; American elm; American sycamore; black walnut; red mulberry; slippery elm
409B: Brayton-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
412B: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatelaef dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
412C: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatelaef dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
412D: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatelaef dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
413C: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatelaef dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
Millsite-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
413E: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatetea leaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
Millsite-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
414: Fredon, cold-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
415C: Millsite-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Westminster-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
415E: Millsite-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
Westminster-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
Rock outcrop-----	---	---	---	---	---
416E: Rock outcrop-----	---	---	---	---	---

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
416E: Westminster-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
416F: Rock outcrop-----	---	---	---	---	---
Westminster-----	beach plum; kinnikinnick; sweet fern	---	---	---	---
417B: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
417C: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
417D: Bice-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternatleaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
418C: Schroon-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
420A: Schroon-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
420B: Schroon-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
421A: Ninigret, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
423A: Sudbury, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
424B: Shelburne-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
424C: Shelburne-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
424D: Shelburne-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
425B: Shelburne-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
425C: Shelburne-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
426D: Shelburne-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
427B: Ashfield-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
427C: Ashfield-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
428A: Ashfield-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
428B: Ashfield-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
428C: Ashfield-----	black raspberry; mapleleaf viburnum; partridgeberry	American witchhazel; beaked hazelnut; sweet birch	American plum; blackhaw; eastern hophornbeam; redbud	alternateteaf dogwood; American beech; bitternut hickory; black oak; flowering dogwood; mountain maple; striped maple; white oak	black cherry; eastern white pine; paper birch; sweetgum
429A: Agawam, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
429B: Agawam, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
429C: Agawam, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
433: Moosilauke-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
434A: Merrimac, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
434B: Merrimac, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
434C: Merrimac, cold-----	lowbush blueberry; mountain laurel	common chokecherry; smooth sumac	deerberry; sassafras	pignut hickory; pin cherry; staghorn sumac	black cherry; eastern white pine; gray birch; northern red oak; quaking aspen
435: Scarboro-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
436: Halsey-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
437: Wonsqueak-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple
438: Bucksport-----	cranberry; holly; leatherleaf	highbush blueberry; swamp birch	Atlantic white cedar; black spruce; coastal sweetpepperbush; tamarack	red spruce	pin oak; red maple
440A: Boscawen-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
440C: Boscawen-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
440E: Boscawen-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
442: Brayton-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple
443: Brayton-----	black chokeberry; common winterberry; inkberry; swamp azalea	common buttonbush; hazel alder; highbush blueberry; southern arrowwood; spicebush	coastal sweetpepperbush; speckled alder	Atlantic white cedar; blackgum	black willow; pin oak; red maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
443: Loonmeadow-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
448B: Hogansburg-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
449B: Hogansburg-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
449C: Hogansburg-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
450B: Pyrities-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
450C: Pyrities-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
450D: Pyrities-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
451B: Pyrities-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple

Table 12.—Conservation and Environmental Plantings—Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
451C: Pyrities-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
451D: Pyrities-----	gray dogwood	redosier dogwood	common pricklyash; hickory; sugar maple	Arnold hawthorn; nannyberry; swamp white oak	balsam poplar; eastern cottonwood; green ash; silver maple
457: Mudgepond-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
458: Mudgepond-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
Alden-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
501: Ondawa-----	huckleberry	roundleaf dogwood	chestnut oak; eastern redcedar; scrub oak	scarlet oak	eastern white pine; gray birch
503: Rumney-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch
508: Medomak-----	elderberry	silky willow	black ash; pussy willow; swamp birch	---	boxelder; green ash; river birch

Table 13.—Recreation (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Very limited Depth to saturated zone Depth to pan	1.00 0.99	Very limited Depth to saturated zone Depth to pan	1.00 0.99	Very limited Depth to saturated zone Depth to pan Gravel content	1.00 0.99 0.18
3: Ridgebury-----	40	Very limited Depth to saturated zone Large stones content Depth to pan	1.00 1.00 0.99	Very limited Depth to saturated zone Large stones content Depth to pan	1.00 1.00 0.99	Very limited Depth to saturated zone Large stones content Depth to pan Gravel content	1.00 1.00 0.99 0.18
Leicester-----	35	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00
Whitman-----	15	Very limited Depth to saturated zone Depth to pan Large stones content Ponding	1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Large stones content Depth to pan Ponding	1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Depth to pan Large stones content Ponding	1.00 1.00 1.00 1.00
4: Leicester-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Gravel content Large stones content	1.00 0.16 0.01
5: Wilbraham-----	80	Very limited Depth to saturated zone Depth to pan	1.00 1.00	Very limited Depth to saturated zone Depth to pan	1.00 1.00	Very limited Depth to saturated zone Depth to pan Gravel content	1.00 1.00 0.18
6: Wilbraham-----	60	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
6: Wilbraham-----		Depth to pan	1.00	Depth to pan	1.00	Depth to pan Gravel content	1.00 0.18
Menlo-----	25	Very limited Depth to saturated zone Large stones content Ponding Depth to pan	1.00 1.00 1.00 1.00 0.71	Very limited Depth to saturated zone Large stones content Ponding Depth to pan	1.00 1.00 1.00 1.00 0.71	Very limited Depth to saturated zone Large stones content Ponding Depth to pan	1.00 1.00 1.00 1.00 0.71
7: Mudgepond-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Gravel content	1.00 0.22
8: Mudgepond-----	45	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content Gravel content	1.00 1.00 0.22
Alden-----	35	Very limited Depth to saturated zone Large stones content Ponding Slow water movement	1.00 1.00 1.00 0.15	Very limited Depth to saturated zone Large stones content Ponding Slow water movement	1.00 1.00 1.00 0.15	Very limited Depth to saturated zone Large stones content Ponding Slow water movement Large stones content	1.00 1.00 1.00 0.15 0.01
9: Scitico-----	40	Very limited Depth to saturated zone Slow water movement	1.00 1.00	Very limited Depth to saturated zone Slow water movement	1.00 1.00	Very limited Depth to saturated zone Slow water movement	1.00 1.00
Shaker-----	30	Very limited Depth to saturated zone Slow water movement	1.00 0.99	Very limited Depth to saturated zone Slow water movement	1.00 0.99	Very limited Depth to saturated zone Slow water movement	1.00 0.99
Maybid-----	15	Very limited Depth to saturated zone Ponding Slow water movement	1.00 1.00 0.99	Very limited Depth to saturated zone Ponding Slow water movement	1.00 1.00 0.99	Very limited Depth to saturated zone Ponding Slow water movement	1.00 1.00 0.99
10: Raynham-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.96	Very limited Depth to saturated zone Slow water movement	1.00 0.96	Very limited Depth to saturated zone Slow water movement	1.00 0.96

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
12: Raypol-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
13: Walpole-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
14: Fredon-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Gravel content	1.00 0.04
15: Scarboro-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
16: Halsey-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
17: Timakwa-----	45	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Natchaug-----	40	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
18: Catden-----	40	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Freetown-----	40	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
20A: Ellington-----	80	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone Slope	0.39 0.12

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
21A: Ninigret-----	60	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone Gravel content Slope	0.39 0.22 0.12
Tisbury-----	25	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone	0.39
22A: Hero-----	85	Somewhat limited Depth to saturated zone Gravel content	0.39 0.18	Somewhat limited Depth to saturated zone Gravel content	0.19 0.18	Very limited Gravel content Depth to saturated zone Large stones content	1.00 0.39 0.01
22B: Hero-----	85	Somewhat limited Depth to saturated zone Gravel content	0.39 0.18	Somewhat limited Depth to saturated zone Gravel content	0.19 0.18	Very limited Gravel content Slope Depth to saturated zone Large stones content	1.00 1.00 0.39 0.01
23A: Sudbury-----	80	Somewhat limited Depth to saturated zone	0.07	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Slope Depth to saturated zone	0.12 0.07
24A: Deerfield-----	80	Somewhat limited Too sandy Depth to saturated zone	0.55 0.07	Somewhat limited Too sandy Depth to saturated zone	0.55 0.03	Somewhat limited Too sandy Depth to saturated zone	0.55 0.07
25A: Brancroft-----	80	Very limited Slow water movement Depth to saturated zone	1.00 0.81	Very limited Slow water movement Depth to saturated zone	1.00 0.48	Very limited Slow water movement Depth to saturated zone	1.00 0.81
25B: Brancroft-----	80	Very limited Slow water movement Depth to saturated zone	1.00 0.81	Very limited Slow water movement Depth to saturated zone	1.00 0.48	Very limited Slow water movement Slope Depth to saturated zone	1.00 1.00 0.81

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
25C: Brancroft-----	80	Very limited Slow water movement Depth to saturated zone Slope	1.00 0.81 0.63	Very limited Slow water movement Slope Depth to saturated zone	1.00 0.63 0.48	Very limited Slope Slow water movement Depth to saturated zone	1.00 1.00 0.81
26A: Berlin-----	80	Very limited Slow water movement Depth to saturated zone	1.00 0.81	Very limited Slow water movement Depth to saturated zone	1.00 0.48	Very limited Slow water movement Depth to saturated zone	1.00 0.81
26B: Berlin-----	80	Very limited Slow water movement Depth to saturated zone	1.00 0.81	Very limited Slow water movement Depth to saturated zone	1.00 0.48	Very limited Slow water movement Slope Depth to saturated zone	1.00 1.00 0.81
27A: Belgrade-----	80	Not limited		Not limited		Somewhat limited Slope	0.12
28A: Elmridge-----	80	Very limited Slow water movement Depth to saturated zone	1.00 0.39	Very limited Slow water movement Depth to saturated zone	1.00 0.19	Very limited Slow water movement Depth to saturated zone	1.00 0.39
28B: Elmridge-----	80	Very limited Slow water movement Depth to saturated zone	1.00 0.39	Very limited Slow water movement Depth to saturated zone	1.00 0.19	Very limited Slow water movement Slope Depth to saturated zone	1.00 1.00 0.39
29A: Agawam-----	80	Not limited		Not limited		Not limited	
29B: Agawam-----	80	Not limited		Not limited		Very limited Slope	1.00
29C: Agawam-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
30A: Branford-----	80	Not limited		Not limited		Somewhat limited Gravel content	0.06

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
30B: Branford-----	80	Not limited		Not limited		Somewhat limited Slope Gravel content	0.72 0.06
30C: Branford-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.06
31A: Copake-----	85	Not limited		Not limited		Somewhat limited Gravel content	0.18
31B: Copake-----	85	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.18
31C: Copake-----	85	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.18
32A: Haven-----	60	Not limited		Not limited		Somewhat limited Gravel content	0.06
Enfield-----	25	Not limited		Not limited		Not limited	
32B: Haven-----	60	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.06
Enfield-----	25	Not limited		Not limited		Very limited Slope	1.00
32C: Haven-----	60	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.06
Enfield-----	25	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
33A: Hartford-----	80	Not limited		Not limited		Somewhat limited Gravel content	0.50
33B: Hartford-----	80	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.50
34A: Merrimac-----	80	Not limited		Not limited		Somewhat limited Gravel content	0.78

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
34B: Merrimac-----	80	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.78
34C: Merrimac-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.78
35A: Penwood-----	80	Somewhat limited Too sandy	0.32	Somewhat limited Too sandy	0.32	Somewhat limited Too sandy	0.32
35B: Penwood-----	80	Somewhat limited Too sandy	0.32	Somewhat limited Too sandy	0.32	Very limited Slope Too sandy	1.00 0.32
36A: Windsor-----	80	Not limited		Not limited		Not limited	
36B: Windsor-----	80	Not limited		Not limited		Very limited Slope	1.00
36C: Windsor-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
37A: Manchester-----	80	Somewhat limited Gravel content	0.68	Somewhat limited Gravel content	0.68	Very limited Gravel content	1.00
37C: Manchester-----	80	Somewhat limited Gravel content Slope	0.68 0.04	Somewhat limited Gravel content Slope	0.68 0.04	Very limited Gravel content Slope	1.00 1.00
37E: Manchester-----	80	Very limited Slope Gravel content	1.00 0.68	Very limited Slope Gravel content	1.00 0.68	Very limited Gravel content Slope	1.00 1.00
38A: Hinckley-----	80	Somewhat limited Gravel content	0.05	Somewhat limited Gravel content	0.05	Very limited Gravel content Large stones content	1.00 0.11
38C: Hinckley-----	80	Somewhat limited Gravel content Slope	0.05 0.04	Somewhat limited Gravel content Slope	0.05 0.04	Very limited Gravel content Slope Large stones content	1.00 1.00 0.11

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
38E: Hinckley-----	80	Very limited Slope Gravel content	1.00 0.05	Very limited Slope Gravel content	1.00 0.05	Very limited Slope Gravel content Large stones content	1.00 1.00 0.11
39A: Groton-----	85	Somewhat limited Gravel content	0.59	Somewhat limited Gravel content	0.59	Very limited Gravel content	1.00
39C: Groton-----	85	Somewhat limited Gravel content Slope	0.59 0.04	Somewhat limited Gravel content Slope	0.59 0.04	Very limited Gravel content Slope	1.00 1.00
39E: Groton-----	85	Very limited Slope Gravel content	1.00 0.59	Very limited Slope Gravel content	1.00 0.59	Very limited Gravel content Slope	1.00 1.00
40A: Ludlow-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Somewhat limited Depth to pan Gravel content Depth to saturated zone	0.80 0.50 0.39
40B: Ludlow-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Very limited Slope Depth to pan Gravel content Depth to saturated zone	1.00 0.80 0.50 0.39
41B: Ludlow-----	80	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.79 0.53 0.39	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.79 0.53 0.19	Somewhat limited Slope Depth to pan Large stones content Gravel content Depth to saturated zone	0.88 0.80 0.53 0.50 0.39
42C: Ludlow-----	80	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.79 0.39 0.04	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.79 0.19 0.04	Very limited Large stones content Slope Depth to pan Gravel content Depth to saturated zone	1.00 1.00 0.80 0.50 0.39

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
43A: Rainbow-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Somewhat limited Depth to pan Depth to saturated zone	0.80 0.39
43B: Rainbow-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.80 0.39
44B: Rainbow-----	80	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.79 0.53 0.39	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.79 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone	1.00 0.80 0.53 0.39
45A: Woodbridge-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.46 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.46 0.19	Somewhat limited Depth to pan Depth to saturated zone Gravel content	0.46 0.39 0.17
45B: Woodbridge-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.46 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.46 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.46 0.39 0.17
45C: Woodbridge-----	80	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.46 0.39	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.46 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.46 0.39 0.17
46B: Woodbridge-----	80	Somewhat limited Large stones content Depth to pan Depth to saturated zone	0.53 0.46 0.39	Somewhat limited Large stones content Depth to pan Depth to saturated zone	0.53 0.46 0.19	Somewhat limited Slope Large stones content Depth to pan Depth to saturated zone Gravel content	0.88 0.53 0.46 0.39 0.17

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
46C: Woodbridge-----	80	Somewhat limited Slope Large stones content Depth to pan Depth to saturated zone	0.63 0.53 0.46 0.39	Somewhat limited Slope Large stones content Depth to pan Depth to saturated zone	0.63 0.53 0.46 0.19	Very limited Slope Large stones content Depth to pan Depth to saturated zone Gravel content	1.00 0.53 0.46 0.39 0.17
47C: Woodbridge-----	80	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.46 0.39 0.04	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.46 0.19 0.04	Very limited Large stones content Slope Depth to pan Depth to saturated zone Gravel content	1.00 1.00 0.46 0.39 0.16
48B: Georgia-----	50	Somewhat limited Slow water movement Depth to saturated zone	0.15 0.07	Somewhat limited Slow water movement Depth to saturated zone	0.15 0.03	Somewhat limited Slope Gravel content Slow water movement Depth to saturated zone	0.88 0.20 0.15 0.07
Amenia-----	35	Somewhat limited Slow water movement Depth to saturated zone	0.15 0.07	Somewhat limited Slow water movement Depth to saturated zone	0.15 0.03	Somewhat limited Slope Gravel content Slow water movement Depth to saturated zone	0.88 0.20 0.15 0.07
48C: Georgia-----	50	Somewhat limited Slope Slow water movement Depth to saturated zone	0.63 0.15 0.07	Somewhat limited Slope Slow water movement Depth to saturated zone	0.63 0.15 0.03	Very limited Slope Gravel content Slow water movement Depth to saturated zone	1.00 0.20 0.15 0.07
Amenia-----	35	Somewhat limited Slope Slow water movement Depth to saturated zone	0.63 0.15 0.07	Somewhat limited Slope Slow water movement Depth to saturated zone	0.63 0.15 0.03	Very limited Slope Gravel content Slow water movement Depth to saturated zone	1.00 0.20 0.15 0.07

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
49B: Georgia-----	50	Very limited Large stones content Slow water movement Depth to saturated zone	1.00 0.15 0.07	Very limited Large stones content Slow water movement Depth to saturated zone	1.00 0.15 0.03	Very limited Slope Large stones content Gravel content Slow water movement Depth to saturated zone	1.00 1.00 0.20 0.15 0.07
Amenia-----	35	Very limited Large stones content Slow water movement Depth to saturated zone	1.00 0.15 0.07	Very limited Large stones content Slow water movement Depth to saturated zone	1.00 0.15 0.03	Very limited Slope Large stones content Gravel content Slow water movement Depth to saturated zone	1.00 1.00 0.20 0.15 0.07
49C: Georgia-----	50	Very limited Large stones content Slope Slow water movement Depth to saturated zone	1.00 0.63 0.15 0.07	Very limited Large stones content Slope Slow water movement Depth to saturated zone	1.00 0.63 0.15 0.03	Very limited Slope Large stones content Gravel content Slow water movement Depth to saturated zone	1.00 1.00 0.20 0.15 0.07
Amenia-----	35	Very limited Large stones content Slope Slow water movement Depth to saturated zone	1.00 0.63 0.15 0.07	Very limited Large stones content Slope Slow water movement Depth to saturated zone	1.00 0.63 0.15 0.03	Very limited Slope Large stones content Gravel content Slow water movement Depth to saturated zone	1.00 1.00 0.20 0.15 0.07
50A: Sutton-----	80	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone Gravel content	0.39 0.18

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
50B: Sutton-----	80	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Very limited Slope	1.00
						Depth to saturated zone	0.39
						Gravel content	0.18
51B: Sutton-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.88
		Depth to saturated zone	0.39	Depth to saturated zone	0.19	Large stones content	0.53
						Depth to saturated zone	0.39
						Gravel content	0.18
52C: Sutton-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Very limited Large stones content	1.00
		Depth to saturated zone	0.39	Depth to saturated zone	0.19	Slope	1.00
		Slope	0.04	Slope	0.04	Depth to saturated zone	0.39
						Gravel content	0.18
53A: Wapping-----	80	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone	0.39
						Gravel content	0.06
53B: Wapping-----	80	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Very limited Slope	1.00
						Depth to saturated zone	0.39
						Gravel content	0.06
54B: Wapping-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.88
		Depth to saturated zone	0.39	Depth to saturated zone	0.19	Large stones content	0.53
						Depth to saturated zone	0.39
						Gravel content	0.06
55A: Watchaug-----	80	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone	0.39
						Gravel content	0.16
						Large stones content	0.01

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
55B: Watchaug-----	80	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Very limited Slope	1.00
						Depth to saturated zone	0.39
						Gravel content	0.16
						Large stones content	0.01
56B: Watchaug-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.88
		Depth to saturated zone	0.39	Depth to saturated zone	0.19	Large stones content	0.53
						Depth to saturated zone	0.39
						Gravel content	0.16
						Large stones content	0.01
57B: Gloucester-----	80	Not limited		Not limited		Very limited Slope	1.00
						Gravel content	0.20
57C: Gloucester-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
						Gravel content	0.20
57D: Gloucester-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
						Gravel content	0.20
58B: Gloucester-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
						Large stones content	0.53
						Gravel content	0.20
58C: Gloucester-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
		Large stones content	0.53	Large stones content	0.53	Large stones content	0.53
						Gravel content	0.20
59C: Gloucester-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Very limited Large stones content	1.00
		Slope	0.04	Slope	0.04	Slope	1.00
						Gravel content	0.20

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
59D: Gloucester-----	80	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content Gravel content	1.00 1.00 0.20
60B: Canton-----	45	Somewhat limited Gravel content	0.01	Somewhat limited Gravel content	0.01	Very limited Slope	1.00
Charlton-----	35	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.43
60C: Canton-----	45	Somewhat limited Slope Gravel content	0.63 0.01	Somewhat limited Slope Gravel content	0.63 0.01	Very limited Slope	1.00
Charlton-----	35	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.43
60D: Canton-----	45	Very limited Slope Gravel content	1.00 0.01	Very limited Slope Gravel content	1.00 0.01	Very limited Slope	1.00
Charlton-----	35	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Gravel content	1.00 0.43
61B: Canton-----	45	Somewhat limited Large stones content Gravel content	0.53 0.01	Somewhat limited Large stones content Gravel content	0.53 0.01	Very limited Slope Large stones content	1.00 0.53
Charlton-----	35	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Slope Large stones content Gravel content	1.00 0.53 0.43
61C: Canton-----	45	Somewhat limited Slope Large stones content Gravel content	0.63 0.53 0.01	Somewhat limited Slope Large stones content Gravel content	0.63 0.53 0.01	Very limited Slope Large stones content	1.00 0.53
Charlton-----	35	Somewhat limited Slope Large stones content	0.63 0.53	Somewhat limited Slope Large stones content	0.63 0.53	Very limited Slope Large stones content Gravel content	1.00 0.53 0.43

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
62C: Canton-----	45	Very limited Large stones content Slope Gravel content	1.00 0.04 0.01	Very limited Large stones content Slope Gravel content	1.00 0.04 0.01	Very limited Large stones content Slope	1.00 1.00
Charlton-----	35	Very limited Large stones content Slope	1.00 0.04	Very limited Large stones content Slope	1.00 0.04	Very limited Large stones content Slope Gravel content	1.00 1.00 0.43
62D: Canton-----	45	Very limited Slope Large stones content Gravel content	1.00 1.00 0.01	Very limited Slope Large stones content Gravel content	1.00 1.00 0.01	Very limited Slope Large stones content	1.00 1.00
Charlton-----	35	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content Gravel content	1.00 1.00 0.43
63B: Cheshire-----	80	Not limited		Not limited		Very limited Slope Gravel content Large stones content	1.00 0.15 0.01
63C: Cheshire-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content Large stones content	1.00 0.15 0.01
63D: Cheshire-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Gravel content Large stones content	1.00 0.15 0.01
64B: Cheshire-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Slope Large stones content Gravel content Large stones content	1.00 0.53 0.15 0.01

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
64C: Cheshire-----	80	Somewhat limited Slope Large stones content	0.63 0.53	Somewhat limited Slope Large stones content	0.63 0.53	Very limited Slope Large stones content Gravel content Large stones content	1.00 0.53 0.15 0.01
65C: Cheshire-----	80	Very limited Large stones content Slope	1.00 0.04	Very limited Large stones content Slope	1.00 0.04	Very limited Large stones content Slope Gravel content Large stones content	1.00 1.00 0.15 0.01
65D: Cheshire-----	80	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content Gravel content Large stones content	1.00 1.00 0.15 0.01
66B: Narragansett-----	80	Not limited		Not limited		Somewhat limited Slope Large stones content	0.88 0.01
66C: Narragansett-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Large stones content	1.00 0.01
67B: Narragansett-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Slope Large stones content Large stones content	1.00 0.53 0.01
67C: Narragansett-----	80	Somewhat limited Slope Large stones content	0.63 0.53	Somewhat limited Slope Large stones content	0.63 0.53	Very limited Slope Large stones content Large stones content	1.00 0.53 0.01

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
68C: Narragansett-----	80	Very limited Large stones content Slope	1.00 0.04	Very limited Large stones content Slope	1.00 0.04	Very limited Large stones content Slope Large stones content	1.00 1.00 0.01
68D: Narragansett-----	80	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content Large stones content	1.00 1.00 0.01
69B: Yalesville-----	75	Not limited		Not limited		Very limited Slope Depth to bedrock	1.00 0.05
69C: Yalesville-----	75	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Depth to bedrock	1.00 0.05
70C: Branford-----	50	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope Gravel content	1.00 0.06
Holyoke-----	30	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53
71C: Brookfield-----	45	Somewhat limited Large stones content Slope	0.53 0.04	Somewhat limited Large stones content Slope	0.53 0.04	Very limited Slope Large stones content	1.00 0.53
Brimfield-----	30	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
71E: Brookfield-----	45	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content	1.00 0.53

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
71E: Brimfield-----	30	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
73C: Charlton-----	45	Somewhat limited Large stones content Slope	0.53 0.04	Somewhat limited Large stones content Slope	0.53 0.04	Very limited Slope Large stones content Gravel content	1.00 0.53 0.43
Chatfield-----	30	Somewhat limited Large stones content Gravel content Slope	0.53 0.25 0.04	Somewhat limited Large stones content Gravel content Slope	0.53 0.25 0.04	Very limited Slope Depth to bedrock Large stones content	1.00 0.54 0.53
73E: Charlton-----	45	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content Gravel content	1.00 0.53 0.43
Chatfield-----	30	Very limited Slope Large stones content Gravel content	1.00 0.53 0.25	Very limited Slope Large stones content Gravel content	1.00 0.53 0.25	Very limited Slope Depth to bedrock Large stones content	1.00 0.54 0.53
74C: Narragansett-----	55	Somewhat limited Large stones content Slope	0.53 0.04	Somewhat limited Large stones content Slope	0.53 0.04	Very limited Slope Large stones content Large stones content	1.00 0.53 0.01
Hollis-----	20	Very limited Depth to bedrock Large stones content Gravel content Slope	1.00 0.53 0.22 0.04	Very limited Depth to bedrock Large stones content Gravel content Slope	1.00 0.53 0.22 0.04	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
75C: Hollis-----	35	Very limited Depth to bedrock	1.00	Very limited Large stones content	1.00	Very limited Depth to bedrock	1.00
		Large stones content	1.00	Depth to bedrock	1.00	Large stones content	1.00
		Gravel content	0.22	Gravel content	0.22	Slope	1.00
		Slope	0.04	Slope	0.04		
Chatfield-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
		Gravel content	0.25	Gravel content	0.25	Depth to bedrock	0.54
		Slope	0.04	Slope	0.04	Large stones content	0.53
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
75E: Hollis-----	35	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Depth to bedrock	1.00	Large stones content	1.00	Depth to bedrock	1.00
		Large stones content	1.00	Depth to bedrock	1.00	Large stones content	1.00
		Gravel content	0.22	Gravel content	0.22		
Chatfield-----	30	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Large stones content	0.53	Large stones content	0.53	Depth to bedrock	0.54
		Gravel content	0.25	Gravel content	0.25	Large stones content	0.53
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
76E: Rock Outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Depth to bedrock	1.00	Very limited Large stones content	1.00	Very limited Depth to bedrock	1.00
		Large stones content	1.00	Depth to bedrock	1.00	Large stones content	1.00
		Slope	1.00	Slope	1.00	Slope	1.00
		Gravel content	0.22	Gravel content	0.22		
76F: Rock Outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Depth to bedrock	1.00	Large stones content	1.00	Depth to bedrock	1.00
		Large stones content	1.00	Depth to bedrock	1.00	Large stones content	1.00
		Gravel content	0.22	Gravel content	0.22		

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
77C: Cheshire-----	45	Somewhat limited Large stones content Slope	0.53 0.04	Somewhat limited Large stones content Slope	0.53 0.04	Very limited Slope Large stones content Gravel content Large stones content	1.00 0.53 0.15 0.01
Holyoke-----	35	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53
77D: Cheshire-----	45	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content Gravel content Large stones content	1.00 0.53 0.15 0.01
Holyoke-----	35	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53
78C: Holyoke-----	50	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Large stones content Slope	1.00 0.53 0.04	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
78E: Holyoke-----	50	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53	Very limited Slope Depth to bedrock Large stones content	1.00 1.00 0.53
78E: Rock Outcrop-----	25	Not rated		Not rated		Not rated	
79E: Rock Outcrop-----	55	Not rated		Not rated		Not rated	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
79E: Holyoke-----	25	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53	Very limited Depth to bedrock Slope Large stones content	1.00 1.00 0.53
80B: Bernardston-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.77	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.43	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.80 0.77 0.18
80C: Bernardston-----	80	Somewhat limited Depth to pan Depth to saturated zone Slope	0.79 0.77 0.63	Somewhat limited Depth to pan Slope Depth to saturated zone	0.79 0.63 0.43	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.80 0.77 0.18
81C: Bernardston-----	80	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.79 0.77 0.04	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.79 0.43 0.04	Very limited Large stones content Slope Depth to pan Depth to saturated zone Gravel content	1.00 1.00 0.80 0.77 0.18
81D: Bernardston-----	80	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.79 0.77	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.79 0.43	Very limited Slope Large stones content Depth to pan Depth to saturated zone Gravel content	1.00 1.00 0.80 0.77 0.18
82B: Broadbrook-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.84 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.84 0.19	Somewhat limited Slope Depth to pan Depth to saturated zone	0.88 0.84 0.39
82C: Broadbrook-----	80	Somewhat limited Depth to pan Slope Depth to saturated zone	0.84 0.63 0.39	Somewhat limited Depth to pan Slope Depth to saturated zone	0.84 0.63 0.19	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.84 0.39

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
82D: Broadbrook-----	80	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.84 0.39	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.84 0.19	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.84 0.39
83B: Broadbrook-----	80	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.84 0.53 0.39	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.84 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone	1.00 0.84 0.53 0.39
83C: Broadbrook-----	80	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.84 0.63 0.53 0.39	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.84 0.63 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone	1.00 0.84 0.53 0.39
84B: Paxton-----	55	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.80 0.39 0.03 0.01
Montauk-----	30	Somewhat limited Depth to pan Depth to saturated zone	0.84 0.07	Somewhat limited Depth to pan Depth to saturated zone	0.84 0.03	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.84 0.07 0.04
84C: Paxton-----	55	Somewhat limited Depth to pan Slope Depth to saturated zone	0.79 0.63 0.39	Somewhat limited Depth to pan Slope Depth to saturated zone	0.79 0.63 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.80 0.39 0.03 0.01
Montauk-----	30	Somewhat limited Depth to pan Slope Depth to saturated zone	0.84 0.63 0.07	Somewhat limited Depth to pan Slope Depth to saturated zone	0.84 0.63 0.03	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.84 0.07 0.04

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
84D: Paxton-----	55	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.79 0.39	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.79 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.80 0.39 0.03 0.01
Montauk-----	30	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.84 0.07	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.84 0.03	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.84 0.07 0.04
85B: Paxton-----	55	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.79 0.53 0.39	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.79 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone Gravel content	1.00 0.80 0.53 0.39 0.03
Montauk-----	30	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.84 0.53 0.07	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.84 0.53 0.03	Very limited Slope Depth to pan Large stones content Depth to saturated zone Gravel content	1.00 0.84 0.53 0.07 0.04
85C: Paxton-----	55	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.79 0.63 0.53 0.39	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.79 0.63 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone Gravel content	1.00 0.80 0.53 0.39 0.03
Montauk-----	30	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.84 0.63 0.53 0.07	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.84 0.63 0.53 0.03	Very limited Slope Depth to pan Large stones content Depth to saturated zone Gravel content	1.00 0.84 0.53 0.07 0.04

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
86C: Paxton-----	55	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.79 0.39 0.04	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.79 0.19 0.04	Very limited Large stones content Slope Depth to pan Depth to saturated zone Gravel content	1.00 1.00 0.80 0.39 0.03
Montauk-----	30	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.84 0.07 0.04	Very limited Large stones content Depth to pan Slope Depth to saturated zone	1.00 0.84 0.04 0.03	Very limited Large stones content Slope Depth to pan Depth to saturated zone Gravel content	1.00 1.00 0.84 0.07 0.04
86D: Paxton-----	55	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.79 0.39	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.79 0.19	Very limited Slope Large stones content Depth to pan Depth to saturated zone Gravel content	1.00 1.00 0.80 0.39 0.03
Montauk-----	30	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.84 0.07	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.84 0.03	Very limited Slope Large stones content Depth to pan Depth to saturated zone Gravel content	1.00 1.00 0.84 0.07 0.04
87B: Wethersfield-----	80	Somewhat limited Depth to pan Depth to saturated zone	0.71 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.71 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.71 0.39 0.16 0.01
87C: Wethersfield-----	80	Somewhat limited Depth to pan Slope Depth to saturated zone	0.71 0.63 0.39	Somewhat limited Depth to pan Slope Depth to saturated zone	0.71 0.63 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.71 0.39 0.16 0.01

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
87D: Wethersfield-----	80	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.71 0.39	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.71 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.71 0.39 0.16 0.01
88B: Wethersfield-----	80	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.71 0.53 0.39	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.71 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone	1.00 0.71 0.53 0.39
88C: Wethersfield-----	80	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.71 0.63 0.53 0.39	Somewhat limited Depth to pan Slope Large stones content Depth to saturated zone	0.71 0.63 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone	1.00 0.71 0.53 0.39
89C: Wethersfield-----	80	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.71 0.39 0.04	Very limited Large stones content Depth to pan Depth to saturated zone Slope	1.00 0.71 0.19 0.04	Very limited Large stones content Slope Depth to pan Depth to saturated zone	1.00 1.00 0.71 0.71 0.39
89D: Wethersfield-----	80	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.71 0.39	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.71 0.19	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 1.00 0.71 0.39
90B: Stockbridge-----	80	Somewhat limited Slow water movement	0.15	Somewhat limited Slow water movement	0.15	Very limited Slope Gravel content Slow water movement	1.00 0.43 0.15

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
90C: Stockbridge-----	80	Somewhat limited Slope Slow water movement	0.63 0.15	Somewhat limited Slope Slow water movement	0.63 0.15	Very limited Slope Gravel content Slow water movement	1.00 0.43 0.15
90D: Stockbridge-----	80	Very limited Slope Slow water movement	1.00 0.15	Very limited Slope Slow water movement	1.00 0.15	Very limited Slope Gravel content Slow water movement	1.00 0.43 0.15
91B: Stockbridge-----	80	Somewhat limited Large stones content Slow water movement	0.53 0.15	Somewhat limited Large stones content Slow water movement	0.53 0.15	Very limited Slope Large stones content Gravel content Slow water movement	1.00 0.53 0.43 0.15
91C: Stockbridge-----	80	Somewhat limited Slope Large stones content Slow water movement	0.63 0.53 0.15	Somewhat limited Slope Large stones content Slow water movement	0.63 0.53 0.15	Very limited Slope Large stones content Gravel content Slow water movement	1.00 0.53 0.43 0.15
91D: Stockbridge-----	80	Very limited Slope Large stones content Slow water movement	1.00 0.53 0.15	Very limited Slope Large stones content Slow water movement	1.00 0.53 0.15	Very limited Slope Large stones content Gravel content Slow water movement	1.00 0.53 0.43 0.15
92B: Nellis-----	85	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.18
92C: Nellis-----	85	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.18
92D: Nellis-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Gravel content	1.00 0.18

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
93C: Nellis-----	85	Somewhat limited Large stones content Slope	0.53 0.04	Somewhat limited Large stones content Slope	0.53 0.04	Very limited Slope Large stones content Gravel content	1.00 0.53 0.18
94C: Farmington-----	40	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope Gravel content	1.00 1.00 0.17
Nellis-----	35	Somewhat limited Large stones content Slope	0.53 0.04	Somewhat limited Large stones content Slope	0.53 0.04	Very limited Slope Large stones content Gravel content	1.00 0.53 0.18
94E: Farmington-----	40	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.17
Nellis-----	35	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content Gravel content	1.00 0.53 0.18
95C: Farmington-----	60	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to bedrock Slope Gravel content	1.00 1.00 0.17
Rock Outcrop-----	20	Not rated		Not rated		Not rated	
95E: Farmington-----	60	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.17
Rock Outcrop-----	20	Not rated		Not rated		Not rated	
96: Ipswich-----	85	Very limited Depth to saturated zone Salinity Flooding Ponding	1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Salinity Flooding	1.00 1.00 1.00 0.60	Very limited Depth to saturated zone Salinity Flooding Ponding	1.00 1.00 1.00 1.00

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
97: Pawcatuck-----	85	Very limited Depth to saturated zone Salinity Flooding Ponding	1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Salinity Flooding	1.00 1.00 1.00 0.60	Very limited Depth to saturated zone Salinity Flooding Ponding	1.00 1.00 1.00 1.00
98: Westbrook-----	80	Very limited Depth to saturated zone Salinity Flooding Ponding	1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Salinity Flooding	1.00 1.00 1.00 0.60	Very limited Depth to saturated zone Salinity Flooding Ponding	1.00 1.00 1.00 1.00
99: Westbrook, low salt-	80	Very limited Depth to saturated zone Flooding Ponding Salinity	1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Salinity Flooding	1.00 1.00 1.00 0.60	Very limited Depth to saturated zone Flooding Ponding Salinity	1.00 1.00 1.00 1.00
100: Suncook-----	80	Very limited Flooding Too sandy	1.00 0.50	Somewhat limited Too sandy	0.50	Somewhat limited Flooding Too sandy	0.60 0.50
101: Occum-----	80	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding Gravel content	0.60 0.06
102: Pootatuck-----	80	Very limited Flooding Depth to saturated zone	1.00 0.39	Somewhat limited Flooding Depth to saturated zone	0.40 0.19	Very limited Flooding Depth to saturated zone Gravel content	1.00 0.39 0.06
103: Rippowam-----	80	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding Gravel content	1.00 1.00 0.06
104: Bash-----	80	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 1.00
105: Hadley-----	80	Very limited Flooding	1.00	Not limited		Somewhat limited Flooding	0.60

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
106: Winooski-----	80	Very limited Flooding Depth to saturated zone	1.00 0.07	Somewhat limited Flooding Depth to saturated zone	0.40 0.03	Very limited Flooding Depth to saturated zone	1.00 0.07
107: Limerick-----	50	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 1.00
Lim-----	30	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 1.00
108: Saco-----	80	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00
109: Fluvaquents, Frequently Flooded-	50	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 1.00
Udifluents, Frequently Flooded-	35	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
221A: Ninigret-----	40	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone Gravel content Slope	0.39 0.22 0.12
Urban Land-----	35	Not rated		Not rated		Not rated	
224A: Deerfield-----	40	Somewhat limited Too sandy Depth to saturated zone	0.55 0.07	Somewhat limited Too sandy Depth to saturated zone	0.55 0.03	Somewhat limited Too sandy Depth to saturated zone	0.55 0.07
Urban Land-----	35	Not rated		Not rated		Not rated	
225B: Brancroft-----	40	Very limited Slow water movement Depth to saturated zone	1.00 0.81	Very limited Slow water movement Depth to saturated zone	1.00 0.48	Very limited Slow water movement Depth to saturated zone Slope	1.00 0.81 0.50

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
225B: Urban Land-----	35	Not rated		Not rated		Not rated	
226B: Berlin-----	40	Very limited Slow water movement Depth to saturated zone	1.00 0.81	Very limited Slow water movement Depth to saturated zone	1.00 0.48	Very limited Slow water movement Depth to saturated zone Slope	1.00 0.81 0.50
Urban Land-----	35	Not rated		Not rated		Not rated	
228B: Elmridge-----	40	Very limited Slow water movement Depth to saturated zone	1.00 0.39	Very limited Slow water movement Depth to saturated zone	1.00 0.19	Very limited Slow water movement Slope Depth to saturated zone	1.00 0.50 0.39
Urban Land-----	35	Not rated		Not rated		Not rated	
229B: Agawam-----	40	Not limited		Not limited		Somewhat limited Slope	0.50
Urban Land-----	35	Not rated		Not rated		Not rated	
229C: Agawam-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Urban Land-----	35	Not rated		Not rated		Not rated	
230B: Branford-----	40	Not limited		Not limited		Somewhat limited Slope Gravel content	0.50 0.06
Urban Land-----	35	Not rated		Not rated		Not rated	
230C: Branford-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.06
Urban Land-----	35	Not rated		Not rated		Not rated	
232B: Haven-----	40	Not limited		Not limited		Somewhat limited Slope Gravel content	0.50 0.06
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
234B: Merrimac-----	40	Not limited		Not limited		Somewhat limited Gravel content	0.78
						Slope	0.50
Urban Land-----	35	Not rated		Not rated		Not rated	
235B: Penwood-----	40	Somewhat limited Too sandy	0.32	Somewhat limited Too sandy	0.32	Somewhat limited Slope Too sandy	0.50 0.32
Urban Land-----	35	Not rated		Not rated		Not rated	
236B: Windsor-----	40	Not limited		Not limited		Somewhat limited Slope	0.50
Urban Land-----	35	Not rated		Not rated		Not rated	
237A: Manchester-----	40	Somewhat limited Gravel content	0.68	Somewhat limited Gravel content	0.68	Very limited Gravel content	1.00
Urban Land-----	35	Not rated		Not rated		Not rated	
237C: Manchester-----	40	Somewhat limited Gravel content Slope	0.68 0.04	Somewhat limited Gravel content Slope	0.68 0.04	Very limited Gravel content Slope	1.00 1.00
Urban Land-----	35	Not rated		Not rated		Not rated	
238A: Hinckley-----	40	Somewhat limited Gravel content	0.05	Somewhat limited Gravel content	0.05	Very limited Gravel content Large stones content	1.00 0.11
Urban Land-----	35	Not rated		Not rated		Not rated	
238C: Hinckley-----	40	Somewhat limited Gravel content Slope	0.05 0.04	Somewhat limited Gravel content Slope	0.05 0.04	Very limited Gravel content Slope Large stones content	1.00 1.00 0.11
Urban Land-----	35	Not rated		Not rated		Not rated	
240B: Ludlow-----	40	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Somewhat limited Depth to pan Gravel content Slope Depth to saturated zone	0.80 0.50 0.50 0.39
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
243B: Rainbow-----	40	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Somewhat limited Depth to pan Slope Depth to saturated zone	0.80 0.50 0.39
Urban Land-----	35	Not rated		Not rated		Not rated	
245B: Woodbridge-----	40	Somewhat limited Depth to pan Depth to saturated zone	0.46 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.46 0.19	Somewhat limited Slope Depth to pan Depth to saturated zone Gravel content	0.50 0.46 0.39 0.17
Urban Land-----	35	Not rated		Not rated		Not rated	
245C: Woodbridge-----	40	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.46 0.39	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.46 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content	1.00 0.46 0.39 0.17
Urban Land-----	35	Not rated		Not rated		Not rated	
248B: Georgia-----	40	Somewhat limited Large stones content Slow water movement Depth to saturated zone	0.53 0.15 0.07	Somewhat limited Large stones content Slow water movement Depth to saturated zone	0.53 0.15 0.03	Somewhat limited Slope Large stones content Gravel content Slow water movement Depth to saturated zone	0.88 0.53 0.20 0.15 0.07
Urban Land-----	35	Not rated		Not rated		Not rated	
250B: Sutton-----	40	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Slope Depth to saturated zone Gravel content	0.50 0.39 0.18
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
253B: Wapping-----	40	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Slope Depth to saturated zone Gravel content	0.50 0.39 0.06
Urban Land-----	35	Not rated		Not rated		Not rated	
255B: Watchaug-----	40	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Slope Depth to saturated zone Gravel content Large stones content	0.50 0.39 0.16 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
260B: Charlton-----	40	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.43
Urban Land-----	35	Not rated		Not rated		Not rated	
260C: Charlton-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.43
Urban Land-----	35	Not rated		Not rated		Not rated	
260D: Charlton-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Gravel content	1.00 0.43
Urban Land-----	35	Not rated		Not rated		Not rated	
263B: Cheshire-----	40	Not limited		Not limited		Very limited Slope Gravel content Large stones content	1.00 0.15 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
263C: Cheshire-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content Large stones content	1.00 0.15 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
266B: Narragansett-----	40	Not limited		Not limited		Very limited Slope Large stones content	1.00 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
269B: Yalesville-----	40	Not limited		Not limited		Very limited Slope Depth to bedrock	1.00 0.05
Urban Land-----	35	Not rated		Not rated		Not rated	
269C: Yalesville-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Depth to bedrock	1.00 0.05
Urban Land-----	35	Not rated		Not rated		Not rated	
273C: Urban Land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope Gravel content	1.00 0.43
Chatfield-----	15	Somewhat limited Large stones content Gravel content Slope	0.53 0.25 0.04	Somewhat limited Large stones content Gravel content Slope	0.53 0.25 0.04	Very limited Slope Depth to bedrock Large stones content	1.00 0.54 0.53
273E: Urban Land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Gravel content	1.00 0.43
Chatfield-----	15	Very limited Slope Large stones content Gravel content	1.00 0.53 0.25	Very limited Slope Large stones content Gravel content	1.00 0.53 0.25	Very limited Slope Depth to bedrock Large stones content	1.00 0.54 0.53

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
275C: Urban Land-----	45	Not rated		Not rated		Not rated	
Chatfield-----	30	Somewhat limited Large stones content Gravel content Slope	0.53 0.25 0.04	Somewhat limited Large stones content Gravel content Slope	0.53 0.25 0.04	Very limited Slope Depth to bedrock Large stones content	1.00 0.54 0.53
275E: Urban Land-----	35	Not rated		Not rated		Not rated	
Chatfield-----	25	Very limited Slope Large stones content Gravel content	1.00 0.53 0.25	Very limited Slope Large stones content Gravel content	1.00 0.53 0.25	Very limited Slope Depth to bedrock Large stones content	1.00 0.54 0.53
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
282B: Broadbrook-----	40	Somewhat limited Depth to pan Depth to saturated zone	0.84 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.84 0.19	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.84 0.39
Urban Land-----	35	Not rated		Not rated		Not rated	
284B: Paxton-----	40	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.79 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.80 0.39 0.03 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
284C: Paxton-----	40	Somewhat limited Depth to pan Slope Depth to saturated zone	0.79 0.63 0.39	Somewhat limited Depth to pan Slope Depth to saturated zone	0.79 0.63 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.80 0.39 0.03 0.01

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
284C: Urban Land-----	35	Not rated		Not rated		Not rated	
284D: Paxton-----	40	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.79 0.39	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.79 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.80 0.39 0.03 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
287B: Wethersfield-----	40	Somewhat limited Depth to pan Depth to saturated zone	0.71 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.71 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.71 0.39 0.16 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
287C: Wethersfield-----	40	Somewhat limited Depth to pan Slope Depth to saturated zone	0.71 0.63 0.39	Somewhat limited Depth to pan Slope Depth to saturated zone	0.71 0.63 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.71 0.39 0.16 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
287D: Wethersfield-----	40	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.71 0.39	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.71 0.19	Very limited Slope Depth to pan Depth to saturated zone Gravel content Large stones content	1.00 0.71 0.39 0.16 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
290B: Stockbridge-----	40	Somewhat limited Slow water movement	0.15	Somewhat limited Slow water movement	0.15	Very limited Slope Gravel content Slow water movement	1.00 0.43 0.15

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
290B: Urban Land-----	35	Not rated		Not rated		Not rated	
290C: Stockbridge-----	40	Somewhat limited Slope Slow water movement	0.63 0.15	Somewhat limited Slope Slow water movement	0.63 0.15	Very limited Slope Gravel content Slow water movement	1.00 0.43 0.15
Urban Land-----	35	Not rated		Not rated		Not rated	
290D: Stockbridge-----	40	Very limited Slope Slow water movement	1.00 0.15	Very limited Slope Slow water movement	1.00 0.15	Very limited Slope Gravel content Slow water movement	1.00 0.43 0.15
Urban Land-----	35	Not rated		Not rated		Not rated	
301: Beaches-----	50	Not rated		Not rated		Not rated	
Udipsamments-----	35	Very limited Flooding Too sandy Slope	1.00 1.00 0.01	Very limited Too sandy Slope	1.00 0.01	Very limited Too sandy Slope Flooding Gravel content	1.00 1.00 0.60 0.06
302: Dumps-----	95	Not rated		Not rated		Not rated	
303: Pits, Quarries-----	90	Not rated		Not rated		Not rated	
304: Udorthents-----	90	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Large stones content	1.00 0.01
305: Udorthents-----	65	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Large stones content	1.00 0.01
Pits-----	25	Not rated		Not rated		Not rated	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
306: Udorthents-----	50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Large stones content	1.00 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
307: Urban Land-----	80	Not rated		Not rated		Not rated	
308: Udorthents-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Large stones content	1.00 0.01
309: Udorthents-----	80	Very limited Flooding Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope Large stones content	1.00 0.01
310: Udorthents, Periodically Flooded-----	85	Very limited Flooding Slope	1.00 1.00	Very limited Slope Flooding	1.00 0.40	Very limited Flooding Slope Large stones content	1.00 1.00 0.01
401C: Macomber-----	55	Somewhat limited Gravel content Slope Large stones content	0.82 0.63 0.19	Somewhat limited Gravel content Slope Large stones content	0.82 0.63 0.19	Very limited Slope Depth to bedrock Large stones content	1.00 0.46 0.19
Taconic-----	30	Very limited Depth to bedrock Gravel content Slope	1.00 0.39 0.01	Very limited Depth to bedrock Gravel content Slope	1.00 0.39 0.01	Very limited Depth to bedrock Slope	1.00 1.00
402D: Macomber-----	50	Very limited Slope Gravel content	1.00 0.82	Very limited Slope Gravel content	1.00 0.82	Very limited Slope Depth to bedrock	1.00 0.46
Taconic-----	25	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.39	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.39	Very limited Slope Depth to bedrock	1.00 1.00
Rock Outcrop-----	15	Not rated		Not rated		Not rated	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
403C: Taconic-----	70	Very limited Depth to bedrock Gravel content Slope	1.00 0.39 0.01	Very limited Depth to bedrock Gravel content Slope	1.00 0.39 0.01	Very limited Depth to bedrock Slope	1.00 1.00
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
403E: Taconic-----	70	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.39	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.39	Very limited Slope Depth to bedrock	1.00 1.00
Rock Outcrop-----	20	Not rated		Not rated		Not rated	
403F: Taconic-----	70	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.39	Very limited Slope Depth to bedrock Gravel content	1.00 1.00 0.39	Very limited Slope Depth to bedrock	1.00 1.00
Rock Outcrop-----	20	Not rated		Not rated		Not rated	
405C: Dummerston-----	85	Somewhat limited Large stones content Gravel content Slope	0.19 0.09 0.01	Somewhat limited Large stones content Gravel content Slope	0.19 0.09 0.01	Very limited Slope Large stones content	1.00 0.19
405E: Dummerston-----	85	Very limited Slope Large stones content Gravel content	1.00 0.19 0.09	Very limited Slope Large stones content Gravel content	1.00 0.19 0.09	Very limited Slope Large stones content	1.00 0.19
407C: Lanesboro-----	85	Somewhat limited Large stones content Depth to pan Depth to saturated zone Slope	0.53 0.46 0.39 0.16	Somewhat limited Large stones content Depth to pan Depth to saturated zone Slope	0.53 0.46 0.19 0.16	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 0.53 0.46 0.39
407E: Lanesboro-----	85	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 0.53 0.46 0.39	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 0.53 0.46 0.19	Very limited Slope Large stones content Depth to pan Depth to saturated zone	1.00 0.53 0.46 0.39

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
408C: Fullam-----	85	Very limited Depth to pan Large stones content Depth to saturated zone Slope	1.00 0.53 0.24 0.16	Very limited Depth to pan Large stones content Slope Depth to saturated zone	1.00 0.53 0.16 0.12	Very limited Slope Depth to pan Large stones content Depth to saturated zone	1.00 1.00 0.53 0.24
409B: Brayton-----	85	Very limited Depth to saturated zone Depth to pan Large stones content	1.00 1.00 0.53	Very limited Depth to saturated zone Depth to pan Large stones content	1.00 1.00 0.53	Very limited Depth to saturated zone Depth to pan Slope Large stones content	1.00 1.00 0.88 0.53
412B: Bice-----	85	Not limited		Not limited		Very limited Slope	1.00
412C: Bice-----	85	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
412D: Bice-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
413C: Bice-----	45	Somewhat limited Slope	0.01	Somewhat limited Slope	0.01	Very limited Slope	1.00
Millsite-----	40	Somewhat limited Slope	0.01	Somewhat limited Slope	0.01	Very limited Slope Depth to bedrock	1.00 0.35
413E: Bice-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Millsite-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Depth to bedrock	1.00 0.35
414: Fredon, cold-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Gravel content	1.00 0.04
415C: Millsite-----	40	Somewhat limited Slope	0.01	Somewhat limited Slope	0.01	Very limited Slope Depth to bedrock	1.00 0.35

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
415C: Westminster-----	40	Very limited Depth to bedrock Slope	1.00 0.01	Very limited Depth to bedrock Slope	1.00 0.01	Very limited Depth to bedrock Slope	1.00 1.00
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
415E: Millsite-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Depth to bedrock	1.00 0.35
Westminster-----	40	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
416E: Rock Outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00
416F: Rock Outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Slope Depth to bedrock	1.00 1.00
417B: Bice-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Slope Large stones content	1.00 0.53
417C: Bice-----	85	Somewhat limited Large stones content Slope	0.53 0.16	Somewhat limited Large stones content Slope	0.53 0.16	Very limited Slope Large stones content	1.00 0.53
417D: Bice-----	85	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content	1.00 0.53	Very limited Slope Large stones content	1.00 0.53
418C: Schroon-----	85	Somewhat limited Large stones content Depth to saturated zone Slope	0.53 0.39 0.04	Somewhat limited Large stones content Depth to saturated zone Slope	0.53 0.19 0.04	Very limited Slope Large stones content Depth to saturated zone	1.00 0.53 0.39

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
420A: Schroon-----	85	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone	0.39
420B: Schroon-----	85	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Slope Depth to saturated zone	0.88 0.39
421A: Ninigret, cold-----	85	Somewhat limited Depth to saturated zone	0.39	Somewhat limited Depth to saturated zone	0.19	Somewhat limited Depth to saturated zone Gravel content	0.39 0.22
423A: Sudbury, cold-----	85	Somewhat limited Depth to saturated zone	0.07	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Depth to saturated zone	0.07
424B: Shelburne-----	85	Somewhat limited Depth to pan Depth to saturated zone	0.74 0.39	Somewhat limited Depth to pan Depth to saturated zone	0.74 0.19	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.74 0.39
424C: Shelburne-----	85	Somewhat limited Depth to pan Slope Depth to saturated zone	0.74 0.63 0.39	Somewhat limited Depth to pan Slope Depth to saturated zone	0.74 0.63 0.19	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.74 0.39
424D: Shelburne-----	85	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.74 0.39	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.74 0.19	Very limited Slope Depth to pan Depth to saturated zone	1.00 0.74 0.39
425B: Shelburne-----	85	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.74 0.53 0.39	Somewhat limited Depth to pan Large stones content Depth to saturated zone	0.74 0.53 0.19	Very limited Slope Depth to pan Large stones content Depth to saturated zone	1.00 0.74 0.53 0.39

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
425C: Shelburne-----	85	Somewhat limited		Somewhat limited		Very limited	
		Depth to pan	0.74	Depth to pan	0.74	Slope	1.00
		Slope	0.63	Slope	0.63	Depth to pan	0.74
		Large stones content	0.53	Large stones content	0.53	Large stones content	0.53
		Depth to saturated zone	0.39	Depth to saturated zone	0.19	Depth to saturated zone	0.39
426D: Shelburne-----	85	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Large stones content	1.00	Large stones content	1.00	Large stones content	1.00
		Depth to pan	0.74	Depth to pan	0.74	Depth to pan	0.74
		Depth to saturated zone	0.39	Depth to saturated zone	0.19	Depth to saturated zone	0.39
427B: Ashfield-----	85	Very limited		Very limited		Very limited	
		Slow water movement	1.00	Slow water movement	1.00	Slow water movement	1.00
		Depth to saturated zone	0.98	Depth to saturated zone	0.75	Slope	1.00
		Depth to pan	0.54	Depth to pan	0.54	Depth to saturated zone	0.98
		Large stones content	0.53	Large stones content	0.53	Depth to pan	0.54
						Large stones content	0.53
427C: Ashfield-----	85	Somewhat limited		Somewhat limited		Very limited	
		Depth to saturated zone	0.98	Depth to saturated zone	0.75	Slope	1.00
		Slope	0.63	Slope	0.63	Depth to saturated zone	0.98
		Depth to pan	0.54	Depth to pan	0.54	Depth to pan	0.54
		Large stones content	0.53	Large stones content	0.53	Large stones content	0.53
428A: Ashfield-----	85	Somewhat limited		Somewhat limited		Somewhat limited	
		Depth to saturated zone	0.98	Depth to saturated zone	0.75	Depth to saturated zone	0.98
		Depth to pan	0.54	Depth to pan	0.54	Depth to pan	0.54
428B: Ashfield-----	85	Somewhat limited		Somewhat limited		Very limited	
		Depth to saturated zone	0.98	Depth to saturated zone	0.75	Slope	1.00
		Depth to pan	0.54	Depth to pan	0.54	Depth to saturated zone	0.98
						Depth to pan	0.54
428C: Ashfield-----	85	Somewhat limited		Somewhat limited		Very limited	
		Depth to saturated zone	0.98	Depth to saturated zone	0.75	Slope	1.00
		Slope	0.63	Slope	0.63	Depth to saturated zone	0.98
		Depth to pan	0.54	Depth to pan	0.54	Depth to pan	0.54

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
429A: Agawam, cold-----	80	Not limited		Not limited		Not limited	
429B: Agawam, cold-----	80	Not limited		Not limited		Very limited Slope	1.00
429C: Agawam, cold-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
433: Moosilauke-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
434A: Merrimac, cold-----	80	Not limited		Not limited		Somewhat limited Gravel content	0.78
434B: Merrimac, cold-----	80	Not limited		Not limited		Very limited Slope Gravel content	1.00 0.78
434C: Merrimac, cold-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope Gravel content	1.00 0.78
435: Scarboro-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
436: Halsey-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
437: Wonsqueak-----	85	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
438: Bucksport-----	85	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
440A: Boscawen-----	80	Not limited		Not limited		Not limited	

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
440C: Boscawen-----	80	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
440E: Boscawen-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
442: Brayton-----	85	Very limited Depth to saturated zone Depth to pan	1.00 0.71	Very limited Depth to saturated zone Depth to pan	1.00 0.71	Very limited Depth to saturated zone Slope Depth to pan	1.00 0.88 0.71
443: Brayton-----	50	Very limited Depth to saturated zone Large stones content Depth to pan	1.00 1.00 0.71	Very limited Depth to saturated zone Large stones content Depth to pan	1.00 1.00 0.71	Very limited Depth to saturated zone Large stones content Slope Depth to pan	1.00 1.00 0.88 0.71
Loonmeadow-----	35	Very limited Depth to saturated zone Large stones content Ponding Slow water movement	1.00 1.00 1.00 0.85	Very limited Depth to saturated zone Large stones content Ponding Slow water movement	1.00 1.00 1.00 0.85	Very limited Depth to saturated zone Large stones content Ponding Slow water movement	1.00 1.00 1.00 0.85
448B: Hogansburg-----	85	Somewhat limited Depth to saturated zone	0.07	Somewhat limited Depth to saturated zone	0.03	Somewhat limited Slope Depth to saturated zone Gravel content	0.88 0.07 0.06
449B: Hogansburg-----	85	Very limited Large stones content Depth to saturated zone	1.00 0.07	Very limited Large stones content Depth to saturated zone	1.00 0.03	Very limited Slope Large stones content Depth to saturated zone Gravel content	1.00 1.00 0.07 0.06
449C: Hogansburg-----	85	Very limited Large stones content Slope Depth to saturated zone	1.00 0.63 0.07	Very limited Large stones content Slope Depth to saturated zone	1.00 0.63 0.03	Very limited Slope Large stones content Depth to saturated zone Gravel content	1.00 1.00 0.07 0.06

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
450B: Pyrities-----	80	Somewhat limited	0.15	Somewhat limited	0.15	Very limited	1.00
		Slow water movement		Slow water movement		Slope	
						Slow water movement	0.15
450C: Pyrities-----	80	Somewhat limited	0.63	Somewhat limited	0.63	Very limited	1.00
		Slope		Slope		Slope	
		Slow water movement	0.15	Slow water movement	0.15	Slow water movement	0.15
450D: Pyrities-----	80	Very limited	1.00	Very limited	1.00	Very limited	1.00
		Slope		Slope		Slope	
		Slow water movement	0.15	Slow water movement	0.15	Slow water movement	0.15
451B: Pyrities-----	80	Somewhat limited	0.53	Somewhat limited	0.53	Very limited	1.00
		Large stones content		Large stones content		Slope	
		Slow water movement	0.15	Slow water movement	0.15	Large stones content	0.53
						Slow water movement	0.15
451C: Pyrities-----	80	Somewhat limited	0.63	Somewhat limited	0.63	Very limited	1.00
		Slope		Slope		Slope	
		Large stones content	0.53	Large stones content	0.53	Large stones content	0.53
		Slow water movement	0.15	Slow water movement	0.15	Slow water movement	0.15
451D: Pyrities-----	80	Very limited	1.00	Very limited	1.00	Very limited	1.00
		Slope		Slope		Slope	
		Large stones content	0.53	Large stones content	0.53	Large stones content	0.53
		Slow water movement	0.15	Slow water movement	0.15	Slow water movement	0.15
457: Mudgepond-----	80	Very limited	1.00	Very limited	1.00	Very limited	1.00
		Depth to saturated zone		Depth to saturated zone		Depth to saturated zone	
						Gravel content	0.22
458: Mudgepond-----	55	Very limited	1.00	Very limited	1.00	Very limited	1.00
		Depth to saturated zone		Depth to saturated zone		Depth to saturated zone	
		Large stones content	1.00	Large stones content	1.00	Large stones content	1.00
						Gravel content	0.22

Table 13.—Recreation (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Camp areas		Picnic areas		Playgrounds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
458: Alden-----	35	Very limited Depth to saturated zone Large stones content Ponding Slow water movement	1.00 1.00 1.00 0.15	Very limited Depth to saturated zone Large stones content Ponding Slow water movement	1.00 1.00 1.00 0.15	Very limited Depth to saturated zone Large stones content Ponding Slow water movement Large stones content	1.00 1.00 1.00 0.15 0.01
501: Ondawa-----	85	Very limited Flooding Slow water movement	1.00 0.15	Somewhat limited Slow water movement	0.15	Very limited Gravel content Flooding Slow water movement	1.00 0.60 0.15
503: Rumney-----	80	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding Gravel content	1.00 1.00 0.22
508: Medomak-----	85	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00

Table 14.—Recreation (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Depth to pan	1.00 0.99
3: Ridgebury-----	40	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Depth to pan	1.00 0.99
Leicester-----	35	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone	1.00
Whitman-----	15	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to pan Depth to saturated zone Ponding	1.00 1.00 1.00
4: Leicester-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Large stones content	1.00 0.01
5: Wilbraham-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Depth to pan	1.00 1.00
6: Wilbraham-----	60	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Depth to pan	1.00 1.00
Menlo-----	25	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Depth to pan	1.00 1.00 0.71

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7: Mudgepond-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
8: Mudgepond-----	45	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone	1.00
Alden-----	35	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Large stones content	1.00 1.00 0.01
9: Scitico-----	40	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Shaker-----	30	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Maybid-----	15	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
10: Raynham-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
12: Raypol-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
13: Walpole-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
14: Fredon-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
15: Scarboro-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Organic matter content Depth to saturated zone Ponding	1.00 1.00 1.00

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
16: Halsey-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
17: Timakwa-----	45	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Organic matter content Depth to saturated zone Ponding	1.00 1.00 1.00
Natchaug-----	40	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
18: Catden-----	40	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Freetown-----	40	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
20A: Ellington-----	80	Not limited		Not limited		Not limited	
21A: Ninigret-----	60	Not limited		Not limited		Not limited	
Tisbury-----	25	Not limited		Not limited		Not limited	
22A: Hero-----	85	Not limited		Not limited		Somewhat limited Gravel content Large stones content	0.18 0.01
22B: Hero-----	85	Not limited		Not limited		Somewhat limited Gravel content Large stones content	0.18 0.01
23A: Sudbury-----	80	Not limited		Not limited		Not limited	
24A: Deerfield-----	80	Somewhat limited Too sandy	0.55	Somewhat limited Too sandy	0.55	Somewhat limited Droughty	0.17
25A: Brancroft-----	80	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Depth to saturated zone	0.11	Not limited	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
25B: Brancroft-----	80	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Depth to saturated zone	0.11	Not limited	
25C: Brancroft-----	80	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Slope	0.63
26A: Berlin-----	80	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Depth to saturated zone	0.11	Not limited	
26B: Berlin-----	80	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Depth to saturated zone	0.11	Not limited	
27A: Belgrade-----	80	Not limited		Not limited		Not limited	
28A: Elmridge-----	80	Not limited		Not limited		Not limited	
28B: Elmridge-----	80	Not limited		Not limited		Not limited	
29A: Agawam-----	80	Not limited		Not limited		Not limited	
29B: Agawam-----	80	Not limited		Not limited		Not limited	
29C: Agawam-----	80	Not limited		Not limited		Somewhat limited Slope	0.63
30A: Branford-----	80	Not limited		Not limited		Not limited	
30B: Branford-----	80	Not limited		Not limited		Not limited	
30C: Branford-----	80	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Somewhat limited Slope	0.63
31A: Copake-----	85	Not limited		Not limited		Not limited	
31B: Copake-----	85	Not limited		Not limited		Not limited	
31C: Copake-----	85	Not limited		Not limited		Somewhat limited Slope	0.63

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
32A:							
Haven-----	60	Not limited		Not limited		Not limited	
Enfield-----	25	Not limited		Not limited		Not limited	
32B:							
Haven-----	60	Not limited		Not limited		Not limited	
Enfield-----	25	Not limited		Not limited		Not limited	
32C:							
Haven-----	60	Not limited		Not limited		Somewhat limited Slope	0.63
Enfield-----	25	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Somewhat limited Slope	0.63
33A:							
Hartford-----	80	Not limited		Not limited		Not limited	
33B:							
Hartford-----	80	Not limited		Not limited		Not limited	
34A:							
Merrimac-----	80	Not limited		Not limited		Not limited	
34B:							
Merrimac-----	80	Not limited		Not limited		Not limited	
34C:							
Merrimac-----	80	Not limited		Not limited		Somewhat limited Slope	0.63
35A:							
Penwood-----	80	Somewhat limited Too sandy	0.32	Somewhat limited Too sandy	0.32	Very limited Droughty	1.00
35B:							
Penwood-----	80	Somewhat limited Too sandy	0.32	Somewhat limited Too sandy	0.32	Very limited Droughty	1.00
36A:							
Windsor-----	80	Not limited		Not limited		Somewhat limited Droughty	0.99
36B:							
Windsor-----	80	Not limited		Not limited		Somewhat limited Droughty	0.99
36C:							
Windsor-----	80	Not limited		Not limited		Somewhat limited Droughty Slope	0.99 0.63
37A:							
Manchester-----	80	Not limited		Not limited		Very limited Droughty Gravel content	1.00 0.68

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
37C: Manchester-----	80	Not limited		Not limited		Very limited Droughty Gravel content Slope	1.00 0.68 0.04
37E: Manchester-----	80	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Slope Droughty Gravel content	1.00 1.00 0.68
38A: Hinckley-----	80	Not limited		Not limited		Very limited Droughty Large stones content Gravel content	1.00 0.11 0.05
38C: Hinckley-----	80	Not limited		Not limited		Very limited Droughty Large stones content Gravel content Slope	1.00 0.11 0.05 0.04
38E: Hinckley-----	80	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Slope Droughty Large stones content Gravel content	1.00 1.00 0.11 0.05
39A: Groton-----	85	Not limited		Not limited		Somewhat limited Droughty Gravel content	0.98 0.59
39C: Groton-----	85	Not limited		Not limited		Somewhat limited Droughty Gravel content Slope	0.98 0.59 0.04
39E: Groton-----	85	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Slope Droughty Gravel content	1.00 0.98 0.59
40A: Ludlow-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.79
40B: Ludlow-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.79

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
41B: Ludlow-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.79
42C: Ludlow-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Depth to pan Slope	0.79 0.04
43A: Rainbow-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.79
43B: Rainbow-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.79
44B: Rainbow-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.79
45A: Woodbridge-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.46
45B: Woodbridge-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.46
45C: Woodbridge-----	80	Not limited		Not limited		Somewhat limited Slope Depth to pan	0.63 0.46
46B: Woodbridge-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.46
46C: Woodbridge-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope Depth to pan	0.63 0.46
47C: Woodbridge-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Depth to pan Slope Large stones content	0.46 0.04 0.01
48B: Georgia-----	50	Not limited		Not limited		Not limited	
Amenia-----	35	Not limited		Not limited		Not limited	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
48C: Georgia-----	50	Not limited		Not limited		Somewhat limited Slope	0.63
Amenia-----	35	Not limited		Not limited		Somewhat limited Slope	0.63
49B: Georgia-----	50	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Not limited	
Amenia-----	35	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Not limited	
49C: Georgia-----	50	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope	0.63
Amenia-----	35	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope	0.63
50A: Sutton-----	80	Not limited		Not limited		Not limited	
50B: Sutton-----	80	Not limited		Not limited		Not limited	
51B: Sutton-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
52C: Sutton-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope	0.04
53A: Wapping-----	80	Not limited		Not limited		Not limited	
53B: Wapping-----	80	Not limited		Not limited		Not limited	
54B: Wapping-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
55A: Watchaug-----	80	Not limited		Not limited		Somewhat limited Large stones content	0.01
55B: Watchaug-----	80	Not limited		Not limited		Somewhat limited Large stones content	0.01

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
56B: Watchaug-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.01
57B: Gloucester-----	80	Not limited		Not limited		Not limited	
57C: Gloucester-----	80	Not limited		Not limited		Somewhat limited Slope	0.63
57D: Gloucester-----	80	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00
58B: Gloucester-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
58C: Gloucester-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.63
59C: Gloucester-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope	0.04
59D: Gloucester-----	80	Very limited Large stones content Slope	1.00 0.50	Very limited Large stones content	1.00	Very limited Slope	1.00
60B: Canton-----	45	Not limited		Not limited		Somewhat limited Gravel content	0.01
Charlton-----	35	Not limited		Not limited		Not limited	
60C: Canton-----	45	Not limited		Not limited		Somewhat limited Slope Gravel content	0.63 0.01
Charlton-----	35	Not limited		Not limited		Somewhat limited Slope	0.63
60D: Canton-----	45	Somewhat limited Slope	0.50	Not limited		Very limited Slope Gravel content	1.00 0.01
Charlton-----	35	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
61B: Canton-----	45	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Gravel content	0.01
Charlton-----	35	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
61C: Canton-----	45	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.63
						Gravel content	0.01
Charlton-----	35	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.63
62C: Canton-----	45	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope	0.04
						Gravel content	0.01
Charlton-----	35	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope	0.04
62D: Canton-----	45	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content	1.00	Very limited Slope	1.00
						Gravel content	0.01
Charlton-----	35	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content	1.00	Very limited Slope	1.00
63B: Cheshire-----	80	Not limited		Not limited		Somewhat limited Large stones content	0.01
63C: Cheshire-----	80	Not limited		Not limited		Somewhat limited Slope Large stones content	0.63 0.01
63D: Cheshire-----	80	Somewhat limited Slope	0.50	Not limited		Very limited Slope Large stones content	1.00 0.01
64B: Cheshire-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.01

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
64C: Cheshire-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope Large stones content	0.63 0.01
65C: Cheshire-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope Large stones content	0.04 0.01
65D: Cheshire-----	80	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content	1.00	Very limited Slope Large stones content	1.00 0.01
66B: Narragansett-----	80	Not limited		Not limited		Somewhat limited Large stones content	0.01
66C: Narragansett-----	80	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Somewhat limited Slope Large stones content	0.63 0.01
67B: Narragansett-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.01
67C: Narragansett-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope Large stones content	0.63 0.01
68C: Narragansett-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope Large stones content	0.04 0.01
68D: Narragansett-----	80	Very limited Large stones content Slope	1.00 0.50	Very limited Large stones content	1.00	Very limited Slope Large stones content	1.00 0.01
69B: Yalesville-----	75	Not limited		Not limited		Somewhat limited Depth to bedrock	0.05

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
69C: Yalesville-----	75	Not limited		Not limited		Somewhat limited Slope	0.63
						Depth to bedrock	0.05
70C: Branford-----	50	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Somewhat limited Slope	0.04
Holyoke-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock	1.00
						Droughty Slope	0.25 0.04
71C: Brookfield-----	45	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.04
Brimfield-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock	1.00
						Droughty Slope	0.93 0.04
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
71E: Brookfield-----	45	Very limited Slope	1.00	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
		Large stones content	0.53	Slope	0.22		
Brimfield-----	30	Very limited Slope	1.00	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock	1.00
		Large stones content	0.53	Slope	0.22	Slope	1.00
						Droughty	0.93
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
73C: Charlton-----	45	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.04
Chatfield-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to bedrock	0.54
						Gravel content	0.25
						Slope	0.04

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
73E: Charlton-----	45	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content Slope	0.53 0.22	Very limited Slope	1.00
Chatfield-----	30	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content Slope	0.53 0.22	Very limited Slope Depth to bedrock Gravel content	1.00 0.54 0.25
74C: Narragansett-----	55	Very limited Water erosion Large stones content	1.00 0.53	Very limited Water erosion Large stones content	1.00 0.53	Somewhat limited Slope Large stones content	0.04 0.01
Hollis-----	20	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.22 0.04
75C: Hollis-----	35	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.22 0.04
Chatfield-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to bedrock Gravel content Slope	0.54 0.25 0.04
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
75E: Hollis-----	35	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content Slope	1.00 0.22	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.22
Chatfield-----	30	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content Slope	0.53 0.22	Very limited Slope Depth to bedrock Gravel content	1.00 0.54 0.25

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
75E: Rock Outcrop-----	15	Not rated		Not rated		Not rated	
76E: Rock Outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Large stones content Slope	1.00 0.98	Very limited Large stones content	1.00	Very limited Depth to bedrock Droughty Slope Gravel content	1.00 1.00 0.22
76F: Rock Outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Slope Large stones content	1.00 1.00	Very limited Slope Large stones content	1.00 1.00	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.22
77C: Cheshire-----	45	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope Large stones content	0.04 0.01
Holyoke-----	35	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock Droughty Slope	1.00 0.25 0.04
77D: Cheshire-----	45	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content	0.53	Very limited Slope Large stones content	1.00 0.01
Holyoke-----	35	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.25
78C: Holyoke-----	50	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock Droughty Slope	1.00 0.25 0.04

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
78C: Rock Outcrop-----	25	Not rated		Not rated		Not rated	
78E: Holyoke-----	50	Very limited Slope	1.00	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock	1.00
		Large stones content	0.53	Slope	0.22	Slope	1.00
						Droughty	0.25
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
79E: Rock Outcrop-----	55	Not rated		Not rated		Not rated	
Holyoke-----	25	Somewhat limited Slope	0.98	Somewhat limited Large stones content	0.53	Very limited Depth to bedrock	1.00
		Large stones content	0.53			Slope	1.00
						Droughty	0.25
80B: Bernardston-----	80	Somewhat limited Depth to saturated zone	0.08	Somewhat limited Depth to saturated zone	0.08	Somewhat limited Depth to pan	0.79
80C: Bernardston-----	80	Somewhat limited Depth to saturated zone	0.08	Somewhat limited Depth to saturated zone	0.08	Somewhat limited Depth to pan	0.79
						Slope	0.63
81C: Bernardston-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.08	Depth to saturated zone	0.08	Slope	0.04
81D: Bernardston-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Very limited Slope	1.00
		Slope	0.50	Depth to saturated zone	0.08	Depth to pan	0.79
		Depth to saturated zone	0.08				
82B: Broadbrook-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.84
82C: Broadbrook-----	80	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Somewhat limited Depth to pan	0.84
						Slope	0.63

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
82D: Broadbrook-----	80	Very limited Water erosion Slope	1.00 0.50	Very limited Water erosion	1.00	Very limited Slope Depth to pan	1.00 0.84
83B: Broadbrook-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.84
83C: Broadbrook-----	80	Very limited Water erosion Large stones content	1.00 0.53	Very limited Water erosion Large stones content	1.00 0.53	Somewhat limited Depth to pan Slope	0.84 0.63
84B: Paxton-----	55	Not limited		Not limited		Somewhat limited Depth to pan Large stones content	0.79 0.01
Montauk-----	30	Not limited		Not limited		Somewhat limited Depth to pan	0.84
84C: Paxton-----	55	Not limited		Not limited		Somewhat limited Depth to pan Slope Large stones content	0.79 0.63 0.01
Montauk-----	30	Not limited		Not limited		Somewhat limited Depth to pan Slope	0.84 0.63
84D: Paxton-----	55	Somewhat limited Slope	0.50	Not limited		Very limited Slope Depth to pan Large stones content	1.00 0.79 0.01
Montauk-----	30	Somewhat limited Slope	0.50	Not limited		Very limited Slope Depth to pan	1.00 0.84
85B: Paxton-----	55	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan Large stones content	0.79 0.01
Montauk-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.84

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
85C: Paxton-----	55	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.79
						Slope	0.63
						Large stones content	0.01
85C: Montauk-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.84
						Slope	0.63
86C: Paxton-----	55	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Depth to pan	0.79
						Slope	0.04
						Large stones content	0.01
Montauk-----	30	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Depth to pan	0.84
						Slope	0.04
86D: Paxton-----	55	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content	1.00	Very limited Slope	1.00
						Depth to pan	0.79
						Large stones content	0.01
Montauk-----	30	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content	1.00	Very limited Slope	1.00
						Depth to pan	0.84
87B: Wethersfield-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.71
						Large stones content	0.01
87C: Wethersfield-----	80	Not limited		Not limited		Somewhat limited Depth to pan	0.71
						Slope	0.63
						Large stones content	0.01
87D: Wethersfield-----	80	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00
						Depth to pan	0.71
						Large stones content	0.01

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
88B: Wethersfield-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.71
88C: Wethersfield-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan Slope	0.71 0.63
89C: Wethersfield-----	80	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Depth to pan Slope	0.71 0.04
89D: Wethersfield-----	80	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content	1.00	Very limited Slope Depth to pan	1.00 0.71
90B: Stockbridge-----	80	Not limited		Not limited		Not limited	
90C: Stockbridge-----	80	Not limited		Not limited		Somewhat limited Slope	0.63
90D: Stockbridge-----	80	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00
91B: Stockbridge-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
91C: Stockbridge-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.63
91D: Stockbridge-----	80	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
92B: Nellis-----	85	Not limited		Not limited		Not limited	
92C: Nellis-----	85	Not limited		Not limited		Somewhat limited Slope	0.63
92D: Nellis-----	85	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
93C: Nellis-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.04
94C: Farmington-----	40	Not limited		Not limited		Very limited Depth to bedrock Droughty Slope	1.00 0.81 0.04
Nellis-----	35	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.04
94E: Farmington-----	40	Very limited Slope	1.00	Not limited		Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.81
Nellis-----	35	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
95C: Farmington-----	60	Not limited		Not limited		Very limited Depth to bedrock Droughty Slope	1.00 0.81 0.04
Rock Outcrop-----	20	Not rated		Not rated		Not rated	
95E: Farmington-----	60	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.81
Rock Outcrop-----	20	Not rated		Not rated		Not rated	
96: Ipswich-----	85	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Ponding Flooding Organic matter content Salinity Sulfur content	1.00 1.00 1.00 1.00 1.00

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
97: Pawcatuck-----	85	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Ponding Flooding Organic matter content Salinity Sulfur content	1.00 1.00 1.00 1.00 1.00
98: Westbrook-----	80	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Ponding Flooding Salinity Sulfur content Depth to saturated zone	1.00 1.00 1.00 1.00 1.00
99: Westbrook, low salt-	80	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.60	Very limited Ponding Flooding Sulfur content Depth to saturated zone Salinity	1.00 1.00 1.00 1.00 1.00
100: Suncook-----	80	Somewhat limited Too sandy	0.50	Somewhat limited Too sandy	0.50	Somewhat limited Droughty Flooding	0.89 0.60
101: Occum-----	80	Not limited		Not limited		Somewhat limited Flooding	0.60
102: Pootatuck-----	80	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
103: Rippowam-----	80	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
104: Bash-----	80	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 0.99
105: Hadley-----	80	Not limited		Not limited		Somewhat limited Flooding	0.60

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
106: Winooski-----	80	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
107: Limerick-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	0.40	Flooding	0.40	Depth to saturated zone	1.00
Lim-----	30	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	0.40	Flooding	0.40	Depth to saturated zone	1.00
108: Saco-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Ponding	1.00	Ponding	1.00	Depth to saturated zone	1.00
		Flooding	0.40	Flooding	0.40	Ponding	1.00
109: Fluvaquents, Frequently Flooded-	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	0.40	Flooding	0.40	Depth to saturated zone	1.00
Udifluvents, Frequently Flooded-	35	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
221A: Ninigret-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
224A: Deerfield-----	40	Somewhat limited Too sandy	0.55	Somewhat limited Too sandy	0.55	Somewhat limited Droughty	0.17
		Urban Land-----	35	Not rated		Not rated	
225B: Brancroft-----	40	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Depth to saturated zone	0.11	Not limited	
		Urban Land-----	35	Not rated		Not rated	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
226B: Berlin-----	40	Somewhat limited Depth to saturated zone	0.11	Somewhat limited Depth to saturated zone	0.11	Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
228B: Elmridge-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
229B: Agawam-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
229C: Agawam-----	40	Not limited		Not limited		Somewhat limited Slope	0.63
Urban Land-----	35	Not rated		Not rated		Not rated	
230B: Branford-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
230C: Branford-----	40	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Somewhat limited Slope	0.63
Urban Land-----	35	Not rated		Not rated		Not rated	
232B: Haven-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
234B: Merrimac-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
235B: Penwood-----	40	Somewhat limited Too sandy	0.32	Somewhat limited Too sandy	0.32	Very limited Droughty	1.00
Urban Land-----	35	Not rated		Not rated		Not rated	
236B: Windsor-----	40	Not limited		Not limited		Somewhat limited Droughty	0.99
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
237A: Manchester-----	40	Not limited		Not limited		Very limited Droughty Gravel content	1.00 0.68
Urban Land-----	35	Not rated		Not rated		Not rated	
237C: Manchester-----	40	Not limited		Not limited		Very limited Droughty Gravel content Slope	1.00 0.68 0.04
Urban Land-----	35	Not rated		Not rated		Not rated	
238A: Hinckley-----	40	Not limited		Not limited		Very limited Droughty Large stones content Gravel content	1.00 0.11 0.05
Urban Land-----	35	Not rated		Not rated		Not rated	
238C: Hinckley-----	40	Not limited		Not limited		Very limited Droughty Large stones content Gravel content Slope	1.00 0.11 0.05 0.04
Urban Land-----	35	Not rated		Not rated		Not rated	
240B: Ludlow-----	40	Not limited		Not limited		Somewhat limited Depth to pan	0.79
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
243B: Rainbow-----	40	Not limited		Not limited		Somewhat limited Depth to pan	0.79
Urban Land-----	35	Not rated		Not rated		Not rated	
245B: Woodbridge-----	40	Not limited		Not limited		Somewhat limited Depth to pan	0.46
Urban Land-----	35	Not rated		Not rated		Not rated	
245C: Woodbridge-----	40	Not limited		Not limited		Somewhat limited Slope Depth to pan	0.63 0.46
Urban Land-----	35	Not rated		Not rated		Not rated	
248B: Georgia-----	40	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
250B: Sutton-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
253B: Wapping-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
255B: Watchaug-----	40	Not limited		Not limited		Somewhat limited Large stones content	0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
260B: Charlton-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
260C: Charlton-----	40	Not limited		Not limited		Somewhat limited Slope	0.63
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
260D: Charlton-----	40	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00
Urban Land-----	35	Not rated		Not rated		Not rated	
263B: Cheshire-----	40	Not limited		Not limited		Somewhat limited Large stones content	0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
263C: Cheshire-----	40	Not limited		Not limited		Somewhat limited Slope Large stones content	0.63 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
266B: Narragansett-----	40	Not limited		Not limited		Somewhat limited Large stones content	0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
269B: Yalesville-----	40	Not limited		Not limited		Somewhat limited Depth to bedrock	0.05
Urban Land-----	35	Not rated		Not rated		Not rated	
269C: Yalesville-----	40	Not limited		Not limited		Somewhat limited Slope Depth to bedrock	0.63 0.05
Urban Land-----	35	Not rated		Not rated		Not rated	
273C: Urban Land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Not limited		Not limited		Somewhat limited Slope	0.04
Chatfield-----	15	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to bedrock Gravel content Slope	0.54 0.25 0.04

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
273E: Urban Land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Slope	1.00
Chatfield-----	15	Very limited Slope	1.00	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
		Large stones content	0.53	Slope	0.22	Depth to bedrock	0.54
						Gravel content	0.25
275C: Urban Land-----	45	Not rated		Not rated		Not rated	
Chatfield-----	30	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to bedrock	0.54
						Gravel content	0.25
						Slope	0.04
275E: Urban Land-----	35	Not rated		Not rated		Not rated	
Chatfield-----	25	Very limited Slope	1.00	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
		Large stones content	0.53	Slope	0.22	Depth to bedrock	0.54
						Gravel content	0.25
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
282B: Broadbrook-----	40	Not limited		Not limited		Somewhat limited Depth to pan	0.84
Urban Land-----	35	Not rated		Not rated		Not rated	
284B: Paxton-----	40	Not limited		Not limited		Somewhat limited Depth to pan	0.79
						Large stones content	0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
284C: Paxton-----	40	Not limited		Not limited		Somewhat limited Depth to pan	0.79
						Slope	0.63
						Large stones content	0.01

Table 14.—Recreation, Part 2—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
284C: Urban Land-----	35	Not rated		Not rated		Not rated	
284D: Paxton-----	40	Somewhat limited Slope	0.50	Not limited		Very limited Slope Depth to pan Large stones content	1.00 0.79 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
287B: Wethersfield-----	40	Not limited		Not limited		Somewhat limited Depth to pan Large stones content	0.71 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
287C: Wethersfield-----	40	Not limited		Not limited		Somewhat limited Depth to pan Slope Large stones content	0.71 0.63 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
287D: Wethersfield-----	40	Somewhat limited Slope	0.50	Not limited		Very limited Slope Depth to pan Large stones content	1.00 0.71 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
290B: Stockbridge-----	40	Not limited		Not limited		Not limited	
Urban Land-----	35	Not rated		Not rated		Not rated	
290C: Stockbridge-----	40	Not limited		Not limited		Somewhat limited Slope	0.63
Urban Land-----	35	Not rated		Not rated		Not rated	
290D: Stockbridge-----	40	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00
Urban Land-----	35	Not rated		Not rated		Not rated	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
301: Beaches-----	50	Not rated		Not rated		Not rated	
Udipsamments-----	35	Very limited Too sandy	1.00	Very limited Too sandy	1.00	Very limited Droughty Flooding Too sandy Slope	1.00 0.60 0.50 0.01
302: Dumps-----	95	Not rated		Not rated		Not rated	
303: Pits, Quarries-----	90	Not rated		Not rated		Not rated	
304: Udorthents-----	90	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Large stones content	1.00 0.01
305: Udorthents-----	65	Somewhat limited Slope	0.18	Not limited		Very limited Slope Large stones content	1.00 0.01
Pits-----	25	Not rated		Not rated		Not rated	
306: Udorthents-----	50	Somewhat limited Slope	0.18	Not limited		Very limited Slope Large stones content	1.00 0.01
Urban Land-----	35	Not rated		Not rated		Not rated	
307: Urban Land-----	80	Not rated		Not rated		Not rated	
308: Udorthents-----	80	Somewhat limited Slope	0.18	Not limited		Very limited Slope Large stones content	1.00 0.01
309: Udorthents-----	80	Somewhat limited Slope	0.18	Not limited		Very limited Slope Large stones content	1.00 0.01

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
310: Udorthents, Periodically Flooded-----	85	Somewhat limited Flooding Slope	0.40 0.18	Somewhat limited Flooding	0.40	Very limited Flooding Slope Large stones content	1.00 1.00 0.01
401C: Macomber-----	55	Somewhat limited Large stones content	0.19	Somewhat limited Large stones content	0.19	Somewhat limited Gravel content Slope Depth to bedrock Droughty	0.82 0.63 0.46 0.01
Taconic-----	30	Not limited		Not limited		Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.39 0.01
402D: Macomber-----	50	Somewhat limited Slope	0.50	Not limited		Very limited Slope Gravel content Depth to bedrock Droughty	1.00 0.82 0.46 0.01
Taconic-----	25	Somewhat limited Slope	0.50	Not limited		Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.39
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
403C: Taconic-----	70	Not limited		Not limited		Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.39 0.01
Rock Outcrop-----	25	Not rated		Not rated		Not rated	
403E: Taconic-----	70	Very limited Slope	1.00	Somewhat limited Slope	0.78	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.39
Rock Outcrop-----	20	Not rated		Not rated		Not rated	

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
403F: Taconic-----	70	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.39
Rock Outcrop-----	20	Not rated		Not rated		Not rated	
405C: Dummerston-----	85	Somewhat limited Large stones content	0.19	Somewhat limited Large stones content	0.19	Somewhat limited Gravel content Slope	0.09 0.01
405E: Dummerston-----	85	Very limited Slope Large stones content	1.00 0.19	Somewhat limited Large stones content	0.19	Very limited Slope Gravel content	1.00 0.09
407C: Lanesboro-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan Slope	0.46 0.16
407E: Lanesboro-----	85	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content Slope	0.53 0.22	Very limited Slope Depth to pan	1.00 0.46
408C: Fullam-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Very limited Depth to pan Slope Droughty	1.00 0.16 0.05
409B: Brayton-----	85	Very limited Depth to saturated zone Large stones content	1.00 0.53	Very limited Depth to saturated zone Large stones content	1.00 0.53	Very limited Depth to pan Depth to saturated zone	1.00 1.00
412B: Bice-----	85	Not limited		Not limited		Not limited	
412C: Bice-----	85	Not limited		Not limited		Somewhat limited Slope	0.16
412D: Bice-----	85	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
413C: Bice-----	45	Not limited		Not limited		Somewhat limited Slope	0.01
Millsite-----	40	Not limited		Not limited		Somewhat limited Depth to bedrock Slope	0.35 0.01
413E: Bice-----	45	Very limited Slope	1.00	Not limited		Very limited Slope	1.00
Millsite-----	40	Very limited Slope	1.00	Not limited		Very limited Slope Depth to bedrock	1.00 0.35
414: Fredon, cold-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
415C: Millsite-----	40	Not limited		Not limited		Somewhat limited Depth to bedrock Slope	0.35 0.01
Westminster-----	40	Not limited		Not limited		Very limited Depth to bedrock Droughty Slope	1.00 0.91 0.01
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
415E: Millsite-----	40	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Slope Depth to bedrock	1.00 0.35
Westminster-----	40	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.91
Rock Outcrop-----	15	Not rated		Not rated		Not rated	
416E: Rock Outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Slope	1.00	Not limited		Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.91

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
416F: Rock Outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.91
417B: Bice-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
417C: Bice-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.16
417D: Bice-----	85	Somewhat limited Large stones content Slope	0.53 0.50	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
418C: Schroon-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.04
420A: Schroon-----	85	Not limited		Not limited		Not limited	
420B: Schroon-----	85	Not limited		Not limited		Not limited	
421A: Ninigret, cold-----	85	Not limited		Not limited		Not limited	
423A: Sudbury, cold-----	85	Not limited		Not limited		Not limited	
424B: Shelburne-----	85	Not limited		Not limited		Somewhat limited Depth to pan	0.74
424C: Shelburne-----	85	Not limited		Not limited		Somewhat limited Depth to pan Slope	0.74 0.63
424D: Shelburne-----	85	Somewhat limited Slope	0.50	Not limited		Very limited Slope Depth to pan	1.00 0.74
425B: Shelburne-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan	0.74

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
425C: Shelburne-----	85	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Depth to pan Slope	0.74 0.63
426D: Shelburne-----	85	Very limited Large stones content Slope	1.00 1.00	Very limited Large stones content	1.00	Very limited Slope Depth to pan	1.00 0.74
427B: Ashfield-----	85	Somewhat limited Large stones content Depth to saturated zone	0.53 0.44	Somewhat limited Large stones content Depth to saturated zone	0.53 0.44	Somewhat limited Depth to pan Depth to saturated zone	0.54 0.08
427C: Ashfield-----	85	Somewhat limited Large stones content Depth to saturated zone	0.53 0.44	Somewhat limited Large stones content Depth to saturated zone	0.53 0.44	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.54 0.08
428A: Ashfield-----	85	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to pan Depth to saturated zone	0.54 0.08
428B: Ashfield-----	85	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to pan Depth to saturated zone	0.54 0.08
428C: Ashfield-----	85	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.54 0.08
429A: Agawam, cold-----	80	Not limited		Not limited		Not limited	
429B: Agawam, cold-----	80	Not limited		Not limited		Not limited	
429C: Agawam, cold-----	80	Not limited		Not limited		Somewhat limited Slope	0.63

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
433: Moosilauke-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
434A: Merrimac, cold-----	80	Not limited		Not limited		Not limited	
434B: Merrimac, cold-----	80	Not limited		Not limited		Not limited	
434C: Merrimac, cold-----	80	Not limited		Not limited		Somewhat limited Slope	0.63
435: Scarboro-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Organic matter content Depth to saturated zone Ponding	1.00 1.00 1.00
436: Halsey-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
437: Wonsqueak-----	85	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
438: Bucksport-----	85	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
440A: Boscawen-----	80	Not limited		Not limited		Very limited Droughty	1.00
440C: Boscawen-----	80	Not limited		Not limited		Very limited Droughty Slope	1.00 0.04
440E: Boscawen-----	85	Very limited Slope	1.00	Somewhat limited Slope	0.22	Very limited Slope Droughty	1.00 1.00
442: Brayton-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Depth to pan	1.00 0.71

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
443: Brayton-----	50	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Depth to pan	1.00 0.71
Loonmeadow-----	35	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
448B: Hogansburg-----	85	Not limited		Not limited		Not limited	
449B: Hogansburg-----	85	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Not limited	
449C: Hogansburg-----	85	Very limited Large stones content	1.00	Very limited Large stones content	1.00	Somewhat limited Slope	0.63
450B: Pyrities-----	80	Not limited		Not limited		Not limited	
450C: Pyrities-----	80	Not limited		Not limited		Somewhat limited Slope	0.63
450D: Pyrities-----	80	Somewhat limited Slope	0.50	Not limited		Very limited Slope	1.00
451B: Pyrities-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Not limited	
451C: Pyrities-----	80	Somewhat limited Large stones content	0.53	Somewhat limited Large stones content	0.53	Somewhat limited Slope	0.63
451D: Pyrities-----	80	Very limited Slope Large stones content	1.00 0.53	Somewhat limited Large stones content	0.53	Very limited Slope	1.00
457: Mudgepond-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00

Table 14.—Recreation (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Paths and trails		Off-road motorcycle trails		Golf fairways	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
458: Mudgepond-----	55	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 1.00	Very limited Depth to saturated zone	1.00
Alden-----	35	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Large stones content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Large stones content	1.00 1.00 0.01
501: Ondawa-----	85	Very limited Gravel content	1.00	Very limited Gravel content	1.00	Somewhat limited Flooding	0.60
503: Rumney-----	80	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
508: Medomak-----	85	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.40	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00

Table 15.--Wildlife Habitat

(See text for definitions of terms used in this table. Absence of an entry indicates that no rating is applicable.)

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
2: Ridgebury-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
3: Ridgebury-----	Very poor	Very poor	Fair	Fair	Fair	Good	Fair	Poor	Fair	Fair
Leicester-----	Very poor	Very poor	Fair	Fair	Fair	Good	Fair	Poor	Fair	Fair
Whitman-----	Very poor	Very poor	Poor	Poor	Poor	Good	Good	Very poor	Poor	Good
4: Leicester-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
5: Wilbraham-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
6: Wilbraham-----	Very poor	Very poor	Fair	Fair	Fair	Good	Fair	Poor	Fair	Fair
Menlo-----	Very poor	Very poor	Poor	Poor	Poor	Good	Good	Very poor	Poor	Good
7: Mudgepond-----	Fair	Fair	Good	Good	Good	Good	Fair	Good	Good	Fair
8: Mudgepond-----	Very poor	Very poor	Good	Good	Good	Good	Fair	Very poor	Good	Fair
Alden-----	Very poor	Very poor	Poor	Poor	Poor	Good	Good	Very poor	Poor	Good
9: Scitico-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
Shaker-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
Maybid-----	Very poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
10: Raynham-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
12: Raypol-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
13: Walpole-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
14: Fredon-----	Poor	Fair	Fair	Fair	Fair	Good	Fair	Fair	Fair	Fair
15: Scarboro-----	Very poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
16: Halsey-----	Very poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
17: Timakwa-----	Poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
Natchaug-----	Poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
18: Catden-----	Poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
Freetown-----	Poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
20A: Ellington-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
21A: Ninigret-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
Tisbury-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
22A: Hero-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
22B: Hero-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
23A: Sudbury-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
24A: Deerfield-----	Poor	Fair	Fair	Poor	Poor	Poor	Poor	Fair	Poor	Poor
25A: Brancroft-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
25B: Brancroft-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
25C: Brancroft-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
26A: Berlin-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
26B: Berlin-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
27A: Belgrade-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
28A: Elmridge-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
28B: Elmridge-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
29A: Agawam-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
29B: Agawam-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
29C: Agawam-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
30A: Branford-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
30B: Branford-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
30C: Branford-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
31A: Copake-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
31B: Copake-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
31C: Copake-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
32A: Haven-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Enfield-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
32B: Haven-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Enfield-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
32C: Haven-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Enfield-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
33A: Hartford-----	Good	Good	Good	Good	Good	Very poor	Very poor	Fair	Fair	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
33B: Hartford-----	Good	Good	Good	Good	Good	Very poor	Very poor	Fair	Fair	Very poor
34A: Merrimac-----	Good	Good	Good	Good	Good	Very poor	Very poor	Fair	Fair	Very poor
34B: Merrimac-----	Good	Good	Good	Good	Good	Very poor	Very poor	Fair	Fair	Very poor
34C: Merrimac-----	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor
35A: Penwood-----	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor
35B: Penwood-----	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor
36A: Windsor-----	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor
36B: Windsor-----	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor
36C: Windsor-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Fair	Poor	Very poor
37A: Manchester-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
37C: Manchester-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
37E: Manchester-----	Very poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
38A: Hinckley-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
38C: Hinckley-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
38E: Hinckley-----	Very poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
39A: Groton-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
39C: Groton-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
39E: Groton-----	Very poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
40A: Ludlow-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
40B: Ludlow-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
41B: Ludlow-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
42C: Ludlow-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
43A: Rainbow-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
43B: Rainbow-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
44B: Rainbow-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
45A: Woodbridge-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
45B: Woodbridge-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
45C: Woodbridge-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
46B: Woodbridge-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
46C: Woodbridge-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
47C: Woodbridge-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
48B: Georgia-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Amenia-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
48C: Georgia-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Amenia-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
49B: Georgia-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
Amenia-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
49C: Georgia-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
Amenia-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
50A: Sutton-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
50B: Sutton-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
51B: Sutton-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
52C: Sutton-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
53A: Wapping-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
53B: Wapping-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
54B: Wapping-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
55A: Watchaug-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
55B: Watchaug-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
56B: Watchaug-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
57B: Gloucester-----	Fair	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
57C: Gloucester-----	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
57D: Gloucester-----	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
58B: Gloucester-----	Very poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
58C: Gloucester-----	Very poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Poor	Very poor
59C: Gloucester-----	Very poor	Very poor	Fair	Fair	Fair	Very poor	Very poor	Very poor	Poor	Very poor
59D: Gloucester-----	Very poor	Very poor	Poor	Fair	Fair	Very poor	Very poor	Very poor	Poor	Very poor
60B: Canton-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Charlton-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
60C: Canton-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Charlton-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
60D: Canton-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
Charlton-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
61B: Canton-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
Charlton-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
61C: Canton-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
Charlton-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
62C: Canton-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
62C: Charlton-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
62D: Canton-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
Charlton-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
63B: Cheshire-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
63C: Cheshire-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
63D: Cheshire-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
64B: Cheshire-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
64C: Cheshire-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
65C: Cheshire-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
65D: Cheshire-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
66B: Narragansett-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
66C: Narragansett-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
67B: Narragansett-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
67C: Narragansett-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
68C: Narragansett-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
68D: Narragansett-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
75E: Hollis-----	Very poor	Very poor	Fair	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor
Chatfield-----	Very poor	Very poor	Good	Fair	Fair	Very poor	Very poor	Very poor	Fair	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
76E: Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
Hollis-----	Very poor	Very poor	Fair	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor
76F: Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
Hollis-----	Very poor	Very poor	Fair	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor
77C: Cheshire-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
Holyoke-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
77D: Cheshire-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
Holyoke-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
78C: Holyoke-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
78E: Holyoke-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
79E: Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
Holyoke-----	Very poor	Very poor	Fair	Poor	Poor	Very poor	Very poor	Very poor	Poor	Very poor
80B: Bernardston-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
80C: Bernardston-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
81C: Bernardston-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
81D: Bernardston-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
82B: Broadbrook-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
82C: Broadbrook-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
82D: Broadbrook-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
83B: Broadbrook-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
83C: Broadbrook-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
84B: Paxton-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Montauk-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
84C: Paxton-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Montauk-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
84D: Paxton-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
Montauk-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
85B: Paxton-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
Montauk-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
85C: Paxton-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
Montauk-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
86C: Paxton-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
Montauk-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
86D: Paxton-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
Montauk-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
87B: Wethersfield-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
87C: Wethersfield-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
87D: Wethersfield-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
88B: Wethersfield-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
88C: Wethersfield-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
89C: Wethersfield-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
89D: Wethersfield-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
90B: Stockbridge-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
90C: Stockbridge-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
90D: Stockbridge-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
91B: Stockbridge-----	Very poor	Poor	Good	Good	Good	Poor	Very poor	Poor	Good	Very poor
91C: Stockbridge-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
91D: Stockbridge-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
92B: Nellis-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
92C: Nellis-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
92D: Nellis-----	Fair	Fair	Fair	Good	Good	Very poor	Very poor	Fair	Good	Very poor
93C: Nellis-----	Very poor	Fair	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
94C: Farmington-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Nellis-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
94E: Farmington-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Nellis-----	Very poor	Fair	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
95C: Farmington-----	Poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
95E: Farmington-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
96: Ipswich-----	Very poor	Very poor	Very poor	Very poor	Very poor	Good	Good	Very poor	Very poor	Good
97: Pawcatuck-----	Very poor	Very poor	Very poor	Very poor	Very poor	Good	Good	Very poor	Very poor	Good
98: Westbrook-----	Very poor	Very poor	Very poor	Very poor	Very poor	Good	Good	Very poor	Very poor	Good
99: Westbrook, low salt-----	Very poor	Very poor	Very poor	Very poor	Very poor	Good	Good	Very poor	Very poor	Good

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
255B: Watchaug-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
260B: Charlton-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
260C: Charlton-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
260D: Charlton-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
263B: Cheshire-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
263C: Cheshire-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
266B: Narragansett-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
269B: Yalesville-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
269C: Yalesville-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
273C: Urban Land-----	---	---	---	---	---	---	---	---	---	---
Charlton-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
Chatfield-----	Very poor	Poor	Good	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
287D: Wethersfield-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
290B: Stockbridge-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
290C: Stockbridge-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
290D: Stockbridge-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
301: Beaches-----	---	---	---	---	---	---	---	---	---	---
Udipsammments-----	Very poor	Very poor	Poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor
302: Dumps-----	---	---	---	---	---	---	---	---	---	---
303: Pits, Quarries-----	---	---	---	---	---	---	---	---	---	---
304: Udorthents-----	Very poor	Very poor	Good	Good	Good	Very poor	Very poor	Poor	Fair	Very poor
305: Udorthents-----	Poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Fair	Very poor
Pits-----	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor	Very poor
306: Udorthents-----	Poor	Poor	Good	Good	Good	Very poor	Very poor	Fair	Fair	Very poor
Urban Land-----	---	---	---	---	---	---	---	---	---	---
307: Urban Land-----	---	---	---	---	---	---	---	---	---	---
308: Udorthents-----	Poor	Poor	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
309: Udorthents-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
310: Udorthents, Periodically Flooded-----	Poor	Poor	Fair	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor
401C: Macomber-----	Poor	Good	Good	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor
Taconic-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
402D: Macomber-----	Very poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Good	Fair	Very poor
Taconic-----	Very poor	Fair	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
403C: Taconic-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
403E: Taconic-----	Very poor	Very poor	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
403F: Taconic-----	Very poor	Very poor	Very poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
405C: Dummerston-----	Poor	Poor	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
405E: Dummerston-----	Very poor	Very poor	Poor	Good	Good	Very poor	Very poor	Good	Good	Very poor
407C: Lanesboro-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
407E: Lanesboro-----	Very poor	Poor	Poor	Good	Good	Very poor	Very poor	Poor	Good	Very poor
408C: Fullam-----	Poor	Poor	Fair	Good	Good	Very poor	Very poor	Good	Fair	Fair
409B: Brayton-----	Very poor	Poor	Fair	Fair	Fair	Poor	Very poor	Poor	Fair	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
412B: Bice-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
412C: Bice-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
412D: Bice-----	Poor	Fair	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
413C: Bice-----	Poor	Fair	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
Millsite-----	Very poor	Poor	Good	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor
413E: Bice-----	Very poor	Poor	Fair	Good	Good	Poor	Very poor	Good	Good	Very poor
Millsite-----	Very poor	Poor	Poor	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor
414: Fredon, cold-----	Poor	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good
415C: Millsite-----	Very poor	Poor	Good	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor
Westminster-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
415E: Millsite-----	Very poor	Poor	Good	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor
Westminster-----	Very poor	Poor	Fair	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
416E: Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
Westminster-----	Very poor	Poor	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
416F: Rock Outcrop-----	---	---	---	---	---	---	---	---	---	---
Westminster-----	Very poor	Very poor	Poor	Poor	Poor	Very poor	Very poor	Poor	Poor	Very poor
417B: Bice-----	Fair	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
417C: Bice-----	Fair	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
417D: Bice-----	Poor	Poor	Fair	Good	Good	Poor	Very poor	Good	Good	Very poor
418C: Schroon-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
420A: Schroon-----	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
420B: Schroon-----	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
421A: Ninigret, cold-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
423A: Sudbury, cold-----	Fair	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor
424B: Shelburne-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
424C: Shelburne-----	Fair	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
424D: Shelburne-----	Poor	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
425B: Shelburne-----	Fair	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
425C: Shelburne-----	Fair	Fair	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
426D: Shelburne-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
427B: Ashfield-----	Fair	Fair	Good	Good	Good	Poor	Very poor	Fair	Good	Poor
427C: Ashfield-----	Fair	Fair	Good	Good	Good	Poor	Very poor	Fair	Good	Poor
428A: Ashfield-----	Good	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Poor

Table 15.--Wildlife Habitat--Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
428B: Ashfield-----	Good	Good	Good	Good	Good	Poor	Very poor	Fair	Good	Poor
428C: Ashfield-----	Fair	Fair	Good	Good	Good	Poor	Very poor	Fair	Good	Poor
429A: Agawam, cold-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
429B: Agawam, cold-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Very poor
429C: Agawam, cold-----	Fair	Fair	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
433: Moosilauke-----	Poor	Poor	Fair	Fair	Fair	Good	Fair	Fair	Fair	Good
434A: Merrimac, cold-----	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
434B: Merrimac, cold-----	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
434C: Merrimac, cold-----	Fair	Fair	Fair	Good	Good	Very poor	Very poor	Good	Good	Very poor
435: Scarboro-----	Very poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
436: Halsey-----	Very poor	Poor	Poor	Poor	Poor	Good	Good	Poor	Poor	Good
437: Wonsqueak-----	Very poor	Very poor	Poor	Very poor	Poor	Good	Good	Poor	Poor	Good
438: Bucksport-----	Very poor	Poor	Poor	Very poor	Poor	Good	Good	Poor	Poor	Good
440A: Boscawen-----	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor
440C: Boscawen-----	Poor	Fair	Fair	Fair	Fair	Very poor	Very poor	Fair	Fair	Very poor
440E: Boscawen-----	Poor	Poor	Poor	Fair	Fair	Very poor	Very poor	Poor	Fair	Very poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
442: Brayton-----	Very poor	Poor	Fair	Fair	Fair	Good	Poor	Poor	Fair	Good
443: Brayton-----	Very poor	Poor	Fair	Poor	Fair	Good	Fair	Poor	Fair	Good
Loonmeadow-----	Very poor	Very poor	Poor	Poor	Fair	Good	Good	Poor	Fair	Good
448B: Hogansburg-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Poor
449B: Hogansburg-----	Poor	Poor	Good	Good	Good	Poor	Very poor	Fair	Good	Poor
449C: Hogansburg-----	Poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
450B: Pyrities-----	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
450C: Pyrities-----	Good	Good	Good	Good	Good	Very poor	Very poor	Good	Good	Very poor
450D: Pyrities-----	Poor	Fair	Good	Good	Good	Very poor	Very poor	Fair	Good	Very poor
451B: Pyrities-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
451C: Pyrities-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
451D: Pyrities-----	Very poor	Poor	Good	Good	Good	Very poor	Very poor	Poor	Good	Very poor
457: Mudgepond-----	Poor	Fair	Good	Fair	Fair	Good	Fair	Fair	Good	Good
458: Mudgepond-----	Poor	Fair	Good	Fair	Fair	Good	Fair	Very poor	Good	Good
Alden-----	Very poor	Very poor	Poor	Poor	Poor	Good	Fair	Very poor	Poor	Good
501: Ondawa-----	Good	Good	Good	Good	Good	Poor	Very poor	Good	Good	Poor

Table 15.—Wildlife Habitat—Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hard- wood trees	Conif- erous plants	Wetland plants	Shallow water areas	Open- land wild- life	Wood- land wild- life	Wetland wild- life
503: Rumney-----	Poor	Fair	Fair	Poor	Poor	Good	Fair	Fair	Fair	Good
508: Medomak-----	Very poor	Poor	Poor	Poor	Poor	Good	Fair	Poor	Poor	Fair

Table 16.—Connecticut Inland Wetlands

The state of Connecticut defines inland wetlands based on soils. The Connecticut Inland Wetlands and Watercourses Act defines wetland soils to include "any of the soil types designated as poorly drained, very poorly drained, alluvial, and floodplain by the National Cooperative Soil Survey, as may be amended from time to time, of the Natural Resources Conservation Service of the United States Department of Agriculture".

Map units may be dominated by Connecticut inland wetland soils, but have minor components of non-wetland soils. Non-wetland map units may contain minor components of Connecticut inland wetland soils. On site investigation is necessary to determine the presence or absence of wetland soils in a particular area.

The following map units meet the definition of Connecticut Inland Wetland Soils:

Map symbol	Map unit name
2	Ridgebury fine sandy loam
3	Ridgebury, Leicester, and Whitman soils, extremely stony
4	Leicester fine sandy loam
5	Wilbraham silt loam
6	Wilbraham and Menlo soils, extremely stony
7	Mudgepond silt loam
8	Mudgepond and Alden soils, extremely stony
9	Scitico, Shaker, and Maybid soils
10	Raynham silt loam
12	Raypol silt loam
13	Walpole sandy loam
14	Fredon silt loam
15	Scarboro muck
16	Halsey silt loam
17	Timakwa and Natchaug soils
18	Catden and Freetown soils
96	Ipswich mucky peat
97	Pawcatuck mucky peat
98	Westbrook mucky peat
99	Westbrook mucky peat, low salt
100	Suncook loamy fine sand
101	Occum fine sandy loam
102	Pootatuck fine sandy loam
103	Rippowam fine sandy loam
104	Bash silt loam
105	Hadley silt loam
106	Winooski silt loam
107	Limerick and Lim soils
108	Saco silt loam
109	Fluvaquents-Udifluents complex, frequently flooded
409B	Brayton mucky silt loam, 0 to 8 percent slopes, very stony
414	Fredon silt loam, cold
433	Moosilauke sandy loam
435	Scarboro muck, cold
436	Halsey silt loam, cold
437	Wonsqueak mucky peat
438	Bucksport muck
442	Brayton loam
443	Brayton-Loonmeadow complex, extremely stony
457	Mudgepond silt loam, cold
458	Mudgepond and Alden soils, extremely stony, cold
501	Ondawa fine sandy loam
503	Rumney fine sandy loam
508	Medomak silt loam

Table 17.—Building Site Development (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
3: Ridgebury-----	40	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Leicester-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Whitman-----	15	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
4: Leicester-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
5: Wilbraham-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
6: Wilbraham-----	60	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Menlo-----	25	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
7: Mudgepond-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
8: Mudgepond-----	45	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Alden-----	35	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
9: Scitico-----	40	Very limited Depth to saturated zone Shrink-swell	1.00 0.99	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 0.99
Shaker-----	30	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Depth to saturated zone	1.00
Maybid-----	15	Very limited Depth to saturated zone Ponding Shrink-swell	1.00 1.00 0.99	Very limited Depth to saturated zone Shrink-swell Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Shrink-swell	1.00 1.00 0.99
10: Raynham-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
12: Raypol-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
13: Walpole-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
14: Fredon-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
15: Scarboro-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
16: Halsey-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
17: Timakwa-----	45	Very limited Flooding Depth to saturated zone Organic matter content	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Subsidence	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Organic matter content	1.00 1.00 1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
17: Natchaug-----	40	Very limited Flooding Depth to saturated zone Subsidence	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Subsidence	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Subsidence	1.00 1.00 1.00
18: Catden-----	40	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00
Freetown-----	40	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00
20A: Ellington-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
21A: Ninigret-----	60	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Tisbury-----	25	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
22A: Hero-----	85	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
22B: Hero-----	85	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
23A: Sudbury-----	80	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
24A: Deerfield-----	80	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
25A: Brancroft-----	80	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
25B: Brancroft-----	80	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope Shrink-swell	0.81 0.50 0.01
25C: Brancroft-----	80	Somewhat limited Depth to saturated zone Slope Shrink-swell	0.81 0.63 0.01	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone Shrink-swell	1.00 0.81 0.01
26A: Berlin-----	80	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01
26B: Berlin-----	80	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope Shrink-swell	0.81 0.50 0.01
27A: Belgrade-----	80	Not limited		Very limited Depth to saturated zone	0.99	Not limited	
28A: Elmridge-----	80	Very limited Shrink-swell Depth to saturated zone	0.99 0.39	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Shrink-swell Depth to saturated zone	0.99 0.39
28B: Elmridge-----	80	Very limited Shrink-swell Depth to saturated zone	0.99 0.39	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Shrink-swell Slope Depth to saturated zone	0.99 0.50 0.39
29A: Agawam-----	80	Not limited		Not limited		Not limited	
29B: Agawam-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
29C: Agawam-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
30A: Branford-----	80	Not limited		Not limited		Not limited	
30B: Branford-----	80	Not limited		Not limited		Somewhat limited Slope	0.03
30C: Branford-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
31A: Copake-----	85	Not limited		Not limited		Not limited	
31B: Copake-----	85	Not limited		Not limited		Somewhat limited Slope	0.50
31C: Copake-----	85	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
32A: Haven-----	60	Not limited		Not limited		Not limited	
Enfield-----	25	Not limited		Not limited		Not limited	
32B: Haven-----	60	Not limited		Not limited		Somewhat limited Slope	0.50
Enfield-----	25	Not limited		Not limited		Somewhat limited Slope	0.50
32C: Haven-----	60	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Enfield-----	25	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
33A: Hartford-----	80	Not limited		Not limited		Not limited	
33B: Hartford-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
34A: Merrimac-----	80	Not limited		Not limited		Not limited	
34B: Merrimac-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
34C: Merrimac-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
35A: Penwood-----	80	Not limited		Not limited		Not limited	
35B: Penwood-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
36A: Windsor-----	80	Not limited		Not limited		Not limited	
36B: Windsor-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
36C: Windsor-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
37A: Manchester-----	80	Not limited		Not limited		Not limited	
37C: Manchester-----	80	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
37E: Manchester-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
38A: Hinckley-----	80	Not limited		Not limited		Not limited	
38C: Hinckley-----	80	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
38E: Hinckley-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
39A: Groton-----	85	Not limited		Not limited		Not limited	
39C: Groton-----	85	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
39E: Groton-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
40A: Ludlow-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
40B: Ludlow-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
41B: Ludlow-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.39 0.12
42C: Ludlow-----	80	Somewhat limited Depth to saturated zone Slope	0.39 0.04	Very limited Depth to saturated zone Slope	1.00 0.04	Very limited Slope Depth to saturated zone	1.00 0.39
43A: Rainbow-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
43B: Rainbow-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
44B: Rainbow-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
45A: Woodbridge-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
45B: Woodbridge-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
45C: Woodbridge-----	80	Somewhat limited Slope Depth to saturated zone	0.63 0.39	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.39
46B: Woodbridge-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.39 0.12

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
46C: Woodbridge-----	80	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.39	Slope	0.63	Depth to saturated zone	0.39
47C: Woodbridge-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.04	Slope	0.04	Depth to saturated zone	0.39
48B: Georgia-----	50	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.12
						Depth to saturated zone	0.07
Amenia-----	35	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.12
						Depth to saturated zone	0.07
48C: Georgia-----	50	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.07	Slope	0.63	Depth to saturated zone	0.07
Amenia-----	35	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.07	Slope	0.63	Depth to saturated zone	0.07
49B: Georgia-----	50	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.50
						Depth to saturated zone	0.07
Amenia-----	35	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.50
						Depth to saturated zone	0.07
49C: Georgia-----	50	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.07	Slope	0.63	Depth to saturated zone	0.07

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
49C: Amenia-----	35	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.07	Slope	0.63	Depth to saturated zone	0.07
50A: Sutton-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
50B: Sutton-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.50
						Depth to saturated zone	0.39
51B: Sutton-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
						Slope	0.12
52C: Sutton-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.04	Slope	0.04	Depth to saturated zone	0.39
53A: Wapping-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
53B: Wapping-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.50
						Depth to saturated zone	0.39
54B: Wapping-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
						Slope	0.12
55A: Watchaug-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
55B: Watchaug-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.50
						Depth to saturated zone	0.39

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
56B: Watchaug-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.39 0.12
57B: Gloucester-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
57C: Gloucester-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
57D: Gloucester-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
58B: Gloucester-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
58C: Gloucester-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
59C: Gloucester-----	80	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
59D: Gloucester-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
60B: Canton-----	45	Not limited		Not limited		Somewhat limited Slope	0.50
Charlton-----	35	Not limited		Not limited		Somewhat limited Slope	0.50
60C: Canton-----	45	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Charlton-----	35	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
60D: Canton-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Charlton-----	35	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
61B: Canton-----	45	Not limited		Not limited		Somewhat limited Slope	0.50
Charlton-----	35	Not limited		Not limited		Somewhat limited Slope	0.50

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
61C: Canton-----	45	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Charlton-----	35	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
62C: Canton-----	45	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Charlton-----	35	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
62D: Canton-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Charlton-----	35	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
63B: Cheshire-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
63C: Cheshire-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
63D: Cheshire-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
64B: Cheshire-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
64C: Cheshire-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
65C: Cheshire-----	80	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
65D: Cheshire-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
66B: Narragansett-----	80	Not limited		Not limited		Somewhat limited Slope	0.12
66C: Narragansett-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
67B: Narragansett-----	80	Not limited		Not limited		Somewhat limited Slope	0.50

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
67C: Narragansett-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
68C: Narragansett-----	80	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
68D: Narragansett-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
69B: Yalesville-----	75	Somewhat limited Depth to hard bedrock	0.05	Very limited Depth to hard bedrock	1.00	Somewhat limited Slope Depth to hard bedrock	0.50 0.05
69C: Yalesville-----	75	Somewhat limited Slope Depth to hard bedrock	0.63 0.05	Very limited Depth to hard bedrock Slope	1.00 0.63	Very limited Slope Depth to hard bedrock	1.00 0.05
70C: Branford-----	50	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Holyoke-----	30	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
71C: Brookfield-----	45	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Brimfield-----	30	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	
71E: Brookfield-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Brimfield-----	30	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
73C:							
Charlton-----	45	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Chatfield-----	30	Somewhat limited Depth to hard bedrock Slope	0.54 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Slope Depth to hard bedrock	1.00 0.54
73E:							
Charlton-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Chatfield-----	30	Very limited Slope Depth to hard bedrock	1.00 0.54	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 0.54
74C:							
Narragansett-----	55	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Hollis-----	20	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
75C:							
Hollis-----	35	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
Chatfield-----	30	Somewhat limited Depth to hard bedrock Slope	0.54 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Slope Depth to hard bedrock	1.00 0.54
Rock outcrop-----	15	Not rated		Not rated		Not rated	
75E:							
Hollis-----	35	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Chatfield-----	30	Very limited Slope Depth to hard bedrock	1.00 0.54	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 0.54
Rock outcrop-----	15	Not rated		Not rated		Not rated	
76E:							
Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
76F: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
77C: Cheshire-----	45	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Holyoke-----	35	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
77D: Cheshire-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Holyoke-----	35	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
78C: Holyoke-----	50	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
Rock outcrop-----	25	Not rated		Not rated		Not rated	
78E: Holyoke-----	50	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Rock outcrop-----	25	Not rated		Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Holyoke-----	25	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00
80B: Bernardston-----	80	Somewhat limited Depth to saturated zone	0.77	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.77 0.50
80C: Bernardston-----	80	Somewhat limited Depth to saturated zone Slope	0.77 0.63	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.77

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
81C: Bernardston-----	80	Somewhat limited Depth to saturated zone Slope	0.77 0.04	Very limited Depth to saturated zone Slope	1.00 0.04	Very limited Slope Depth to saturated zone	1.00 0.77
81D: Bernardston-----	80	Very limited Slope Depth to saturated zone	1.00 0.77	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.77
82B: Broadbrook-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.39 0.12
82C: Broadbrook-----	80	Somewhat limited Slope Depth to saturated zone	0.63 0.39	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.39
82D: Broadbrook-----	80	Very limited Slope Depth to saturated zone	1.00 0.39	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.39
83B: Broadbrook-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
83C: Broadbrook-----	80	Somewhat limited Slope Depth to saturated zone	0.63 0.39	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.39
84B: Paxton-----	55	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
Montauk-----	30	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.07

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
84C: Paxton-----	55	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.39	Slope	0.63	Depth to saturated zone	0.39
Montauk-----	30	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.07	Slope	0.63	Depth to saturated zone	0.07
84D: Paxton-----	55	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.39	Depth to saturated zone	1.00	Depth to saturated zone	0.39
Montauk-----	30	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.07	Depth to saturated zone	1.00	Depth to saturated zone	0.07
85B: Paxton-----	55	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.50
						Depth to saturated zone	0.39
Montauk-----	30	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.50
						Depth to saturated zone	0.07
85C: Paxton-----	55	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.39	Slope	0.63	Depth to saturated zone	0.39
Montauk-----	30	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.07	Slope	0.63	Depth to saturated zone	0.07
86C: Paxton-----	55	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.04	Slope	0.04	Depth to saturated zone	0.39

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
86C: Montauk-----	30	Somewhat limited Depth to saturated zone Slope	0.07 0.04	Very limited Depth to saturated zone Slope	1.00 0.04	Very limited Slope Depth to saturated zone	1.00 0.07
86D: Paxton-----	55	Very limited Slope Depth to saturated zone	1.00 0.39	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.39
Montauk-----	30	Very limited Slope Depth to saturated zone	1.00 0.07	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.07
87B: Wethersfield-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
87C: Wethersfield-----	80	Somewhat limited Slope Depth to saturated zone	0.63 0.39	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.39
87D: Wethersfield-----	80	Very limited Slope Depth to saturated zone	1.00 0.39	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.39
88B: Wethersfield-----	80	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
88C: Wethersfield-----	80	Somewhat limited Slope Depth to saturated zone	0.63 0.39	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.39
89C: Wethersfield-----	80	Somewhat limited Depth to saturated zone Slope	0.39 0.04	Very limited Depth to saturated zone Slope	1.00 0.04	Very limited Slope Depth to saturated zone	1.00 0.39

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
89D: Wethersfield-----	80	Very limited Slope Depth to saturated zone	1.00 0.39	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.39
90B: Stockbridge-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
90C: Stockbridge-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
90D: Stockbridge-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
91B: Stockbridge-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
91C: Stockbridge-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
91D: Stockbridge-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
92B: Nellis-----	85	Not limited		Not limited		Somewhat limited Slope	0.50
92C: Nellis-----	85	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
92D: Nellis-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
93C: Nellis-----	85	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
94C: Farmington-----	40	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
Nellis-----	35	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
94E: Farmington-----	40	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
94E: Nellis-----	35	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
95C: Farmington-----	60	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 1.00
Rock outcrop-----	20	Not rated		Not rated		Not rated	
95E: Farmington-----	60	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Rock outcrop-----	20	Not rated		Not rated		Not rated	
96: Ipswich-----	85	Very limited Ponding Subsidence Flooding	1.00 1.00 1.00	Very limited Ponding Subsidence Flooding	1.00 1.00 1.00	Very limited Ponding Subsidence Flooding	1.00 1.00 1.00
97: Pawcatuck-----	85	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00
98: Westbrook-----	80	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00
99: Westbrook, low salt-	80	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00
100: Suncook-----	80	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.09	Very limited Flooding	1.00
101: Occum-----	80	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.09	Very limited Flooding	1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
102: Pootatuck-----	80	Very limited Flooding Depth to saturated zone	1.00 0.39	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.39
103: Rippowam-----	80	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
104: Bash-----	80	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
105: Hadley-----	80	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.03	Very limited Flooding	1.00
106: Winooski-----	80	Very limited Flooding Depth to saturated zone	1.00 0.07	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 0.07
107: Limerick-----	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Lim-----	30	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
108: Saco-----	80	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00
109: Fluvaquents, Frequently Flooded-	50	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
Udifluvents, Frequently Flooded-	35	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
221A: Ninigret-----	40	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
224A: Deerfield-----	40	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
Urban land-----	35	Not rated		Not rated		Not rated	
225B: Brancroft-----	40	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01
Urban land-----	35	Not rated		Not rated		Not rated	
226B: Berlin-----	40	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Shrink-swell	0.81 0.01
Urban land-----	35	Not rated		Not rated		Not rated	
228B: Elmridge-----	40	Very limited Shrink-swell	0.99	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Shrink-swell	0.99
Urban land-----	35	Not rated		Not rated		Not rated	
229B: Agawam-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
229C: Agawam-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
230B: Branford-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
230C: Branford-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
232B:							
Haven-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
234B:							
Merrimac-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
235B:							
Penwood-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
236B:							
Windsor-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
237A:							
Manchester-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
237C:							
Manchester-----	40	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
238A:							
Hinckley-----	40	Not limited		Not limited		Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
238C:							
Hinckley-----	40	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
240B:							
Ludlow-----	40	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
243B:							
Rainbow-----	40	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
245B:							
Woodbridge-----	40	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
245B: Urban land-----	35	Not rated		Not rated		Not rated	
245C: Woodbridge-----	40	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.39	Slope	0.63	Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
248B: Georgia-----	40	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.12
						Depth to saturated zone	0.07
Urban land-----	35	Not rated		Not rated		Not rated	
250B: Sutton-----	40	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
253B: Wapping-----	40	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
255B: Watchaug-----	40	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
260B: Charlton-----	40	Not limited		Not limited		Somewhat limited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
260C: Charlton-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
260D: Charlton-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
263B: Cheshire-----	40	Not limited		Not limited		Somewhat limited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
263C: Cheshire-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
266B: Narragansett-----	40	Not limited		Not limited		Somewhat limited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
269B: Yalesville-----	40	Somewhat limited Depth to hard bedrock	0.05	Very limited Depth to hard bedrock	1.00	Somewhat limited Slope	0.50
						Depth to hard bedrock	0.05
Urban land-----	35	Not rated		Not rated		Not rated	
269C: Yalesville-----	40	Somewhat limited Slope	0.63	Very limited Depth to hard bedrock	1.00	Very limited Slope	1.00
		Depth to hard bedrock	0.05	Slope	0.63	Depth to hard bedrock	0.05
Urban land-----	35	Not rated		Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
Chatfield-----	15	Somewhat limited Depth to hard bedrock	0.54	Very limited Depth to hard bedrock	1.00	Very limited Slope	1.00
		Slope	0.04	Slope	0.04	Depth to hard bedrock	0.54
273E: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Chatfield-----	15	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Depth to hard bedrock	0.54	Depth to hard bedrock	1.00	Depth to hard bedrock	0.54

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
275C:							
Urban land-----	45	Not rated		Not rated		Not rated	
Chatfield-----	30	Somewhat limited		Very limited		Very limited	
		Depth to hard bedrock	0.54	Depth to hard bedrock	1.00	Slope	1.00
		Slope	0.04	Slope	0.04	Depth to hard bedrock	0.54
275E:							
Urban land-----	35	Not rated		Not rated		Not rated	
Chatfield-----	25	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Depth to hard bedrock	0.54	Depth to hard bedrock	1.00	Depth to hard bedrock	0.54
Rock outcrop-----	15	Not rated		Not rated		Not rated	
282B:							
Broadbrook-----	40	Somewhat limited		Very limited		Somewhat limited	
		Depth to saturated zone	0.39	Depth to saturated zone	1.00	Slope	0.50
						Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
284B:							
Paxton-----	40	Somewhat limited		Very limited		Somewhat limited	
		Depth to saturated zone	0.39	Depth to saturated zone	1.00	Slope	0.50
						Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
284C:							
Paxton-----	40	Somewhat limited		Very limited		Very limited	
		Slope	0.63	Depth to saturated zone	1.00	Slope	1.00
		Depth to saturated zone	0.39	Slope	0.63	Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
284D:							
Paxton-----	40	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00
		Depth to saturated zone	0.39	Depth to saturated zone	1.00	Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
287B:							
Wethersfield-----	40	Somewhat limited		Very limited		Somewhat limited	
		Depth to saturated zone	0.39	Depth to saturated zone	1.00	Slope	0.50
						Depth to saturated zone	0.39

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
287B: Urban land-----	35	Not rated		Not rated		Not rated	
287C: Wethersfield-----	40	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.39	Slope	0.63	Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
287D: Wethersfield-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Depth to saturated zone	0.39	Depth to saturated zone	1.00	Depth to saturated zone	0.39
Urban land-----	35	Not rated		Not rated		Not rated	
290B: Stockbridge-----	40	Not limited		Not limited		Somewhat limited Slope	0.50
Urban land-----	35	Not rated		Not rated		Not rated	
290C: Stockbridge-----	40	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
290D: Stockbridge-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
301: Beaches-----	50	Not rated		Not rated		Not rated	
Udipsamments-----	35	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
		Slope	0.01	Depth to saturated zone	0.16	Slope	1.00
				Slope	0.01		
302: Dumps-----	95	Not rated		Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated		Not rated	
304: Udorthents-----	90	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Depth to saturated zone	0.18		

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
305: Udorthents-----	65	Very limited Slope	1.00	Very limited Slope Depth to saturated zone	1.00 0.89	Very limited Slope	1.00
Pits-----	25	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
306: Udorthents-----	50	Very limited Slope	1.00	Very limited Slope Depth to saturated zone	1.00 0.18	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
307: Urban land-----	80	Not rated		Not rated		Not rated	
308: Udorthents-----	80	Very limited Slope	1.00	Very limited Slope Depth to saturated zone	1.00 0.89	Very limited Slope	1.00
309: Udorthents-----	80	Very limited Flooding Slope	1.00 1.00	Very limited Flooding Slope Depth to saturated zone	1.00 1.00 0.89	Very limited Flooding Slope	1.00 1.00
310: Udorthents, Periodically Flooded-----	85	Very limited Flooding Slope	1.00 1.00	Very limited Flooding Slope Depth to saturated zone	1.00 1.00 0.89	Very limited Flooding Slope	1.00 1.00
401C: Macomber-----	55	Somewhat limited Slope Depth to hard bedrock	0.63 0.46	Very limited Depth to hard bedrock Slope	1.00 0.63	Very limited Slope Depth to hard bedrock	1.00 0.46
Taconic-----	30	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 1.00
402D: Macomber-----	50	Very limited Slope Depth to hard bedrock	1.00 0.46	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 0.46

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
402D: Taconic-----	25	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	
403C: Taconic-----	70	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 1.00
Rock outcrop-----	25	Not rated		Not rated		Not rated	
403E: Taconic-----	70	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Rock outcrop-----	20	Not rated		Not rated		Not rated	
403F: Taconic-----	70	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Rock outcrop-----	20	Not rated		Not rated		Not rated	
405C: Dummerston-----	85	Somewhat limited Slope	0.01	Somewhat limited Slope	0.01	Very limited Slope	1.00
405E: Dummerston-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
407C: Lanesboro-----	85	Somewhat limited Depth to saturated zone Slope	0.39 0.16	Very limited Depth to saturated zone Slope	1.00 0.16	Very limited Slope Depth to saturated zone	1.00 0.39
407E: Lanesboro-----	85	Very limited Slope Depth to saturated zone	1.00 0.39	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.39
408C: Fullam-----	85	Somewhat limited Depth to saturated zone Slope	0.24 0.16	Very limited Depth to saturated zone Slope	1.00 0.16	Very limited Slope Depth to saturated zone	1.00 0.24

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
409B: Brayton-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Slope	1.00 0.12
412B: Bice-----	85	Not limited		Not limited		Somewhat limited Slope	0.50
412C: Bice-----	85	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
412D: Bice-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
413C: Bice-----	45	Somewhat limited Slope	0.01	Somewhat limited Slope	0.01	Very limited Slope	1.00
Millsite-----	40	Somewhat limited Depth to hard bedrock Slope	0.35 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Slope Depth to hard bedrock	1.00 0.35
413E: Bice-----	45	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
Millsite-----	40	Very limited Slope Depth to hard bedrock	1.00 0.35	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 0.35
414: Fredon, cold-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
415C: Millsite-----	40	Somewhat limited Depth to hard bedrock Slope	0.35 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Slope Depth to hard bedrock	1.00 0.35
Westminster-----	40	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	
415E: Millsite-----	40	Very limited Slope Depth to hard bedrock	1.00 0.35	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 0.35

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
415E: Westminster-----	40	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
Rock outcrop-----	15	Not rated		Not rated		Not rated	
416E: Rock outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
416F: Rock outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00	Very limited Slope Depth to hard bedrock	1.00 1.00
417B: Bice-----	85	Not limited		Not limited		Somewhat limited Slope	0.50
417C: Bice-----	85	Somewhat limited Slope	0.16	Somewhat limited Slope	0.16	Very limited Slope	1.00
417D: Bice-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
418C: Schroon-----	85	Somewhat limited Depth to saturated zone Slope	0.39 0.04	Very limited Depth to saturated zone Slope	1.00 0.04	Very limited Slope Depth to saturated zone	1.00 0.39
420A: Schroon-----	85	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39
420B: Schroon-----	85	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.39 0.12
421A: Ninigret, cold-----	85	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.39

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
423A: Sudbury, cold-----	85	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.07
424B: Shelburne-----	85	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
424C: Shelburne-----	85	Somewhat limited Slope Depth to saturated zone	0.63 0.39	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.39
424D: Shelburne-----	85	Very limited Slope Depth to saturated zone	1.00 0.39	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.39
425B: Shelburne-----	85	Somewhat limited Depth to saturated zone	0.39	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.39
425C: Shelburne-----	85	Somewhat limited Slope Depth to saturated zone	0.63 0.39	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.39
426D: Shelburne-----	85	Very limited Slope Depth to saturated zone	1.00 0.39	Very limited Slope Depth to saturated zone	1.00 1.00	Very limited Slope Depth to saturated zone	1.00 0.39
427B: Ashfield-----	85	Somewhat limited Depth to saturated zone	0.98	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.98 0.50
427C: Ashfield-----	85	Somewhat limited Depth to saturated zone Slope	0.98 0.63	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.98
428A: Ashfield-----	85	Somewhat limited Depth to saturated zone	0.98	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.98

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
428B: Ashfield-----	85	Somewhat limited Depth to saturated zone	0.98	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone Slope	0.98 0.50
428C: Ashfield-----	85	Somewhat limited Depth to saturated zone Slope	0.98 0.63	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.98
429A: Agawam, cold-----	80	Not limited		Not limited		Not limited	
429B: Agawam, cold-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
429C: Agawam, cold-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
433: Moosilauke-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
434A: Merrimac, cold-----	80	Not limited		Not limited		Not limited	
434B: Merrimac, cold-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
434C: Merrimac, cold-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
435: Scarboro-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
436: Halsey-----	80	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
437: Wonsqueak-----	85	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
438: Bucksport-----	85	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00	Very limited Subsidence Flooding Depth to saturated zone	1.00 1.00 1.00
440A: Boscawen-----	80	Not limited		Not limited		Not limited	
440C: Boscawen-----	80	Somewhat limited Slope	0.04	Somewhat limited Slope	0.04	Very limited Slope	1.00
440E: Boscawen-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
442: Brayton-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Slope	1.00 0.12
443: Brayton-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Slope	1.00 0.12
Loonmeadow-----	35	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
448B: Hogansburg-----	85	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.12 0.07
449B: Hogansburg-----	85	Somewhat limited Depth to saturated zone	0.07	Very limited Depth to saturated zone	1.00	Somewhat limited Slope Depth to saturated zone	0.50 0.07
449C: Hogansburg-----	85	Somewhat limited Slope Depth to saturated zone	0.63 0.07	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Depth to saturated zone	1.00 0.07
450B: Pyrities-----	80	Not limited		Not limited		Somewhat limited Slope	0.50

Table 17.—Building Site Development (Part 1)—Continued

Map symbol and soil name	Pct. of map unit	Dwellings without basements		Dwellings with basements		Small commercial buildings	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
450C: Pyrities-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
450D: Pyrities-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
451B: Pyrities-----	80	Not limited		Not limited		Somewhat limited Slope	0.50
451C: Pyrities-----	80	Somewhat limited Slope	0.63	Somewhat limited Slope	0.63	Very limited Slope	1.00
451D: Pyrities-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
457: Mudgepond-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
458: Mudgepond-----	55	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
Alden-----	35	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
501: Ondawa-----	85	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.09	Very limited Flooding	1.00
503: Rumney-----	80	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00
508: Medomak-----	85	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Ponding	1.00 1.00 1.00

Table 18.—Building Site Development (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Very limited Depth to saturated zone Depth to pan	1.00 0.99
3: Ridgebury-----	40	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Very limited Depth to saturated zone Depth to pan	1.00 0.99
Leicester-----	35	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
Whitman-----	15	Very limited Depth to saturated zone Frost action Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Dense layer	1.00 1.00 1.00 0.50	Very limited Depth to pan Depth to saturated zone Ponding	1.00 1.00 1.00
4: Leicester-----	80	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone Large stones content	1.00 0.01
5: Wilbraham-----	80	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 1.00 0.50	Very limited Depth to saturated zone Depth to pan	1.00 0.99
6: Wilbraham-----	60	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 1.00 0.50	Very limited Depth to saturated zone Depth to pan	1.00 0.99
Menlo-----	25	Very limited Depth to saturated zone Frost action Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Dense layer Cutbanks cave	1.00 1.00 1.00 0.50 0.10	Very limited Depth to saturated zone Ponding Depth to pan	1.00 1.00 0.71

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7: Mudgepond-----	85	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
8: Mudgepond-----	45	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
Alden-----	35	Very limited Depth to saturated zone Frost action Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Large stones content	1.00 1.00 0.01
9: Scitico-----	40	Very limited Depth to saturated zone Frost action Low strength Shrink-swell	1.00 1.00 1.00 0.99	Very limited Depth to saturated zone Cutbanks cave Too clayey	1.00 0.10 0.02	Very limited Depth to saturated zone	1.00
Shaker-----	30	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Too clayey Cutbanks cave	1.00 0.12 0.10	Very limited Depth to saturated zone	1.00
Maybid-----	15	Very limited Depth to saturated zone Frost action Low strength Ponding Shrink-swell	1.00 1.00 1.00 1.00 0.99	Very limited Depth to saturated zone Ponding Cutbanks cave	1.00 1.00 0.10	Very limited Depth to saturated zone Ponding	1.00 1.00
10: Raynham-----	80	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 0.10	Very limited Depth to saturated zone	1.00
12: Raypol-----	80	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
13: Walpole-----	80	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
14: Fredon-----	85	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
15: Scarboro-----	80	Very limited Depth to saturated zone Ponding Frost action	1.00 1.00 0.50	Very limited Depth to saturated zone Cutbanks cave Ponding	1.00 1.00 1.00	Very limited Organic matter content Depth to saturated zone Ponding	1.00 1.00 1.00
16: Halsey-----	80	Very limited Depth to saturated zone Frost action Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
17: Timakwa-----	45	Very limited Depth to saturated zone Frost action Shrink-swell Subsidence Ponding	1.00 1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Ponding Organic matter content	1.00 1.00 1.00 1.00	Very limited Organic matter content Depth to saturated zone Ponding	1.00 1.00 1.00
Natchaug-----	40	Very limited Depth to saturated zone Frost action Subsidence Ponding Flooding	1.00 1.00 1.00 1.00 0.40	Very limited Depth to saturated zone Ponding Organic matter content Cutbanks cave	1.00 1.00 1.00 0.10	Very limited Depth to saturated zone Ponding	1.00 1.00
18: Catden-----	40	Very limited Depth to saturated zone Subsidence Frost action Shrink-swell Ponding	1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Organic matter content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
Freetown-----	40	Very limited Depth to saturated zone Subsidence Frost action Shrink-swell Ponding	1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Organic matter content Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
20A: Ellington-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
21A: Ninigret-----	60	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
Tisbury-----	25	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
22A: Hero-----	85	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Gravel content	0.18
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
22B: Hero-----	85	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Gravel content	0.18
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
23A: Sudbury-----	80	Somewhat limited Depth to saturated zone	0.03	Very limited Depth to saturated zone	1.00	Not limited	
				Cutbanks cave	1.00		
24A: Deerfield-----	80	Somewhat limited Depth to saturated zone	0.03	Very limited Depth to saturated zone	1.00	Somewhat limited Droughty	0.17
				Cutbanks cave	1.00		
25A: Brancroft-----	80	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Low strength	1.00	Cutbanks cave	0.10		
		Depth to saturated zone	0.48				
		Shrink-swell	0.01				

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
25B: Brancroft-----	80	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Low strength	1.00	Cutbanks cave	0.10		
		Depth to saturated zone	0.48				
		Shrink-swell	0.01				
25C: Brancroft-----	80	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Low strength	1.00	Slope	0.63		
		Slope	0.63	Cutbanks cave	0.10		
		Depth to saturated zone	0.48				
		Shrink-swell	0.01				
26A: Berlin-----	80	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Low strength	1.00	Cutbanks cave	0.10		
		Depth to saturated zone	0.48				
		Shrink-swell	0.01				
26B: Berlin-----	80	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Low strength	1.00	Cutbanks cave	0.10		
		Depth to saturated zone	0.48				
		Shrink-swell	0.01				
27A: Belgrade-----	80	Very limited Frost action	1.00	Very limited Depth to saturated zone	0.99	Not limited	
				Cutbanks cave	0.10		
28A: Elmridge-----	80	Very limited Low strength	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Shrink-swell	0.99	Too clayey	0.24		
		Frost action	0.50	Cutbanks cave	0.10		
		Depth to saturated zone	0.19				
28B: Elmridge-----	80	Very limited Low strength	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Shrink-swell	0.99	Too clayey	0.24		
		Frost action	0.50	Cutbanks cave	0.10		
		Depth to saturated zone	0.19				

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
29A: Agawam-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
29B: Agawam-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
29C: Agawam-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
30A: Branford-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
30B: Branford-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
30C: Branford-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
31A: Copake-----	85	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
31B: Copake-----	85	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
31C: Copake-----	85	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
32A: Haven-----	60	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Enfield-----	25	Very limited Frost action	1.00	Very limited Cutbanks cave	1.00	Not limited	
32B: Haven-----	60	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Enfield-----	25	Very limited Frost action	1.00	Very limited Cutbanks cave	1.00	Not limited	
32C: Haven-----	60	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
Enfield-----	25	Very limited Frost action Slope	1.00 0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
33A: Hartford-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
33B: Hartford-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
34A: Merrimac-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
34B: Merrimac-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
34C: Merrimac-----	80	Somewhat limited Slope	0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
35A: Penwood-----	80	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty	1.00
35B: Penwood-----	80	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty	1.00
36A: Windsor-----	80	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.99
36B: Windsor-----	80	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.99
36C: Windsor-----	80	Somewhat limited Slope	0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Droughty Slope	0.99 0.63
37A: Manchester-----	80	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty Gravel content	1.00 0.68
37C: Manchester-----	80	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Very limited Droughty Gravel content Slope	1.00 0.68 0.04
37E: Manchester-----	80	Very limited Slope	1.00	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Droughty Gravel content	1.00 1.00 0.68

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
38A: Hinckley-----	80	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty Large stones content Gravel content	1.00 0.11 0.05
38C: Hinckley-----	80	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Very limited Droughty Large stones content Gravel content Slope	1.00 0.11 0.05 0.04
38E: Hinckley-----	80	Very limited Slope	1.00	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Droughty Large stones content Gravel content	1.00 1.00 0.11 0.05
39A: Groton-----	85	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty Gravel content	0.98 0.59
39C: Groton-----	85	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Droughty Gravel content Slope	0.98 0.59 0.04
39E: Groton-----	85	Very limited Slope	1.00	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Droughty Gravel content	1.00 0.98 0.59
40A: Ludlow-----	80	Somewhat limited Frost action Depth to saturated zone	0.50 0.19	Very limited Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 0.50	Somewhat limited Depth to pan	0.79
40B: Ludlow-----	80	Somewhat limited Frost action Depth to saturated zone	0.50 0.19	Very limited Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 0.50	Somewhat limited Depth to pan	0.79

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
41B: Ludlow-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
42C: Ludlow-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Slope	0.04
		Slope	0.04	Dense layer	0.50		
				Slope	0.04		
43A: Rainbow-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
43B: Rainbow-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
44B: Rainbow-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
45A: Woodbridge-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.46
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
45B: Woodbridge-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.46
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
45C: Woodbridge-----	80	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00	Depth to pan	0.46
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		
46B: Woodbridge-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.46
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
46C: Woodbridge-----	80	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00	Depth to pan	0.46
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		
47C: Woodbridge-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.46
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Slope	0.04
		Slope	0.04	Dense layer	0.50	Large stones content	0.01
				Slope	0.04		
48B: Georgia-----	50	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
Amenia-----	35	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
48C: Georgia-----	50	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00		
		Depth to saturated zone	0.03	Slope	0.63		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
48C: Amenia-----	35	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00		
		Depth to saturated zone	0.03	Slope	0.63		
49B: Georgia-----	50	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
Amenia-----	35	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
49C: Georgia-----	50	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00		
		Depth to saturated zone	0.03	Slope	0.63		
49C: Amenia-----	35	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00		
		Depth to saturated zone	0.03	Slope	0.63		
50A: Sutton-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
50B: Sutton-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
51B: Sutton-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
52C: Sutton-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.04
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
		Slope	0.04	Slope	0.04		
53A: Wapping-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
53B: Wapping-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
54B: Wapping-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
55A: Watchaug-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Large stones content	0.01
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
55B: Watchaug-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Large stones content	0.01
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
56B: Watchaug-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Large stones content	0.01
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
57B: Gloucester-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
57C: Gloucester-----	80	Somewhat limited Slope	0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
57D: Gloucester-----	80	Very limited Slope	1.00	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
58B: Gloucester-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
58C: Gloucester-----	80	Somewhat limited Slope	0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
59C: Gloucester-----	80	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope	0.04
59D: Gloucester-----	80	Very limited Slope	1.00	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
60B: Canton-----	45	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Gravel content	0.01
60B: Charlton-----	35	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
60C: Canton-----	45	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Gravel content	0.63 0.01
Charlton-----	35	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
60D: Canton-----	45	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Gravel content	1.00 0.01
Charlton-----	35	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
61B: Canton-----	45	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Gravel content	0.01
Charlton-----	35	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
61C: Canton-----	45	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Gravel content	0.63 0.01

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
61C: Charlton-----	35	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
62C: Canton-----	45	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope Gravel content	0.04 0.01
Charlton-----	35	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope	0.04
62D: Canton-----	45	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Gravel content	1.00 0.01
Charlton-----	35	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
63B: Cheshire-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Large stones content	0.01
63C: Cheshire-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Large stones content	0.63 0.01
63D: Cheshire-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Large stones content	1.00 0.01
64B: Cheshire-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Large stones content	0.01
64C: Cheshire-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Large stones content	0.63 0.01
65C: Cheshire-----	80	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope Large stones content	0.04 0.01
65D: Cheshire-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Large stones content	1.00 0.01

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
66B: Narragansett-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Large stones content	0.01
66C: Narragansett-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Large stones content	0.63 0.01
67B: Narragansett-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Large stones content	0.01
67C: Narragansett-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Large stones content	0.63 0.01
68C: Narragansett-----	80	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope Large stones content	0.04 0.01
68D: Narragansett-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Large stones content	1.00 0.01
69B: Yalesville-----	75	Somewhat limited Frost action Depth to hard bedrock	0.50 0.05	Very limited Depth to hard bedrock Cutbanks cave	1.00 1.00	Somewhat limited Depth to bedrock	0.05
69C: Yalesville-----	75	Somewhat limited Slope Frost action Depth to hard bedrock	0.63 0.50 0.05	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 1.00 0.63	Somewhat limited Slope Depth to bedrock	0.63 0.05
70C: Branford-----	50	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope	0.04
Holyoke-----	30	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to bedrock Droughty Slope	1.00 0.25 0.04

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
71C:							
Brookfield-----	45	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope	0.04
Brimfield-----	30	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to bedrock Droughty Slope	1.00 0.93 0.04
Rock outcrop-----	15	Not rated		Not rated		Not rated	
71E:							
Brookfield-----	45	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
Brimfield-----	30	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.93
Rock outcrop-----	15	Not rated		Not rated		Not rated	
73C:							
Charlton-----	45	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope	0.04
Chatfield-----	30	Somewhat limited Depth to hard bedrock Frost action Slope	0.54 0.50 0.04	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 1.00 0.04	Somewhat limited Depth to bedrock Gravel content Slope	0.54 0.25 0.04
73E:							
Charlton-----	45	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
Chatfield-----	30	Very limited Slope Depth to hard bedrock Frost action	1.00 0.54 0.50	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 1.00 1.00	Very limited Slope Depth to bedrock Gravel content	1.00 0.54 0.25
74C:							
Narragansett-----	55	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope Large stones content	0.04 0.01
Hollis-----	20	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.22 0.04

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
75C: Hollis-----	35	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.22 0.04
75C: Chatfield-----	30	Somewhat limited Depth to hard bedrock Frost action Slope	0.54 0.50 0.04	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 1.00 0.04	Somewhat limited Depth to bedrock Gravel content Slope	0.54 0.25 0.04
Rock outcrop-----	15	Not rated		Not rated		Not rated	
75E: Hollis-----	35	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.22
Chatfield-----	30	Very limited Slope Depth to hard bedrock Frost action	1.00 0.54 0.50	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 1.00 1.00	Very limited Slope Depth to bedrock Gravel content	1.00 0.54 0.25
Rock outcrop-----	15	Not rated		Not rated		Not rated	
76E: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Droughty Slope Gravel content	1.00 1.00 1.00 0.22
76F: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Hollis-----	25	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.22
77C: Cheshire-----	45	Somewhat limited Frost action Slope	0.50 0.04	Very limited Cutbanks cave Slope	1.00 0.04	Somewhat limited Slope Large stones content	0.04 0.01

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
77C: Holyoke-----	35	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to bedrock Droughty Slope	1.00 0.25 0.04
77D: Cheshire-----	45	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Large stones content	1.00 0.01
Holyoke-----	35	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.25
78C: Holyoke-----	50	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope	1.00 0.04	Very limited Depth to bedrock Droughty Slope	1.00 0.25 0.04
Rock outcrop-----	25	Not rated		Not rated		Not rated	
78E: Holyoke-----	50	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.25
Rock outcrop-----	25	Not rated		Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated		Not rated	
Holyoke-----	25	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.25
80B: Bernardston-----	80	Somewhat limited Frost action Depth to saturated zone	0.50 0.43	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Somewhat limited Depth to pan	0.79
80C: Bernardston-----	80	Somewhat limited Slope Frost action Depth to saturated zone	0.63 0.50 0.43	Very limited Depth to saturated zone Slope Dense layer Cutbanks cave	1.00 0.63 0.50 0.10	Somewhat limited Depth to pan Slope	0.79 0.63

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
81C: Bernardston-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.43	Dense layer	0.50	Slope	0.04
		Slope	0.04	Cutbanks cave	0.10		
				Slope	0.04		
81D: Bernardston-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.79
		Depth to saturated zone	0.43	Dense layer	0.50		
				Cutbanks cave	0.10		
82B: Broadbrook-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
82C: Broadbrook-----	80	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		
82D: Broadbrook-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.84
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
83B: Broadbrook-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
83C: Broadbrook-----	80	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
84B: Paxton-----	55	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
Montauk-----	30	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
				Dense layer	0.50		
84C: Paxton-----	55	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63	Large stones content	0.01
				Dense layer	0.50		
Montauk-----	30	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.03	Slope	0.63		
				Dense layer	0.50		
84D: Paxton-----	55	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
Montauk-----	30	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.84
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
				Dense layer	0.50		
85B: Paxton-----	55	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
85B: Montauk-----	30	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
				Dense layer	0.50		
85C: Paxton-----	55	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63	Large stones content	0.01
				Dense layer	0.50		
Montauk-----	30	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.03	Slope	0.63		
				Dense layer	0.50		
86C: Paxton-----	55	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Slope	0.04
		Slope	0.04	Dense layer	0.50	Large stones content	0.01
				Slope	0.04		
Montauk-----	30	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Slope	0.04	Cutbanks cave	1.00	Slope	0.04
		Depth to saturated zone	0.03	Dense layer	0.50		
				Slope	0.04		
86D: Paxton-----	55	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
86D: Montauk-----	30	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.84
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
				Dense layer	0.50		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
87B: Wethersfield-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.71
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
87C: Wethersfield-----	80	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.71
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63	Large stones content	0.01
				Dense layer	0.50		
87D: Wethersfield-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.71
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
88B: Wethersfield-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.71
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
88C: Wethersfield-----	80	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.71
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		
89C: Wethersfield-----	80	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.71
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Slope	0.04
		Slope	0.04	Dense layer	0.50		
				Slope	0.04		
89D: Wethersfield-----	80	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.71
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
90B: Stockbridge-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
90C: Stockbridge-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
90D: Stockbridge-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
91B: Stockbridge-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
91C: Stockbridge-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
91D: Stockbridge-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
92B: Nellis-----	85	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
92C: Nellis-----	85	Somewhat limited Slope Frost action	0.63 0.50	Somewhat limited Slope Cutbanks cave	0.63 0.10	Somewhat limited Slope	0.63
92D: Nellis-----	85	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
93C: Nellis-----	85	Somewhat limited Frost action Slope	0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04
94C: Farmington-----	40	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 0.10 0.04	Very limited Depth to bedrock Droughty Slope	1.00 0.81 0.04
Nellis-----	35	Somewhat limited Frost action Slope	0.50 0.04	Somewhat limited Cutbanks cave Slope	0.10 0.04	Somewhat limited Slope	0.04

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
94E: Farmington-----	40	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.81
Nellis-----	35	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
95C: Farmington-----	60	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 0.10 0.04	Very limited Depth to bedrock Droughty Slope	1.00 0.81 0.04
Rock outcrop-----	20	Not rated		Not rated		Not rated	
95E: Farmington-----	60	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Depth to bedrock Slope Droughty	1.00 1.00 0.81
Rock outcrop-----	20	Not rated		Not rated		Not rated	
96: Ipswich-----	85	Very limited Ponding Depth to saturated zone Subsidence Frost action Flooding	1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone Organic matter content	1.00 1.00 1.00 1.00	Very limited Ponding Flooding Organic matter content Salinity Sulfur content	1.00 1.00 1.00 1.00 1.00
97: Pawcatuck-----	85	Very limited Ponding Depth to saturated zone Frost action Flooding Shrink-swell	1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone Cutbanks cave Organic matter content	1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Organic matter content Salinity Sulfur content	1.00 1.00 1.00 1.00 1.00
98: Westbrook-----	80	Very limited Ponding Depth to saturated zone Frost action Flooding Shrink-swell	1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone Organic matter content	1.00 1.00 1.00 1.00	Very limited Ponding Flooding Salinity Sulfur content Depth to saturated zone	1.00 1.00 1.00 1.00 1.00

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
99: Westbrook, low salt-	80	Very limited Ponding Depth to saturated zone Frost action Flooding Shrink-swell	1.00 1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Depth to saturated zone Organic matter content	1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Sulfur content Depth to saturated zone Salinity	1.00 1.00 1.00 1.00 1.00 1.00
100: Suncook-----	80	Very limited Flooding	1.00	Very limited Cutbanks cave Flooding Depth to saturated zone	1.00 0.60 0.09	Somewhat limited Droughty Flooding	0.89 0.60
101: Occum-----	80	Very limited Flooding Frost action	1.00 0.50	Very limited Cutbanks cave Flooding Depth to saturated zone	1.00 0.60 0.09	Somewhat limited Flooding	0.60
102: Pootatuck-----	80	Very limited Flooding Frost action Depth to saturated zone	1.00 0.50 0.19	Very limited Depth to saturated zone Cutbanks cave Flooding	1.00 1.00 0.80	Very limited Flooding	1.00
103: Rippowam-----	80	Very limited Depth to saturated zone Frost action Flooding	1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Flooding	1.00 1.00 0.80	Very limited Flooding Depth to saturated zone	1.00 1.00
104: Bash-----	80	Very limited Flooding Depth to saturated zone Frost action	1.00 0.99 0.50	Very limited Depth to saturated zone Flooding Cutbanks cave	1.00 0.80 0.10	Very limited Flooding Depth to saturated zone	1.00 0.99
105: Hadley-----	80	Very limited Frost action Flooding	1.00 1.00	Very limited Cutbanks cave Flooding Depth to saturated zone	1.00 0.60 0.03	Somewhat limited Flooding	0.60

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
106: Winooski-----	80	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Flooding	1.00	Flooding	0.80		
		Depth to saturated zone	0.03	Cutbanks cave	0.10		
107: Limerick-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Frost action	1.00	Flooding	0.80	Depth to saturated zone	1.00
		Flooding	1.00	Cutbanks cave	0.10		
Lim-----	30	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Frost action	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Flooding	0.80		
108: Saco-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Frost action	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Ponding	1.00	Ponding	1.00
		Ponding	1.00	Flooding	0.80		
109: Fluvaquents, Frequently Flooded-	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Flooding	1.00
		Frost action	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Flooding	0.80		
Udifluents, Frequently Flooded-	35	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Very limited Flooding	1.00
		Frost action	0.50	Flooding	0.80		
221A: Ninigret-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
Urban land-----	35	Not rated		Not rated		Not rated	
224A: Deerfield-----	40	Somewhat limited Depth to saturated zone	0.03	Very limited Depth to saturated zone	1.00	Somewhat limited Droughty	0.17
				Cutbanks cave	1.00		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
224A: Urban land-----	35	Not rated		Not rated		Not rated	
225B: Brancroft-----	40	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Low strength Depth to saturated zone	1.00 0.48	Cutbanks cave	0.10		
		Shrink-swell	0.01				
Urban land-----	35	Not rated		Not rated		Not rated	
226B: Berlin-----	40	Very limited Frost action	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Low strength Depth to saturated zone	1.00 0.48	Cutbanks cave	0.10		
		Shrink-swell	0.01				
Urban land-----	35	Not rated		Not rated		Not rated	
228B: Elmridge-----	40	Very limited Low strength	1.00	Very limited Depth to saturated zone	1.00	Not limited	
		Shrink-swell	0.99	Too clayey	0.24		
		Frost action	0.50	Cutbanks cave	0.10		
		Depth to saturated zone	0.19				
Urban land-----	35	Not rated		Not rated		Not rated	
229B: Agawam-----	40	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
229C: Agawam-----	40	Somewhat limited Slope	0.63	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Slope	0.63		
Urban land-----	35	Not rated		Not rated		Not rated	
230B: Branford-----	40	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
230C: Branford-----	40	Somewhat limited Slope	0.63	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Slope	0.63		
Urban land-----	35	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
232B: Haven-----	40	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
234B: Merrimac-----	40	Not limited		Very limited Cutbanks cave	1.00	Not limited	
234B: Urban land-----	35	Not rated		Not rated		Not rated	
235B: Penwood-----	40	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
236B: Windsor-----	40	Not limited		Very limited Cutbanks cave	1.00	Somewhat limited Droughty	0.99
Urban land-----	35	Not rated		Not rated		Not rated	
237A: Manchester-----	40	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty Gravel content	1.00 0.68
Urban land-----	35	Not rated		Not rated		Not rated	
237C: Manchester-----	40	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Very limited Droughty Gravel content Slope	1.00 0.68 0.04
Urban land-----	35	Not rated		Not rated		Not rated	
238A: Hinckley-----	40	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty Large stones content Gravel content	1.00 0.11 0.05
Urban land-----	35	Not rated		Not rated		Not rated	
238C: Hinckley-----	40	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Very limited Droughty Large stones content Gravel content Slope	1.00 0.11 0.05 0.04
Urban land-----	35	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
240B: Ludlow-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
243B: Rainbow-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
245B: Woodbridge-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.46
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
245C: Woodbridge-----	40	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00	Depth to pan	0.46
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
248B: Georgia-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.03	Cutbanks cave	1.00		
Urban land-----	35	Not rated		Not rated		Not rated	
250B: Sutton-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
Urban land-----	35	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
253B: Wapping-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
Urban land-----	35	Not rated		Not rated		Not rated	
255B: Watchaug-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Large stones content	0.01
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
Urban land-----	35	Not rated		Not rated		Not rated	
260B: Charlton-----	40	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
260C: Charlton-----	40	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
Urban land-----	35	Not rated		Not rated		Not rated	
260D: Charlton-----	40	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
Urban land-----	35	Not rated		Not rated		Not rated	
263B: Cheshire-----	40	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Large stones content	0.01
Urban land-----	35	Not rated		Not rated		Not rated	
263C: Cheshire-----	40	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope Large stones content	0.63 0.01
Urban land-----	35	Not rated		Not rated		Not rated	
266B: Narragansett-----	40	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Large stones content	0.01
Urban land-----	35	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
269B: Yalesville-----	40	Somewhat limited Frost action	0.50	Very limited Depth to hard bedrock	1.00	Somewhat limited Depth to bedrock	0.05
		Depth to hard bedrock	0.05	Cutbanks cave	1.00		
269B: Urban land-----	35	Not rated		Not rated		Not rated	
269C: Yalesville-----	40	Somewhat limited Slope	0.63	Very limited Depth to hard bedrock	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Cutbanks cave	1.00	Depth to bedrock	0.05
		Depth to hard bedrock	0.05	Slope	0.63		
Urban land-----	35	Not rated		Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.04
		Slope	0.04	Slope	0.04		
Chatfield-----	15	Somewhat limited Depth to hard bedrock	0.54	Very limited Depth to hard bedrock	1.00	Somewhat limited Depth to bedrock	0.54
		Frost action	0.50	Cutbanks cave	1.00	Gravel content	0.25
		Slope	0.04	Slope	0.04	Slope	0.04
273E: Urban land-----	35	Not rated		Not rated		Not rated	
Charlton-----	25	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Cutbanks cave	1.00		
Chatfield-----	15	Very limited Slope	1.00	Very limited Depth to hard bedrock	1.00	Very limited Slope	1.00
		Depth to hard bedrock	0.54	Slope	1.00	Depth to bedrock	0.54
		Frost action	0.50	Cutbanks cave	1.00	Gravel content	0.25
275C: Urban land-----	45	Not rated		Not rated		Not rated	
Chatfield-----	30	Somewhat limited Depth to hard bedrock	0.54	Very limited Depth to hard bedrock	1.00	Somewhat limited Depth to bedrock	0.54
		Frost action	0.50	Cutbanks cave	1.00	Gravel content	0.25
		Slope	0.04	Slope	0.04	Slope	0.04
275E: Urban land-----	35	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
275E: Chatfield-----	25	Very limited Slope	1.00	Very limited Depth to hard bedrock	1.00	Very limited Slope	1.00
		Depth to hard bedrock	0.54	Slope	1.00	Depth to bedrock	0.54
		Frost action	0.50	Cutbanks cave	1.00	Gravel content	0.25
Rock outcrop-----	15	Not rated		Not rated		Not rated	
282B: Broadbrook-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.84
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
284B: Paxton-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
284C: Paxton-----	40	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.79
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63	Large stones content	0.01
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
284D: Paxton-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.79
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
287B: Wethersfield-----	40	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.71
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
287C: Wethersfield-----	40	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.71
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63	Large stones content	0.01
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
287D: Wethersfield-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.71
		Depth to saturated zone	0.19	Cutbanks cave	1.00	Large stones content	0.01
				Dense layer	0.50		
Urban land-----	35	Not rated		Not rated		Not rated	
290B: Stockbridge-----	40	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
Urban land-----	35	Not rated		Not rated		Not rated	
290C: Stockbridge-----	40	Somewhat limited Slope	0.63	Very limited Cutbanks cave	1.00	Somewhat limited Slope	0.63
		Frost action	0.50	Slope	0.63		
Urban land-----	35	Not rated		Not rated		Not rated	
290D: Stockbridge-----	40	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Cutbanks cave	1.00		
Urban land-----	35	Not rated		Not rated		Not rated	
301: Beaches-----	50	Not rated		Not rated		Not rated	
Udipsamments-----	35	Very limited Flooding	1.00	Very limited Cutbanks cave	1.00	Very limited Droughty	1.00
		Slope	0.01	Flooding	0.60	Flooding	0.60
				Depth to saturated zone	0.16	Too sandy	0.50
				Slope	0.01	Slope	0.01
302: Dumps-----	95	Not rated		Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
304: Udorthents-----	90	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave Depth to saturated zone	1.00 1.00 0.18	Very limited Slope Large stones content	1.00 0.01
305: Udorthents-----	65	Very limited Slope Frost action	1.00 0.50	Very limited Cutbanks cave Slope Depth to saturated zone	1.00 1.00 0.89	Very limited Slope Large stones content	1.00 0.01
Pits-----	25	Very limited Slope	1.00	Very limited Cutbanks cave Slope	1.00 1.00	Very limited Droughty Gravel content Slope Too sandy Large stones content	1.00 1.00 1.00 0.50 0.16
306: Udorthents-----	50	Very limited Slope Frost action	1.00 0.50	Very limited Cutbanks cave Slope Depth to saturated zone	1.00 1.00 0.18	Very limited Slope Large stones content	1.00 0.01
Urban land-----	35	Not rated		Not rated		Not rated	
307: Urban land-----	80	Not rated		Not rated		Not rated	
308: Udorthents-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Cutbanks cave Slope Depth to saturated zone	1.00 1.00 0.89	Very limited Slope Large stones content	1.00 0.01
309: Udorthents-----	80	Very limited Slope Frost action Flooding	1.00 0.50 0.40	Very limited Cutbanks cave Slope Depth to saturated zone	1.00 1.00 0.89	Very limited Slope Large stones content	1.00 0.01

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
310: Udorthents, Periodically Flooded-----	85	Very limited Flooding Slope Frost action	1.00 1.00 0.50	Very limited Cutbanks cave Slope Depth to saturated zone Flooding	1.00 1.00 0.89 0.80	Very limited Flooding Slope Large stones content	1.00 1.00 0.01
401C: Macomber-----	55	Somewhat limited Slope Frost action Depth to hard bedrock	0.63 0.50 0.46	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 0.63 0.10	Somewhat limited Gravel content Slope Depth to bedrock Droughty	0.82 0.63 0.46 0.01
Taconic-----	30	Very limited Depth to hard bedrock Frost action Slope	1.00 0.50 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.39 0.01
402D: Macomber-----	50	Very limited Slope Frost action Depth to hard bedrock	1.00 0.50 0.46	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Slope Gravel content Depth to bedrock Droughty	1.00 0.82 0.46 0.01
Taconic-----	25	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 0.50	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.39
Rock outcrop-----	15	Not rated		Not rated		Not rated	
403C: Taconic-----	70	Very limited Depth to hard bedrock Frost action Slope	1.00 0.50 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to bedrock Droughty Gravel content Slope	1.00 1.00 0.39 0.01
Rock outcrop-----	25	Not rated		Not rated		Not rated	
403E: Taconic-----	70	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 0.50	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.39

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
403E: Rock outcrop-----	20	Not rated		Not rated		Not rated	
403F: Taconic-----	70	Very limited Depth to hard bedrock Slope Frost action	1.00 1.00 0.50	Very limited Depth to hard bedrock Slope	1.00 1.00	Very limited Depth to bedrock Slope Droughty Gravel content	1.00 1.00 1.00 0.39
Rock outcrop-----	20	Not rated		Not rated		Not rated	
405C: Dummerston-----	85	Somewhat limited Frost action Slope	0.50 0.01	Very limited Cutbanks cave Slope	1.00 0.01	Somewhat limited Gravel content Slope	0.09 0.01
405E: Dummerston-----	85	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Gravel content	1.00 0.09
407C: Lanesboro-----	85	Somewhat limited Frost action Depth to saturated zone Slope	0.50 0.19 0.16	Very limited Depth to saturated zone Dense layer Slope Cutbanks cave	1.00 0.50 0.16 0.10	Somewhat limited Depth to pan Slope	0.46 0.16
407E: Lanesboro-----	85	Very limited Slope Frost action Depth to saturated zone	1.00 0.50 0.19	Very limited Slope Depth to saturated zone Dense layer Cutbanks cave	1.00 1.00 0.50 0.10	Very limited Slope Depth to pan	1.00 0.46
408C: Fullam-----	85	Somewhat limited Frost action Slope Depth to saturated zone	0.50 0.16 0.12	Very limited Depth to saturated zone Cutbanks cave Dense layer Slope	1.00 1.00 0.50 0.16	Very limited Depth to pan Slope Droughty	0.99 0.16 0.05
409B: Brayton-----	85	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 1.00 0.50	Very limited Depth to pan Depth to saturated zone	1.00 1.00
412B: Bice-----	85	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
412C: Bice-----	85	Somewhat limited Frost action Slope	0.50 0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Slope	0.16
412D: Bice-----	85	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
413C: Bice-----	45	Somewhat limited Frost action Slope	0.50 0.01	Very limited Cutbanks cave Slope	1.00 0.01	Somewhat limited Slope	0.01
Millsite-----	40	Somewhat limited Frost action Depth to hard bedrock Slope	0.50 0.35 0.01	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 0.10 0.01	Somewhat limited Depth to bedrock Slope	0.35 0.01
413E: Bice-----	45	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
Millsite-----	40	Very limited Slope Frost action Depth to hard bedrock	1.00 0.50 0.35	Very limited Depth to hard bedrock Slope Cutbanks cave	1.00 1.00 0.10	Very limited Slope Depth to bedrock	1.00 0.35
414: Fredon, cold-----	85	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
415C: Millsite-----	40	Somewhat limited Frost action Depth to hard bedrock Slope	0.50 0.35 0.01	Very limited Depth to hard bedrock Cutbanks cave Slope	1.00 0.10 0.01	Somewhat limited Depth to bedrock Slope	0.35 0.01
Westminster-----	40	Very limited Depth to hard bedrock Frost action Slope	1.00 1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 0.01	Very limited Depth to bedrock Droughty Slope	1.00 0.91 0.01
Rock outcrop-----	15	Not rated		Not rated		Not rated	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
415E: Millsite-----	40	Very limited Slope	1.00	Very limited Depth to hard bedrock	1.00	Very limited Slope	1.00
		Frost action Depth to hard bedrock	0.50 0.35	Slope Cutbanks cave	1.00 0.10	Depth to bedrock	0.35
Westminster-----	40	Very limited Depth to hard bedrock	1.00	Very limited Depth to hard bedrock	1.00	Very limited Depth to bedrock	1.00
		Slope Frost action	1.00 1.00	Slope	1.00	Slope Droughty	1.00 0.91
Rock outcrop-----	15	Not rated		Not rated		Not rated	
416E: Rock outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Depth to hard bedrock	1.00	Very limited Depth to hard bedrock	1.00	Very limited Depth to bedrock	1.00
		Frost action Slope	1.00 1.00	Slope	1.00	Slope Droughty	1.00 0.91
416F: Rock outcrop-----	70	Not rated		Not rated		Not rated	
Westminster-----	20	Very limited Depth to hard bedrock	1.00	Very limited Depth to hard bedrock	1.00	Very limited Depth to bedrock	1.00
		Slope Frost action	1.00 1.00	Slope	1.00	Slope Droughty	1.00 0.91
417B: Bice-----	85	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
417C: Bice-----	85	Somewhat limited Frost action Slope	0.50 0.16	Very limited Cutbanks cave Slope	1.00 0.16	Somewhat limited Slope	0.16
417D: Bice-----	85	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope	1.00
418C: Schroon-----	85	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.04
		Depth to saturated zone Slope	0.19 0.04	Cutbanks cave Slope	0.10 0.04		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
420B: Schroon-----	85	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	0.10		
421A: Ninigret, cold-----	85	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Not limited	
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
423A: Sudbury, cold-----	85	Somewhat limited Depth to saturated zone	0.03	Very limited Depth to saturated zone	1.00	Not limited	
				Cutbanks cave	1.00		
424B: Shelburne-----	85	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.74
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
424C: Shelburne-----	85	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.74
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		
424D: Shelburne-----	85	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Frost action	0.50	Depth to saturated zone	1.00	Depth to pan	0.74
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
425B: Shelburne-----	85	Somewhat limited Frost action	0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.74
		Depth to saturated zone	0.19	Cutbanks cave	1.00		
				Dense layer	0.50		
425C: Shelburne-----	85	Somewhat limited Slope	0.63	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to pan	0.74
		Frost action	0.50	Cutbanks cave	1.00	Slope	0.63
		Depth to saturated zone	0.19	Slope	0.63		
				Dense layer	0.50		

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
426D: Shelburne-----	85	Very limited Slope Frost action Depth to saturated zone	1.00 0.50 0.19	Very limited Slope Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 1.00 0.50	Very limited Slope Depth to pan	1.00 0.74
427B: Ashfield-----	85	Somewhat limited Depth to saturated zone Frost action	0.75 0.50	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Somewhat limited Depth to pan Depth to saturated zone	0.54 0.08
427C: Ashfield-----	85	Somewhat limited Depth to saturated zone Slope Frost action	0.75 0.63 0.50	Very limited Depth to saturated zone Slope Dense layer Cutbanks cave	1.00 0.63 0.50 0.10	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.54 0.08
428A: Ashfield-----	85	Somewhat limited Depth to saturated zone Frost action	0.75 0.50	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Somewhat limited Depth to pan Depth to saturated zone	0.54 0.08
428B: Ashfield-----	85	Somewhat limited Depth to saturated zone Frost action	0.75 0.50	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Somewhat limited Depth to pan Depth to saturated zone	0.54 0.08
428C: Ashfield-----	85	Somewhat limited Depth to saturated zone Slope Frost action	0.75 0.63 0.50	Very limited Depth to saturated zone Slope Dense layer Cutbanks cave	1.00 0.63 0.50 0.10	Somewhat limited Slope Depth to pan Depth to saturated zone	0.63 0.54 0.08
429A: Agawam, cold-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	
429B: Agawam, cold-----	80	Somewhat limited Frost action	0.50	Very limited Cutbanks cave	1.00	Not limited	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
429C: Agawam, cold-----	80	Somewhat limited Slope Frost action	0.63 0.50	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
433: Moosilauke-----	80	Very limited Depth to saturated zone Frost action	1.00 0.50	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
434A: Merrimac, cold-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
434B: Merrimac, cold-----	80	Not limited		Very limited Cutbanks cave	1.00	Not limited	
434C: Merrimac, cold-----	80	Somewhat limited Slope	0.63	Very limited Cutbanks cave Slope	1.00 0.63	Somewhat limited Slope	0.63
435: Scarboro-----	80	Very limited Depth to saturated zone Ponding Frost action	1.00 1.00 0.50	Very limited Depth to saturated zone Cutbanks cave Ponding	1.00 1.00 1.00	Very limited Organic matter content Depth to saturated zone Ponding	1.00 1.00 1.00
436: Halsey-----	80	Very limited Depth to saturated zone Frost action Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
437: Wonsqueak-----	85	Very limited Depth to saturated zone Frost action Ponding Flooding	1.00 1.00 1.00 0.40	Very limited Depth to saturated zone Cutbanks cave Ponding Organic matter content	1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
438: Bucksport-----	85	Very limited Depth to saturated zone Subsidence Frost action Shrink-swell Ponding	1.00 1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Ponding Organic matter content	1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
440A: Boscawen-----	80	Not limited		Very limited Cutbanks cave	1.00	Very limited Droughty	1.00

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
440C: Boscawen-----	80	Somewhat limited Slope	0.04	Very limited Cutbanks cave Slope	1.00 0.04	Very limited Droughty Slope	1.00 0.04
440E: Boscawen-----	85	Very limited Slope	1.00	Very limited Slope Cutbanks cave	1.00 1.00	Very limited Slope Droughty	1.00 1.00
442: Brayton-----	85	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 1.00 0.50	Very limited Depth to saturated zone Depth to pan	1.00 0.71
443: Brayton-----	50	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Dense layer	1.00 1.00 1.00 0.50	Very limited Depth to saturated zone Depth to pan	1.00 0.71
Loonmeadow-----	35	Very limited Depth to saturated zone Frost action Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Ponding Dense layer	1.00 1.00 1.00 1.00 0.50	Very limited Depth to saturated zone Ponding	1.00 1.00
448B: Hogansburg-----	85	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Not limited	
449B: Hogansburg-----	85	Somewhat limited Frost action Depth to saturated zone	0.50 0.03	Very limited Depth to saturated zone Dense layer Cutbanks cave	1.00 0.50 0.10	Not limited	
449C: Hogansburg-----	85	Somewhat limited Slope Frost action Depth to saturated zone	0.63 0.50 0.03	Very limited Depth to saturated zone Slope Dense layer Cutbanks cave	1.00 0.63 0.50 0.10	Somewhat limited Slope	0.63
450B: Pyrities-----	80	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
450C: Pyrities-----	80	Somewhat limited Slope Frost action	0.63 0.50	Somewhat limited Slope Cutbanks cave	0.63 0.10	Somewhat limited Slope	0.63
450D: Pyrities-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
451B: Pyrities-----	80	Somewhat limited Frost action	0.50	Somewhat limited Cutbanks cave	0.10	Not limited	
451C: Pyrities-----	80	Somewhat limited Slope Frost action	0.63 0.50	Somewhat limited Slope Cutbanks cave	0.63 0.10	Somewhat limited Slope	0.63
451D: Pyrities-----	80	Very limited Slope Frost action	1.00 0.50	Very limited Slope Cutbanks cave	1.00 0.10	Very limited Slope	1.00
457: Mudgepond-----	80	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
458: Mudgepond-----	55	Very limited Depth to saturated zone Frost action	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave	1.00 1.00	Very limited Depth to saturated zone	1.00
Alden-----	35	Very limited Depth to saturated zone Frost action Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Large stones content	1.00 1.00 0.01
501: Ondawa-----	85	Very limited Flooding Frost action	1.00 0.50	Somewhat limited Flooding Cutbanks cave Depth to saturated zone	0.60 0.10 0.09	Somewhat limited Flooding	0.60
503: Rumney-----	80	Very limited Depth to saturated zone Frost action Flooding	1.00 1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Flooding	1.00 1.00 0.80	Very limited Flooding Depth to saturated zone	1.00 1.00

Table 18.—Building Site Development (Part 2)—Continued

Map symbol and soil name	Pct. of map unit	Local roads and streets		Shallow excavations		Lawns and landscaping	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
508: Medomak-----	85	Very limited		Very limited		Very limited	
		Depth to saturated zone	1.00	Depth to saturated zone	1.00	Flooding	1.00
		Frost action	1.00	Cutbanks cave	1.00	Depth to saturated zone	1.00
		Flooding	1.00	Ponding	1.00	Ponding	1.00
		Ponding	1.00	Flooding	0.80		

Table 19.--Sewage Disposal

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00
3: Ridgebury-----	40	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Seepage	1.00 1.00
Leicester-----	35	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00
Whitman-----	15	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Seepage Ponding	1.00 1.00 1.00
4: Leicester-----	80	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00
5: Wilbraham-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Seepage	1.00 0.50
6: Wilbraham-----	60	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Seepage	1.00 0.50
Menlo-----	25	Very limited Depth to saturated zone Ponding Slow water movement	1.00 1.00 0.50	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 0.50
7: Mudgepond-----	85	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Very limited Depth to saturated zone Seepage	1.00 1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
8: Mudgepond-----	45	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Very limited Depth to saturated zone Seepage	1.00 1.00
Alden-----	35	Very limited Depth to saturated zone Slow water movement Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 0.50
9: Scitico-----	40	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone	1.00
Shaker-----	30	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 1.00
Maybid-----	15	Very limited Slow water movement Depth to saturated zone Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
10: Raynham-----	80	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Seepage	1.00 0.32
12: Raypol-----	80	Very limited Depth to saturated zone Seepage Slow water movement	1.00 1.00 0.50	Very limited Seepage Depth to saturated zone	1.00 1.00
13: Walpole-----	80	Very limited Depth to saturated zone Seepage Filtering capacity	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
14: Fredon-----	85	Very limited Depth to saturated zone Seepage Slow water movement	1.00 1.00 0.50	Very limited Depth to saturated zone Seepage	1.00 1.00
15: Scarboro-----	80	Very limited Depth to saturated zone Seepage Filtering capacity Ponding	1.00 1.00 1.00 1.00	Very limited Seepage Depth to saturated zone Ponding Organic matter content	1.00 1.00 1.00 1.00
16: Halsey-----	80	Very limited Depth to saturated zone Seepage Ponding	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone Ponding	1.00 1.00 1.00
17: Timakwa-----	45	Very limited Depth to saturated zone Seepage Filtering capacity Ponding Flooding	1.00 1.00 1.00 1.00 0.40	Very limited Seepage Depth to saturated zone Ponding Organic matter content Flooding	1.00 1.00 1.00 1.00 0.40
Natchaug-----	40	Very limited Depth to saturated zone Ponding Slow water movement Flooding	1.00 1.00 0.68 0.40	Very limited Depth to saturated zone Seepage Ponding Organic matter content Flooding	1.00 1.00 1.00 1.00 1.00 0.40
18: Catden-----	40	Very limited Depth to saturated zone Subsidence Seepage Ponding Flooding	1.00 1.00 1.00 1.00 0.40	Very limited Organic matter content Depth to saturated zone Seepage Ponding Flooding	1.00 1.00 1.00 1.00 0.40

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
18: Freetown-----	40	Very limited Depth to saturated zone Subsidence Seepage Ponding Flooding	1.00 1.00 1.00 1.00 0.40	Very limited Organic matter content Depth to saturated zone Seepage Ponding Flooding	1.00 1.00 1.00 1.00 0.40
20A: Ellington-----	80	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone Slope	1.00 1.00 0.08
21A: Ninigret-----	60	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone Slope	1.00 1.00 0.08
Tisbury-----	25	Very limited Depth to saturated zone Seepage Slow water movement	1.00 1.00 0.50	Very limited Seepage Depth to saturated zone	1.00 1.00
22A: Hero-----	85	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
22B: Hero-----	85	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone Slope	1.00 1.00 0.92
23A: Sudbury-----	80	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone Slope	1.00 1.00 0.08

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
24A: Deerfield-----	80	Very limited Depth to saturated zone Seepage Filtering capacity	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
25A: Brancroft-----	80	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone	1.00
25B: Brancroft-----	80	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Slope	1.00 0.92
25C: Brancroft-----	80	Very limited Slow water movement Depth to saturated zone Slope	1.00 1.00 0.63	Very limited Slope Depth to saturated zone	1.00 1.00
26A: Berlin-----	80	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone	1.00
26B: Berlin-----	80	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Slope	1.00 0.92
27A: Belgrade-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Very limited Depth to saturated zone Seepage Slope	1.00 0.50 0.08
28A: Elmridge-----	80	Very limited Slow water movement Depth to saturated zone	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 0.75

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
28B: Elmridge-----	80	Very limited Slow water movement	1.00	Very limited Seepage	1.00
		Depth to saturated zone	1.00	Slope	0.92
				Depth to saturated zone	0.75
29A: Agawam-----	80	Very limited Seepage	1.00	Very limited Seepage	1.00
29B: Agawam-----	80	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
29C: Agawam-----	80	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
30A: Branford-----	80	Very limited Seepage	1.00	Very limited Seepage	1.00
30B: Branford-----	80	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.50
30C: Branford-----	80	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
31A: Copake-----	85	Very limited Seepage	1.00	Very limited Seepage	1.00
31B: Copake-----	85	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
31C: Copake-----	85	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
32A: Haven-----	60	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage	1.00
Enfield-----	25	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage	1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
32B: Haven-----	60	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage Slope	1.00 0.92
Enfield-----	25	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage Slope	1.00 0.92
32C: Haven-----	60	Very limited Seepage Slope Slow water movement	1.00 0.63 0.50	Very limited Slope Seepage	1.00 1.00
Enfield-----	25	Very limited Seepage Slope Slow water movement	1.00 0.63 0.50	Very limited Slope Seepage	1.00 1.00
33A: Hartford-----	80	Very limited Seepage	1.00	Very limited Seepage	1.00
33B: Hartford-----	80	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
34A: Merrimac-----	80	Very limited Seepage	1.00	Very limited Seepage	1.00
34B: Merrimac-----	80	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
34C: Merrimac-----	80	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
35A: Penwood-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00
35B: Penwood-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage Slope	1.00 0.92

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
36A: Windsor-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00
36B: Windsor-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage Slope	1.00 0.92
36C: Windsor-----	80	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.63	Very limited Slope Seepage	1.00 1.00
37A: Manchester-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00
37C: Manchester-----	80	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 1.00
37E: Manchester-----	80	Very limited Slope Seepage Filtering capacity	1.00 1.00 1.00	Very limited Slope Seepage	1.00 1.00
38A: Hinckley-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00
38C: Hinckley-----	80	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 1.00
38E: Hinckley-----	80	Very limited Slope Seepage Filtering capacity	1.00 1.00 1.00	Very limited Slope Seepage	1.00 1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
39A: Groton-----	85	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00
39C: Groton-----	85	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 1.00
39E: Groton-----	85	Very limited Slope Seepage Filtering capacity	1.00 1.00 1.00	Very limited Slope Seepage	1.00 1.00
40A: Ludlow-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Depth to saturated zone Seepage	0.75 0.50
40B: Ludlow-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Slope Depth to saturated zone Seepage	0.92 0.75 0.50
41B: Ludlow-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Depth to saturated zone Slope Seepage	0.75 0.68 0.50
42C: Ludlow-----	80	Very limited Depth to saturated zone Slow water movement Slope	1.00 0.50 0.04	Very limited Slope Depth to saturated zone Seepage	1.00 0.75 0.50
43A: Rainbow-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Depth to saturated zone Seepage	0.75 0.50

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
43B: Rainbow-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone Seepage	0.75 0.50
44B: Rainbow-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone Seepage	0.75 0.50
45A: Woodbridge-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
45B: Woodbridge-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone Seepage	0.75 0.50
45C: Woodbridge-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
46B: Woodbridge-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.75
		Slow water movement	0.50	Slope	0.68
				Seepage	0.50
46C: Woodbridge-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
47C: Woodbridge-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	0.50	Depth to saturated zone	0.75
		Slope	0.04	Seepage	0.50
48B: Georgia-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Slow water movement	1.00	Slope	0.68
				Seepage	0.50
Amenia-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Slow water movement	1.00	Slope	0.68
				Seepage	0.50
48C: Georgia-----	50	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Depth to saturated zone	1.00
		Slope	0.63	Seepage	0.50
Amenia-----	35	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Depth to saturated zone	1.00
		Slope	0.63	Seepage	0.50
49B: Georgia-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Slow water movement	1.00	Slope	0.92
				Seepage	0.50
Amenia-----	35	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Slow water movement	1.00	Slope	0.92
				Seepage	0.50
49C: Georgia-----	50	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Depth to saturated zone	1.00
		Slope	0.63	Seepage	0.50

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
49C: Amenia-----	35	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Depth to saturated zone	1.00
		Slope	0.63	Seepage	0.50
50A: Sutton-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
50B: Sutton-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
				Slope	0.92
51B: Sutton-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
				Slope	0.68
52C: Sutton-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
		Slope	0.04	Slope	1.00
53A: Wapping-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
53B: Wapping-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
				Slope	0.92
54B: Wapping-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
				Slope	0.68
55A: Watchaug-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
		Slow water movement	0.50		

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
55B: Watchaug-----	80	Very limited Depth to saturated zone Seepage Slow water movement	1.00 1.00 0.50	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.92
56B: Watchaug-----	80	Very limited Depth to saturated zone Seepage Slow water movement	1.00 1.00 0.50	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.68
57B: Gloucester-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage Slope	1.00 0.92
57C: Gloucester-----	80	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.63	Very limited Slope Seepage	1.00 1.00
57D: Gloucester-----	80	Very limited Slope Seepage Filtering capacity	1.00 1.00 1.00	Very limited Slope Seepage	1.00 1.00
58B: Gloucester-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage Slope	1.00 0.92
58C: Gloucester-----	80	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.63	Very limited Slope Seepage	1.00 1.00
59C: Gloucester-----	80	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
59D: Gloucester-----	80	Very limited Slope Seepage Filtering capacity	1.00 1.00 1.00	Very limited Slope Seepage	1.00 1.00
60B: Canton-----	45	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
Charlton-----	35	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
60C: Canton-----	45	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
Charlton-----	35	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
60D: Canton-----	45	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Charlton-----	35	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
61B: Canton-----	45	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
Charlton-----	35	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
61C: Canton-----	45	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
Charlton-----	35	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
62C: Canton-----	45	Very limited Seepage Slope	1.00 0.04	Very limited Seepage Slope	1.00 1.00
Charlton-----	35	Very limited Seepage Slope	1.00 0.04	Very limited Slope Seepage	1.00 1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
62D: Canton-----	45	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Charlton-----	35	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
63B: Cheshire-----	80	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
63C: Cheshire-----	80	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
63D: Cheshire-----	80	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
64B: Cheshire-----	80	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
64C: Cheshire-----	80	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
65C: Cheshire-----	80	Very limited Seepage Slope	1.00 0.04	Very limited Slope Seepage	1.00 1.00
65D: Cheshire-----	80	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
66B: Narragansett-----	80	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage Slope	1.00 0.68
66C: Narragansett-----	80	Very limited Seepage Slope Slow water movement	1.00 0.63 0.50	Very limited Slope Seepage	1.00 1.00
67B: Narragansett-----	80	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage Slope	1.00 0.92

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
67C: Narragansett-----	80	Very limited Seepage Slope Slow water movement	1.00 0.63 0.50	Very limited Slope Seepage	1.00 1.00
68C: Narragansett-----	80	Very limited Seepage Slow water movement Slope	1.00 0.50 0.04	Very limited Seepage Slope	1.00 1.00
68D: Narragansett-----	80	Very limited Slope Seepage Slow water movement	1.00 1.00 0.50	Very limited Slope Seepage	1.00 1.00
69B: Yalesville-----	75	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to hard bedrock Seepage Slope	1.00 1.00 0.92
69C: Yalesville-----	75	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.63	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
70C: Branford-----	50	Very limited Seepage Slope	1.00 0.04	Very limited Seepage Slope	1.00 1.00
Holyoke-----	30	Very limited Depth to bedrock Slope	1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 0.50
71C: Brookfield-----	45	Very limited Seepage Slope	1.00 0.04	Very limited Slope Seepage	1.00 1.00
Brimfield-----	30	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
Rock outcrop-----	15	Not rated		Not rated	

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
71E: Brookfield-----	45	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
71E: Brimfield-----	30	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
Rock outcrop-----	15	Not rated		Not rated	
73C: Charlton-----	45	Very limited Seepage Slope	1.00 0.04	Very limited Slope Seepage	1.00 1.00
Chatfield-----	30	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
73E: Charlton-----	45	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Chatfield-----	30	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
74C: Narragansett-----	55	Very limited Seepage Slow water movement Slope	1.00 0.50 0.04	Very limited Seepage Slope	1.00 1.00
Hollis-----	20	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
75C: Hollis-----	35	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
Chatfield-----	30	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
75C: Rock outcrop-----	15	Not rated		Not rated	
75E: Hollis-----	35	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
Chatfield-----	30	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
Rock outcrop-----	15	Not rated		Not rated	
76E: Rock outcrop-----	55	Not rated		Not rated	
Hollis-----	25	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
76F: Rock outcrop-----	55	Not rated		Not rated	
Hollis-----	25	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
77C: Cheshire-----	45	Very limited Seepage	1.00	Very limited Slope	1.00
		Slope	0.04	Seepage	1.00
Holyoke-----	35	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	0.04	Slope	1.00
				Seepage	0.50
77D: Cheshire-----	45	Very limited Slope	1.00	Very limited Slope	1.00
		Seepage	1.00	Seepage	1.00
Holyoke-----	35	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
				Seepage	0.50

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
78C: Holyoke-----	50	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	0.04	Slope Seepage	1.00 0.50
Rock outcrop-----	25	Not rated		Not rated	
78E: Holyoke-----	50	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope Seepage	1.00 0.50
Rock outcrop-----	25	Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated	
Holyoke-----	25	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope Seepage	1.00 0.50
80B: Bernardston-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone Seepage	0.92 0.50
80C: Bernardston-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Depth to saturated zone Seepage	0.92 0.50
		Slow water movement	0.50		
81C: Bernardston-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	0.50	Depth to saturated zone Seepage	0.92 0.50
		Slope	0.04		
81D: Bernardston-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to saturated zone Seepage	0.92 0.50
		Slow water movement	0.50		

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
82B: Broadbrook-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Depth to saturated zone Slope Seepage	0.75 0.68 0.50
82C: Broadbrook-----	80	Very limited Depth to saturated zone Slope Slow water movement	1.00 0.63 0.50	Very limited Slope Depth to saturated zone Seepage	1.00 0.75 0.50
82D: Broadbrook-----	80	Very limited Depth to saturated zone Slope Slow water movement	1.00 1.00 0.50	Very limited Slope Depth to saturated zone Seepage	1.00 0.75 0.50
83B: Broadbrook-----	80	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Slope Depth to saturated zone Seepage	0.92 0.75 0.50
83C: Broadbrook-----	80	Very limited Depth to saturated zone Slope Slow water movement	1.00 0.63 0.50	Very limited Slope Depth to saturated zone Seepage	1.00 0.75 0.50
84B: Paxton-----	55	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Slope Depth to saturated zone Seepage	0.92 0.75 0.50
Montauk-----	30	Very limited Depth to saturated zone	1.00	Very limited Seepage Slope Depth to saturated zone	1.00 0.92 0.44

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
84C: Paxton-----	55	Very limited		Very limited	
		Depth to saturated zone	1.00	Slope	1.00
		Slope	0.63	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
Montauk-----	30	Very limited		Very limited	
		Depth to saturated zone	1.00	Slope	1.00
		Slope	0.63	Seepage	1.00
				Depth to saturated zone	0.44
84D: Paxton-----	55	Very limited		Very limited	
		Depth to saturated zone	1.00	Slope	1.00
		Slope	1.00	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
Montauk-----	30	Very limited		Very limited	
		Depth to saturated zone	1.00	Slope	1.00
		Slope	1.00	Seepage	1.00
				Depth to saturated zone	0.44
85B: Paxton-----	55	Very limited		Somewhat limited	
		Depth to saturated zone	1.00	Slope	0.92
		Slow water movement	0.50	Depth to saturated zone	0.75
				Seepage	0.50
85B: Montauk-----	30	Very limited		Very limited	
		Depth to saturated zone	1.00	Seepage	1.00
				Slope	0.92
				Depth to saturated zone	0.44
85C: Paxton-----	55	Very limited		Very limited	
		Depth to saturated zone	1.00	Slope	1.00
		Slope	0.63	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
85C: Montauk-----	30	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage Depth to saturated zone	1.00 0.44
86C: Paxton-----	55	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	0.50	Depth to saturated zone	0.75
		Slope	0.04	Seepage	0.50
Montauk-----	30	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.04	Seepage Depth to saturated zone	1.00 0.44
86D: Paxton-----	55	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
Montauk-----	30	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Seepage Depth to saturated zone	1.00 0.44
87B: Wethersfield-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone	0.75
				Seepage	0.50
87C: Wethersfield-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
87D: Wethersfield-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
88B: Wethersfield-----	80	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone	0.75
				Seepage	0.50
88C: Wethersfield-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
89C: Wethersfield-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	0.50	Depth to saturated zone	0.75
		Slope	0.04	Seepage	0.50
89D: Wethersfield-----	80	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
90B: Stockbridge-----	80	Very limited Slow water movement	1.00	Somewhat limited Slope	0.92
				Seepage	0.50
90C: Stockbridge-----	80	Very limited Slow water movement	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage	0.50
90D: Stockbridge-----	80	Very limited Slope	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Seepage	0.50

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
91B: Stockbridge-----	80	Very limited Slow water movement	1.00	Somewhat limited Slope Seepage	0.92 0.50
91C: Stockbridge-----	80	Very limited Slow water movement Slope	1.00 0.63	Very limited Slope Seepage	1.00 0.50
91D: Stockbridge-----	80	Very limited Slope Slow water movement	1.00 1.00	Very limited Slope Seepage	1.00 0.50
92B: Nellis-----	85	Not limited		Very limited Seepage Slope	0.99 0.92
92C: Nellis-----	85	Somewhat limited Slope	0.63	Very limited Slope Seepage	1.00 0.99
92D: Nellis-----	85	Very limited Slope	1.00	Very limited Slope Seepage	1.00 0.99
93C: Nellis-----	85	Somewhat limited Slope	0.04	Very limited Slope Seepage	1.00 0.99
94C: Farmington-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
Nellis-----	35	Somewhat limited Slope	0.04	Very limited Slope Seepage	1.00 0.99
94E: Farmington-----	40	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
Nellis-----	35	Very limited Slope	1.00	Very limited Slope Seepage	1.00 0.99

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
95C: Farmington-----	60	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Seepage Slope	1.00 0.04	Slope Seepage	1.00 1.00
Rock outcrop-----	20	Not rated		Not rated	
95E: Farmington-----	60	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope Seepage	1.00 1.00	Slope Seepage	1.00 1.00
Rock outcrop-----	20	Not rated		Not rated	
96: Ipswich-----	85	Very limited Flooding Ponding Depth to saturated zone Filtering capacity Subsidence	1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Organic matter content Depth to saturated zone Seepage	1.00 1.00 1.00 1.00 1.00
97: Pawcatuck-----	85	Very limited Flooding Ponding Depth to saturated zone Seepage Filtering capacity	1.00 1.00 1.00 1.00 1.00	Very limited Ponding Flooding Seepage Depth to saturated zone Organic matter content	1.00 1.00 1.00 1.00 1.00
98: Westbrook-----	80	Very limited Flooding Ponding Depth to saturated zone Subsidence Slow water movement	1.00 1.00 1.00 1.00 0.82	Very limited Ponding Flooding Depth to saturated zone Seepage Organic matter content	1.00 1.00 1.00 1.00 1.00
99: Westbrook, low salt-	80	Very limited Flooding Ponding Depth to saturated zone Subsidence Slow water movement	1.00 1.00 1.00 1.00 0.82	Very limited Ponding Flooding Depth to saturated zone Seepage Organic matter content	1.00 1.00 1.00 1.00 1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
100: Suncook-----	80	Very limited Flooding Seepage Filtering capacity Depth to saturated zone	1.00 1.00 1.00 0.24	Very limited Flooding Seepage	1.00 1.00
101: Occum-----	80	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 0.24	Very limited Flooding Seepage	1.00 1.00
102: Pootatuck-----	80	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
103: Rippowam-----	80	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
104: Bash-----	80	Very limited Flooding Depth to saturated zone Slow water movement	1.00 1.00 0.68	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 0.50
105: Hadley-----	80	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 0.08	Very limited Flooding Seepage	1.00 1.00
106: Winooski-----	80	Very limited Flooding Depth to saturated zone Seepage Slow water movement	1.00 1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
107: Limerick-----	50	Very limited Flooding Depth to saturated zone Slow water movement	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 0.50
Lim-----	30	Very limited Flooding Depth to saturated zone Seepage Slow water movement	1.00 1.00 1.00 0.50	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
108: Saco-----	80	Very limited Flooding Depth to saturated zone Seepage Ponding Slow water movement	1.00 1.00 1.00 1.00 0.50	Very limited Flooding Seepage Depth to saturated zone Ponding	1.00 1.00 1.00 1.00
109: Fluvaquents, Frequently Flooded-	50	Very limited Flooding Depth to saturated zone Seepage Slow water movement	1.00 1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00
Udifluents, Frequently Flooded-	35	Very limited Flooding Filtering capacity Seepage	1.00 1.00 1.00	Very limited Flooding Seepage	1.00 1.00
221A: Ninigret-----	40	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone Slope	1.00 1.00 0.08
Urban land-----	35	Not rated		Not rated	

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
224A: Deerfield-----	40	Very limited Depth to saturated zone	1.00	Very limited Seepage	1.00
		Seepage	1.00	Depth to saturated zone	1.00
		Filtering capacity	1.00		
Urban land-----	35	Not rated		Not rated	
225B: Brancroft-----	40	Very limited Slow water movement	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Slope	0.32
Urban land-----	35	Not rated		Not rated	
226B: Berlin-----	40	Very limited Slow water movement	1.00	Very limited Depth to saturated zone	1.00
		Depth to saturated zone	1.00	Slope	0.32
Urban land-----	35	Not rated		Not rated	
228B: Elmridge-----	40	Very limited Slow water movement	1.00	Very limited Seepage	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	0.75
				Slope	0.32
Urban land-----	35	Not rated		Not rated	
229B: Agawam-----	40	Very limited Seepage	1.00	Very limited Seepage	1.00
				Slope	0.32
Urban land-----	35	Not rated		Not rated	
229C: Agawam-----	40	Very limited Seepage	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage	1.00
Urban land-----	35	Not rated		Not rated	
230B: Branford-----	40	Very limited Seepage	1.00	Very limited Seepage	1.00
				Slope	0.32
Urban land-----	35	Not rated		Not rated	

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
230C: Branford-----	40	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
Urban land-----	35	Not rated		Not rated	
232B: Haven-----	40	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage Slope	1.00 0.32
Urban land-----	35	Not rated		Not rated	
234B: Merrimac-----	40	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.32
Urban land-----	35	Not rated		Not rated	
235B: Penwood-----	40	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage Slope	1.00 0.32
Urban land-----	35	Not rated		Not rated	
236B: Windsor-----	40	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage Slope	1.00 0.32
Urban land-----	35	Not rated		Not rated	
237A: Manchester-----	40	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00
Urban land-----	35	Not rated		Not rated	
237C: Manchester-----	40	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 1.00
Urban land-----	35	Not rated		Not rated	
238A: Hinckley-----	40	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
238A: Urban land-----	35	Not rated		Not rated	
238C: Hinckley-----	40	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 1.00
Urban land-----	35	Not rated		Not rated	
240B: Ludlow-----	40	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Depth to saturated zone Seepage Slope	0.75 0.50 0.32
Urban land-----	35	Not rated		Not rated	
243B: Rainbow-----	40	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Depth to saturated zone Seepage Slope	0.75 0.50 0.32
Urban land-----	35	Not rated		Not rated	
245B: Woodbridge-----	40	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Depth to saturated zone Seepage Slope	0.75 0.50 0.32
Urban land-----	35	Not rated		Not rated	
245C: Woodbridge-----	40	Very limited Depth to saturated zone Slope Slow water movement	1.00 0.63 0.50	Very limited Slope Depth to saturated zone Seepage	1.00 0.75 0.50
Urban land-----	35	Not rated		Not rated	
248B: Georgia-----	40	Very limited Depth to saturated zone Slow water movement	1.00 1.00	Very limited Depth to saturated zone Slope Seepage	1.00 0.68 0.50

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
248B: Urban land-----	35	Not rated		Not rated	
250B: Sutton-----	40	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.32
250B: Urban land-----	35	Not rated		Not rated	
253B: Wapping-----	40	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.32
Urban land-----	35	Not rated		Not rated	
255B: Watchaug-----	40	Very limited Depth to saturated zone Seepage Slow water movement	1.00 1.00 0.50	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.32
Urban land-----	35	Not rated		Not rated	
260B: Charlton-----	40	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
Urban land-----	35	Not rated		Not rated	
260C: Charlton-----	40	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
Urban land-----	35	Not rated		Not rated	
260D: Charlton-----	40	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Urban land-----	35	Not rated		Not rated	
263B: Cheshire-----	40	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
Urban land-----	35	Not rated		Not rated	

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
263C: Cheshire-----	40	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
Urban land-----	35	Not rated		Not rated	
266B: Narragansett-----	40	Very limited Seepage Slow water movement	1.00 0.50	Very limited Seepage Slope	1.00 0.92
Urban land-----	35	Not rated		Not rated	
269B: Yalesville-----	40	Very limited Depth to bedrock Seepage	1.00 1.00	Very limited Depth to hard bedrock Seepage Slope	1.00 1.00 0.92
Urban land-----	35	Not rated		Not rated	
269C: Yalesville-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.63	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
Urban land-----	35	Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Very limited Seepage Slope	1.00 0.04	Very limited Slope Seepage	1.00 1.00
Chatfield-----	15	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.04	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
273E: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Chatfield-----	15	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
275C: Urban land-----	45	Not rated		Not rated	
275C: Chatfield-----	30	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Seepage	1.00	Slope	1.00
		Slope	0.04	Seepage	1.00
275E: Urban land-----	35	Not rated		Not rated	
Chatfield-----	25	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
Rock outcrop-----	15	Not rated		Not rated	
282B: Broadbrook-----	40	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone	0.75
				Seepage	0.50
Urban land-----	35	Not rated		Not rated	
284B: Paxton-----	40	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone	0.75
				Seepage	0.50
Urban land-----	35	Not rated		Not rated	
284C: Paxton-----	40	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
Urban land-----	35	Not rated		Not rated	
284D: Paxton-----	40	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
284D: Urban land-----	35	Not rated		Not rated	
287B: Wethersfield-----	40	Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.92
		Slow water movement	0.50	Depth to saturated zone Seepage	0.75 0.50
Urban land-----	35	Not rated		Not rated	
287C: Wethersfield-----	40	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Depth to saturated zone Seepage	0.75 0.50
		Slow water movement	0.50		
Urban land-----	35	Not rated		Not rated	
287D: Wethersfield-----	40	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to saturated zone Seepage	0.75 0.50
		Slow water movement	0.50		
Urban land-----	35	Not rated		Not rated	
290B: Stockbridge-----	40	Very limited Slow water movement	1.00	Somewhat limited Slope	0.92
				Seepage	0.50
Urban land-----	35	Not rated		Not rated	
290C: Stockbridge-----	40	Very limited Slow water movement	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage	0.50
Urban land-----	35	Not rated		Not rated	
290D: Stockbridge-----	40	Very limited Slope	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Seepage	0.50
Urban land-----	35	Not rated		Not rated	

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
301: Beaches-----	50	Not rated		Not rated	
301: Udipsamments-----	35	Very limited Flooding Seepage Filtering capacity Depth to saturated zone Slope	1.00 1.00 1.00 0.43 0.01	Very limited Flooding Seepage Slope	1.00 1.00 1.00
302: Dumps-----	95	Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated	
304: Udorthents-----	90	Very limited Slope Seepage Depth to saturated zone	1.00 1.00 0.50	Very limited Slope Seepage	1.00 1.00
305: Udorthents-----	65	Very limited Depth to saturated zone Slope Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Slope Seepage	1.00 1.00 1.00
Pits-----	25	Very limited Filtering capacity Seepage Slope	1.00 1.00 1.00	Very limited Seepage Slope	1.00 1.00
306: Udorthents-----	50	Very limited Slope Seepage Depth to saturated zone	1.00 1.00 0.50	Very limited Slope Seepage	1.00 1.00
Urban land-----	35	Not rated		Not rated	
307: Urban land-----	80	Not rated		Not rated	
308: Udorthents-----	80	Very limited Depth to saturated zone Slope Seepage	1.00 1.00 1.00	Very limited Depth to saturated zone Slope Seepage	1.00 1.00 1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
309: Udorthents-----	80	Very limited Depth to saturated zone Slope Seepage Flooding	1.00 1.00 1.00 0.40	Very limited Depth to saturated zone Slope Seepage Flooding	1.00 1.00 1.00 0.40
310: Udorthents, Periodically Flooded-----	85	Very limited Flooding Depth to saturated zone Slope Seepage	1.00 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Slope Seepage	1.00 1.00 1.00 1.00
401C: Macomber-----	55	Very limited Depth to bedrock Slope Slow water movement	1.00 0.63 0.50	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 0.50
Taconic-----	30	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 1.00
402D: Macomber-----	50	Very limited Depth to bedrock Slope Slow water movement	1.00 1.00 0.50	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 0.50
Taconic-----	25	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope	1.00 1.00
Rock outcrop-----	15	Not rated		Not rated	
403C: Taconic-----	70	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.01	Very limited Depth to hard bedrock Slope	1.00 1.00
Rock outcrop-----	25	Not rated		Not rated	

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
403E: Taconic-----	70	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00		
Rock outcrop-----	20	Not rated		Not rated	
403F: Taconic-----	70	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00		
Rock outcrop-----	20	Not rated		Not rated	
405C: Dummerston-----	85	Somewhat limited Slow water movement	0.50	Very limited Slope	1.00
		Slope	0.01	Seepage	0.50
405E: Dummerston-----	85	Very limited Slope	1.00	Very limited Slope	1.00
		Slow water movement	0.50	Seepage	0.50
407C: Lanesboro-----	85	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slow water movement	0.50	Depth to saturated zone	0.75
		Slope	0.16	Seepage	0.50
407E: Lanesboro-----	85	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Depth to saturated zone	0.75
		Slow water movement	0.50	Seepage	0.50
408C: Fullam-----	85	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.16	Depth to saturated zone	0.64
				Seepage	0.50
409B: Brayton-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
				Seepage	1.00
				Slope	0.68

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
412B: Bice-----	85	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
412C: Bice-----	85	Very limited Seepage Slope	1.00 0.16	Very limited Slope Seepage	1.00 1.00
412D: Bice-----	85	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
413C: Bice-----	45	Very limited Seepage Slope	1.00 0.01	Very limited Slope Seepage	1.00 1.00
Millsite-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.01	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
413E: Bice-----	45	Very limited Slope Seepage	1.00 1.00	Very limited Slope Seepage	1.00 1.00
Millsite-----	40	Very limited Depth to bedrock Slope Seepage	1.00 1.00 1.00	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
414: Fredon, cold-----	85	Very limited Depth to saturated zone Seepage Slow water movement	1.00 1.00 0.50	Very limited Depth to saturated zone Seepage	1.00 1.00
415C: Millsite-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.01	Very limited Depth to hard bedrock Slope Seepage	1.00 1.00 1.00
415C: Westminster-----	40	Very limited Depth to bedrock Seepage Slope	1.00 1.00 0.01	Very limited Depth to hard bedrock Seepage Slope	1.00 1.00 1.00
Rock outcrop-----	15	Not rated		Not rated	

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
415E: Millsite-----	40	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
Westminster-----	40	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
Rock outcrop-----	15	Not rated		Not rated	
416E: Rock outcrop-----	70	Not rated		Not rated	
Westminster-----	20	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
416F: Rock outcrop-----	70	Not rated		Not rated	
Westminster-----	20	Very limited Depth to bedrock	1.00	Very limited Depth to hard bedrock	1.00
		Slope	1.00	Slope	1.00
		Seepage	1.00	Seepage	1.00
417B: Bice-----	85	Very limited Seepage	1.00	Very limited Seepage	1.00
				Slope	0.92
417C: Bice-----	85	Very limited Seepage	1.00	Very limited Slope	1.00
		Slope	0.16	Seepage	1.00
417D: Bice-----	85	Very limited Slope	1.00	Very limited Slope	1.00
		Seepage	1.00	Seepage	1.00
418C: Schroon-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00
		Slope	0.04	Slope	1.00
420A: Schroon-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Seepage	1.00	Seepage	1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
420B: Schroon-----	85	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.68
421A: Ninigret, cold-----	85	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
423A: Sudbury, cold-----	85	Very limited Depth to saturated zone Seepage	1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
424B: Shelburne-----	85	Very limited Depth to saturated zone	1.00	Very limited Seepage Slope Depth to saturated zone	1.00 0.92 0.75
424C: Shelburne-----	85	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Seepage Depth to saturated zone	1.00 1.00 0.75
424D: Shelburne-----	85	Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope Seepage Depth to saturated zone	1.00 1.00 0.75
425B: Shelburne-----	85	Very limited Depth to saturated zone	1.00	Very limited Seepage Slope Depth to saturated zone	1.00 0.92 0.75
425C: Shelburne-----	85	Very limited Depth to saturated zone Slope	1.00 0.63	Very limited Slope Seepage Depth to saturated zone	1.00 1.00 0.75

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
426D: Shelburne-----	85	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	1.00	Seepage Depth to saturated zone	1.00 0.75
427B: Ashfield-----	85	Very limited Slow water movement	1.00	Very limited Seepage	1.00
		Depth to saturated zone	1.00	Depth to saturated zone	0.99
				Slope	0.92
427C: Ashfield-----	85	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage Depth to saturated zone	1.00 0.99
428A: Ashfield-----	85	Very limited Depth to saturated zone	1.00	Very limited Seepage	1.00
				Depth to saturated zone	0.99
428B: Ashfield-----	85	Very limited Depth to saturated zone	1.00	Very limited Seepage	1.00
				Depth to saturated zone	0.99
				Slope	0.92
428C: Ashfield-----	85	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage Depth to saturated zone	1.00 0.99
429A: Agawam, cold-----	80	Very limited Seepage	1.00	Very limited Seepage	1.00
429B: Agawam, cold-----	80	Very limited Seepage	1.00	Very limited Seepage	1.00
				Slope	0.92
429C: Agawam, cold-----	80	Very limited Seepage	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage	1.00

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
433: Moosilauke-----	80	Very limited Depth to saturated zone Seepage Filtering capacity	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone	1.00 1.00
434A: Merrimac, cold-----	80	Very limited Seepage	1.00	Very limited Seepage	1.00
434B: Merrimac, cold-----	80	Very limited Seepage	1.00	Very limited Seepage Slope	1.00 0.92
434C: Merrimac, cold-----	80	Very limited Seepage Slope	1.00 0.63	Very limited Slope Seepage	1.00 1.00
435: Scarboro-----	80	Very limited Depth to saturated zone Seepage Filtering capacity Ponding	1.00 1.00 1.00 1.00	Very limited Seepage Depth to saturated zone Ponding Organic matter content	1.00 1.00 1.00 1.00
436: Halsey-----	80	Very limited Depth to saturated zone Seepage Ponding	1.00 1.00 1.00	Very limited Seepage Depth to saturated zone Ponding	1.00 1.00 1.00
437: Wonsqueak-----	85	Very limited Depth to saturated zone Ponding Slow water movement Flooding	1.00 1.00 0.68 0.40	Very limited Depth to saturated zone Seepage Ponding Organic matter content Flooding	1.00 1.00 1.00 1.00 1.00 0.40
438: Bucksport-----	85	Very limited Depth to saturated zone Subsidence Seepage Ponding Flooding	1.00 1.00 1.00 1.00 0.40	Very limited Depth to saturated zone Seepage Ponding Organic matter content Flooding	1.00 1.00 1.00 1.00 1.00 0.40

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
440A: Boscawen-----	80	Very limited Seepage Filtering capacity	1.00 1.00	Very limited Seepage	1.00
440C: Boscawen-----	80	Very limited Seepage Filtering capacity Slope	1.00 1.00 0.04	Very limited Seepage Slope	1.00 1.00
440E: Boscawen-----	85	Very limited Slope Seepage Filtering capacity	1.00 1.00 1.00	Very limited Slope Seepage	1.00 1.00
442: Brayton-----	85	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.68
443: Brayton-----	50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Seepage Slope	1.00 1.00 0.68
Loonmeadow-----	35	Very limited Depth to saturated zone Slow water movement Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Seepage Ponding	1.00 1.00 1.00
448B: Hogansburg-----	85	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Slope Seepage Depth to saturated zone	0.68 0.50 0.44
449B: Hogansburg-----	85	Very limited Depth to saturated zone Slow water movement	1.00 0.50	Somewhat limited Slope Seepage Depth to saturated zone	0.92 0.50 0.44

Table 19.—Sewage Disposal—Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
449C: Hogansburg-----	85	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage	0.50
		Slow water movement	0.50	Depth to saturated zone	0.44
450B: Pyrities-----	80	Very limited Slow water movement	1.00	Somewhat limited Slope	0.92
				Seepage	0.50
450C: Pyrities-----	80	Very limited Slow water movement	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage	0.50
450D: Pyrities-----	80	Very limited Slope	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Seepage	0.50
451B: Pyrities-----	80	Very limited Slow water movement	1.00	Somewhat limited Slope	0.92
				Seepage	0.50
451C: Pyrities-----	80	Very limited Slow water movement	1.00	Very limited Slope	1.00
		Slope	0.63	Seepage	0.50
451D: Pyrities-----	80	Very limited Slope	1.00	Very limited Slope	1.00
		Slow water movement	1.00	Seepage	0.50
457: Mudgepond-----	80	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Slow water movement	0.50	Seepage	1.00
458: Mudgepond-----	55	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
		Slow water movement	0.50	Seepage	1.00

Table 19.--Sewage Disposal--Continued

Map symbol and soil name	Pct. of map unit	Septic tank absorption fields		Sewage lagoons	
		Rating class and limiting features	Value	Rating class and limiting features	Value
458: Alden-----	35	Very limited Depth to saturated zone Slow water movement Ponding	1.00 1.00 1.00	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 0.50
501: Ondawa-----	85	Very limited Flooding Slow water movement Depth to saturated zone	1.00 1.00 0.24	Very limited Flooding Seepage	1.00 0.50
503: Rumney-----	80	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Flooding Seepage Depth to saturated zone	1.00 1.00 1.00
508: Medomak-----	85	Very limited Flooding Depth to saturated zone Seepage Ponding Slow water movement	1.00 1.00 1.00 1.00 0.50	Very limited Flooding Seepage Depth to saturated zone Ponding	1.00 1.00 1.00 1.00

Table 20.—Source of Sand and Gravel

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
2: Ridgebury-----	80	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
3: Ridgebury-----	40	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
Leicester-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
Whitman-----	15	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
4: Leicester-----	80	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
5: Wilbraham-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
6: Wilbraham-----	60	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
Menlo-----	25	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
7: Mudgepond-----	85	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.21	Thickest layer	0.00
8: Mudgepond-----	45	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.21	Thickest layer	0.00
9: Scitico-----	40	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
9: Shaker-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Maybid-----	15	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Alden-----	35	Fair Bottom layer Thickest layer	0.00 0.14	Poor Bottom layer Thickest layer	0.00 0.00
10: Raynham-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
12: Raypol-----	80	Fair Thickest layer Bottom layer	0.14 0.14	Fair Thickest layer Bottom layer	0.09 0.10
13: Walpole-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Fair Bottom layer Thickest layer	0.15 0.15
14: Fredon-----	85	Fair Thickest layer Bottom layer	0.14 0.14	Fair Thickest layer Bottom layer	0.08 0.43
15: Scarboro-----	80	Fair Thickest layer Bottom layer	0.00 0.14	Fair Thickest layer Bottom layer	0.31 0.36
16: Halsey-----	80	Fair Thickest layer Bottom layer	0.00 0.36	Fair Thickest layer Bottom layer	0.00 0.31
17: Timakwa-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Natchaug-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
18: Catden-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Organic matter content Thickest layer Bottom layer	0.00 0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
18: Freetown-----	40	Poor Bottom layer	0.00	Poor Organic matter content	0.00
		Thickest layer	0.00	Thickest layer	0.00
				Bottom layer	0.00
20A: Ellington-----	80	Fair Thickest layer	0.00	Fair Thickest layer	0.00
		Bottom layer	0.14	Bottom layer	0.11
21A: Ninigret-----	60	Poor Thickest layer	0.00	Fair Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.03
Tisbury-----	25	Poor Thickest layer	0.00	Fair Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.10
22A: Hero-----	85	Fair Thickest layer	0.29	Fair Thickest layer	0.00
		Bottom layer	0.50	Bottom layer	0.11
22B: Hero-----	85	Fair Thickest layer	0.29	Fair Thickest layer	0.00
		Bottom layer	0.50	Bottom layer	0.11
23A: Sudbury-----	80	Fair Thickest layer	0.00	Fair Thickest layer	0.00
		Bottom layer	0.43	Bottom layer	0.10
24A: Deerfield-----	80	Poor Bottom layer	0.00	Fair Thickest layer	0.10
		Thickest layer	0.00	Bottom layer	0.13
25A: Brancroft-----	80	Poor Bottom layer	0.00	Poor Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
25B: Brancroft-----	80	Poor Bottom layer	0.00	Poor Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
25C: Brancroft-----	80	Poor Bottom layer	0.00	Poor Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
26A: Berlin-----	80	Poor Bottom layer	0.00	Poor Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
26B: Berlin-----	80	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
27A: Belgrade-----	80	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
28A: Elmridge-----	80	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
28B: Elmridge-----	80	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
29A: Agawam-----	80	Poor Thickest layer Bottom layer	 0.00 0.00	Fair Thickest layer Bottom layer	 0.00 0.12
29B: Agawam-----	80	Poor Thickest layer Bottom layer	 0.00 0.00	Fair Thickest layer Bottom layer	 0.00 0.12
29C: Agawam-----	80	Poor Thickest layer Bottom layer	 0.00 0.00	Fair Thickest layer Bottom layer	 0.00 0.12
30A: Branford-----	80	Poor Thickest layer Bottom layer	 0.00 0.07	Fair Thickest layer Bottom layer	 0.00 0.12
30B: Branford-----	80	Poor Thickest layer Bottom layer	 0.00 0.07	Fair Thickest layer Bottom layer	 0.00 0.12
30C: Branford-----	80	Poor Thickest layer Bottom layer	 0.00 0.07	Fair Thickest layer Bottom layer	 0.00 0.12
31A: Copake-----	85	Fair Thickest layer Bottom layer	 0.43 0.43	Fair Bottom layer Thickest layer	 0.09 0.29
31B: Copake-----	85	Fair Thickest layer Bottom layer	 0.43 0.43	Fair Bottom layer Thickest layer	 0.09 0.29

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
31C: Copake-----	85	Fair		Fair	
		Thickest layer	0.43	Bottom layer	0.09
		Bottom layer	0.43	Thickest layer	0.29
32A: Haven-----	60	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.39
Enfield-----	25	Poor		Fair	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.50
32B: Haven-----	60	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.39
Enfield-----	25	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.50
32C: Haven-----	60	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.39
Enfield-----	25	Poor		Fair	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.50
33A: Hartford-----	80	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.10
33B: Hartford-----	80	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.10
34A: Merrimac-----	80	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.29	Bottom layer	0.12
34B: Merrimac-----	80	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.29	Bottom layer	0.12
34C: Merrimac-----	80	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.29	Bottom layer	0.12
35A: Penwood-----	80	Poor		Fair	
		Bottom layer	0.00	Bottom layer	0.15
		Thickest layer	0.00	Thickest layer	0.15

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
35B: Penwood-----	80	Poor		Fair	
		Bottom layer	0.00	Bottom layer	0.15
		Thickest layer	0.00	Thickest layer	0.15
36A: Windsor-----	80	Poor		Fair	
		Bottom layer	0.00	Thickest layer	0.06
		Thickest layer	0.00	Bottom layer	0.10
36B: Windsor-----	80	Poor		Fair	
		Bottom layer	0.00	Thickest layer	0.06
		Thickest layer	0.00	Bottom layer	0.10
36C: Windsor-----	80	Poor		Fair	
		Bottom layer	0.00	Thickest layer	0.06
		Thickest layer	0.00	Bottom layer	0.10
37A: Manchester-----	80	Good		Fair	
		Thickest layer	0.14	Thickest layer	0.06
		Bottom layer	0.93	Bottom layer	0.11
37C: Manchester-----	80	Good		Fair	
		Thickest layer	0.14	Thickest layer	0.06
		Bottom layer	0.93	Bottom layer	0.11
37E: Manchester-----	80	Good		Fair	
		Thickest layer	0.14	Thickest layer	0.06
		Bottom layer	0.93	Bottom layer	0.11
38A: Hinckley-----	80	Good		Fair	
		Thickest layer	0.21	Bottom layer	0.09
		Bottom layer	0.64	Thickest layer	0.09
38C: Hinckley-----	80	Good		Fair	
		Thickest layer	0.21	Bottom layer	0.09
		Bottom layer	0.64	Thickest layer	0.09
38E: Hinckley-----	80	Good		Fair	
		Thickest layer	0.21	Bottom layer	0.09
		Bottom layer	0.64	Thickest layer	0.09
39A: Groton-----	85	Good		Fair	
		Thickest layer	0.37	Bottom layer	0.08
		Bottom layer	0.93	Thickest layer	0.08
39C: Groton-----	85	Good		Fair	
		Thickest layer	0.37	Bottom layer	0.08
		Bottom layer	0.93	Thickest layer	0.08

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
39E: Groton-----	85	Good		Fair	
		Thickest layer	0.37	Thickest layer	0.08
		Bottom layer	0.93	Bottom layer	0.08
40A: Ludlow-----	80	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
40B: Ludlow-----	80	Poor		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
41B: Ludlow-----	80	Poor		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
42C: Ludlow-----	80	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
43A: Rainbow-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
43B: Rainbow-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
44B: Rainbow-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
45A: Woodbridge-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
45B: Woodbridge-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
45C: Woodbridge-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
46B: Woodbridge-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
46C: Woodbridge-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
47C: Woodbridge-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
48B: Georgia-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Amenia-----	35	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
48C: Georgia-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Amenia-----	35	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
49B: Georgia-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Amenia-----	35	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
49C: Georgia-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
49C: Amenia-----	35	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
50A: Sutton-----	80	Poor Bottom layer Thickest layer	0.07 0.07	Poor Bottom layer Thickest layer	0.00 0.00
50B: Sutton-----	80	Poor Bottom layer Thickest layer	0.07 0.07	Poor Bottom layer Thickest layer	0.00 0.00
51B: Sutton-----	80	Poor Bottom layer Thickest layer	0.07 0.07	Poor Bottom layer Thickest layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
52C: Sutton-----	80	Poor Bottom layer Thickest layer	 0.07 0.07	Poor Bottom layer Thickest layer	 0.00 0.00
53A: Wapping-----	80	Fair Thickest layer Bottom layer	 0.00 0.36	Fair Thickest layer Bottom layer	 0.00 0.09
53B: Wapping-----	80	Fair Thickest layer Bottom layer	 0.00 0.36	Fair Thickest layer Bottom layer	 0.00 0.09
54B: Wapping-----	80	Fair Thickest layer Bottom layer	 0.00 0.36	Fair Thickest layer Bottom layer	 0.00 0.09
55A: Watchaug-----	80	Poor Bottom layer Thickest layer	 0.00 0.07	Poor Bottom layer Thickest layer	 0.00 0.00
55B: Watchaug-----	80	Poor Bottom layer Thickest layer	 0.00 0.07	Poor Bottom layer Thickest layer	 0.00 0.00
56B: Watchaug-----	80	Poor Bottom layer Thickest layer	 0.00 0.07	Poor Bottom layer Thickest layer	 0.00 0.00
57B: Gloucester-----	80	Good Thickest layer Bottom layer	 0.47 0.83	Fair Thickest layer Bottom layer	 0.15 0.22
57C: Gloucester-----	80	Good Thickest layer Bottom layer	 0.47 0.83	Fair Thickest layer Bottom layer	 0.15 0.22
57D: Gloucester-----	80	Good Thickest layer Bottom layer	 0.47 0.83	Fair Thickest layer Bottom layer	 0.15 0.22
58B: Gloucester-----	80	Good Thickest layer Bottom layer	 0.47 0.83	Fair Thickest layer Bottom layer	 0.15 0.22
58C: Gloucester-----	80	Good Thickest layer Bottom layer	 0.47 0.83	Fair Thickest layer Bottom layer	 0.15 0.22

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
59C: Gloucester-----	80	Good		Fair	
		Thickest layer	0.47	Thickest layer	0.15
		Bottom layer	0.83	Bottom layer	0.22
59D: Gloucester-----	80	Good		Fair	
		Thickest layer	0.47	Thickest layer	0.15
		Bottom layer	0.83	Bottom layer	0.22
60B: Canton-----	45	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Charlton-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
60C: Canton-----	45	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Charlton-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
60D: Canton-----	45	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Charlton-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
61B: Canton-----	45	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Charlton-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
61C: Canton-----	45	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Charlton-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
62C: Canton-----	45	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Charlton-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
62D: Canton-----	45	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Charlton-----	35	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
63B: Cheshire-----	80	Fair		Poor	
		Bottom layer	0.14	Bottom layer	0.00
		Thickest layer	0.14	Thickest layer	0.00
63C: Cheshire-----	80	Fair		Poor	
		Bottom layer	0.14	Bottom layer	0.00
		Thickest layer	0.14	Thickest layer	0.00
63D: Cheshire-----	80	Fair		Poor	
		Bottom layer	0.14	Bottom layer	0.00
		Thickest layer	0.14	Thickest layer	0.00
64B: Cheshire-----	80	Fair		Poor	
		Bottom layer	0.14	Bottom layer	0.00
		Thickest layer	0.14	Thickest layer	0.00
64C: Cheshire-----	80	Fair		Poor	
		Bottom layer	0.14	Bottom layer	0.00
		Thickest layer	0.14	Thickest layer	0.00
65C: Cheshire-----	80	Fair		Poor	
		Bottom layer	0.14	Bottom layer	0.00
		Thickest layer	0.14	Thickest layer	0.00
65D: Cheshire-----	80	Fair		Poor	
		Bottom layer	0.14	Bottom layer	0.00
		Thickest layer	0.14	Thickest layer	0.00
66B: Narragansett-----	80	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.07
66C: Narragansett-----	80	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.07
67B: Narragansett-----	80	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.07

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
67C: Narragansett-----	80	Poor Thickest layer Bottom layer	0.00 0.07	Fair Thickest layer Bottom layer	0.00 0.07
68C: Narragansett-----	80	Poor Thickest layer Bottom layer	0.00 0.07	Fair Thickest layer Bottom layer	0.00 0.07
68D: Narragansett-----	80	Poor Thickest layer Bottom layer	0.00 0.07	Fair Thickest layer Bottom layer	0.00 0.07
69B: Yalesville-----	75	Fair Thickest layer Bottom layer	0.00 0.36	Poor Thickest layer Bottom layer	0.00 0.00
69C: Yalesville-----	75	Fair Thickest layer Bottom layer	0.00 0.36	Poor Thickest layer Bottom layer	0.00 0.00
70C: Branford-----	50	Poor Thickest layer Bottom layer	0.00 0.07	Fair Thickest layer Bottom layer	0.00 0.12
70C: Holyoke-----	30	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
71C: Brookfield-----	45	Fair Thickest layer Bottom layer	0.00 0.14	Fair Thickest layer Bottom layer	0.00 0.01
Brimfield-----	30	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rock outcrop-----	15	Not rated		Not rated	
71E: Brookfield-----	45	Fair Thickest layer Bottom layer	0.00 0.14	Fair Thickest layer Bottom layer	0.00 0.01
Brimfield-----	30	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rock outcrop-----	15	Not rated		Not rated	

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
73C: Charlton-----	45	Fair		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.14	Bottom layer	0.00
Chatfield-----	30	Fair		Poor	
		Thickest layer	0.09	Bottom layer	0.00
		Bottom layer	0.29	Thickest layer	0.00
73E: Charlton-----	45	Fair		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.14	Bottom layer	0.00
Chatfield-----	30	Fair		Poor	
		Thickest layer	0.09	Bottom layer	0.00
		Bottom layer	0.29	Thickest layer	0.00
74C: Narragansett-----	55	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.07
Hollis-----	20	Poor		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
75C: Hollis-----	35	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
75C: Chatfield-----	30	Fair		Poor	
		Thickest layer	0.09	Thickest layer	0.00
		Bottom layer	0.29	Bottom layer	0.00
Rock outcrop-----	15	Not rated		Not rated	
75E: Hollis-----	35	Poor		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
Chatfield-----	30	Fair		Poor	
		Thickest layer	0.09	Thickest layer	0.00
		Bottom layer	0.29	Bottom layer	0.00
Rock outcrop-----	15	Not rated		Not rated	
76E: Rock outcrop-----	55	Not rated		Not rated	
Hollis-----	25	Poor		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
76F: Rock outcrop-----	55	Not rated		Not rated	
Hollis-----	25	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
77C: Cheshire-----	45	Fair Bottom layer Thickest layer	0.14 0.14	Poor Thickest layer Bottom layer	0.00 0.00
Holyoke-----	35	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
77D: Cheshire-----	45	Fair Bottom layer Thickest layer	0.14 0.14	Poor Thickest layer Bottom layer	0.00 0.00
Holyoke-----	35	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
78C: Holyoke-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
78C: Rock outcrop-----	25	Not rated		Not rated	
78E: Holyoke-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
Rock outcrop-----	25	Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated	
Holyoke-----	25	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
80B: Bernardston-----	80	Poor Thickest layer Bottom layer	0.00 0.09	Poor Thickest layer Bottom layer	0.00 0.00
80C: Bernardston-----	80	Poor Thickest layer Bottom layer	0.00 0.09	Poor Thickest layer Bottom layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
81C: Bernardston-----	80	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.09	Bottom layer	0.00
81D: Bernardston-----	80	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.09	Bottom layer	0.00
82B: Broadbrook-----	80	Poor		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
82C: Broadbrook-----	80	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
82D: Broadbrook-----	80	Poor		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
83B: Broadbrook-----	80	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
83C: Broadbrook-----	80	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
84B: Paxton-----	55	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
Montauk-----	30	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.04
84C: Paxton-----	55	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
Montauk-----	30	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.04
84D: Paxton-----	55	Poor		Poor	
		Bottom layer	0.00	Bottom layer	0.00
		Thickest layer	0.00	Thickest layer	0.00
Montauk-----	30	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.04

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
85B: Paxton-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Montauk-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.04
85C: Paxton-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
Montauk-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.04
86C: Paxton-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Montauk-----	30	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.04
86D: Paxton-----	55	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
Montauk-----	30	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.04
87B: Wethersfield-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
87C: Wethersfield-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
87D: Wethersfield-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
88B: Wethersfield-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
88C: Wethersfield-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
89C: Wethersfield-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
89D: Wethersfield-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
90B: Stockbridge-----	80	Fair Thickest layer Bottom layer	0.19 0.50	Poor Bottom layer Thickest layer	0.00 0.00
90C: Stockbridge-----	80	Fair Thickest layer Bottom layer	0.19 0.50	Poor Bottom layer Thickest layer	0.00 0.00
90D: Stockbridge-----	80	Fair Thickest layer Bottom layer	0.19 0.50	Poor Bottom layer Thickest layer	0.00 0.00
91B: Stockbridge-----	80	Fair Thickest layer Bottom layer	0.19 0.50	Poor Bottom layer Thickest layer	0.00 0.00
91C: Stockbridge-----	80	Fair Thickest layer Bottom layer	0.19 0.50	Poor Thickest layer Bottom layer	0.00 0.00
91D: Stockbridge-----	80	Fair Thickest layer Bottom layer	0.19 0.50	Poor Thickest layer Bottom layer	0.00 0.00
92B: Nellis-----	85	Fair Thickest layer Bottom layer	0.21 0.36	Poor Thickest layer Bottom layer	0.00 0.00
92C: Nellis-----	85	Fair Thickest layer Bottom layer	0.21 0.36	Poor Thickest layer Bottom layer	0.00 0.00
92D: Nellis-----	85	Fair Thickest layer Bottom layer	0.21 0.36	Poor Bottom layer Thickest layer	0.00 0.00
93C: Nellis-----	85	Fair Thickest layer Bottom layer	0.21 0.36	Poor Thickest layer Bottom layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
94C: Farmington-----	40	Poor Thickest layer Bottom layer	0.00 0.07	Poor Bottom layer Thickest layer	0.00 0.00
Nellis-----	35	Fair Thickest layer Bottom layer	0.21 0.36	Poor Thickest layer Bottom layer	0.00 0.00
94E: Farmington-----	40	Poor Thickest layer Bottom layer	0.00 0.07	Poor Thickest layer Bottom layer	0.00 0.00
Nellis-----	35	Fair Thickest layer Bottom layer	0.21 0.36	Poor Thickest layer Bottom layer	0.00 0.00
95C: Farmington-----	60	Poor Thickest layer Bottom layer	0.00 0.07	Poor Thickest layer Bottom layer	0.00 0.00
95C: Rock outcrop-----	20	Not rated		Not rated	
95E: Farmington-----	60	Poor Thickest layer Bottom layer	0.00 0.07	Poor Thickest layer Bottom layer	0.00 0.00
Rock outcrop-----	20	Not rated		Not rated	
96: Ipswich-----	85	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer Organic matter content	0.00 0.00 0.00
97: Pawcatuck-----	85	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.04
98: Westbrook-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
99: Westbrook, low salt-	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
100: Suncook-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.10 0.10

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
101: Occum-----	80	Poor Thickest layer Bottom layer	0.00 0.07	Fair Thickest layer Bottom layer	0.00 0.36
102: Pootatuck-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.36
103: Rippowam-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.50
104: Bash-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
105: Hadley-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
106: Winooski-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
107: Limerick-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Lim-----	30	Poor Thickest layer Bottom layer	0.04 0.07	Fair Thickest layer Bottom layer	0.04 0.09
108: Saco-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.10
109: Fluvaquents, Frequently Flooded-	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
Udifluents, Frequently Flooded-	35	Fair Bottom layer Thickest layer	0.29 0.29	Fair Thickest layer Bottom layer	0.08 0.17
221A: Ninigret-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.03
Urban land-----	35	Not rated		Not rated	

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
224A: Deerfield-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.10 0.13
Urban land-----	35	Not rated		Not rated	
225B: Brancroft-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Urban land-----	35	Not rated		Not rated	
226B: Berlin-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
Urban land-----	35	Not rated		Not rated	
228B: Elmridge-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Urban land-----	35	Not rated		Not rated	
229B: Agawam-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.12
Urban land-----	35	Not rated		Not rated	
229C: Agawam-----	40	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.12
Urban land-----	35	Not rated		Not rated	
230B: Branford-----	40	Poor Thickest layer Bottom layer	0.00 0.07	Fair Thickest layer Bottom layer	0.00 0.12
Urban land-----	35	Not rated		Not rated	
230C: Branford-----	40	Poor Thickest layer Bottom layer	0.00 0.07	Fair Thickest layer Bottom layer	0.00 0.12
Urban land-----	35	Not rated		Not rated	

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
232B:					
Haven-----	40	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.39
Urban land-----	35	Not rated		Not rated	
234B:					
Merrimac-----	40	Fair		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.29	Bottom layer	0.12
Urban land-----	35	Not rated		Not rated	
235B:					
Penwood-----	40	Poor		Fair	
		Bottom layer	0.00	Bottom layer	0.15
		Thickest layer	0.00	Thickest layer	0.15
Urban land-----	35	Not rated		Not rated	
236B:					
Windsor-----	40	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.06
		Bottom layer	0.00	Bottom layer	0.10
Urban land-----	35	Not rated		Not rated	
237A:					
Manchester-----	40	Good		Fair	
		Thickest layer	0.14	Thickest layer	0.06
		Bottom layer	0.93	Bottom layer	0.11
Urban land-----	35	Not rated		Not rated	
237C:					
Manchester-----	40	Good		Fair	
		Thickest layer	0.14	Thickest layer	0.06
		Bottom layer	0.93	Bottom layer	0.11
Urban land-----	35	Not rated		Not rated	
238A:					
Hinckley-----	40	Good		Fair	
		Thickest layer	0.21	Bottom layer	0.09
		Bottom layer	0.64	Thickest layer	0.09
Urban land-----	35	Not rated		Not rated	
238C:					
Hinckley-----	40	Good		Fair	
		Thickest layer	0.21	Bottom layer	0.09
		Bottom layer	0.64	Thickest layer	0.09
Urban land-----	35	Not rated		Not rated	

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
240B: Ludlow-----	40	Poor Thickest layer	0.00	Poor Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
Urban land-----	35	Not rated		Not rated	
243B: Rainbow-----	40	Poor Thickest layer	0.00	Poor Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
245B: Woodbridge-----	40	Poor Bottom layer	0.00	Poor Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
245C: Woodbridge-----	40	Poor Bottom layer	0.00	Poor Thickest layer	0.00
		Thickest layer	0.00	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
248B: Georgia-----	40	Poor Thickest layer	0.00	Poor Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
250B: Sutton-----	40	Poor Bottom layer	0.07	Poor Thickest layer	0.00
		Thickest layer	0.07	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
253B: Wapping-----	40	Fair Thickest layer	0.00	Fair Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.09
Urban land-----	35	Not rated		Not rated	
255B: Watchaug-----	40	Poor Bottom layer	0.00	Poor Thickest layer	0.00
		Thickest layer	0.07	Bottom layer	0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
255B: Urban land-----	35	Not rated		Not rated	
260B: Charlton-----	40	Fair		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.14	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
260C: Charlton-----	40	Fair		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
Urban land-----	35	Not rated		Not rated	
260D: Charlton-----	40	Fair		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.14	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
263B: Cheshire-----	40	Fair		Poor	
		Bottom layer	0.14	Thickest layer	0.00
		Thickest layer	0.14	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	
263C: Cheshire-----	40	Fair		Poor	
		Thickest layer	0.14	Bottom layer	0.00
		Bottom layer	0.14	Thickest layer	0.00
Urban land-----	35	Not rated		Not rated	
266B: Narragansett-----	40	Poor		Fair	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.07	Bottom layer	0.07
Urban land-----	35	Not rated		Not rated	
269B: Yalesville-----	40	Fair		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.36	Bottom layer	0.00
Urban land-----	35	Not rated		Not rated	

Table 20.--Source of Sand and Gravel--Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
269C: Yalesville-----	40	Fair Thickest layer Bottom layer	 0.00 0.36	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Fair Thickest layer Bottom layer	 0.00 0.14	Poor Thickest layer Bottom layer	 0.00 0.00
Chatfield-----	15	Fair Thickest layer Bottom layer	 0.09 0.29	Poor Thickest layer Bottom layer	 0.00 0.00
273E: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Fair Thickest layer Bottom layer	 0.00 0.14	Poor Thickest layer Bottom layer	 0.00 0.00
Chatfield-----	15	Fair Thickest layer Bottom layer	 0.09 0.29	Poor Thickest layer Bottom layer	 0.00 0.00
275C: Urban land-----	45	Not rated		Not rated	
Chatfield-----	30	Fair Thickest layer Bottom layer	 0.09 0.29	Poor Thickest layer Bottom layer	 0.00 0.00
275E: Urban land-----	35	Not rated		Not rated	
Chatfield-----	25	Fair Thickest layer Bottom layer	 0.09 0.29	Poor Thickest layer Bottom layer	 0.00 0.00
Rock outcrop-----	15	Not rated		Not rated	
282B: Broadbrook-----	40	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
284B: Paxton-----	40	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
284C: Paxton-----	40	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
284D: Paxton-----	40	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
287B: Wethersfield-----	40	Poor Thickest layer Bottom layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
287C: Wethersfield-----	40	Poor Thickest layer Bottom layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
287D: Wethersfield-----	40	Poor Thickest layer Bottom layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
290B: Stockbridge-----	40	Fair Thickest layer Bottom layer	 0.19 0.50	Poor Bottom layer Thickest layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
290C: Stockbridge-----	40	Fair Thickest layer Bottom layer	 0.19 0.50	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
290D: Stockbridge-----	40	Fair Thickest layer Bottom layer	 0.19 0.50	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
301: Beaches-----	50	Not rated		Not rated	
Udipsamments-----	35	Poor Thickest layer Bottom layer	 0.00 0.00	Fair Thickest layer Bottom layer	 0.10 0.76
302: Dumps-----	95	Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated	
304: Udorthents-----	90	Fair Thickest layer Bottom layer	 0.43 0.43	Poor Thickest layer Bottom layer	 0.00 0.00
305: Udorthents-----	65	Fair Thickest layer Bottom layer	 0.43 0.43	Poor Bottom layer Thickest layer	 0.00 0.00
Pits-----	25	Good Bottom layer Thickest layer	 1.00 1.00	Fair Bottom layer Thickest layer	 0.71 0.71
306: Udorthents-----	50	Fair Bottom layer Thickest layer	 0.43 0.43	Poor Thickest layer Bottom layer	 0.00 0.00
Urban land-----	35	Not rated		Not rated	
307: Urban land-----	80	Not rated		Not rated	
308: Udorthents-----	80	Fair Bottom layer Thickest layer	 0.43 0.43	Poor Bottom layer Thickest layer	 0.00 0.00
309: Udorthents-----	80	Fair Bottom layer Thickest layer	 0.43 0.43	Poor Bottom layer Thickest layer	 0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
310: Udorthents, Periodically Flooded-----	85	Fair Thickest layer Bottom layer	0.43 0.43	Poor Bottom layer Thickest layer	0.00 0.00
401C: Macomber-----	55	Good Thickest layer Bottom layer	0.00 1.00	Poor Thickest layer Bottom layer	0.00 0.00
Taconic-----	30	Good Thickest layer Bottom layer	0.00 0.77	Poor Bottom layer Thickest layer	0.00 0.00
402D: Macomber-----	50	Good Thickest layer Bottom layer	0.00 1.00	Poor Bottom layer Thickest layer	0.00 0.00
Taconic-----	25	Good Thickest layer Bottom layer	0.00 0.77	Poor Bottom layer Thickest layer	0.00 0.00
Rock outcrop-----	15	Not rated		Not rated	
403C: Taconic-----	70	Good Thickest layer Bottom layer	0.00 0.77	Poor Bottom layer Thickest layer	0.00 0.00
Rock outcrop-----	25	Not rated		Not rated	
403E: Taconic-----	70	Good Thickest layer Bottom layer	0.00 0.77	Poor Thickest layer Bottom layer	0.00 0.00
403E: Rock outcrop-----	20	Not rated		Not rated	
403F: Taconic-----	70	Good Thickest layer Bottom layer	0.00 0.77	Poor Bottom layer Thickest layer	0.00 0.00
Rock outcrop-----	20	Not rated		Not rated	
405C: Dummerston-----	85	Poor Thickest layer Bottom layer	0.07 0.07	Poor Bottom layer Thickest layer	0.00 0.00
405E: Dummerston-----	85	Poor Thickest layer Bottom layer	0.07 0.07	Poor Bottom layer Thickest layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
407C: Lanesboro-----	85	Fair Thickest layer Bottom layer	0.00 0.14	Poor Thickest layer Bottom layer	0.00 0.00
407E: Lanesboro-----	85	Fair Thickest layer Bottom layer	0.00 0.14	Poor Thickest layer Bottom layer	0.00 0.00
408C: Fullam-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
409B: Brayton-----	85	Fair Thickest layer Bottom layer	0.00 0.14	Poor Bottom layer Thickest layer	0.00 0.00
412B: Bice-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
412C: Bice-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
412D: Bice-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
413C: Bice-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
Millsite-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
413E: Bice-----	45	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
Millsite-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
414: Fredon, cold-----	85	Fair Thickest layer Bottom layer	0.14 0.14	Fair Thickest layer Bottom layer	0.08 0.43
415C: Millsite-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
415C: Westminster-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rock outcrop-----	15	Not rated		Not rated	
415E: Millsite-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Westminster-----	40	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
Rock outcrop-----	15	Not rated		Not rated	
416E: Rock outcrop-----	70	Not rated		Not rated	
Westminster-----	20	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
416F: Rock outcrop-----	70	Not rated		Not rated	
416F: Westminster-----	20	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
417B: Bice-----	85	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
417C: Bice-----	85	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
417D: Bice-----	85	Poor Thickest layer Bottom layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
418C: Schroon-----	85	Fair Bottom layer Thickest layer	0.07 0.14	Fair Thickest layer Bottom layer	0.00 0.03
420A: Schroon-----	85	Fair Bottom layer Thickest layer	0.07 0.14	Fair Thickest layer Bottom layer	0.00 0.03

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
420B: Schroon-----	85	Fair Bottom layer Thickest layer	 0.07 0.14	Fair Thickest layer Bottom layer	 0.00 0.03
421A: Ninigret, cold-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Fair Thickest layer Bottom layer	 0.00 0.03
423A: Sudbury, cold-----	85	Fair Thickest layer Bottom layer	 0.00 0.43	Fair Thickest layer Bottom layer	 0.00 0.10
424B: Shelburne-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
424C: Shelburne-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
424D: Shelburne-----	85	Poor Thickest layer Bottom layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
425B: Shelburne-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
425C: Shelburne-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
426D: Shelburne-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00
427B: Ashfield-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
427C: Ashfield-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Bottom layer Thickest layer	 0.00 0.00
428A: Ashfield-----	85	Poor Bottom layer Thickest layer	 0.00 0.00	Poor Thickest layer Bottom layer	 0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
428B: Ashfield-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
428C: Ashfield-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
429A: Agawam, cold-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.12
429B: Agawam, cold-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.12
429C: Agawam, cold-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.12
433: Moosilauke-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.12 0.15
434A: Merrimac, cold-----	80	Fair Thickest layer Bottom layer	0.00 0.29	Fair Thickest layer Bottom layer	0.00 0.12
434B: Merrimac, cold-----	80	Fair Thickest layer Bottom layer	0.00 0.29	Fair Thickest layer Bottom layer	0.00 0.12
434C: Merrimac, cold-----	80	Fair Thickest layer Bottom layer	0.00 0.29	Fair Thickest layer Bottom layer	0.00 0.12
435: Scarboro-----	80	Fair Thickest layer Bottom layer	0.00 0.14	Fair Thickest layer Bottom layer	0.31 0.36
436: Halsey-----	80	Fair Thickest layer Bottom layer	0.00 0.36	Fair Thickest layer Bottom layer	0.00 0.31
437: Wonsqueak-----	85	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
438: Bucksport-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.00 0.10
440A: Boscawen-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.22 0.22
440C: Boscawen-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.22 0.22
440E: Boscawen-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Fair Thickest layer Bottom layer	0.22 0.22
442: Brayton-----	85	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
443: Brayton-----	50	Poor Bottom layer Thickest layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00
443: Loonmeadow-----	35	Poor Thickest layer Bottom layer	0.07 0.07	Poor Thickest layer Bottom layer	0.00 0.00
448B: Hogansburg-----	85	Fair Thickest layer Bottom layer	0.07 0.14	Poor Thickest layer Bottom layer	0.00 0.00
449B: Hogansburg-----	85	Fair Thickest layer Bottom layer	0.07 0.14	Poor Thickest layer Bottom layer	0.00 0.00
449C: Hogansburg-----	85	Fair Thickest layer Bottom layer	0.07 0.14	Poor Thickest layer Bottom layer	0.00 0.00
450B: Pyrities-----	80	Poor Bottom layer Thickest layer	0.00 0.00	Poor Thickest layer Bottom layer	0.00 0.00
450C: Pyrities-----	80	Poor Thickest layer Bottom layer	0.00 0.00	Poor Bottom layer Thickest layer	0.00 0.00

Table 20.—Source of Sand and Gravel—Continued

Map symbol and soil name	Pct. of map unit	Potential source of gravel		Potential source of sand	
		Rating class	Value	Rating class	Value
450D: Pyrities-----	80	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
451B: Pyrities-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
451C: Pyrities-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
451D: Pyrities-----	80	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
457: Mudgepond-----	80	Fair		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.21	Bottom layer	0.00
458: Mudgepond-----	55	Fair		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.21	Bottom layer	0.00
458: Alden-----	35	Fair		Poor	
		Bottom layer	0.00	Thickest layer	0.00
		Thickest layer	0.14	Bottom layer	0.00
501: Ondawa-----	85	Poor		Poor	
		Thickest layer	0.00	Bottom layer	0.00
		Bottom layer	0.00	Thickest layer	0.00
503: Rumney-----	80	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00
508: Medomak-----	85	Poor		Poor	
		Thickest layer	0.00	Thickest layer	0.00
		Bottom layer	0.00	Bottom layer	0.00

Table 21.—Construction Materials

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments	0.00 0.00 0.03
3: Ridgebury-----	40	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments	0.00 0.00 0.03
Leicester-----	35	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.12 0.68
Whitman-----	15	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth	0.00 0.00
4: Leicester-----	80	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.12 0.68
5: Wilbraham-----	80	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (dense layer) Rock fragments Too acid	0.00 0.00 0.28 0.98
6: Wilbraham-----	60	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (dense layer) Rock fragments Too acid	0.00 0.00 0.28 0.98
Menlo-----	25	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (dense layer) Rock fragments	0.00 0.29 0.50

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
7: Mudgepond-----	85	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments	0.00 0.88
8: Mudgepond-----	45	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments	0.00 0.88
Alden-----	35	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (rock fragments)	0.00 0.88
9: Scitico-----	40	Poor Wetness depth Low strength Shrink-swell	0.00 0.00 0.58	Poor Wetness depth Too clayey	0.00 0.11
Shaker-----	30	Poor Wetness depth Low strength Shrink-swell	0.00 0.00 0.92	Poor Wetness depth	0.00
Maybid-----	15	Poor Wetness depth Low strength Shrink-swell	0.00 0.00 0.50	Poor Wetness depth Too clayey	0.00 0.04
10: Raynham-----	80	Poor Wetness depth	0.00	Poor Wetness depth	0.00
12: Raypol-----	80	Poor Wetness depth	0.00	Poor Too sandy Wetness depth Rock fragments Hard to reclaim (rock fragments) Too acid	0.00 0.00 0.00 0.68 0.88
13: Walpole-----	80	Poor Wetness depth	0.00	Poor Too sandy Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.82
14: Fredon-----	85	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments Too sandy Hard to reclaim (rock fragments)	0.00 0.02 0.22 0.97

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
15: Scarboro-----	80	Poor Wetness depth	0.00	Poor Too sandy Wetness depth Hard to reclaim (rock fragments) Rock fragments Too acid	0.00 0.00 0.82 0.97 0.98
16: Halsey-----	80	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (rock fragments) Rock fragments	0.00 0.50 0.97
17: Timakwa-----	45	Poor Wetness depth Shrink-swell	0.00 0.44	Poor Wetness depth Organic matter content high Too acid	0.00 0.00 0.88
Natchaug-----	40	Poor Wetness depth Shrink-swell	0.00 0.95	Poor Wetness depth Rock fragments	0.00 0.88
18: Catden-----	40	Poor Wetness depth Shrink-swell	0.00 0.00	Poor Wetness depth Organic matter content high Too acid	0.00 0.00 0.88
Freetown-----	40	Poor Wetness depth Shrink-swell	0.00 0.00	Poor Wetness depth Organic matter content high Too acid	0.00 0.00 0.12
20A: Ellington-----	80	Fair Wetness depth	0.53	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Wetness depth Too acid	0.00 0.00 0.50 0.53 0.98
21A: Ninigret-----	60	Fair Wetness depth	0.53	Poor Too sandy Rock fragments Wetness depth Hard to reclaim (rock fragments)	0.00 0.00 0.53 0.82

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
21A: Tisbury-----	25	Fair Wetness depth	0.53	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Wetness depth Too acid	0.00 0.00 0.50 0.53 0.98
22A: Hero-----	85	Fair Wetness depth	0.53	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Wetness depth	0.00 0.00 0.00 0.53
22B: Hero-----	85	Fair Wetness depth	0.53	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Wetness depth	0.00 0.00 0.00 0.53
23A: Sudbury-----	80	Fair Wetness depth	0.76	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Wetness depth	0.00 0.00 0.08 0.76
24A: Deerfield-----	80	Fair Wetness depth	0.76	Poor Too sandy Wetness depth	0.00 0.76
25A: Brancroft-----	80	Poor Low strength Wetness depth	0.00 0.29	Fair Wetness depth	0.29
25B: Brancroft-----	80	Poor Low strength Wetness depth	0.00 0.29	Fair Wetness depth	0.29
25C: Brancroft-----	80	Poor Low strength Wetness depth	0.00 0.29	Fair Wetness depth Slope	0.29 0.37
26A: Berlin-----	80	Poor Low strength Wetness depth	0.00 0.29	Fair Wetness depth Too clayey	0.29 0.55

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
26B: Berlin-----	80	Poor Low strength Wetness depth	0.00 0.29	Fair Wetness depth Too clayey	0.29 0.55
27A: Belgrade-----	80	Fair Wetness depth	0.91	Fair Wetness depth	0.91
28A: Elmridge-----	80	Poor Low strength Wetness depth Shrink-swell	0.00 0.53 0.86	Poor Too clayey Wetness depth	0.00 0.53
28B: Elmridge-----	80	Poor Low strength Wetness depth Shrink-swell	0.00 0.53 0.86	Poor Too clayey Wetness depth	0.00 0.53
29A: Agawam-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.50
29B: Agawam-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.50
29C: Agawam-----	80	Good		Poor Too sandy Rock fragments Slope Hard to reclaim (rock fragments)	0.00 0.00 0.37 0.50
30A: Branford-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Too acid	0.00 0.00 0.32 0.98
30B: Branford-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Too acid	0.00 0.00 0.32 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
30C: Branford-----	80	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.32
				Slope	0.37
				Too acid	0.98
31A: Copake-----	85	Good		Poor	
				Hard to reclaim (rock fragments)	0.00
				Rock fragments	0.12
31B: Copake-----	85	Good		Poor	
				Hard to reclaim (rock fragments)	0.00
				Rock fragments	0.12
31C: Copake-----	85	Good		Poor	
				Hard to reclaim (rock fragments)	0.00
				Rock fragments	0.12
				Slope	0.37
32A: Haven-----	60	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.02
				Too acid	0.98
Enfield-----	25	Good		Fair	
				Hard to reclaim (rock fragments)	0.68
32B: Haven-----	60	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.02
				Too acid	0.98
Enfield-----	25	Good		Fair	
				Hard to reclaim (rock fragments)	0.68
32C: Haven-----	60	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.02
				Slope	0.37
				Too acid	0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
32C: Enfield-----	25	Good		Fair Slope	0.37
				Hard to reclaim (rock fragments)	0.68
33A: Hartford-----	80	Good		Poor Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.50
				Too acid	0.98
33B: Hartford-----	80	Good		Poor Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.50
				Too acid	0.98
34A: Merrimac-----	80	Good		Poor Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.00
34B: Merrimac-----	80	Good		Poor Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.00
34C: Merrimac-----	80	Good		Poor Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.00
				Slope	0.37
35A: Penwood-----	80	Good		Poor Too sandy	0.00
				Too acid	0.98
35B: Penwood-----	80	Good		Poor Too sandy	0.00
				Too acid	0.98
36A: Windsor-----	80	Good		Poor Too sandy	0.00
				Rock fragments	0.97

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
36B: Windsor-----	80	Good		Poor Too sandy Rock fragments	0.00 0.97
36C: Windsor-----	80	Good		Poor Too sandy Slope Rock fragments	0.00 0.37 0.97
37A: Manchester-----	80	Good		Poor Too sandy Hard to reclaim (rock fragments) Rock fragments Too acid	0.00 0.00 0.00 0.98
37C: Manchester-----	80	Good		Poor Too sandy Hard to reclaim (rock fragments) Rock fragments Slope Too acid	0.00 0.00 0.00 0.96 0.98
37E: Manchester-----	80	Poor Slope	0.00	Poor Slope Too sandy Hard to reclaim (rock fragments) Rock fragments Too acid	0.00 0.00 0.00 0.00 0.98
38A: Hinckley-----	80	Good		Poor Hard to reclaim (rock fragments) Rock fragments Too sandy Too acid	0.00 0.00 0.32 0.76
38C: Hinckley-----	80	Good		Poor Hard to reclaim (rock fragments) Rock fragments Too sandy Too acid Slope	0.00 0.00 0.32 0.76 0.96
38E: Hinckley-----	80	Poor Slope	0.00	Poor Slope Hard to reclaim (rock fragments) Rock fragments Too sandy Too acid	0.00 0.00 0.00 0.32 0.76

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
39A: Groton-----	85	Good		Poor Rock fragments Hard to reclaim (rock fragments)	0.00 0.00
39C: Groton-----	85	Good		Poor Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.00 0.96
39E: Groton-----	85	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00
40A: Ludlow-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
40B: Ludlow-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
41B: Ludlow-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
42C: Ludlow-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Slope Too acid	0.00 0.28 0.53 0.96 0.98
43A: Rainbow-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.20 0.53 0.97 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
43B: Rainbow-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.20 0.53 0.97 0.98
44B: Rainbow-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.20 0.53 0.97 0.98
45A: Woodbridge-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
45B: Woodbridge-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
45C: Woodbridge-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Slope Wetness depth Too acid	0.00 0.28 0.37 0.53 0.98
46B: Woodbridge-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
46C: Woodbridge-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Slope Wetness depth Too acid	0.00 0.28 0.37 0.53 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
47C: Woodbridge-----	80	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Slope Too acid	0.00 0.28 0.53 0.96 0.98
48B: Georgia-----	50	Fair Wetness depth	0.76	Fair Rock fragments Wetness depth	0.12 0.76
Amenia-----	35	Fair Wetness depth	0.76	Fair Rock fragments Wetness depth	0.12 0.76
48C: Georgia-----	50	Fair Wetness depth	0.76	Fair Rock fragments Slope Wetness depth	0.12 0.37 0.76
Amenia-----	35	Fair Wetness depth	0.76	Fair Rock fragments Slope Wetness depth	0.12 0.37 0.76
49B: Georgia-----	50	Fair Wetness depth	0.76	Fair Rock fragments Wetness depth	0.12 0.76
Amenia-----	35	Fair Wetness depth	0.76	Fair Rock fragments Wetness depth	0.12 0.76
49C: Georgia-----	50	Fair Wetness depth	0.76	Fair Rock fragments Slope Wetness depth	0.12 0.37 0.76
Amenia-----	35	Fair Wetness depth	0.76	Fair Rock fragments Slope Wetness depth	0.12 0.37 0.76
50A: Sutton-----	80	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth Too acid	0.03 0.53 0.98
50B: Sutton-----	80	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth Too acid	0.03 0.53 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
51B: Sutton-----	80	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth Too acid	0.03 0.53 0.98
52C: Sutton-----	80	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth Slope Too acid	0.03 0.53 0.96 0.98
53A: Wapping-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (rock fragments) Wetness depth Rock fragments	0.18 0.53 0.97
53B: Wapping-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (rock fragments) Wetness depth Rock fragments	0.18 0.53 0.97
54B: Wapping-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (rock fragments) Wetness depth Rock fragments	0.18 0.53 0.97
55A: Watchaug-----	80	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth	0.28 0.53
55B: Watchaug-----	80	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth	0.28 0.53
56B: Watchaug-----	80	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth	0.28 0.53
57B: Gloucester-----	80	Fair Stone content	0.99	Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
57C: Gloucester-----	80	Fair Stone content	0.99	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.00 0.00 0.37
57D: Gloucester-----	80	Fair Slope Stone content	0.50 0.99	Poor Slope Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.00
58B: Gloucester-----	80	Fair Stone content	0.99	Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00
58C: Gloucester-----	80	Fair Stone content	0.99	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.00 0.00 0.37
59C: Gloucester-----	80	Fair Stone content	0.99	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.00 0.00 0.96
59D: Gloucester-----	80	Fair Slope Stone content	0.50 0.99	Poor Slope Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.00
60B: Canton-----	45	Good		Fair Rock fragments Hard to reclaim (rock fragments) Too acid	0.12 0.32 0.88
Charlton-----	35	Good		Poor Rock fragments Hard to reclaim (rock fragments)	0.00 0.68

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
60C: Canton-----	45	Good		Fair Rock fragments Hard to reclaim (rock fragments) Slope Too acid	0.12 0.32 0.37 0.88
Charlton-----	35	Good		Poor Rock fragments Slope Hard to reclaim (rock fragments)	0.00 0.37 0.68
60D: Canton-----	45	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments) Too acid	0.00 0.12 0.32 0.88
Charlton-----	35	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.68
61B: Canton-----	45	Good		Fair Rock fragments Hard to reclaim (rock fragments) Too acid	0.12 0.32 0.88
Charlton-----	35	Good		Poor Rock fragments Hard to reclaim (rock fragments)	0.00 0.68
61C: Canton-----	45	Good		Fair Rock fragments Hard to reclaim (rock fragments) Slope Too acid	0.12 0.32 0.37 0.88
Charlton-----	35	Good		Poor Rock fragments Slope Hard to reclaim (rock fragments)	0.00 0.37 0.68
62C: Canton-----	45	Good		Fair Rock fragments Hard to reclaim (rock fragments) Too acid Slope	0.12 0.32 0.88 0.96

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
62C: Charlton-----	35	Good		Poor Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.68 0.96
62D: Canton-----	45	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments) Too acid	0.00 0.12 0.32 0.88
Charlton-----	35	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.68
63B: Cheshire-----	80	Good		Fair Rock fragments Too sandy Too acid	0.12 0.98 0.98
63C: Cheshire-----	80	Good		Fair Rock fragments Slope Too sandy Too acid	0.12 0.37 0.98 0.98
63D: Cheshire-----	80	Fair Slope	0.50	Poor Slope Rock fragments Too sandy Too acid	0.00 0.12 0.98 0.98
64B: Cheshire-----	80	Good		Fair Rock fragments Too sandy Too acid	0.12 0.98 0.98
64C: Cheshire-----	80	Good		Fair Rock fragments Slope Too sandy Too acid	0.12 0.37 0.98 0.98
65C: Cheshire-----	80	Good		Fair Rock fragments Slope Too sandy Too acid	0.12 0.96 0.98 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
65D: Cheshire-----	80	Poor Slope	0.00	Poor Slope Rock fragments Too sandy Too acid	0.00 0.12 0.98 0.98
66B: Narragansett-----	80	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments) Too acid	0.00 0.04 0.68 0.76
66C: Narragansett-----	80	Good		Poor Rock fragments Too sandy Slope Hard to reclaim (rock fragments) Too acid	0.00 0.04 0.37 0.68 0.76
67B: Narragansett-----	80	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments) Too acid	0.00 0.04 0.68 0.76
67C: Narragansett-----	80	Good		Poor Rock fragments Too sandy Slope Hard to reclaim (rock fragments) Too acid	0.00 0.04 0.37 0.68 0.76
68C: Narragansett-----	80	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments) Too acid Slope	0.00 0.04 0.68 0.76 0.96
68D: Narragansett-----	80	Fair Slope	0.50	Poor Slope Rock fragments Too sandy Hard to reclaim (rock fragments) Too acid	0.00 0.00 0.04 0.68 0.76

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
69B: Yalesville-----	75	Poor Depth to bedrock	0.00	Fair Rock fragments Depth to bedrock Too acid	0.12 0.95 0.98
69C: Yalesville-----	75	Poor Depth to bedrock	0.00	Fair Rock fragments Slope Depth to bedrock Too acid	0.12 0.37 0.95 0.98
70C: Branford-----	50	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Slope Too acid	0.00 0.00 0.32 0.96 0.98
Holyoke-----	30	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.12 0.96
71C: Brookfield-----	45	Good		Fair Rock fragments Hard to reclaim (rock fragments) Slope Too acid	0.03 0.82 0.96 0.98
Brimfield-----	30	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope Too acid	0.00 0.12 0.96 0.98
Rock outcrop-----	15	Not rated		Not rated	
71E: Brookfield-----	45	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments) Too acid	0.00 0.03 0.82 0.98
Brimfield-----	30	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments Too acid	0.00 0.00 0.12 0.98
Rock outcrop-----	15	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
73C: Charlton-----	45	Good		Poor Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.68 0.96
Chatfield-----	30	Poor Depth to bedrock	0.00	Fair Rock fragments Depth to bedrock Slope	0.12 0.46 0.96
73E: Charlton-----	45	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.68
Chatfield-----	30	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Rock fragments Depth to bedrock	0.00 0.12 0.46
74C: Narragansett-----	55	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments) Too acid Slope	0.00 0.04 0.68 0.76 0.96
Hollis-----	20	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.28 0.96
75C: Hollis-----	35	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.28 0.96
Chatfield-----	30	Poor Depth to bedrock	0.00	Fair Rock fragments Depth to bedrock Slope	0.12 0.46 0.96
Rock outcrop-----	15	Not rated		Not rated	
75E: Hollis-----	35	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments	0.00 0.00 0.28

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
75E: Chatfield-----	30	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Rock fragments Depth to bedrock	0.00 0.12 0.46
Rock outcrop-----	15	Not rated		Not rated	
76E: Rock outcrop-----	55	Not rated		Not rated	
Hollis-----	25	Poor Depth to bedrock Slope	0.00 0.02	Poor Depth to bedrock Slope Rock fragments	0.00 0.00 0.28
76F: Rock outcrop-----	55	Not rated		Not rated	
Hollis-----	25	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments	0.00 0.00 0.28
77C: Cheshire-----	45	Good		Fair Rock fragments Slope Too sandy Too acid	0.12 0.96 0.98 0.98
Holyoke-----	35	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.12 0.96
77D: Cheshire-----	45	Poor Slope	0.00	Poor Slope Rock fragments Too sandy Too acid	0.00 0.12 0.98 0.98
Holyoke-----	35	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments	0.00 0.00 0.12
78C: Holyoke-----	50	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.12 0.96
Rock outcrop-----	25	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
78E: Holyoke-----	50	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments	0.00 0.00 0.12
Rock outcrop-----	25	Not rated		Not rated	
79E: Rock outcrop-----	55	Not rated		Not rated	
Holyoke-----	25	Poor Depth to bedrock Slope	0.00 0.02	Poor Depth to bedrock Slope Rock fragments	0.00 0.00 0.12
80B: Bernardston-----	80	Fair Wetness depth	0.32	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.10 0.32 0.32 0.98
80C: Bernardston-----	80	Fair Wetness depth	0.32	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Slope Too acid	0.10 0.32 0.32 0.37 0.98
81C: Bernardston-----	80	Fair Wetness depth	0.32	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Slope Too acid	0.10 0.32 0.32 0.96 0.98
81D: Bernardston-----	80	Fair Wetness depth Slope	0.32 0.50	Poor Slope Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.00 0.10 0.32 0.32 0.98
82B: Broadbrook-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.16 0.53 0.97 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
82C: Broadbrook-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Slope Wetness depth Rock fragments Too acid	0.16 0.37 0.53 0.97 0.98
82D: Broadbrook-----	80	Fair Slope Wetness depth	0.50 0.53	Poor Slope Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.00 0.16 0.53 0.97 0.98
83B: Broadbrook-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.16 0.53 0.97 0.98
83C: Broadbrook-----	80	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Slope Wetness depth Rock fragments Too acid	0.16 0.37 0.53 0.97 0.98
84B: Paxton-----	55	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
Montauk-----	30	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.50 0.76 0.98
84C: Paxton-----	55	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Slope Wetness depth Too acid	0.00 0.28 0.37 0.53 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
84C: Montauk-----	30	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Slope Rock fragments Wetness depth Too acid	0.00 0.37 0.50 0.76 0.98
84D: Paxton-----	55	Fair Slope Wetness depth	0.50 0.53	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.00 0.28 0.53 0.98
Montauk-----	30	Fair Slope Wetness depth	0.50 0.76	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.00 0.50 0.76 0.98
85B: Paxton-----	55	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
Montauk-----	30	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.50 0.76 0.98
85C: Paxton-----	55	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Slope Wetness depth Too acid	0.00 0.28 0.37 0.53 0.98
Montauk-----	30	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Slope Rock fragments Wetness depth Too acid	0.00 0.37 0.50 0.76 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
86C: Paxton-----	55	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Slope Too acid	0.00 0.28 0.53 0.96 0.98
Montauk-----	30	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Slope Too acid	0.00 0.50 0.76 0.96 0.98
86D: Paxton-----	55	Poor Slope Wetness depth	0.00 0.53	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.00 0.28 0.53 0.98
Montauk-----	30	Poor Slope Wetness depth	0.00 0.76	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.00 0.50 0.76 0.98
87B: Wethersfield-----	80	Fair Wetness depth	0.53	Fair Rock fragments Hard to reclaim (dense layer) Wetness depth Too acid	0.28 0.29 0.53 0.98
87C: Wethersfield-----	80	Fair Wetness depth	0.53	Fair Rock fragments Hard to reclaim (dense layer) Slope Wetness depth Too acid	0.28 0.29 0.37 0.53 0.98
87D: Wethersfield-----	80	Fair Slope Wetness depth	0.50 0.53	Poor Slope Rock fragments Hard to reclaim (dense layer) Wetness depth Too acid	0.00 0.28 0.29 0.53 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
88B: Wethersfield-----	80	Fair Wetness depth	0.53	Fair Rock fragments Hard to reclaim (dense layer) Wetness depth Too acid	0.28 0.29 0.53 0.98
88C: Wethersfield-----	80	Fair Wetness depth	0.53	Fair Rock fragments Hard to reclaim (dense layer) Slope Wetness depth Too acid	0.28 0.29 0.37 0.53 0.98
89C: Wethersfield-----	80	Fair Wetness depth	0.53	Fair Rock fragments Hard to reclaim (dense layer) Wetness depth Slope Too acid	0.28 0.29 0.53 0.96 0.98
89D: Wethersfield-----	80	Poor Slope Wetness depth	0.00 0.53	Poor Slope Rock fragments Hard to reclaim (dense layer) Wetness depth Too acid	0.00 0.28 0.29 0.53 0.98
90B: Stockbridge-----	80	Good		Fair Rock fragments Hard to reclaim (rock fragments)	0.03 0.68
90C: Stockbridge-----	80	Good		Fair Rock fragments Slope Hard to reclaim (rock fragments)	0.03 0.37 0.68
90D: Stockbridge-----	80	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.03 0.68
91B: Stockbridge-----	80	Good		Fair Rock fragments Hard to reclaim (rock fragments)	0.03 0.68

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
91C: Stockbridge-----	80	Good		Fair Rock fragments Slope Hard to reclaim (rock fragments)	0.03 0.37 0.68
91D: Stockbridge-----	80	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.03 0.68
92B: Nellis-----	85	Good		Poor Rock fragments Hard to reclaim (rock fragments)	0.00 0.82
92C: Nellis-----	85	Good		Poor Rock fragments Slope Hard to reclaim (rock fragments)	0.00 0.37 0.82
92D: Nellis-----	85	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.82
93C: Nellis-----	85	Good		Poor Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.82 0.96
94C: Farmington-----	40	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.12 0.96
Nellis-----	35	Good		Poor Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.82 0.96
94E: Farmington-----	40	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments	0.00 0.00 0.00 0.12

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
94E: Nellis-----	35	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.82
95C: Farmington-----	60	Poor Depth to bedrock	0.00	Poor Depth to bedrock Rock fragments Slope	0.00 0.12 0.96
Rock outcrop-----	20	Not rated		Not rated	
95E: Farmington-----	60	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments	0.00 0.00 0.12
Rock outcrop-----	20	Not rated		Not rated	
96: Ipswich-----	85	Poor Wetness depth Shrink-swell	0.00 0.00	Poor Wetness depth Organic matter content high Salinity	0.00 0.00 0.00
97: Pawcatuck-----	85	Poor Wetness depth Shrink-swell	0.00 0.09	Poor Wetness depth Salinity Organic matter content high	0.00 0.00 0.00
98: Westbrook-----	80	Poor Wetness depth Shrink-swell	0.00 0.02	Poor Wetness depth Salinity Organic matter content high	0.00 0.00 0.00
99: Westbrook, low salt-	80	Poor Wetness depth Shrink-swell	0.00 0.02	Poor Wetness depth Organic matter content high Salinity	0.00 0.00 0.00
100: Suncook-----	80	Good		Poor Too sandy Rock fragments	0.00 0.97

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
101: Occum-----	80	Good		Fair Hard to reclaim (rock fragments) Rock fragments	0.68 0.97
102: Pootatuck-----	80	Fair Wetness depth	0.53	Fair Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.53 0.97 0.98
103: Rippowam-----	80	Poor Wetness depth	0.00	Poor Too sandy Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.12 0.88
104: Bash-----	80	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments Too acid	0.00 0.28 0.98
105: Hadley-----	80	Good		Good	
106: Winooski-----	80	Fair Wetness depth	0.76	Fair Wetness depth	0.76
107: Limerick-----	50	Poor Wetness depth	0.00	Poor Wetness depth	0.00
Lim-----	30	Poor Wetness depth	0.00	Poor Too sandy Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.68
108: Saco-----	80	Poor Wetness depth	0.00	Poor Wetness depth	0.00
109: Fluvaquents, Frequently Flooded-	50	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth	0.00 0.00

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
109: Udifluvents, Frequently Flooded-	35	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments)	0.00 0.32 0.68
221A: Ninigret-----	40	Fair Wetness depth	0.53	Poor Too sandy Rock fragments Wetness depth Hard to reclaim (rock fragments)	0.00 0.00 0.53 0.82
Urban land-----	35	Not rated		Not rated	
224A: Deerfield-----	40	Fair Wetness depth	0.76	Poor Too sandy Wetness depth	0.00 0.76
Urban land-----	35	Not rated		Not rated	
225B: Brancroft-----	40	Poor Low strength Wetness depth	0.00 0.29	Fair Wetness depth	0.29
Urban land-----	35	Not rated		Not rated	
226B: Berlin-----	40	Poor Low strength Wetness depth	0.00 0.29	Fair Wetness depth Too clayey	0.29 0.55
Urban land-----	35	Not rated		Not rated	
228B: Elmridge-----	40	Poor Low strength Wetness depth Shrink-swell	0.00 0.53 0.86	Poor Too clayey Wetness depth	0.00 0.53
Urban land-----	35	Not rated		Not rated	
229B: Agawam-----	40	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.50
Urban land-----	35	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
229C: Agawam-----	40	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Slope	0.37
				Hard to reclaim (rock fragments)	0.50
Urban land-----	35	Not rated		Not rated	
230B: Branford-----	40	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.32
				Too acid	0.98
Urban land-----	35	Not rated		Not rated	
230C: Branford-----	40	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.32
				Slope	0.37
				Too acid	0.98
Urban land-----	35	Not rated		Not rated	
232B: Haven-----	40	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.02
				Too acid	0.98
Urban land-----	35	Not rated		Not rated	
234B: Merrimac-----	40	Good		Poor	
				Too sandy	0.00
				Rock fragments	0.00
				Hard to reclaim (rock fragments)	0.00
Urban land-----	35	Not rated		Not rated	
235B: Penwood-----	40	Good		Poor	
				Too sandy	0.00
				Too acid	0.98
Urban land-----	35	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
236B: Windsor-----	40	Good		Poor Too sandy Rock fragments	0.00 0.97
Urban land-----	35	Not rated		Not rated	
237A: Manchester-----	40	Good		Poor Too sandy Hard to reclaim (rock fragments) Rock fragments Too acid	0.00 0.00 0.00 0.98
Urban land-----	35	Not rated		Not rated	
237C: Manchester-----	40	Good		Poor Too sandy Hard to reclaim (rock fragments) Rock fragments Slope Too acid	0.00 0.00 0.00 0.96 0.98
Urban land-----	35	Not rated		Not rated	
238A: Hinckley-----	40	Good		Poor Hard to reclaim (rock fragments) Rock fragments Too sandy Too acid	0.00 0.00 0.32 0.76
Urban land-----	35	Not rated		Not rated	
238C: Hinckley-----	40	Good		Poor Hard to reclaim (rock fragments) Rock fragments Too sandy Too acid Slope	0.00 0.00 0.32 0.76 0.96
Urban land-----	35	Not rated		Not rated	
240B: Ludlow-----	40	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
240B: Urban land-----	35	Not rated		Not rated	
243B: Rainbow-----	40	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.20 0.53 0.97 0.98
Urban land-----	35	Not rated		Not rated	
245B: Woodbridge-----	40	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
Urban land-----	35	Not rated		Not rated	
245C: Woodbridge-----	40	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Slope Wetness depth Too acid	0.00 0.28 0.37 0.53 0.98
Urban land-----	35	Not rated		Not rated	
248B: Georgia-----	40	Fair Wetness depth	0.76	Fair Rock fragments Wetness depth	0.12 0.76
Urban land-----	35	Not rated		Not rated	
250B: Sutton-----	40	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth Too acid	0.03 0.53 0.98
Urban land-----	35	Not rated		Not rated	
253B: Wapping-----	40	Fair Wetness depth	0.53	Fair Hard to reclaim (rock fragments) Wetness depth Rock fragments	0.18 0.53 0.97

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
253B: Urban land-----	35	Not rated		Not rated	
255B: Watchaug-----	40	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth	0.28 0.53
Urban land-----	35	Not rated		Not rated	
260B: Charlton-----	40	Good		Poor Rock fragments Hard to reclaim (rock fragments)	0.00 0.68
Urban land-----	35	Not rated		Not rated	
260C: Charlton-----	40	Good		Poor Rock fragments Slope Hard to reclaim (rock fragments)	0.00 0.37 0.68
Urban land-----	35	Not rated		Not rated	
260D: Charlton-----	40	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.68
Urban land-----	35	Not rated		Not rated	
263B: Cheshire-----	40	Good		Fair Rock fragments Too sandy Too acid	0.12 0.98 0.98
Urban land-----	35	Not rated		Not rated	
263C: Cheshire-----	40	Good		Fair Rock fragments Slope Too sandy Too acid	0.12 0.37 0.98 0.98
Urban land-----	35	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
266B: Narragansett-----	40	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments) Too acid	0.00 0.04 0.68 0.76
Urban land-----	35	Not rated		Not rated	
269B: Yalesville-----	40	Poor Depth to bedrock	0.00	Fair Rock fragments Depth to bedrock Too acid	0.12 0.95 0.98
Urban land-----	35	Not rated		Not rated	
269C: Yalesville-----	40	Poor Depth to bedrock	0.00	Fair Rock fragments Slope Depth to bedrock Too acid	0.12 0.37 0.95 0.98
Urban land-----	35	Not rated		Not rated	
273C: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Good		Poor Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.68 0.96
Chatfield-----	15	Poor Depth to bedrock	0.00	Fair Rock fragments Depth to bedrock Slope	0.12 0.46 0.96
273E: Urban land-----	35	Not rated		Not rated	
Charlton-----	25	Poor Slope	0.00	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.68
Chatfield-----	15	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Rock fragments Depth to bedrock	0.00 0.12 0.46

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
275C: Urban land-----	45	Not rated		Not rated	
Chatfield-----	30	Poor Depth to bedrock	0.00	Fair Rock fragments Depth to bedrock Slope	0.12 0.46 0.96
275E: Urban land-----	35	Not rated		Not rated	
Chatfield-----	25	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Rock fragments Depth to bedrock	0.00 0.12 0.46
Rock outcrop-----	15	Not rated		Not rated	
282B: Broadbrook-----	40	Fair Wetness depth	0.53	Fair Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.16 0.53 0.97 0.98
Urban land-----	35	Not rated		Not rated	
284B: Paxton-----	40	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.28 0.53 0.98
Urban land-----	35	Not rated		Not rated	
284C: Paxton-----	40	Fair Wetness depth	0.53	Poor Hard to reclaim (dense layer) Rock fragments Slope Wetness depth Too acid	0.00 0.28 0.37 0.53 0.98
Urban land-----	35	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
284D: Paxton-----	40	Fair Slope Wetness depth	0.50 0.53	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.00 0.28 0.53 0.98
Urban land-----	35	Not rated		Not rated	
287B: Wethersfield-----	40	Fair Wetness depth	0.53	Fair Rock fragments Hard to reclaim (dense layer) Wetness depth Too acid	0.28 0.29 0.53 0.98
Urban land-----	35	Not rated		Not rated	
287C: Wethersfield-----	40	Fair Wetness depth	0.53	Fair Rock fragments Hard to reclaim (dense layer) Slope Wetness depth Too acid	0.28 0.29 0.37 0.53 0.98
Urban land-----	35	Not rated		Not rated	
287D: Wethersfield-----	40	Fair Slope Wetness depth	0.50 0.53	Poor Slope Rock fragments Hard to reclaim (dense layer) Wetness depth Too acid	0.00 0.28 0.29 0.53 0.98
Urban land-----	35	Not rated		Not rated	
290B: Stockbridge-----	40	Good		Fair Rock fragments Hard to reclaim (rock fragments)	0.03 0.68
Urban land-----	35	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
290C: Stockbridge-----	40	Good		Fair Rock fragments Slope Hard to reclaim (rock fragments)	0.03 0.37 0.68
Urban land-----	35	Not rated		Not rated	
290D: Stockbridge-----	40	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.03 0.68
Urban land-----	35	Not rated		Not rated	
301: Beaches-----	50	Not rated		Poor Too sandy Wetness depth Salinity Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.00 0.82
Udipsamments-----	35	Good		Poor Too sandy Rock fragments	0.00 0.97
302: Dumps-----	95	Not rated		Not rated	
303: Pits, quarries-----	90	Not rated		Not rated	
304: Udorthents-----	90	Poor Slope Stone content	0.00 0.73	Poor Slope Hard to reclaim (dense layer) Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.50
305: Udorthents-----	65	Fair Stone content Slope	0.73 0.82	Poor Hard to reclaim (dense layer) Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.50

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
305: Pits-----	25	Poor Slope	0.00	Poor Too sandy Hard to reclaim (rock fragments) Rock fragments Slope	0.00 0.00 0.00 0.00
306: Udorthents-----	50	Fair Stone content Slope	0.73 0.82	Poor Hard to reclaim (dense layer) Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.50
Urban land-----	35	Not rated		Not rated	
307: Urban land-----	80	Not rated		Not rated	
308: Udorthents-----	80	Fair Stone content Slope	0.73 0.82	Poor Hard to reclaim (dense layer) Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.50
309: Udorthents-----	80	Fair Stone content Slope	0.73 0.82	Poor Hard to reclaim (dense layer) Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.50
310: Udorthents, Periodically Flooded-----	85	Fair Stone content Slope	0.73 0.82	Poor Hard to reclaim (dense layer) Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.50
401C: Macomber-----	55	Poor Depth to bedrock	0.00	Poor Rock fragments Slope Depth to bedrock Too acid	0.00 0.37 0.54 0.92

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
401C: Taconic-----	30	Poor Depth to bedrock	0.00	Poor Rock fragments Depth to bedrock Too acid	0.00 0.00 0.88
402D: Macomber-----	50	Poor Depth to bedrock Slope	0.00 0.50	Poor Slope Rock fragments Depth to bedrock Too acid	0.00 0.00 0.54 0.92
Taconic-----	25	Poor Depth to bedrock Slope	0.00 0.50	Poor Slope Rock fragments Depth to bedrock Too acid	0.00 0.00 0.00 0.88
Rock outcrop-----	15	Not rated		Not rated	
403C: Taconic-----	70	Poor Depth to bedrock	0.00	Poor Rock fragments Depth to bedrock Too acid	0.00 0.00 0.88
Rock outcrop-----	25	Not rated		Not rated	
403E: Taconic-----	70	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Rock fragments Depth to bedrock Too acid	0.00 0.00 0.00 0.88
Rock outcrop-----	20	Not rated		Not rated	
403F: Taconic-----	70	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Rock fragments Depth to bedrock Too acid	0.00 0.00 0.00 0.88
Rock outcrop-----	20	Not rated		Not rated	
405C: Dummerston-----	85	Fair Stone content	0.96	Fair Rock fragments Hard to reclaim (rock fragments) Too acid	0.03 0.98 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
405E: Dummerston-----	85	Poor Slope Stone content	0.00 0.96	Poor Slope Rock fragments Hard to reclaim (rock fragments) Too acid	0.00 0.03 0.98 0.98
407C: Lanesboro-----	85	Fair Wetness depth Stone content	0.53 0.94	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Slope Too acid	0.00 0.00 0.53 0.84 0.98
407E: Lanesboro-----	85	Poor Slope Wetness depth Stone content	0.00 0.53 0.94	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.00 0.00 0.53 0.98
408C: Fullam-----	85	Fair Wetness depth Stone content	0.62 0.92	Poor Hard to reclaim (dense layer) Rock fragments Wetness depth Slope Too acid	0.00 0.00 0.62 0.84 0.98
409B: Brayton-----	85	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments	0.00 0.00 0.12
412B: Bice-----	85	Good		Fair Rock fragments Hard to reclaim (rock fragments)	0.50 0.95
412C: Bice-----	85	Good		Fair Rock fragments Slope Hard to reclaim (rock fragments)	0.50 0.84 0.95
412D: Bice-----	85	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.50 0.95

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
413C: Bice-----	45	Good		Fair	
				Rock fragments	0.50
				Hard to reclaim (rock fragments)	0.95
Millsite-----	40	Poor		Fair	
		Depth to bedrock	0.00	Rock fragments	0.28
		Stone content	0.99	Depth to bedrock	0.65
413E: Bice-----	45	Poor		Poor	
		Slope	0.00	Slope	0.00
				Rock fragments	0.50
				Hard to reclaim (rock fragments)	0.95
Millsite-----	40	Poor		Poor	
		Depth to bedrock	0.00	Slope	0.00
		Slope	0.00	Rock fragments	0.28
		Stone content	0.99	Depth to bedrock	0.65
414: Fredon, cold-----	85	Poor		Poor	
		Wetness depth	0.00	Wetness depth	0.00
				Rock fragments	0.02
				Too sandy	0.22
				Hard to reclaim (rock fragments)	0.97
415C: Millsite-----	40	Poor		Fair	
		Depth to bedrock	0.00	Rock fragments	0.28
		Stone content	0.99	Depth to bedrock	0.65
Westminster-----	40	Poor		Poor	
		Depth to bedrock	0.00	Depth to bedrock	0.00
				Rock fragments	0.50
				Too acid	0.76
Rock outcrop-----	15	Not rated		Not rated	
415E: Millsite-----	40	Poor		Poor	
		Depth to bedrock	0.00	Slope	0.00
		Slope	0.00	Rock fragments	0.28
		Stone content	0.99	Depth to bedrock	0.65
Westminster-----	40	Poor		Poor	
		Depth to bedrock	0.00	Slope	0.00
		Slope	0.00	Depth to bedrock	0.00
				Rock fragments	0.50
				Too acid	0.76
Rock outcrop-----	15	Not rated		Not rated	

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
416E: Rock outcrop-----	70	Not rated		Not rated	
Westminster-----	20	Poor Depth to bedrock Slope	0.00 0.00	Poor Depth to bedrock Slope Rock fragments Too acid	0.00 0.00 0.50 0.76
416F: Rock outcrop-----	70	Not rated		Not rated	
Westminster-----	20	Poor Depth to bedrock Slope	0.00 0.00	Poor Slope Depth to bedrock Rock fragments Too acid	0.00 0.00 0.50 0.76
417B: Bice-----	85	Good		Fair Rock fragments Hard to reclaim (rock fragments)	0.50 0.95
417C: Bice-----	85	Good		Fair Rock fragments Slope Hard to reclaim (rock fragments)	0.50 0.84 0.95
417D: Bice-----	85	Fair Slope	0.50	Poor Slope Rock fragments Hard to reclaim (rock fragments)	0.00 0.50 0.95
418C: Schroon-----	85	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth Slope	0.12 0.53 0.96
420A: Schroon-----	85	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth	0.12 0.53
420B: Schroon-----	85	Fair Wetness depth	0.53	Fair Rock fragments Wetness depth	0.12 0.53

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
421A: Ninigret, cold-----	85	Fair Wetness depth	0.53	Poor Too sandy Rock fragments Wetness depth Hard to reclaim (rock fragments)	0.00 0.00 0.53 0.82
423A: Sudbury, cold-----	85	Fair Wetness depth	0.76	Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Wetness depth	0.00 0.00 0.08 0.76
424B: Shelburne-----	85	Fair Wetness depth Stone content	0.53 0.92	Fair Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.26 0.50 0.53 0.98
424C: Shelburne-----	85	Fair Wetness depth Stone content	0.53 0.92	Fair Hard to reclaim (dense layer) Slope Rock fragments Wetness depth Too acid	0.26 0.37 0.50 0.53 0.98
424D: Shelburne-----	85	Fair Slope Wetness depth Stone content	0.50 0.53 0.92	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.26 0.50 0.53 0.98
425B: Shelburne-----	85	Fair Wetness depth Stone content	0.53 0.92	Fair Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.26 0.50 0.53 0.98
425C: Shelburne-----	85	Fair Wetness depth Stone content	0.53 0.92	Fair Hard to reclaim (dense layer) Slope Rock fragments Wetness depth Too acid	0.26 0.37 0.50 0.53 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
426D: Shelburne-----	85	Poor Slope Wetness depth Stone content	0.00 0.53 0.92	Poor Slope Hard to reclaim (dense layer) Rock fragments Wetness depth Too acid	0.00 0.26 0.50 0.53 0.98
427B: Ashfield-----	85	Fair Wetness depth	0.14	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments	0.00 0.14 0.50
427C: Ashfield-----	85	Fair Wetness depth	0.14	Poor Hard to reclaim (dense layer) Wetness depth Slope Rock fragments Too acid	0.00 0.14 0.37 0.50 0.88
428A: Ashfield-----	85	Fair Wetness depth	0.14	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.00 0.14 0.50 0.88
428B: Ashfield-----	85	Fair Wetness depth	0.14	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments Too acid	0.00 0.14 0.50 0.88
428C: Ashfield-----	85	Fair Wetness depth	0.14	Poor Hard to reclaim (dense layer) Wetness depth Slope Rock fragments Too acid	0.00 0.14 0.37 0.50 0.88
429A: Agawam, cold-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.50

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
429B: Agawam, cold-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.50
429C: Agawam, cold-----	80	Good		Poor Too sandy Rock fragments Slope Hard to reclaim (rock fragments)	0.00 0.00 0.37 0.50
433: Moosilauke-----	80	Poor Wetness depth	0.00	Poor Too sandy Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00 0.82
434A: Merrimac, cold-----	80	Good		Poor Rock fragments Too sandy Hard to reclaim (rock fragments)	0.00 0.00 0.00
434B: Merrimac, cold-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments)	0.00 0.00 0.00
434C: Merrimac, cold-----	80	Good		Poor Too sandy Rock fragments Hard to reclaim (rock fragments) Slope	0.00 0.00 0.00 0.37
435: Scarboro-----	80	Poor Wetness depth	0.00	Poor Too sandy Wetness depth Hard to reclaim (rock fragments) Rock fragments Too acid	0.00 0.00 0.82 0.97 0.98
436: Halsey-----	80	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (rock fragments) Rock fragments	0.00 0.50 0.97

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
437: Wonsqueak-----	85	Poor Wetness depth Shrink-swell	0.00 0.98	Poor Wetness depth Rock fragments	0.00 0.98
438: Bucksport-----	85	Poor Wetness depth Shrink-swell	0.00 0.00	Poor Wetness depth Organic matter content high Too acid	0.00 0.00 0.88
440A: Boscawen-----	80	Good		Poor Too sandy Rock fragments Too acid	0.00 0.00 0.99
440C: Boscawen-----	80	Good		Poor Too sandy Rock fragments Slope Too acid	0.00 0.00 0.96 0.99
440E: Boscawen-----	85	Poor Slope	0.00	Poor Slope Too sandy Rock fragments Too acid	0.00 0.00 0.00 0.99
442: Brayton-----	85	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth Too acid Rock fragments	0.00 0.00 0.88 0.88
443: Brayton-----	50	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth Too acid Rock fragments	0.00 0.00 0.88 0.88
Loonmeadow-----	35	Poor Wetness depth	0.00	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments	0.00 0.00 0.28
448B: Hogansburg-----	85	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.76 0.97 0.98

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
449B: Hogansburg-----	85	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.76 0.97 0.98
449C: Hogansburg-----	85	Fair Wetness depth	0.76	Poor Hard to reclaim (dense layer) Slope Wetness depth Rock fragments Hard to reclaim (rock fragments)	0.00 0.37 0.76 0.97 0.98
450B: Pyrities-----	80	Good		Fair Rock fragments	0.88
450C: Pyrities-----	80	Good		Fair Slope Rock fragments	0.37 0.88
450D: Pyrities-----	80	Fair Slope	0.50	Poor Slope Rock fragments	0.00 0.88
451B: Pyrities-----	80	Good		Fair Rock fragments	0.88
451C: Pyrities-----	80	Good		Fair Slope Rock fragments	0.37 0.88
451D: Pyrities-----	80	Poor Slope	0.00	Poor Slope Rock fragments	0.00 0.88
457: Mudgepond-----	80	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments	0.00 0.88
458: Mudgepond-----	55	Poor Wetness depth	0.00	Poor Wetness depth Rock fragments	0.00 0.88
Alden-----	35	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (rock fragments)	0.00 0.88

Table 21.—Construction Materials—Continued

Map symbol and soil name	Pct. of map unit	Potential source of roadfill		Potential source of topsoil	
		Rating class and limiting features	Value	Rating class and limiting features	Value
501: Ondawa-----	85	Good		Fair Rock fragments Too acid	0.88 0.88
503: Rumney-----	80	Poor Wetness depth	0.00	Poor Wetness depth Hard to reclaim (rock fragments) Rock fragments	0.00 0.32 0.88
508: Medomak-----	85	Poor Wetness depth	0.00	Poor Wetness depth	0.00

Table 22.—Ponds and Embankments

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Thin layer Seepage	1.00 0.99 0.01	Very limited Depth to water	1.00
3: Ridgebury-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone Thin layer Seepage	1.00 0.99 0.01	Very limited Depth to water	1.00
Leicester-----	35	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.01	Very limited Cutbanks cave	1.00
Whitman-----	15	Not limited		Very limited Depth to saturated zone Thin layer Ponding	1.00 1.00 1.00	Very limited Depth to water	1.00
4: Leicester-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.01	Very limited Cutbanks cave	1.00
5: Wilbraham-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping Thin layer	1.00 1.00 0.99	Very limited Depth to water	1.00
6: Wilbraham-----	60	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping Thin layer	1.00 1.00 0.99	Very limited Depth to water	1.00
Menlo-----	25	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping Ponding Thin layer	1.00 1.00 1.00 0.93	Very limited Depth to water	1.00

Table 22.--Ponds and Embankments--Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7: Mudgepond-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	1.00 1.00	Very limited Cutbanks cave	1.00
8: Mudgepond-----	45	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	1.00 1.00	Very limited Cutbanks cave	1.00
Alden-----	35	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping Ponding	1.00 1.00 1.00	Very limited Cutbanks cave Slow refill	1.00 0.30
9: Scitico-----	40	Not limited		Very limited Depth to saturated zone Piping	1.00 0.16	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
Shaker-----	30	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	1.00 0.96	Somewhat limited Cutbanks cave	0.10
Maybid-----	15	Not limited		Very limited Depth to saturated zone Ponding Piping	1.00 1.00 0.20	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
10: Raynham-----	80	Somewhat limited Seepage	0.57	Very limited Depth to saturated zone Piping	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.43 0.10
12: Raypol-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.10	Very limited Cutbanks cave	1.00
13: Walpole-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.15	Very limited Cutbanks cave	1.00
14: Fredon-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.43	Very limited Cutbanks cave	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
15: Scarboro-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 0.36	Very limited Cutbanks cave	1.00
16: Halsey-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 0.31	Very limited Cutbanks cave	1.00
17: Timakwa-----	45	Very limited Seepage	1.00	Very limited Organic matter content Depth to saturated zone Piping Ponding	1.00 1.00 1.00 1.00	Very limited Cutbanks cave	1.00
Natchaug-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping Ponding	1.00 1.00 1.00	Somewhat limited Cutbanks cave	0.10
18: Catden-----	40	Very limited Seepage	1.00	Not rated		Somewhat limited Cutbanks cave	0.10
Freetown-----	40	Very limited Seepage	1.00	Not rated		Somewhat limited Cutbanks cave	0.10
20A: Ellington-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.11	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
21A: Ninigret-----	60	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
Tisbury-----	25	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.10	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
22A: Hero-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.11	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
22B: Hero-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.11	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
23A: Sudbury-----	80	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage	0.95 0.10	Very limited Cutbanks cave Depth to saturated zone	1.00 0.02
24A: Deerfield-----	80	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage	0.95 0.13	Very limited Cutbanks cave Depth to saturated zone	1.00 0.02
25A: Brancroft-----	80	Somewhat limited Seepage	0.01	Very limited Depth to saturated zone Piping	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.99 0.10
25B: Brancroft-----	80	Somewhat limited Seepage	0.01	Very limited Depth to saturated zone Piping	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.99 0.10
25C: Brancroft-----	80	Somewhat limited Seepage Slope	0.01 0.01	Very limited Depth to saturated zone Piping	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.99 0.10
26A: Berlin-----	80	Somewhat limited Seepage	0.01	Very limited Depth to saturated zone Piping	1.00 0.97	Very limited Slow refill Cutbanks cave	1.00 0.10
26B: Berlin-----	80	Somewhat limited Seepage	0.01	Very limited Depth to saturated zone Piping	1.00 0.97	Very limited Slow refill Cutbanks cave	1.00 0.10

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
27A: Belgrade-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone	1.00 0.84	Somewhat limited Slow refill Cutbanks cave Depth to saturated zone	0.30 0.10 0.07
28A: Elmridge-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	0.99 0.84	Very limited Depth to water	1.00
28B: Elmridge-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	0.99 0.84	Very limited Depth to water	1.00
29A: Agawam-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
29B: Agawam-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
29C: Agawam-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
30A: Branford-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
30B: Branford-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
30C: Branford-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
31A: Copake-----	85	Very limited Seepage	1.00	Somewhat limited Seepage	0.43	Very limited Depth to water	1.00
31B: Copake-----	85	Very limited Seepage	1.00	Somewhat limited Seepage	0.43	Very limited Depth to water	1.00
31C: Copake-----	85	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.43	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
32A: Haven-----	60	Very limited Seepage	1.00	Somewhat limited Seepage	0.39	Very limited Depth to water	1.00
Enfield-----	25	Very limited Seepage	1.00	Somewhat limited Seepage	0.50	Very limited Depth to water	1.00
32B: Haven-----	60	Very limited Seepage	1.00	Somewhat limited Seepage	0.39	Very limited Depth to water	1.00
Enfield-----	25	Very limited Seepage	1.00	Somewhat limited Seepage	0.50	Very limited Depth to water	1.00
32C: Haven-----	60	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.39	Very limited Depth to water	1.00
Enfield-----	25	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.50	Very limited Depth to water	1.00
33A: Hartford-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.10	Very limited Depth to water	1.00
33B: Hartford-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.10	Very limited Depth to water	1.00
34A: Merrimac-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
34B: Merrimac-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
34C: Merrimac-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
35A: Penwood-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.15	Very limited Depth to water	1.00
35B: Penwood-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.15	Very limited Depth to water	1.00
36A: Windsor-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.10	Very limited Depth to water	1.00
36B: Windsor-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.10	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
36C: Windsor-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.10	Very limited Depth to water	1.00
37A: Manchester-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.11	Very limited Depth to water	1.00
37C: Manchester-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.11	Very limited Depth to water	1.00
37E: Manchester-----	80	Very limited Seepage Slope	1.00 0.50	Somewhat limited Seepage	0.11	Very limited Depth to water	1.00
38A: Hinckley-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
38C: Hinckley-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
38E: Hinckley-----	80	Very limited Seepage Slope	1.00 0.50	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
39A: Groton-----	85	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
39C: Groton-----	85	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
39E: Groton-----	85	Very limited Seepage Slope	1.00 0.50	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
40A: Ludlow-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
40B: Ludlow-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
41B: Ludlow-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
42C: Ludlow-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
43A: Rainbow-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
43B: Rainbow-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
44B: Rainbow-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
45A: Woodbridge-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.86	Very limited Depth to water	1.00
45B: Woodbridge-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.86	Very limited Depth to water	1.00
45C: Woodbridge-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Depth to saturated zone Thin layer	0.99 0.86	Very limited Depth to water	1.00
46B: Woodbridge-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.86	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
46C: Woodbridge-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
		Slope	0.01	Thin layer	0.86		
47C: Woodbridge-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
				Thin layer	0.86		
48B: Georgia-----	50	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Cutbanks cave	1.00
				Depth to saturated zone	0.95	Slow refill	0.95
						Depth to saturated zone	0.02
Amenia-----	35	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Cutbanks cave	1.00
				Depth to saturated zone	0.95	Slow refill	0.95
						Depth to saturated zone	0.02
48C: Georgia-----	50	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Cutbanks cave	1.00
		Slope	0.01	Depth to saturated zone	0.95	Slow refill	0.95
						Depth to saturated zone	0.02
Amenia-----	35	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Cutbanks cave	1.00
		Slope	0.01	Depth to saturated zone	0.95	Slow refill	0.95
						Depth to saturated zone	0.02
49B: Georgia-----	50	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Cutbanks cave	1.00
				Depth to saturated zone	0.95	Slow refill	0.95
						Depth to saturated zone	0.02
Amenia-----	35	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Cutbanks cave	1.00
				Depth to saturated zone	0.95	Slow refill	0.95
						Depth to saturated zone	0.02

Table 22.--Ponds and Embankments--Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
49C: Georgia-----	50	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Slow refill Depth to saturated zone	1.00 0.95 0.02
Amenia-----	35	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Slow refill Depth to saturated zone	1.00 0.95 0.02
50A: Sutton-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
50B: Sutton-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
51B: Sutton-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
52C: Sutton-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
53A: Wapping-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.09	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
53B: Wapping-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.09	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
54B: Wapping-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.09	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
55A: Watchaug-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
55B: Watchaug-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
56B: Watchaug-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
57B: Gloucester-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
57C: Gloucester-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
57D: Gloucester-----	80	Very limited Seepage Slope	1.00 0.12	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
58B: Gloucester-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
58C: Gloucester-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
59C: Gloucester-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
59D: Gloucester-----	80	Very limited Seepage Slope	1.00 0.12	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00

Table 22.--Ponds and Embankments--Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
60B:							
Canton-----	45	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Charlton-----	35	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
60C:							
Canton-----	45	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Charlton-----	35	Very limited Seepage Slope	1.00 0.01	Not limited		Very limited Depth to water	1.00
60D:							
Canton-----	45	Very limited Seepage Slope	1.00 0.12	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Charlton-----	35	Very limited Seepage Slope	1.00 0.12	Not limited		Very limited Depth to water	1.00
61B:							
Canton-----	45	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Charlton-----	35	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
61C:							
Canton-----	45	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Charlton-----	35	Very limited Seepage Slope	1.00 0.01	Not limited		Very limited Depth to water	1.00
62C:							
Canton-----	45	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Charlton-----	35	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
62D:							
Canton-----	45	Very limited Seepage Slope	1.00 0.28	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Charlton-----	35	Very limited Seepage Slope	1.00 0.28	Not limited		Very limited Depth to water	1.00
63B:							
Cheshire-----	80	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
63C: Cheshire-----	80	Very limited Seepage Slope	1.00 0.01	Not limited		Very limited Depth to water	1.00
63D: Cheshire-----	80	Very limited Seepage Slope	1.00 0.12	Not limited		Very limited Depth to water	1.00
64B: Cheshire-----	80	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
64C: Cheshire-----	80	Very limited Seepage Slope	1.00 0.01	Not limited		Very limited Depth to water	1.00
65C: Cheshire-----	80	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
65D: Cheshire-----	80	Very limited Seepage Slope	1.00 0.28	Not limited		Very limited Depth to water	1.00
66B: Narragansett-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
66C: Narragansett-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
67B: Narragansett-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
67C: Narragansett-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
68C: Narragansett-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
68D: Narragansett-----	80	Very limited Seepage Slope	1.00 0.12	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
69B: Yalesville-----	75	Very limited Seepage Depth to bedrock	1.00 0.65	Somewhat limited Thin layer Seepage	0.65 0.01	Very limited Depth to water	1.0

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
69C: Yalesville-----	75	Very limited Seepage Depth to bedrock Slope	1.00 0.65 0.01	Somewhat limited Thin layer Seepage	0.65 0.01	Very limited Depth to water	1.00
70C: Branford-----	50	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
Holyoke-----	30	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
71C: Brookfield-----	45	Very limited Seepage	1.00	Somewhat limited Seepage	0.01	Very limited Depth to water	1.00
Brimfield-----	30	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock	1.00	Not rated		Not rated	
71E: Brookfield-----	45	Very limited Seepage Slope	1.00 0.50	Somewhat limited Seepage	0.01	Very limited Depth to water	1.00
Brimfield-----	30	Very limited Depth to bedrock Slope	1.00 0.50	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock Slope	1.00 0.50	Not rated		Not rated	
73C: Charlton-----	45	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
Chatfield-----	30	Very limited Seepage Depth to bedrock	1.00 0.88	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
73E: Charlton-----	45	Very limited Seepage Slope	1.00 0.50	Not limited		Very limited Depth to water	1.00
Chatfield-----	30	Very limited Seepage Depth to bedrock Slope	1.00 0.88 0.50	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
74C: Narragansett-----	55	Very limited Seepage	1.00	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
Hollis-----	20	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
75C: Hollis-----	35	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Chatfield-----	30	Very limited Seepage Depth to bedrock	1.00 0.88	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock	1.00	Not rated		Not rated	
75E: Hollis-----	35	Very limited Depth to bedrock Slope	1.00 0.50	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Chatfield-----	30	Very limited Seepage Depth to bedrock Slope	1.00 0.88 0.50	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock Slope	1.00 0.50	Not rated		Not rated	
76E: Rock outcrop-----	55	Very limited Depth to bedrock Slope	1.00 0.24	Not rated		Not rated	
Hollis-----	25	Very limited Depth to bedrock Slope	1.00 0.24	Very limited Thin layer	1.00	Very limited Depth to water	1.00
76F: Rock outcrop-----	55	Very limited Depth to bedrock Slope	1.00 1.00	Not rated		Not rated	
Hollis-----	25	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
77C: Cheshire-----	45	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
Holyoke-----	35	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
77D: Cheshire-----	45	Very limited Seepage Slope	1.00 0.28	Not limited		Very limited Depth to water	1.00
Holyoke-----	35	Very limited Depth to bedrock Slope	1.00 0.28	Very limited Thin layer	1.00	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
78C: Holyoke-----	50	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	25	Very limited Depth to bedrock	1.00	Not rated		Not rated	
78E: Holyoke-----	50	Very limited Depth to bedrock Slope	1.00 0.50	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	25	Very limited Depth to bedrock Slope	1.00 0.50	Not rated		Not rated	
79E: Rock outcrop-----	55	Very limited Depth to bedrock Slope	1.00 0.24	Not rated		Not rated	
Holyoke-----	25	Very limited Depth to bedrock Slope	1.00 0.24	Very limited Thin layer	1.00	Very limited Depth to water	1.00
80B: Bernardston-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping Thin layer	1.00 1.00 0.95	Very limited Depth to water	1.00
80C: Bernardston-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Depth to saturated zone Piping Thin layer	1.00 1.00 0.95	Very limited Depth to water	1.00
81C: Bernardston-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping Thin layer	1.00 1.00 0.95	Very limited Depth to water	1.00
81D: Bernardston-----	80	Somewhat limited Seepage Slope	0.70 0.12	Very limited Depth to saturated zone Piping Thin layer	1.00 1.00 0.95	Very limited Depth to water	1.00
82B: Broadbrook-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.96	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
82C: Broadbrook-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.96	Very limited Depth to water	1.00
82D: Broadbrook-----	80	Somewhat limited Seepage Slope	0.70 0.12	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.96	Very limited Depth to water	1.00
83B: Broadbrook-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.96	Very limited Depth to water	1.00
83C: Broadbrook-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.96	Very limited Depth to water	1.00
84B: Paxton-----	55	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00
Montauk-----	30	Very limited Seepage	1.00	Somewhat limited Thin layer Depth to saturated zone Seepage	0.96 0.95 0.04	Very limited Depth to water	1.00
84C: Paxton-----	55	Somewhat limited Seepage Slope	0.70 0.01	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00
Montauk-----	30	Very limited Seepage Slope	1.00 0.01	Somewhat limited Thin layer Depth to saturated zone Seepage	0.96 0.95 0.04	Very limited Depth to water	1.00
84D: Paxton-----	55	Somewhat limited Seepage Slope	0.70 0.12	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
84D: Montauk-----	30	Very limited Seepage Slope	1.00 0.12	Somewhat limited Thin layer Depth to saturated zone Seepage	0.96 0.95 0.04	Very limited Depth to water	1.00
85B: Paxton-----	55	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00
Montauk-----	30	Very limited Seepage	1.00	Somewhat limited Thin layer Depth to saturated zone Seepage	0.96 0.95 0.04	Very limited Depth to water	1.00
85C: Paxton-----	55	Somewhat limited Seepage Slope	0.70 0.01	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00
Montauk-----	30	Very limited Seepage Slope	1.00 0.01	Somewhat limited Thin layer Depth to saturated zone Seepage	0.96 0.95 0.04	Very limited Depth to water	1.00
86C: Paxton-----	55	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00
Montauk-----	30	Very limited Seepage	1.00	Somewhat limited Thin layer Depth to saturated zone Seepage	0.96 0.95 0.04	Very limited Depth to water	1.00
86D: Paxton-----	55	Somewhat limited Seepage Slope	0.70 0.28	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00
Montauk-----	30	Very limited Seepage Slope	1.00 0.28	Somewhat limited Thin layer Depth to saturated zone Seepage	0.96 0.95 0.04	Very limited Depth to water	1.00
87B: Wethersfield-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.93	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
87C: Wethersfield-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.93	Very limited Depth to water	1.00
87D: Wethersfield-----	80	Somewhat limited Seepage Slope	0.70 0.12	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.93	Very limited Depth to water	1.00
88B: Wethersfield-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.93	Very limited Depth to water	1.00
88C: Wethersfield-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.93	Very limited Depth to water	1.00
89C: Wethersfield-----	80	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.93	Very limited Depth to water	1.00
89D: Wethersfield-----	80	Somewhat limited Seepage Slope	0.70 0.28	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.93	Very limited Depth to water	1.00
90B: Stockbridge-----	80	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00
90C: Stockbridge-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping	1.00	Very limited Depth to water	1.00
90D: Stockbridge-----	80	Somewhat limited Seepage Slope	0.70 0.12	Very limited Piping	1.00	Very limited Depth to water	1.00
91B: Stockbridge-----	80	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
91C: Stockbridge-----	80	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping	1.00	Very limited Depth to water	1.00
91D: Stockbridge-----	80	Somewhat limited Seepage Slope	0.70 0.28	Very limited Piping	1.00	Very limited Depth to water	1.00
92B: Nellis-----	85	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
92C: Nellis-----	85	Very limited Seepage Slope	1.00 0.01	Not limited		Very limited Depth to water	1.00
92D: Nellis-----	85	Very limited Seepage Slope	1.00 0.12	Not limited		Very limited Depth to water	1.00
93C: Nellis-----	85	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
94C: Farmington-----	40	Very limited Depth to bedrock	1.00	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00
Nellis-----	35	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
94E: Farmington-----	40	Very limited Depth to bedrock Slope	1.00 0.28	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00
Nellis-----	35	Very limited Seepage Slope	1.00 0.28	Not limited		Very limited Depth to water	1.00
95C: Farmington-----	60	Very limited Depth to bedrock	1.00	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00
Rock outcrop-----	20	Very limited Depth to bedrock	1.00	Not rated		Not rated	
95E: Farmington-----	60	Very limited Depth to bedrock Slope	1.00 0.50	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00
Rock outcrop-----	20	Very limited Depth to bedrock Slope	1.00 0.50	Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
96: Ipswich-----	85	Very limited Seepage	1.00	Not rated		Very limited Salinity and saturated zone Cutbanks cave	1.00 0.10
97: Pawcatuck-----	85	Very limited Seepage	1.00	Very limited Organic matter content Ponding Depth to saturated zone Salinity Piping	1.00 1.00 1.00 1.00 1.00	Very limited Cutbanks cave Salinity and saturated zone	1.00 1.00
98: Westbrook-----	80	Very limited Seepage	1.00	Very limited Organic matter content Ponding Depth to saturated zone Salinity Piping	1.00 1.00 1.00 1.00 1.00	Very limited Salinity and saturated zone Cutbanks cave	1.00 0.10
99: Westbrook, low salt-	80	Very limited Seepage	1.00	Very limited Organic matter content Ponding Depth to saturated zone Piping Salinity	1.00 1.00 1.00 1.00 0.03	Somewhat limited Salinity and saturated zone Cutbanks cave	0.35 0.10
100: Suncook-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.10	Very limited Depth to water	1.00
101: Occum-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.36	Very limited Depth to water	1.00
102: Pootatuck-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.36	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
103: Rippowam-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.50	Very limited Cutbanks cave	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
104: Bash-----	80	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
105: Hadley-----	80	Very limited Seepage	1.00	Very limited Piping	1.00	Very limited Depth to water	1.00
106: Winooski-----	80	Very limited Seepage	1.00	Very limited Piping Depth to saturated zone	1.00 0.95	Somewhat limited Cutbanks cave Depth to saturated zone	0.10 0.02
107: Limerick-----	50	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.30 0.10
Lim-----	30	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.09	Very limited Cutbanks cave	1.00
108: Saco-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping Ponding Seepage	1.00 1.00 1.00 0.10	Very limited Cutbanks cave	1.00
109: Fluvaquents, Frequently Flooded-	50	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.01	Very limited Cutbanks cave	1.00
Udifluents, Frequently Flooded-	35	Very limited Seepage	1.00	Somewhat limited Seepage	0.17	Very limited Depth to water	1.00
221A: Ninigret-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
Urban land-----	35	Not limited		Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
224A: Deerfield-----	40	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage	0.95 0.13	Very limited Cutbanks cave Depth to saturated zone	1.00 0.02
Urban land-----	35	Not limited		Not rated		Not rated	
225B: Brancroft-----	40	Somewhat limited Seepage	0.01	Very limited Depth to saturated zone Piping	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.99 0.10
Urban land-----	35	Not limited		Not rated		Not rated	
226B: Berlin-----	40	Somewhat limited Seepage	0.01	Very limited Depth to saturated zone Piping	1.00 0.97	Very limited Slow refill Cutbanks cave	1.00 0.10
Urban land-----	35	Not limited		Not rated		Not rated	
228B: Elmridge-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	0.99 0.84	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
229B: Agawam-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
229C: Agawam-----	40	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	
230B: Branford-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
230C: Branford-----	40	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
232B: Haven-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.39	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
234B: Merrimac-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
235B: Penwood-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.15	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
236B: Windsor-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.10	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
237A: Manchester-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.11	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
237C: Manchester-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.11	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
238A: Hinckley-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
238C: Hinckley-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.09	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
240B: Ludlow-----	40	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
243B: Rainbow-----	40	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.95	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
245B: Woodbridge-----	40	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.86	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
245C: Woodbridge-----	40	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.86	Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	
248B: Georgia-----	40	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone	1.00 0.95	Very limited Cutbanks cave Slow refill Depth to saturated zone	1.00 0.95 0.02
Urban land-----	35	Not limited		Not rated		Not rated	
250B: Sutton-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
Urban land-----	35	Not limited		Not rated		Not rated	
253B: Wapping-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.09	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
Urban land-----	35	Not limited		Not rated		Not rated	
255B: Watchaug-----	40	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
255B: Urban land-----	35	Not limited		Not rated		Not rated	
260B: Charlton-----	40	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
260C: Charlton-----	40	Very limited Seepage Slope	1.00 0.01	Not limited		Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	
260D: Charlton-----	40	Very limited Seepage Slope	1.00 0.12	Not limited		Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.12	Not rated		Not rated	
263B: Cheshire-----	40	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
263C: Cheshire-----	40	Very limited Seepage Slope	1.00 0.01	Not limited		Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	
266B: Narragansett-----	40	Very limited Seepage	1.00	Somewhat limited Seepage	0.07	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
269B: Yalesville-----	40	Very limited Seepage Depth to bedrock	1.00 0.65	Somewhat limited Thin layer Seepage	0.65 0.01	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
269C: Yalesville-----	40	Very limited Seepage Depth to bedrock Slope	1.00 0.65 0.01	Somewhat limited Thin layer Seepage	0.65 0.01	Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
273C:							
Urban land-----	35	Not limited		Not rated		Not rated	
Charlton-----	25	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
Chatfield-----	15	Very limited Seepage Depth to bedrock	1.00 0.88	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
273E:							
Urban land-----	35	Somewhat limited Slope	0.50	Not rated		Not rated	
Charlton-----	25	Very limited Seepage Slope	1.00 0.50	Not limited		Very limited Depth to water	1.00
Chatfield-----	15	Very limited Seepage Depth to bedrock Slope	1.00 0.88 0.50	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
275C:							
Urban land-----	45	Not limited		Not rated		Not rated	
Chatfield-----	30	Very limited Seepage Depth to bedrock	1.00 0.88	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
275E:							
Urban land-----	35	Somewhat limited Slope	0.50	Not rated		Not rated	
Chatfield-----	25	Very limited Seepage Depth to bedrock Slope	1.00 0.88 0.50	Somewhat limited Thin layer Seepage	0.88 0.01	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock Slope	1.00 0.50	Not rated		Not rated	
282B:							
Broadbrook-----	40	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.96	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
284B:							
Paxton-----	40	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Thin layer	0.99 0.95	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
284C: Paxton-----	40	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
		Slope	0.01	Thin layer	0.95		
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	
284D: Paxton-----	40	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
		Slope	0.12	Thin layer	0.95		
Urban land-----	35	Somewhat limited Slope	0.12	Not rated		Not rated	
287B: Wethersfield-----	40	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00
				Depth to saturated zone	0.99		
				Thin layer	0.93		
Urban land-----	35	Not limited		Not rated		Not rated	
287C: Wethersfield-----	40	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00
		Slope	0.01	Depth to saturated zone	0.99		
				Thin layer	0.93		
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	
287D: Wethersfield-----	40	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00
		Slope	0.12	Depth to saturated zone	0.99		
				Thin layer	0.93		
Urban land-----	35	Somewhat limited Slope	0.12	Not rated		Not rated	
290B: Stockbridge-----	40	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00
Urban land-----	35	Not limited		Not rated		Not rated	
290C: Stockbridge-----	40	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00
		Slope	0.01				
Urban land-----	35	Somewhat limited Slope	0.01	Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
290D: Stockbridge-----	40	Somewhat limited Seepage Slope	0.70 0.12	Very limited Piping	1.00	Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.12	Not rated		Not rated	
301: Beaches-----	50	Very limited Seepage	1.00	Not rated		Not rated	
Udipsamments-----	35	Very limited Seepage	1.00	Somewhat limited Seepage	0.76	Very limited Depth to water	1.00
302: Dumps-----	95	Somewhat limited Seepage	0.81	Not rated		Not rated	
303: Pits, quarries-----	90	Very limited Depth to bedrock Slope	1.00 0.28	Not rated		Not rated	
304: Udorthents-----	90	Very limited Seepage Slope	1.00 0.99	Very limited Piping	1.00	Very limited Depth to water	1.00
305: Udorthents-----	65	Very limited Seepage Slope	1.00 0.08	Very limited Piping Depth to saturated zone	1.00 0.22	Very limited Cutbanks cave Depth to saturated zone	1.00 0.40
Pits-----	25	Very limited Seepage Slope	1.00 0.88	Somewhat limited Seepage	0.93	Very limited Depth to water	1.00
306: Udorthents-----	50	Very limited Seepage Slope	1.00 0.08	Very limited Piping	1.00	Very limited Depth to water	1.00
Urban land-----	35	Somewhat limited Slope	0.08	Not rated		Not rated	
307: Urban land-----	80	Somewhat limited Slope	0.21	Not rated		Not rated	
308: Udorthents-----	80	Very limited Seepage Slope	1.00 0.08	Very limited Piping Depth to saturated zone	1.00 0.22	Very limited Cutbanks cave Depth to saturated zone	1.00 0.40

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
309: Udorthents-----	80	Very limited Seepage Slope	1.00 0.08	Very limited Piping Depth to saturated zone	1.00 0.22	Very limited Cutbanks cave Depth to saturated zone	1.00 0.40
310: Udorthents, Periodically Flooded-----	85	Very limited Seepage Slope	1.00 0.08	Very limited Piping Depth to saturated zone	1.00 0.22	Very limited Cutbanks cave Depth to saturated zone	1.00 0.40
401C: Macomber-----	55	Somewhat limited Depth to bedrock Seepage Slope	0.86 0.70 0.01	Somewhat limited Thin layer Seepage	0.86 0.12	Very limited Depth to water	1.00
Taconic-----	30	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
402D: Macomber-----	50	Somewhat limited Depth to bedrock Seepage Slope	0.86 0.70 0.12	Somewhat limited Thin layer Seepage	0.86 0.12	Very limited Depth to water	1.00
Taconic-----	25	Very limited Depth to bedrock Slope	1.00 0.12	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock Slope	1.00 0.12	Not rated		Not rated	
403C: Taconic-----	70	Very limited Depth to bedrock	1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	25	Very limited Depth to bedrock	1.00	Not rated		Not rated	
403E: Taconic-----	70	Very limited Depth to bedrock Slope	1.00 0.72	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	20	Very limited Depth to bedrock Slope	1.00 0.72	Not rated		Not rated	
403F: Taconic-----	70	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Thin layer	1.00	Very limited Depth to water	1.00
Rock outcrop-----	20	Very limited Depth to bedrock Slope	1.00 1.00	Not rated		Not rated	

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
405C: Dummerston-----	85	Somewhat limited Seepage	0.70	Not limited		Very limited Depth to water	1.00
405E: Dummerston-----	85	Somewhat limited Seepage Slope	0.70 0.28	Not limited		Very limited Depth to water	1.00
407C: Lanesboro-----	85	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.86	Very limited Depth to water	1.00
407E: Lanesboro-----	85	Somewhat limited Seepage Slope	0.70 0.50	Very limited Piping Depth to saturated zone Thin layer	1.00 0.99 0.86	Very limited Depth to water	1.00
408C: Fullam-----	85	Somewhat limited Seepage	0.70	Very limited Piping Thin layer Depth to saturated zone	1.00 0.99 0.99	Very limited Depth to water	1.00
409B: Brayton-----	85	Not limited		Very limited Depth to saturated zone Piping Thin layer	1.00 1.00 1.00	Very limited Depth to water	1.00
412B: Bice-----	85	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
412C: Bice-----	85	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
412D: Bice-----	85	Very limited Seepage Slope	1.00 0.12	Not limited		Very limited Depth to water	1.00
413C: Bice-----	45	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
Millsite-----	40	Very limited Seepage Depth to bedrock	1.00 0.83	Somewhat limited Thin layer	0.83	Very limited Depth to water	1.00

Table 22.--Ponds and Embankments--Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
413E: Bice-----	45	Very limited Seepage Slope	1.00 0.28	Not limited		Very limited Depth to water	1.00
Millsite-----	40	Very limited Seepage Depth to bedrock Slope	1.00 0.83 0.28	Somewhat limited Thin layer	0.83	Very limited Depth to water	1.00
414: Fredon, cold-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.43	Very limited Cutbanks cave	1.00
415C: Millsite-----	40	Very limited Seepage Depth to bedrock	1.00 0.83	Somewhat limited Thin layer	0.83	Very limited Depth to water	1.00
Westminster-----	40	Very limited Depth to bedrock	1.00	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock	1.00	Not rated		Not rated	
415E: Millsite-----	40	Very limited Seepage Depth to bedrock Slope	1.00 0.83 0.50	Somewhat limited Thin layer	0.83	Very limited Depth to water	1.00
Westminster-----	40	Very limited Depth to bedrock Slope	1.00 0.50	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00
Rock outcrop-----	15	Very limited Depth to bedrock Slope	1.00 0.50	Not rated		Not rated	
416E: Rock outcrop-----	70	Very limited Depth to bedrock Slope	1.00 0.28	Not rated		Not rated	
Westminster-----	20	Very limited Depth to bedrock Slope	1.00 0.28	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00
416F: Rock outcrop-----	70	Very limited Depth to bedrock Slope	1.00 1.00	Not rated		Not rated	
Westminster-----	20	Very limited Depth to bedrock Slope	1.00 1.00	Very limited Thin layer Piping	1.00 1.00	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
417B: Bice-----	85	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
417C: Bice-----	85	Very limited Seepage	1.00	Not limited		Very limited Depth to water	1.00
417D: Bice-----	85	Very limited Seepage Slope	1.00 0.12	Not limited		Very limited Depth to water	1.00
418C: Schroon-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.03	Somewhat limited Cutbanks cave Depth to saturated zone	0.10 0.01
420A: Schroon-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.03	Somewhat limited Cutbanks cave Depth to saturated zone	0.10 0.01
420B: Schroon-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.03	Somewhat limited Cutbanks cave Depth to saturated zone	0.10 0.01
421A: Ninigret, cold-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	0.99 0.03	Very limited Cutbanks cave Depth to saturated zone	1.00 0.01
423A: Sudbury, cold-----	85	Very limited Seepage	1.00	Somewhat limited Depth to saturated zone Seepage	0.95 0.10	Very limited Cutbanks cave Depth to saturated zone	1.00 0.02
424B: Shelburne-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Thin layer	0.99 0.94	Very limited Depth to water	1.00
424C: Shelburne-----	85	Very limited Seepage Slope	1.00 0.01	Very limited Depth to saturated zone Thin layer	0.99 0.94	Very limited Depth to water	1.00

Table 22.--Ponds and Embankments--Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
424D: Shelburne-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
		Slope	0.12	Thin layer	0.94		
425B: Shelburne-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
				Thin layer	0.94		
425C: Shelburne-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
		Slope	0.01	Thin layer	0.94		
426D: Shelburne-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	0.99	Very limited Depth to water	1.00
		Slope	0.28	Thin layer	0.94		
427B: Ashfield-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to water	1.00
427C: Ashfield-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to water	1.00
		Slope	0.01	Piping	1.00		
				Thin layer	0.88		
428A: Ashfield-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to water	1.00
				Piping	1.00		
				Thin layer	0.88		
428B: Ashfield-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to water	1.00
				Piping	1.00		
				Thin layer	0.88		
428C: Ashfield-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to water	1.00
		Slope	0.01	Piping	1.00		
				Thin layer	0.88		
429A: Agawam, cold-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
429B: Agawam, cold-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
429C: Agawam, cold-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
433: Moosilauke-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Seepage	1.00 0.15	Very limited Cutbanks cave	1.00
434A: Merrimac, cold-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
434B: Merrimac, cold-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
434C: Merrimac, cold-----	80	Very limited Seepage Slope	1.00 0.01	Somewhat limited Seepage	0.12	Very limited Depth to water	1.00
435: Scarboro-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 0.36	Very limited Cutbanks cave	1.00
436: Halsey-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding Seepage	1.00 1.00 0.31	Very limited Cutbanks cave	1.00
437: Wonsqueak-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping Ponding	1.00 1.00 1.00	Very limited Cutbanks cave	1.00
438: Bucksport-----	85	Very limited Seepage	1.00	Very limited Organic matter content Depth to saturated zone Piping Ponding Seepage	1.00 1.00 1.00 1.00 0.10	Very limited Cutbanks cave	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
440A: Boscawen-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
440C: Boscawen-----	80	Very limited Seepage	1.00	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
440E: Boscawen-----	85	Very limited Seepage Slope	1.00 0.50	Somewhat limited Seepage	0.22	Very limited Depth to water	1.00
442: Brayton-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Thin layer	1.00 0.93	Very limited Depth to water	1.00
443: Brayton-----	50	Very limited Seepage	1.00	Very limited Depth to saturated zone Thin layer	1.00 0.93	Very limited Depth to water	1.00
Loonmeadow-----	35	Very limited Seepage	1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Cutbanks cave	1.00
448B: Hogansburg-----	85	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.95 0.35	Very limited Depth to water	1.00
449B: Hogansburg-----	85	Somewhat limited Seepage	0.70	Very limited Piping Depth to saturated zone Thin layer	1.00 0.95 0.35	Very limited Depth to water	1.00
449C: Hogansburg-----	85	Somewhat limited Seepage Slope	0.70 0.01	Very limited Piping Depth to saturated zone Thin layer	1.00 0.95 0.35	Very limited Depth to water	1.00
450B: Pyrities-----	80	Somewhat limited Seepage	0.05	Very limited Piping	1.00	Very limited Depth to water	1.00
450C: Pyrities-----	80	Somewhat limited Seepage Slope	0.05 0.01	Very limited Piping	1.00	Very limited Depth to water	1.00

Table 22.—Ponds and Embankments—Continued

Map symbol and soil name	Pct. of map unit	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
450D: Pyrities-----	80	Somewhat limited Slope Seepage	0.12 0.05	Very limited Piping	1.00	Very limited Depth to water	1.00
451B: Pyrities-----	80	Somewhat limited Seepage	0.05	Very limited Piping	1.00	Very limited Depth to water	1.00
451C: Pyrities-----	80	Somewhat limited Seepage Slope	0.05 0.01	Very limited Piping	1.00	Very limited Depth to water	1.00
451D: Pyrities-----	80	Somewhat limited Slope Seepage	0.28 0.05	Very limited Piping	1.00	Very limited Depth to water	1.00
457: Mudgepond-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	1.00 1.00	Very limited Cutbanks cave	1.00
458: Mudgepond-----	55	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping	1.00 1.00	Very limited Cutbanks cave	1.00
Alden-----	35	Somewhat limited Seepage	0.70	Very limited Depth to saturated zone Piping Ponding	1.00 1.00 1.00	Very limited Cutbanks cave Slow refill	1.00 0.30
501: Ondawa-----	85	Somewhat limited Seepage	0.70	Very limited Piping	1.00	Very limited Depth to water	1.00
503: Rumney-----	80	Very limited Seepage	1.00	Very limited Depth to saturated zone	1.00	Very limited Cutbanks cave	1.00
508: Medomak-----	85	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping Ponding	1.00 1.00 1.00	Very limited Cutbanks cave	1.00

Table 23.--Engineering Properties

(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
2: Ridgebury-----	0-5	Fine sandy loam	SM	A-2, A-4	0-5	0-5	85-95	80-90	65-85	25-45	0-20	NP-5
	5-14	Fine sandy loam, sandy loam, gravelly loam	SM	A-1, A-4, A-2	0-5	0-5	70-100	55-90	40-90	20-50	0-20	NP-5
	14-21	Fine sandy loam, sandy loam, gravelly loam	SM	A-2, A-4, A-1	0-5	0-5	70-100	55-90	40-90	20-50	0-20	NP-5
	21-60	Sandy loam, gravelly loam	SM	A-2, A-4, A-1	0-5	0-5	70-100	55-90	40-90	20-50	0-20	NP-5
3: Ridgebury-----	0-5	Fine sandy loam	SM	A-4, A-2	0-5	0-5	85-95	80-90	65-85	25-45	0-20	NP-5
	5-14	Fine sandy loam, sandy loam, gravelly loam	SM	A-2, A-4, A-1	0-5	0-5	70-100	55-90	40-90	20-50	0-20	NP-5
	14-21	Fine sandy loam, sandy loam, gravelly loam	SM	A-1, A-2, A-4	0-5	0-5	70-100	55-90	40-90	20-50	0-20	NP-5
	21-60	Sandy loam, gravelly loam	SM	A-1, A-2, A-4	0-5	0-5	70-100	55-90	40-90	20-50	0-20	NP-5
3: Leicester-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	SM, SC-SM	A-4	0-3	0-10	85-95	80-90	65-90	35-45	5-25	NP-5
	7-10	Fine sandy loam, loam, gravelly sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	0-5	0-15	75-100	60-90	45-90	25-65	5-25	NP-5
	10-18	Fine sandy loam, loam, gravelly sandy loam	SM, SC-SM, ML, CL-ML	A-4, A-2	0-5	0-15	75-100	60-90	45-90	25-65	5-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
3: Leicester-----	18-24	Fine sandy loam, loam, gravelly sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	0-5	0-15	75-100	60-90	45-90	25-65	5-25	NP-5
	24-43	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM, SC-SM	A-2, A-4	0-5	0-10	75-100	60-90	45-85	25-50	5-20	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, very gravelly sandy loam	SM, SC-SM	A-2, A-4	0-10	0-20	60-100	40-90	35-85	20-50	5-20	NP-5
Whitman-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-9	Fine sandy loam	SM, SC-SM	A-4, A-2	1-5	1-10	95-100	85-95	75-90	30-50	15-25	NP-5
	9-16	Fine sandy loam, gravelly fine sandy loam, gravelly loam	SM, SC-SM, ML, CL-ML	A-2, A-4	1-10	1-15	80-100	65-95	55-90	25-60	15-25	NP-5
	16-22	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	1-10	1-15	80-100	65-95	50-95	25-55	15-20	NP-5
	22-60	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	1-10	1-15	80-100	65-95	50-95	25-55	15-20	NP-5
4: Leicester-----	0-7	Fine sandy loam	SM, SC-SM	A-4	0-3	0-10	85-95	80-90	65-90	35-45	5-25	NP-5
	7-10	Fine sandy loam, loam, gravelly sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	0-5	0-15	75-100	60-90	45-90	25-65	5-25	NP-5
	10-18	Fine sandy loam, loam, gravelly sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	0-5	0-15	75-100	60-90	45-90	25-65	5-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
4: Leicester-----	18-24	Fine sandy loam, loam, gravelly sandy loam	SM, SC-SM, ML, CL-ML	A-2, A-4	0-5	0-15	75-100	60-90	45-90	25-65	5-25	NP-5
	24-43	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM, SC-SM	A-4, A-2	0-5	0-10	75-100	60-90	45-85	25-50	5-20	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, very gravelly sandy loam	SM, SC-SM	A-2, A-4	0-10	0-20	60-100	40-90	35-85	20-50	5-20	NP-5
5: Wilbraham-----	0-4	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	55-85	15-25	NP-5
	4-8	Silt loam, gravelly loam, very fine sandy loam	SM, ML, CL-ML	A-4	0-5	0-5	85-100	65-90	60-90	40-85	15-25	NP-10
	8-20	Silt loam, gravelly loam, very fine sandy loam	CL-ML, ML, SM	A-4	0-5	0-5	85-100	65-90	60-90	40-85	15-25	NP-10
	20-65	Gravelly loam, silt loam, very fine sandy loam	CL-ML, ML, SM	A-4	0-5	0-10	80-100	65-95	60-95	35-90	15-25	NP-10
6: Wilbraham-----	0-4	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	55-85	15-25	NP-5
	4-8	Silt loam, gravelly loam, very fine sandy loam	ML, SM, CL-ML	A-4	0-5	0-5	85-100	65-90	60-90	40-85	15-25	NP-10
	8-20	Silt loam, gravelly loam, very fine sandy loam	SM, CL-ML, ML	A-4	0-5	0-5	85-100	65-90	60-90	40-85	15-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
6: Wilbraham-----	20-65	Gravelly loam, silt loam, very fine sandy loam	CL-ML, ML, SM	A-4	0-5	0-10	80-100	65-95	60-95	35-90	15-25	NP-10
6: Menlo-----	0-5	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	5-16	Mucky silt loam	ML, SM	A-4	0-5	0-5	75-95	65-95	60-95	40-90	15-25	NP-5
	16-22	Flaggy very fine sandy loam, silt loam, gravelly fine sandy loam, loam	SM, ML, SC- SM, CL-ML	A-4, A-2	0-5	0-20	95	80-95	70-95	30-90	15-25	NP-10
	22-27	Flaggy fine sandy loam, silt loam, loam, gravelly very fine sandy loam	ML, SM, SC- SM, CL-ML	A-2, A-4	0-5	0-20	95	80-95	70-95	30-90	15-25	NP-10
	27-40	Fine sandy loam, silt loam, loam, gravelly fine sandy loam, gravelly very fine sandy loam	ML, SM, SC- SM, CL-ML	A-4, A-2	0-5	0-20	80-90	65-90	55-90	25-90	15-25	NP-10
	40-60	Fine sandy loam, silt loam, loam, gravelly fine sandy loam, gravelly very fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0-5	0-20	80-90	65-90	55-90	25-85	15-25	NP-10
7: Mudgepond-----	0-11	Silt loam	ML	A-4	0-5	0	90-95	80-90	75-85	60-75	20-30	NP-5
	11-16	Loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
					Pct	Pct						
7: Mudgepond-----	16-26	Fine sandy loam, loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	26-35	Gravelly fine sandy loam, very fine sandy loam, silt loam, loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	35-65	Gravelly fine sandy loam, very fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-85	25-65	20-25	NP-10
8: Mudgepond-----	0-11	Silt loam	ML	A-4	0-5	0	90-95	80-90	75-85	60-75	20-30	NP-5
	11-16	Loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	16-26	Fine sandy loam, loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	26-35	Gravelly fine sandy loam, very fine sandy loam, silt loam, loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	35-65	Gravelly fine sandy loam, very fine sandy loam, sandy loam, loam	ML, SM	A-4, A-2	0-5	5-10	65-90	60-90	55-85	25-65	20-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
8: Alden-----	0-4	Mucky silt loam	ML	A-4	0-5	0-5	90-100	80-100	75-100	60-95	25-35	NP-5
	4-13	Silt loam, very fine sandy loam	ML	A-4	0-5	0-5	90-100	80-100	75-100	50-95	25-35	NP-5
	13-23	Silt loam, very fine sandy loam	ML	A-4	0-5	0-5	90-100	80-100	75-100	50-95	25-35	NP-5
	23-29	Silt loam, very fine sandy loam	ML, SM	A-4	0-5	0-5	90-100	80-100	75-100	45-95	25-35	NP-5
	29-43	Gravelly loam, silt loam, loam	CL, SC	A-6	0-5	0-5	60-100	55-90	50-90	40-80	25-35	10-15
	43-60	Loam, gravelly loam, silt loam	CL, SC	A-6	0-5	0-15	70-100	60-90	50-90	40-80	25-35	10-15
9: Scitico-----	0-8	Silt loam	ML	A-4	0	0	100	95-100	90-100	65-100	25-35	NP-5
	8-11	Silt loam, silty clay loam, silty clay	CL	A-7, A-6	0	0	100	95-100	90-100	85-100	30-50	10-25
	11-18	Silty clay loam, silt loam, silty clay	CL	A-7, A-6	0	0	100	95-100	90-100	85-100	30-50	10-25
	18-30	Silty clay loam, silty clay, clay	CL, CH	A-7	0	0	100	95-100	95-100	90-100	40-55	15-30
	30-38	Silty clay, silty clay loam, clay	CL, CH	A-7	0	0	100	95-100	95-100	90-100	40-55	15-30
	38-52	Silty clay loam, silty clay, clay	CH, CL	A-7	0	0	100	95-100	95-100	90-100	40-55	15-30
	52-65	Silty clay, clay, silty clay loam	CH, CL	A-7	0	0	100	95-100	95-100	90-100	40-55	15-30

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
9: Shaker-----	In											
	0-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-6	Fine sandy loam	SM	A-4	0	0	100	90-100	85-100	40-50	15-25	NP
	6-20	Sandy loam, fine sandy loam, loam	ML, SM	A-4	0	0	100	90-100	70-100	40-60	15-25	NP-5
	20-30	Sandy loam, fine sandy loam, loam	ML, SM	A-4	0	0	100	90-100	70-100	40-60	15-25	NP-5
	30-65	Silty clay, silty clay loam, clay	CH, CL	A-7	0	0	100	95-100	95-100	85-100	40-55	15-25
Maybid-----	0-9	Silt loam	ML	A-4	0	0	100	100	95-100	85-100	25-35	NP-5
	9-18	Silty clay loam, silty clay	CL	A-7, A-6	0	0	100	100	95-100	90-95	30-50	10-25
	18-26	Silty clay loam, silty clay, clay	CL	A-7, A-6	0	0	100	100	95-100	90-95	30-50	10-25
	26-36	Silty clay loam, silty clay, clay	CL	A-7, A-6	0	0	100	100	100	85-95	35-50	15-25
	36-60	Silty clay loam, silty clay, clay	CL	A-7, A-6	0	0	100	100	100	85-95	35-50	15-25
10: Raynham-----	0-10	Silt loam	ML	A-4	0	0	95-100	95-100	90-100	60-95	15-25	NP
	10-16	Silt loam, very fine sandy loam	ML	A-4	0	0	95-100	95-100	90-100	55-85	15-25	NP-5
	16-26	Silt loam, very fine sandy loam	ML	A-4	0	0	95-100	95-100	90-100	55-85	15-25	NP-5
	26-34	Very fine sandy loam, silt loam	ML	A-4	0	0	95-100	95-100	90-100	55-85	15-25	NP-5
	34-47	Silt loam, very fine sandy loam	ML	A-4	0	0	95-100	95-100	90-100	55-85	15-25	NP-5
	47-60	Silt loam, very fine sandy loam	ML	A-4	0	0	95-100	95-100	90-100	55-85	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
12: Raypol-----	0-8	Silt loam	ML	A-4	0	0	90-100	85-100	70-100	50-90	15-25	NP
	8-12	Very fine sandy loam, silt loam, loam	ML, SM	A-4	0	0	90-100	85-100	75-100	45-90	15-25	NP-5
	12-20	Silt loam, very fine sandy loam, loam	ML, SM	A-4	0	0	90-100	85-100	75-100	45-90	15-25	NP-5
	20-26	Silt loam, very fine sandy loam, loam	ML, SM	A-4	0	0	90-100	85-100	75-100	45-90	15-25	NP-5
	26-29	Very fine sandy loam, silt loam, loam	ML, SM	A-4	0	0	90-100	85-100	75-100	45-90	15-25	NP-5
	29-52	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-15	60-100	30-100	15-95	5-30	0-15	NP
	52-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-15	60-100	30-100	15-95	5-30	0-15	NP
	13: Walpole-----	0-1	Moderately decomposed plant material	OL		0	0	100	100	100	100	---
	1-7	Sandy loam	ML, SM	A-2, A-4	0	0-5	90-100	80-100	55-90	25-70	0-15	NP
	7-21	Sandy loam, fine sandy loam, gravelly sandy loam	SM	A-1, A-2, A-4	0	0-10	85-100	65-100	45-90	20-50	0-15	NP
	21-25	Gravelly sandy loam, sandy loam, fine sandy loam	SM	A-1, A-2, A-4	0	0-10	85-100	65-100	45-90	20-50	0-15	NP
	25-41	Stratified very gravelly coarse sand to loamy fine sand	SP, SM, SW-SM, SW	A-2, A-3, A-1	0	0-10	70-100	35-100	10-95	1-20	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
13: Walpole-----	41-65	Stratified very gravelly coarse sand to loamy fine sand	SP, SW, SW- SM, SM	A-1, A-2, A-3	0	0-10	70-100	35-100	10-95	1-20	0-15	NP
14: Fredon-----	0-8	Silt loam	ML, SM	A-4	0	0	85-100	75-100	70-100	45-95	15-25	NP-5
	8-17	Loam, very fine sandy loam, silt loam, fine sandy loam	SM, ML	A-4	0	0-5	90-100	80-100	75-100	35-85	15-25	NP-5
	17-24	Loam, fine sandy loam, very fine sandy loam, silt loam	ML, SM	A-4	0	0-5	90-100	80-100	75-100	35-85	15-25	NP-5
	24-29	Stratified gravelly sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-90	60-85	30-80	5-25	15-20	NP-5
	29-48	Stratified gravelly sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-90	60-85	30-80	5-25	15-20	NP-5
	48-60	Stratified gravelly sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-90	60-85	30-80	5-25	15-20	NP-5
15: Scarboro-----	0-12	Muck	PT	A-8	0	0	100	100	100	100	---	---
	12-17	Loamy sand, sandy loam, fine sandy loam, loamy fine sand	SM	A-2	0	0	90-100	85-100	70-90	20-30	0-20	NP
	17-31	Stratified sand to loamy fine sand	SM, SP-SM	A-2	0	0	85-100	75-100	45-95	5-30	0-15	NP
	31-72	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-15	70-100	40-100	20-95	5-30	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
16: Halsey-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Silt loam	CL-ML, ML	A-4	0	0-5	90-100	80-100	75-100	50-85	20-25	NP-5
	8-16	Silt loam, loam, very fine sandy loam, gravelly loam	SC-SM, SM, CL-ML, ML	A-4	0	0-5	80-100	75-100	70-100	45-85	20-25	NP-10
	16-28	Fine sandy loam, very fine sandy loam, gravelly fine sandy loam, gravelly very fine sandy loam	SM, ML	A-4	0	0-5	90-100	75-100	65-100	35-70	20-25	NP-5
	28-38	Loamy sand, gravelly loamy sand, very gravelly loamy sand, sand	SW-SM, SM	A-1, A-2, A-3	0	0-15	55-100	40-85	25-65	5-25	15-20	NP-5
	38-60	Sand, loamy sand, gravelly loamy sand, very gravelly loamy sand	SM, SW-SM	A-1, A-3, A-2	0	0-15	55-90	40-85	25-65	5-25	15-20	NP-5
17: Timakwa-----	0-10	Muck	PT	A-8	0	0	100	100	100	100	---	---
	10-21	Muck	PT	A-8	0	0	100	100	100	100	---	---
	21-24	Muck	PT	A-8	0	0	100	100	100	100	---	---
	24-37	Muck	PT	A-8	0	0	100	100	100	100	---	---
	37-47	Very gravelly loamy coarse sand, sand, loamy sand, coarse sand, loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	60-100	50-100	25-95	5-25	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
17: Timakwa-----	47-60	Gravelly loamy very fine sand, loamy sand, loamy fine sand, coarse sand, loamy coarse sand, gravelly sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	60-100	50-100	25-95	5-25	0-15	NP
Natchaug-----	0-2	Peat	PT	A-8	0	0	100	100	100	100	---	---
	2-4	Peat	PT	A-8	0	0	100	100	100	100	---	---
	4-6	Muck	PT	A-8	0	0	100	100	100	100	---	---
	6-11	Muck	PT	A-8	0	0	100	100	100	100	---	---
	11-18	Muck	PT	A-8	0	0	100	100	100	100	---	---
	18-24	Muck	PT	A-8	0	0	100	100	100	100	---	---
	24-33	Fine sandy loam, loam, very fine sandy loam, gravelly sandy loam	ML, SM	A-2, A-4	0-2	0-10	85-100	75-100	55-95	30-75	20-25	NP-5
	33-36	Fine sandy loam, sandy loam, silt loam, gravelly sandy loam	ML, SM	A-4, A-2	0-2	0-10	85-100	75-100	55-95	30-75	20-25	NP-5
	36-80	Loam, sandy loam, fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-2	0-10	85-100	75-100	55-95	30-75	20-25	NP-5
18: Catden-----	0-2	Muck	PT	A-8	0	0	100	100	100	100	---	---
	2-18	Muck	PT	A-8	0	0	100	100	100	100	---	---
	18-47	Muck	PT	A-8	0	0	100	100	100	100	---	---
	47-49	Muck	PT	A-8	0	0	100	100	100	100	---	---
	49-61	Muck	PT	A-8	0	0	100	100	100	100	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
18: Freetown-----	0-4	Peat	PT	A-8	0	0	100	100	100	100	---	---
	4-10	Peat	PT	A-8	0	0	100	100	100	100	---	---
	10-22	Muck	PT	A-8	0	0	100	100	100	100	---	---
	22-35	Muck	PT	A-8	0	0	100	100	100	100	---	---
	35-41	Muck	PT	A-8	0	0	100	100	100	100	---	---
	41-55	Muck	PT	A-8	0	0	100	100	100	100	---	---
	55-71	Muck	PT	A-8	0	0	100	100	100	100	---	---
	71-91	Muck	PT	A-8	0	0	100	100	100	100	---	---
20A: Ellington-----	0-8	Silt loam	ML	A-4	0	0-5	85-100	80-100	75-100	55-90	10-20	NP
	8-18	Silt loam, very fine sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-10	70-100	60-100	55-100	25-85	10-20	NP
	18-26	Very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	70-100	60-100	55-100	30-85	10-20	NP
	26-65	Stratified loamy fine sand to very gravelly coarse sand	SW-SM, SM	A-1, A-3, A-2	0-10	0-25	60-100	45-100	25-90	5-35	0-15	NP
21A: Ninigret-----	0-8	Fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	70-100	60-100	25-55	0-35	NP-5
	8-16	Fine sandy loam, very fine sandy loam, silt loam	ML, SM	A-2, A-4	0	0	95-100	75-100	70-100	30-80	0-25	NP-5
	16-26	Fine sandy loam, very fine sandy loam, silt loam	ML, SM	A-2, A-4	0	0	95-100	75-100	70-100	30-80	0-25	NP-5
	26-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0-10	0-15	70-100	35-100	5-95	1-30	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
21A: Tisbury-----	0-8	Silt loam	ML	A-4	0	0	100	90-100	80-100	60-95	0-35	NP-5
	8-18	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	100	90-100	85-100	45-95	0-35	NP-5
	18-26	Silt loam, very fine sandy loam	SM, ML	A-4	0	0	100	90-100	85-100	45-95	0-35	NP-5
	26-60	Stratified very gravelly sand to loamy sand	SM, SP	A-1, A-3	0-10	0-25	65-100	35-100	20-80	2-30	0-15	NP
22A: Hero-----	0-9	Gravelly loam	SM	A-2, A-4	0-5	0-10	65-85	55-75	45-75	25-50	15-25	NP-5
	9-18	Gravelly silt loam, silt loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-95	55-90	50-90	20-75	15-25	NP-5
	18-24	Gravelly silt loam, silt loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-10	65-95	55-90	50-90	20-75	15-25	NP-5
	24-27	Gravelly sandy loam, silt loam, loam, gravelly fine sandy loam, gravelly silt loam	ML, SM	A-4, A-2	0-5	0-10	65-95	55-90	40-90	20-75	15-25	NP-5
	27-60	Stratified extremely gravelly coarse sand to gravelly loamy fine sand	GM, GP-GM, SM, SP-SM	A-2, A-3, A-1	0-10	0-15	50-85	25-75	10-65	5-15	15-25	NP-5
22B: Hero-----	0-9	Gravelly loam	SM	A-2, A-4	0-5	0-10	65-85	55-75	45-75	25-50	15-25	NP-5
	9-18	Gravelly silt loam, silt loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-95	55-90	50-90	20-75	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
22B: Hero-----	18-24	Gravelly silt loam, silt loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-95	55-90	50-90	20-75	15-25	NP-5
	24-27	Gravelly sandy loam, silt loam, loam, gravelly fine sandy loam, gravelly silt loam	ML, SM	A-4, A-2	0-5	0-10	65-95	55-90	40-90	20-75	15-25	NP-5
	27-60	Stratified extremely gravelly coarse sand to gravelly loamy fine sand	GM, GP-GM, SM, SP-SM	A-2, A-3, A-1	0-10	0-15	50-85	25-75	10-65	5-15	15-25	NP-5
23A: Sudbury-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-5	Sandy loam	ML, SM	A-2, A-4	0	0-10	90-100	80-100	50-95	30-65	0-15	NP
	5-17	Gravelly sandy loam, fine sandy loam, sandy loam	SM	A-2	0	0-20	70-100	60-100	35-95	25-35	0-15	NP
	17-25	Sandy loam, fine sandy loam, gravelly sandy loam	SM	A-2	0	0-20	70-100	60-100	35-95	25-35	0-15	NP
	25-60	Stratified gravel to sand	SP-SM, SM, GM, GP-GM	A-2, A-3, A-1	0-20	0-40	40-100	20-90	15-75	5-20	0-15	NP
24A: Deerfield-----	0-8	Loamy fine sand	SM	A-2	0	0	100	85-100	70-90	20-30	0-20	NP
	8-16	Loamy sand, fine sand, loamy fine sand	SM	A-2	0	0	100	85-100	55-90	15-25	0-20	NP
	16-28	Loamy sand, fine sand, loamy fine sand	SM	A-2	0	0	100	85-100	55-90	15-25	0-20	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
24A: Deerfield-----	28-34	Fine sand, sand, loamy sand	SM	A-2	0	0	100	85-100	70-80	15-25	0-15	NP
	34-60	Fine sand, loamy sand, coarse sand	SP-SM, SM, SP	A-1, A-2	0	0	100	80-100	20-85	2-30	0-15	NP
25A: Brancroft-----	0-6	Silt loam	ML	A-4	0	0	100	95-100	90-100	70-95	25-35	NP-5
	6-17	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	17-22	Silty clay loam, silt loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	22-32	Silt loam, silty clay loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	32-43	Silty clay loam, silt loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	43-66	Silt loam, silty clay loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
25B: Brancroft-----	0-6	Silt loam	ML	A-4	0	0	100	95-100	90-100	70-95	25-35	NP-5
	6-17	Silt loam, silty clay loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	17-22	Silty clay loam, silt loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	22-32	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	32-43	Silty clay loam, silt loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	43-66	Silt loam, silty clay loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
25C: Brancroft-----	0-6	Silt loam	ML	A-4	0	0	100	95-100	90-100	70-95	25-35	NP-5
	6-17	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	17-22	Silty clay loam, silt loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	22-32	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	32-43	Silty clay loam, silt loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	43-66	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
26A: Berlin-----	0-6	Silt loam	ML	A-4	0	0	95-100	85-100	80-100	60-95	25-35	NP-5
	6-12	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	95-100	85-100	80-100	75-95	25-40	5-15
	12-20	Silty clay loam, silt loam	CL, CL-ML	A-6, A-4	0	0	95-100	85-100	80-100	75-95	25-40	5-15
	20-34	Silty clay loam, silt loam, silty clay	CL, CL-ML	A-7, A-4, A- 6, A-5	0	0	95-100	85-100	85-100	80-95	25-45	5-20
	34-48	Silty clay loam, silt loam, silty clay	CL, CL-ML	A-5, A-6, A- 7, A-4	0	0	95-100	85-100	85-100	80-95	25-45	5-20
	48-65	Silty clay loam, silt loam, silty clay	CL-ML, CL	A-7, A-4, A- 6, A-5	0	0	95-100	85-100	85-100	80-95	25-45	5-20
26B: Berlin-----	0-6	Silt loam	ML	A-4	0	0	95-100	85-100	80-100	60-95	25-35	NP-5
	6-12	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	95-100	85-100	80-100	75-95	25-40	5-15
	12-20	Silty clay loam, silt loam	CL, CL-ML	A-4, A-6	0	0	95-100	85-100	80-100	75-95	25-40	5-15

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
26B: Berlin-----	20-34	Silty clay loam, silt loam, silty clay	CL, CL-ML	A-5, A-4, A-6, A-7	0	0	95-100	85-100	85-100	80-95	25-45	5-20
	34-48	Silty clay loam, silt loam, silty clay	CL, CL-ML	A-4, A-6, A-7, A-5	0	0	95-100	85-100	85-100	80-95	25-45	5-20
	48-65	Silty clay loam, silt loam, silty clay	CL-ML, CL	A-4, A-6, A-7, A-5	0	0	95-100	85-100	85-100	80-95	25-45	5-20
27A: Belgrade-----	0-8	Silt loam	ML	A-4	0	0	100	100	90-100	70-85	15-25	NP-5
	8-16	Silt loam, very fine sandy loam	ML	A-4	0	0	100	100	95-100	60-90	15-25	NP-5
	16-27	Silt loam, very fine sandy loam	ML	A-4	0	0	100	100	95-100	60-90	15-25	NP-5
	27-45	Silt loam, fine sandy loam	ML	A-4	0	0	100	100	95-100	60-90	15-30	NP-10
	45-60	Silt loam, fine sandy loam	ML	A-4	0	0	100	100	95-100	60-90	15-30	NP-10
28A: Elmridge-----	0-6	Fine sandy loam	ML, SM	A-4	0	0	100	90-100	85-95	35-55	15-20	NP
	6-10	Fine sandy loam, sandy loam, loam	SM, CL-ML, ML	A-4	0	0	100	90-100	65-100	45-65	15-20	NP-5
	10-18	Fine sandy loam, sandy loam, loam	SM, ML, CL-ML	A-4	0	0	100	90-100	65-90	45-65	15-20	NP-5
	18-25	Sandy loam, fine sandy loam, loam	SM, CL-ML, ML	A-4	0	0	100	90-100	65-90	45-65	15-20	NP-5
	25-65	Silty clay, silty clay loam, clay	CL, CH	A-7	0	0	100	95-100	95-100	85-95	40-55	15-30

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
28B: Elmridge-----	0-6	Fine sandy loam	SM, ML	A-4	0	0	100	90-100	85-95	35-55	15-20	NP
	6-10	Fine sandy loam, sandy loam, loam	CL-ML, SM, ML	A-4	0	0	100	90-100	65-100	45-65	15-20	NP-5
	10-18	Fine sandy loam, sandy loam, loam	SM, ML, CL-ML	A-4	0	0	100	90-100	65-90	45-65	15-20	NP-5
	18-25	Sandy loam, fine sandy loam, loam	SM, CL-ML, ML	A-4	0	0	100	90-100	65-90	45-65	15-20	NP-5
	25-65	Silty clay, silty clay loam, clay	CH, CL	A-7	0	0	100	95-100	95-100	85-95	40-55	15-30
29A: Agawam-----	0-8	Fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5
	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP
29B: Agawam-----	0-8	Fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5
	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
29C: Agawam-----	0-8	Fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5
	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP
30A: Branford-----	0-8	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-85	15-25	NP-5
	8-18	Loam, silt loam, very fine sandy loam, gravelly fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	35-85	15-25	NP-5
	18-24	Gravelly loam, gravelly fine sandy loam, loam, silt loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	75-100	55-100	55-100	25-85	15-25	NP-5
	24-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-3, A-1, A-2	0	0-10	75-90	35-85	10-75	0-30	0-15	NP
30B: Branford-----	0-8	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-85	15-25	NP-5
	8-18	Loam, silt loam, very fine sandy loam, gravelly fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	35-85	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
30B: Branford-----	18-24	Gravelly loam, gravelly fine sandy loam, loam, silt loam, very fine sandy loam	ML, SM	A-4, A-2	0	0	75-100	55-100	55-100	25-85	15-25	NP-5
	24-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0-10	75-90	35-85	10-75	0-30	0-15	NP
30C: Branford-----	0-8	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-85	15-25	NP-5
	8-18	Loam, silt loam, very fine sandy loam, gravelly fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	35-85	15-25	NP-5
	18-24	Gravelly loam, gravelly fine sandy loam, loam, silt loam, very fine sandy loam	ML, SM	A-4, A-2	0	0	75-100	55-100	55-100	25-85	15-25	NP-5
	24-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0-10	75-90	35-85	10-75	0-30	0-15	NP
31A: Copake-----	0-6	Fine sandy loam	SM	A-4	0	0-5	90-100	80-90	70-90	35-45	15-25	NP-5
	6-13	Gravelly fine sandy loam, fine sandy loam, loam	ML, SM	A-2, A-4	0	0-10	75-100	70-90	60-85	25-60	15-25	NP-5
	13-21	Gravelly fine sandy loam, loam, silt loam	SM, CL-ML, ML, SC-SM	A-4, A-2	0	0-15	70-100	60-90	50-90	20-70	15-25	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
31A: Copake-----	21-31	Gravelly fine sandy loam, silt loam, loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-15	70-100	60-90	50-90	20-70	15-25	NP-10
	31-56	Very gravelly coarse sand, fine sand, gravelly sand, extremely gravelly coarse sand, loamy fine sand	SP-SM, GP-GM, GM, SM	A-1, A-2, A-3	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	56-65	Gravelly sand, loamy fine sand, fine sand, coarse sand, extremely gravelly coarse sand	GM, GP-GM, SM, SW-SM	A-2, A-1, A-3	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	65-75	Gravelly sand, coarse sand, loamy fine sand, fine sand, extremely gravelly coarse sand	SW-SM, GM, GP-GM, SM	A-2, A-1, A-3	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	75-80	Gravelly sand, coarse sand, fine sand, extremely gravelly coarse sand, loamy fine sand	SW-SM, SM, GM, GP-GM	A-3, A-1, A-2	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
31B: Copake-----	0-6	Fine sandy loam	SM	A-4	0	0-5	90-100	80-90	70-90	35-45	15-25	NP-5
	6-13	Gravelly fine sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0	0-10	75-100	70-90	60-85	25-60	15-25	NP-5
	13-21	Gravelly fine sandy loam, loam, silt loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-15	70-100	60-90	50-90	20-70	15-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
31B: Copake-----	21-31	Gravelly fine sandy loam, silt loam, loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-15	70-100	60-90	50-90	20-70	15-25	NP-10
	31-56	Very gravelly coarse sand, fine sand, gravelly sand, extremely gravelly coarse sand, loamy fine sand	GP-GM, GM, SM, SP-SM	A-3, A-1, A-2	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	56-65	Extremely gravelly coarse sand, loamy fine sand, gravelly sand, fine sand, coarse sand	GM, GP-GM, SM, SW-SM	A-1, A-3, A-2	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	65-75	Gravelly sand, coarse sand, loamy fine sand, fine sand, extremely gravelly coarse sand	SW-SM, GM, GP-GM, SM	A-1, A-3, A-2	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	75-80	Gravelly sand, coarse sand, fine sand, extremely gravelly coarse sand, loamy fine sand	SM, SW-SM, GM, GP-GM	A-1, A-3, A-2	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
31C: Copake-----	0-6	Fine sandy loam	SM	A-4	0	0-5	90-100	80-90	70-90	35-45	15-25	NP-5
	6-13	Gravelly fine sandy loam, fine sandy loam, loam	ML, SM	A-2, A-4	0	0-10	75-100	70-90	60-85	25-60	15-25	NP-5
	13-21	Gravelly fine sandy loam, loam, silt loam	SC-SM, SM, CL-ML, ML	A-4, A-2	0	0-15	70-100	60-90	50-90	20-70	15-25	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
31C: Copake-----	21-31	Gravelly fine sandy loam, silt loam, loam	SC-SM, CL-ML, ML, SM	A-4, A-2	0	0-15	70-100	60-90	50-90	20-70	15-25	NP-10
	31-56	Very gravelly coarse sand, fine sand, gravelly sand, extremely gravelly coarse sand, loamy fine sand	GP-GM, GM, SM, SP-SM	A-2, A-3, A-1	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	56-65	Loamy fine sand, extremely gravelly coarse sand, gravelly sand, coarse sand, fine sand	SW-SM, GM, GP-GM, SM	A-3, A-2, A-1	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	65-75	Gravelly sand, coarse sand, loamy fine sand, fine sand, extremely gravelly coarse sand	SW-SM, SM, GM, GP-GM	A-3, A-2, A-1	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
	75-80	Gravelly sand, coarse sand, fine sand, extremely gravelly coarse sand, loamy fine sand	SM, GP-GM, GM, SW-SM	A-3, A-2, A-1	0-10	0-50	40-100	20-90	10-85	5-30	0-20	NP-5
32A: Haven-----	0-7	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-100	15-25	NP-5
	7-14	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	14-20	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
32A: Haven-----	0-7	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-100	15-25	NP-5
	7-14	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	14-20	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	20-24	Fine sandy loam, very fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	75-100	70-95	30-65	15-25	NP-5
	24-60	Stratified very gravelly sand to gravelly fine sand	SM, SW-SM, SW	A-1, A-2, A-3	0	0-15	75-90	30-75	10-70	0-15	10-20	NP
Enfield-----	0-3	Slightly decomposed plant material	PT		0	0	100	100	100	100	---	---
	3-4	Moderately decomposed plant material	PT		0	0	100	100	100	100	---	---
	4-12	Silt loam	ML	A-4	0	0	100	90-100	90-100	60-95	15-25	NP-5
	12-20	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	20-26	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	26-30	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	30-37	Stratified coarse sand to very gravelly loamy sand	SP-SM, SM	A-1, A-2, A-3	0	0	70-100	25-100	10-95	5-25	15-25	NP-5
	37-65	Stratified very gravelly coarse sand to loamy sand	SM, SP	A-1, A-3	0	0-15	75-100	30-100	5-95	1-35	10-20	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
32B: Haven-----	0-7	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-100	15-25	NP-5
	7-14	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	14-20	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	20-24	Fine sandy loam, very fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	75-100	70-95	30-65	15-25	NP-5
	24-60	Stratified very gravelly sand to gravelly fine sand	SM, SW-SM, SW	A-1, A-2, A-3	0	0-15	75-90	30-75	10-70	0-15	10-20	NP
Enfield-----	0-3	Slightly decomposed plant material	PT		0	0	100	100	100	100	---	---
	3-4	Moderately decomposed plant material	PT		0	0	100	100	100	100	---	---
	4-12	Silt loam	ML	A-4	0	0	100	90-100	90-100	60-95	15-25	NP-5
	12-20	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	20-26	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	26-30	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	30-37	Stratified coarse sand to very gravelly loamy sand	SP-SM, SM	A-1, A-2, A-3	0	0	70-100	25-100	10-95	5-25	15-25	NP-5
	37-65	Stratified very gravelly coarse sand to loamy sand	SM, SP	A-3, A-1	0	0-15	75-100	30-100	5-95	1-35	10-20	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
32C: Haven-----	0-7	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-100	15-25	NP-5
	7-14	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	14-20	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	20-24	Fine sandy loam, very fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	75-100	70-95	30-65	15-25	NP-5
	24-60	Stratified very gravelly sand to gravelly fine sand	SM, SW-SM, SW	A-1, A-2, A-3	0	0-15	75-90	30-75	10-70	0-15	10-20	NP
Enfield-----	0-3	Slightly decomposed plant material	PT		0	0	100	100	100	100	---	---
	3-4	Moderately decomposed plant material	PT		0	0	100	100	100	100	---	---
	4-12	Silt loam	ML	A-4	0	0	100	90-100	90-100	60-95	15-25	NP-5
	12-20	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	20-26	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	26-30	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	40-95	15-25	NP-5
	30-37	Stratified coarse sand to very gravelly loamy sand	SM, SP-SM	A-1, A-2, A-3	0	0	70-100	25-100	10-95	5-25	15-25	NP-5
	37-65	Stratified very gravelly coarse sand to loamy sand	SM, SP	A-3, A-1	0	0-15	75-100	30-100	5-95	1-35	10-20	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
33A: Hartford-----	0-8	Sandy loam	SM	A-4, A-2	0	0	85-100	75-90	60-85	30-50	0-15	NP
	8-20	Sandy loam, gravelly sandy loam	SM	A-2, A-4	0	0-10	70-100	60-90	45-80	25-50	0-15	NP
	20-26	Loamy sand, sandy loam, gravelly sandy loam	SM	A-2	0	0-10	70-100	60-90	50-65	25-35	0-15	NP
	26-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0-5	0-15	55-90	40-85	20-80	5-30	0-15	NP
33B: Hartford-----	0-8	Sandy loam	SM	A-4, A-2	0	0	85-100	75-90	60-85	30-50	0-15	NP
	8-20	Sandy loam, gravelly sandy loam	SM	A-4, A-2	0	0-10	70-100	60-90	45-80	25-50	0-15	NP
	20-26	Loamy sand, sandy loam, gravelly sandy loam	SM	A-2	0	0-10	70-100	60-90	50-65	25-35	0-15	NP
	26-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0-5	0-15	55-90	40-85	20-80	5-30	0-15	NP
34A: Merrimac-----	0-9	Sandy loam	ML, SM	A-2, A-4	0	0	90-100	70-90	45-85	25-60	0-20	NP
	9-16	Sandy loam, fine sandy loam	SM	A-2, A-4	0	0	90-100	75-90	45-90	25-50	0-25	NP
	16-24	Gravelly sandy loam, sandy loam	SM	A-2, A-4	0	0	80-100	55-90	35-70	15-50	0-25	NP
	24-60	Stratified very gravelly coarse sand to gravelly sand	SP, SP-SM	A-1	0	10-20	65-85	35-60	35-60	1-15	0-25	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
34B: Merrimac-----	0-9	Sandy loam	ML, SM	A-2, A-4	0	0	90-100	70-90	45-85	25-60	0-20	NP
	9-16	Sandy loam, fine sandy loam	SM	A-2, A-4	0	0	90-100	75-90	45-90	25-50	0-25	NP
	16-24	Gravelly sandy loam, sandy loam	SM	A-2, A-4	0	0	80-100	55-90	35-70	15-50	0-25	NP
	24-60	Stratified very gravelly coarse sand to gravelly sand	SP, SP-SM	A-1	0	10-20	65-85	35-60	35-60	1-15	0-25	NP
34C: Merrimac-----	0-9	Sandy loam	ML, SM	A-2, A-4	0	0	90-100	70-90	45-85	25-60	0-20	NP
	9-16	Sandy loam, fine sandy loam	SM	A-2, A-4	0	0	90-100	75-90	45-90	25-50	0-25	NP
	16-24	Gravelly sandy loam, sandy loam	SM	A-2, A-4	0	0	80-100	55-90	35-70	15-50	0-25	NP
	24-60	Stratified very gravelly coarse sand to gravelly sand	SP, SP-SM	A-1	0	10-20	65-85	35-60	35-60	1-15	0-25	NP
35A: Penwood-----	0-8	Loamy sand	SM	A-2	0	0	90-100	85-100	50-75	25-30	0-15	NP
	8-18	Loamy sand, loamy fine sand	SM	A-2	0	0	90-100	85-100	55-95	25-30	0-15	NP
	18-30	Sand, fine sand	SM, SP-SM	A-1, A-2	0	0	90-100	85-100	45-90	10-20	0-15	NP
	30-60	Sand, fine sand	SM, SP-SM	A-2, A-1	0	0	85-100	75-100	40-90	10-20	0-15	NP
35B: Penwood-----	0-8	Loamy sand	SM	A-2	0	0	90-100	85-100	50-75	25-30	0-15	NP
	8-18	Loamy sand, loamy fine sand	SM	A-2	0	0	90-100	85-100	55-95	25-30	0-15	NP
	18-30	Sand, fine sand	SM, SP-SM	A-2, A-1	0	0	90-100	85-100	45-90	10-20	0-15	NP
	30-60	Sand, fine sand	SM, SP-SM	A-1, A-2	0	0	85-100	75-100	40-90	10-20	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
36A: Windsor-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Loamy sand	SM	A-2	0	0	100	85-100	50-80	15-35	0-15	NP
	3-9	Loamy sand, loamy fine sand	SM	A-2	0	0	100	85-100	50-95	15-30	0-15	NP
	9-21	Loamy sand, loamy fine sand	SM	A-2	0	0	100	85-100	50-95	15-30	0-15	NP
	21-25	Sand, fine sand, loamy sand	SW-SM, SM	A-2, A-3	0	0	100	85-100	60-70	5-30	0-15	NP
	25-65	Sand, fine sand, loamy sand	SM, SW-SM	A-2, A-3	0	0	95-100	75-100	60-70	5-30	0-15	NP
36B: Windsor-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Loamy sand	SM	A-2	0	0	100	85-100	50-80	15-35	0-15	NP
	3-9	Loamy sand, loamy fine sand	SM	A-2	0	0	100	85-100	50-95	15-30	0-15	NP
	9-21	Loamy sand, loamy fine sand	SM	A-2	0	0	100	85-100	50-95	15-30	0-15	NP
	21-25	Sand, fine sand, loamy sand	SM, SW-SM	A-2, A-3	0	0	100	85-100	60-70	5-30	0-15	NP
	25-65	Sand, fine sand, loamy sand	SW-SM, SM	A-3, A-2	0	0	95-100	75-100	60-70	5-30	0-15	NP
36C: Windsor-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Loamy sand	SM	A-2	0	0	100	85-100	50-80	15-35	0-15	NP
	3-9	Loamy sand, loamy fine sand	SM	A-2	0	0	100	85-100	50-95	15-30	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
36C: Windsor-----	9-21	Loamy sand, loamy fine sand	SM	A-2	0	0	100	85-100	50-95	15-30	0-15	NP
	21-25	Sand, fine sand, loamy sand	SM, SW-SM	A-2, A-3	0	0	100	85-100	60-70	5-30	0-15	NP
	25-65	Sand, fine sand, loamy sand	SM, SW-SM	A-2, A-3	0	0	95-100	75-100	60-70	5-30	0-15	NP
37A: Manchester-----	0-9	Gravelly sandy loam	SM	A-4, A-2	0	0	75-85	50-70	35-70	20-45	0-15	NP
	9-18	Gravelly loamy sand, very gravelly loamy sand, very gravelly sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	60-85	35-75	20-60	5-25	0-15	NP
	18-65	Stratified extremely gravelly coarse sand to very gravelly loamy sand	GM, SM, SW-SM	A-1	0	0-15	45-60	20-50	10-40	0-20	0-15	NP
37C: Manchester-----	0-9	Gravelly sandy loam	SM	A-2, A-4	0	0	75-85	50-70	35-70	20-45	0-15	NP
	9-18	Gravelly loamy sand, very gravelly loamy sand, very gravelly sand	SM, SP-SM	A-2, A-3, A-1	0	0-10	60-85	35-75	20-60	5-25	0-15	NP
	18-65	Stratified extremely gravelly coarse sand to very gravelly loamy sand	GM, SM, SW-SM	A-1	0	0-15	45-60	20-50	10-40	0-20	0-15	NP
37E: Manchester-----	0-9	Gravelly sandy loam	SM	A-4, A-2	0	0	75-85	50-70	35-70	20-45	0-15	NP
	9-18	Gravelly loamy sand, very gravelly loamy sand, very gravelly sand	SM, SP-SM	A-3, A-2, A-1	0	0-10	60-85	35-75	20-60	5-25	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
37E: Manchester-----	18-65	Stratified extremely gravelly coarse sand to very gravelly loamy sand	GM, SM, SW-SM	A-1	0	0-15	45-60	20-50	10-40	0-20	0-15	NP
38A: Hinckley-----	0-8	Gravelly sandy loam	SM	A-2, A-4	0-5	1-15	65-85	60-75	45-70	25-50	0-20	NP
	8-20	Very gravelly loamy sand, gravelly loamy sand, loamy fine sand, very gravelly loamy coarse sand	SM, GM	A-1, A-2	0-10	1-15	50-100	40-90	25-90	15-25	0-20	NP
	20-27	Very gravelly sand, gravelly sand	SM, GM	A-1	1-10	5-20	50-90	40-80	25-50	5-15	0-20	NP
	27-42	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
	42-60	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
38C: Hinckley-----	0-8	Gravelly sandy loam	SM	A-2, A-4	0-5	1-15	65-85	60-75	45-70	25-50	0-20	NP
	8-20	Very gravelly loamy sand, gravelly loamy sand, loamy fine sand, very gravelly loamy coarse sand	SM, GM	A-1, A-2	0-10	1-15	50-100	40-90	25-90	15-25	0-20	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
38C: Hinckley-----	20-27	Very gravelly sand, gravelly sand	SM, GM	A-1	1-10	5-20	50-90	40-80	25-50	5-15	0-20	NP
	27-42	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
	42-60	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
38E: Hinckley-----	0-8	Gravelly sandy loam	SM	A-2, A-4	0-5	1-15	65-85	60-75	45-70	25-50	0-20	NP
	8-20	Very gravelly loamy sand, gravelly loamy sand, loamy fine sand, very gravelly loamy coarse sand	SM, GM	A-1, A-2	0-10	1-15	50-100	40-90	25-90	15-25	0-20	NP
	20-27	Very gravelly sand, gravelly sand	SM, GM	A-1	1-10	5-20	50-90	40-80	25-50	5-15	0-20	NP
	27-42	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
	42-60	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
39A: Groton-----	0-8	Gravelly sandy loam	SM	A-1, A-2	0	0-5	65-75	50-70	30-50	15-35	0-15	NP
	8-18	Very gravelly sandy loam, gravelly fine sandy loam, gravelly sandy loam	GM, SM	A-2, A-1	0	0-10	45-70	30-70	15-65	10-35	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
39A: Groton-----	18-24	Very gravelly loamy sand, very gravelly sandy loam, gravelly loamy sand	GM, GW-GM	A-1	0	0-10	40-50	30-50	15-50	5-25	0-15	NP
	24-30	Very gravelly loamy sand, very gravelly sandy loam, gravelly loamy sand, very gravelly loamy fine sand	GM, GW-GM	A-1	0	0-10	40-50	20-50	10-35	5-25	0-15	NP
	30-52	Stratified extremely gravelly coarse sand to very gravelly loamy fine sand	GW-GM, GM	A-1	0	0-10	40-50	20-50	5-50	0-15	0-15	NP
	52-72	Stratified extremely gravelly coarse sand to gravelly loamy fine sand	GM, GW-GM	A-3, A-1	0	0-10	40-65	20-65	5-65	0-15	0-15	NP
39C: Groton-----	0-8	Gravelly sandy loam	SM	A-2, A-1	0	0-5	65-75	50-70	30-50	15-35	0-15	NP
	8-18	Very gravelly sandy loam, gravelly fine sandy loam, gravelly sandy loam	GM, SM	A-1, A-2	0	0-10	45-70	30-70	15-65	10-35	0-15	NP
	18-24	Very gravelly loamy sand, very gravelly sandy loam, gravelly loamy sand	GM, GW-GM	A-1	0	0-10	40-50	30-50	15-50	5-25	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
39C: Groton-----	24-30	Very gravelly loamy sand, very gravelly sandy loam, gravelly loamy sand, very gravelly loamy fine sand	GM, GW-GM	A-1	0	0-10	40-50	20-50	10-35	5-25	0-15	NP
	30-52	Stratified extremely gravelly coarse sand to very gravelly loamy fine sand	GM, GW-GM	A-1	0	0-10	40-50	20-50	5-50	0-15	0-15	NP
	52-72	Stratified extremely gravelly coarse sand to gravelly loamy fine sand	GW-GM, GM	A-1, A-3	0	0-10	40-65	20-65	5-65	0-15	0-15	NP
39E: Groton-----	0-8	Gravelly sandy loam	SM	A-1, A-2	0	0-5	65-75	50-70	30-50	15-35	0-15	NP
	8-18	Very gravelly sandy loam, gravelly fine sandy loam, gravelly sandy loam	GM, SM	A-2, A-1	0	0-10	45-70	30-70	15-65	10-35	0-15	NP
	18-24	Very gravelly loamy sand, very gravelly sandy loam, gravelly loamy sand	GW-GM, GM	A-1	0	0-10	40-50	30-50	15-50	5-25	0-15	NP
	24-30	Very gravelly loamy sand, very gravelly sandy loam, gravelly loamy sand, very gravelly loamy fine sand	GW-GM, GM	A-1	0	0-10	40-50	20-50	10-35	5-25	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
39E: Groton-----	30-52	Stratified extremely gravelly coarse sand to very gravelly loamy fine sand	GM, GW-GM	A-1	0	0-10	40-50	20-50	5-50	0-15	0-15	NP
	52-72	Stratified extremely gravelly coarse sand to gravelly loamy fine sand	GM, GW-GM	A-3, A-1	0	0-10	40-65	20-65	5-65	0-15	0-15	NP
40A: Ludlow-----	0-8	Silt loam	ML	A-4	0	0	85-100	75-90	70-90	50-80	15-25	NP-5
	8-20	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	20-26	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	26-65	Gravelly loam, gravelly silt loam, silt loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-15	70-100	65-90	60-90	30-75	15-25	NP-5
40B: Ludlow-----	0-8	Silt loam	ML	A-4	0	0	85-100	75-90	70-90	50-80	15-25	NP-5
	8-20	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	20-26	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	26-65	Gravelly loam, gravelly silt loam, silt loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-15	70-100	65-90	60-90	30-75	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
41B: Ludlow-----	0-8	Silt loam	ML	A-4	0	0	85-100	75-90	70-90	50-80	15-25	NP-5
	8-20	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	20-26	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	26-65	Gravelly loam, gravelly silt loam, silt loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-15	70-100	65-90	60-90	30-75	15-25	NP-5
42C: Ludlow-----	0-8	Silt loam	ML	A-4	0	0	85-100	75-90	70-90	50-80	15-25	NP-5
	8-20	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	20-26	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	26-65	Gravelly loam, gravelly silt loam, silt loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-15	70-100	65-90	60-90	30-75	15-25	NP-5
43A: Rainbow-----	0-6	Silt loam	ML	A-4	0	0-5	90-100	80-100	70-100	50-90	15-25	NP-5
	6-18	Silt loam, very fine sandy loam, gravelly loam	ML, SM	A-2, A-4	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	18-26	Silt loam, very fine sandy loam, gravelly loam	ML, SM	A-2, A-4	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	26-65	Gravelly fine sandy loam, gravelly sandy loam, loam	SM, ML	A-4, A-2	0-5	0-10	75-100	60-90	45-80	20-60	10-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
				Pct	Pct					Pct		
43B: Rainbow-----	In											
	0-6	Silt loam	ML	A-4	0	0-5	90-100	80-100	70-100	50-90	15-25	NP-5
	6-18	Silt loam, very fine sandy loam, gravelly loam	SM, ML	A-4, A-2	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	18-26	Silt loam, very fine sandy loam, gravelly loam	ML, SM	A-2, A-4	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	26-65	Gravelly fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-5	0-10	75-100	60-90	45-80	20-60	10-25	NP-5
44B: Rainbow-----	0-6	Silt loam	ML	A-4	0	0-5	90-100	80-100	70-100	50-90	15-25	NP-5
	6-18	Silt loam, very fine sandy loam, gravelly loam	ML, SM	A-2, A-4	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	18-26	Silt loam, very fine sandy loam, gravelly loam	ML, SM	A-2, A-4	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	26-65	Gravelly fine sandy loam, gravelly sandy loam, loam	SM, ML	A-2, A-4	0-5	0-10	75-100	60-90	45-80	20-60	10-25	NP-5
45A: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
45A: Woodbridge-----	30-43	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
45B: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	30-43	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
45C: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
45C: Woodbridge-----	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	30-43	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
46B: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	30-43	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
46B: Woodbridge-----	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
46C: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	30-43	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
47C: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
47C: Woodbridge-----	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	30-43	Gravelly fine sandy loam, fine sandy loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
48B: Georgia-----	0-8	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	8-14	Loam, silt loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	14-24	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	24-60	Gravelly fine sandy loam, gravelly loam, loam, silt loam	ML, SM	A-4, A-2	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
Amenia-----	0-9	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	9-16	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	16-25	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	25-60	Gravelly loam, gravelly silt loam, silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
48C: Georgia-----	0-8	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	8-14	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	14-24	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	24-60	Gravelly fine sandy loam, gravelly loam, loam, silt loam	SM, ML	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
Amenia-----	0-9	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	9-16	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	16-25	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	25-60	Gravelly loam, gravelly silt loam, silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
49B: Georgia-----	0-8	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	8-14	Loam, silt loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	14-24	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	24-60	Gravelly fine sandy loam, gravelly loam, loam, silt loam	ML, SM	A-4, A-2	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
49B: Amenia-----	0-9	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	9-16	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	16-25	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	25-60	Gravelly loam, gravelly silt loam, silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
49C: Georgia-----	0-8	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	8-14	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	14-24	Loam, silt loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	24-60	Gravelly fine sandy loam, gravelly loam, loam, silt loam	SM, ML	A-4, A-2	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
Amenia-----	0-9	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	9-16	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	16-25	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5
	25-60	Gravelly loam, gravelly silt loam, silt loam	ML, SM	A-4	0-5	0-5	70-100	60-90	55-90	40-75	20-30	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
50A: Sutton-----	0-6	Fine sandy loam	SC-SM, SM	A-4, A-2	0	0-5	90-100	80-90	70-90	30-45	15-25	NP-5
	6-12	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, SC-SM, CL-ML	A-2, A-4	0	0-5	60-100	55-90	40-90	20-55	15-25	NP-5
	12-24	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	SM, ML, CL-ML, SC-SM	A-4, A-2	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	24-28	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	28-36	Gravelly fine sandy loam, gravelly sandy loam, sandy loam	SM, SC-SM	A-2	0-5	0-10	65-100	60-90	55-85	30-40	15-20	NP-5
	36-65	Gravelly sandy loam, gravelly fine sandy loam, sandy loam	SC-SM, SM	A-2, A-4	0-5	0-10	65-100	60-90	45-85	25-50	15-20	NP-5
50B: Sutton-----	0-6	Fine sandy loam	SC-SM, SM	A-4, A-2	0	0-5	90-100	80-90	70-90	30-45	15-25	NP-5
	6-12	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	CL-ML, SM, ML, SC-SM	A-4, A-2	0	0-5	60-100	55-90	40-90	20-55	15-25	NP-5
	12-24	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	24-28	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
50B: Sutton-----	28-36	Gravelly fine sandy loam, gravelly sandy loam, sandy loam	SC-SM, SM	A-2	0-5	0-10	65-100	60-90	55-85	30-40	15-20	NP-5
	36-65	Gravelly sandy loam, gravelly fine sandy loam, sandy loam	SM, SC-SM	A-2, A-4	0-5	0-10	65-100	60-90	45-85	25-50	15-20	NP-5
51B: Sutton-----	0-6	Fine sandy loam	SM, SC-SM	A-4, A-2	0	0-5	90-100	80-90	70-90	30-45	15-25	NP-5
	6-12	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	SM, SC-SM, CL-ML, ML	A-4, A-2	0	0-5	60-100	55-90	40-90	20-55	15-25	NP-5
	12-24	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, CL- ML, SC-SM	A-2, A-4	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	24-28	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, CL- ML, SC-SM	A-2, A-4	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	28-36	Gravelly fine sandy loam, gravelly sandy loam, sandy loam	SC-SM, SM	A-2	0-5	0-10	65-100	60-90	55-85	30-40	15-20	NP-5
	36-65	Gravelly sandy loam, gravelly fine sandy loam, sandy loam	SC-SM, SM	A-2, A-4	0-5	0-10	65-100	60-90	45-85	25-50	15-20	NP-5
52C: Sutton-----	0-6	Fine sandy loam	SC-SM, SM	A-4, A-2	0	0-5	90-100	80-90	70-90	30-45	15-25	NP-5
	6-12	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, SC- SM, CL-ML	A-4, A-2	0	0-5	60-100	55-90	40-90	20-55	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
52C: Sutton-----	12-24	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	SC-SM, CL-ML, SM, ML	A-4, A-2	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	24-28	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, CL- ML, SC-SM	A-4, A-2	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	28-36	Gravelly fine sandy loam, gravelly sandy loam, sandy loam	SC-SM, SM	A-2	0-5	0-10	65-100	60-90	55-85	30-40	15-20	NP-5
	36-65	Gravelly sandy loam, gravelly fine sandy loam, sandy loam	SC-SM, SM	A-2, A-4	0-5	0-10	65-100	60-90	45-85	25-50	15-20	NP-5
53A: Wapping-----	0-11	Very fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	70-100	40-75	15-25	NP-5
	11-16	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	16-20	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	20-28	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-85	55-75	40-70	20-35	15-25	NP-5
	28-36	Gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-1	0-5	0-5	70-85	55-75	35-65	15-20	0-20	NP-5
	36-80	Gravelly loamy sand, very gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM, SW-SM	A-1, A-2	0-10	0-15	60-85	40-75	30-65	10-20	0-20	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
53B: Wapping-----	0-11	Very fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	70-100	40-75	15-25	NP-5
	11-16	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	16-20	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	20-28	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-85	55-75	40-70	20-35	15-25	NP-5
	28-36	Gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-1	0-5	0-5	70-85	55-75	35-65	15-20	0-20	NP-5
	36-80	Gravelly loamy sand, very gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM, SW-SM	A-2, A-1	0-10	0-15	60-85	40-75	30-65	10-20	0-20	NP-5
54B: Wapping-----	0-11	Very fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	70-100	40-75	15-25	NP-5
	11-16	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	16-20	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	20-28	Gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-1	0-5	0-5	70-85	55-75	40-70	20-35	15-25	NP-5
	28-36	Gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-1	0-5	0-5	70-85	55-75	35-65	15-20	0-20	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
54B: Wapping-----	36-80	Gravelly loamy sand, very gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM, SW-SM	A-2, A-1	0-10	0-15	60-85	40-75	30-65	10-20	0-20	NP-5
55A: Watchaug-----	0-8	Fine sandy loam	SM	A-4	0	0-10	90-100	80-90	75-85	35-45	15-25	NP
	8-18	Fine sandy loam, silt loam, gravelly loam	ML, SM	A-2, A-4	0	0-10	65-100	55-90	45-90	25-70	15-25	NP-5
	18-24	Fine sandy loam, silt loam, gravelly loam	ML, SM	A-2, A-4	0-10	0-15	70-100	60-90	55-90	25-70	15-25	NP-5
	24-65	Gravelly sandy loam, gravelly fine sandy loam, fine sandy loam	SM	A-2, A-4	0-10	0-15	70-100	65-90	45-80	30-40	15-25	NP-5
55B: Watchaug-----	0-8	Fine sandy loam	SM	A-4	0	0-10	90-100	80-90	75-85	35-45	15-25	NP
	8-18	Fine sandy loam, silt loam, gravelly loam	ML, SM	A-2, A-4	0	0-10	65-100	55-90	45-90	25-70	15-25	NP-5
	18-24	Fine sandy loam, silt loam, gravelly loam	ML, SM	A-2, A-4	0-10	0-15	70-100	60-90	55-90	25-70	15-25	NP-5
	24-65	Gravelly sandy loam, gravelly fine sandy loam, fine sandy loam	SM	A-2, A-4	0-10	0-15	70-100	65-90	45-80	30-40	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
56B: Watchaug-----	0-8	Fine sandy loam	SM	A-4	0	0-10	90-100	80-90	75-85	35-45	15-25	NP
	8-18	Fine sandy loam, silt loam, gravelly loam	ML, SM	A-2, A-4	0	0-10	65-100	55-90	45-90	25-70	15-25	NP-5
	18-24	Fine sandy loam, silt loam, gravelly loam	ML, SM	A-4, A-2	0-10	0-15	70-100	60-90	55-90	25-70	15-25	NP-5
	24-65	Gravelly sandy loam, gravelly fine sandy loam, fine sandy loam	SM	A-2, A-4	0-10	0-15	70-100	65-90	45-80	30-40	15-25	NP-5
57B: Gloucester-----	0-4	Gravelly sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	55-75	30-50	0-20	NP
	4-12	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-80	55-70	40-65	20-35	0-20	NP
	12-25	Very gravelly loamy sand, very gravelly loamy coarse sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-15	10-15	50-60	35-55	15-45	5-10	0-15	NP
	25-35	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
	35-60	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
57C: Gloucester-----	0-4	Gravelly sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	55-75	30-50	0-20	NP
	4-12	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-80	55-70	40-65	20-35	0-20	NP
	12-25	Very gravelly loamy sand, very gravelly loamy coarse sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-15	10-15	50-60	35-55	15-45	5-10	0-15	NP
	25-35	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
	35-60	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
57D: Gloucester-----	0-4	Gravelly sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	55-75	30-50	0-20	NP
	4-12	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-80	55-70	40-65	20-35	0-20	NP
	12-25	Very gravelly loamy sand, very gravelly loamy coarse sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-15	10-15	50-60	35-55	15-45	5-10	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
57D: Gloucester-----	25-35	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
	35-60	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
58B: Gloucester-----	0-4	Gravelly sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	55-75	30-50	0-20	NP
	4-12	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-80	55-70	40-65	20-35	0-20	NP
	12-25	Very gravelly loamy sand, very gravelly loamy coarse sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-15	10-15	50-60	35-55	15-45	5-10	0-15	NP
	25-35	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
	35-60	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
58C: Gloucester-----	In											
	0-4	Gravelly sandy loam	SM	A-4, A-2	0-5	0-5	95-100	80-90	55-75	30-50	0-20	NP
	4-12	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-80	55-70	40-65	20-35	0-20	NP
	12-25	Very gravelly loamy sand, very gravelly loamy coarse sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-15	10-15	50-60	35-55	15-45	5-10	0-15	NP
	25-35	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
	35-60	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
59C: Gloucester-----	0-4	Gravelly sandy loam	SM	A-4, A-2	0-5	0-5	95-100	80-90	55-75	30-50	0-20	NP
	4-12	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-80	55-70	40-65	20-35	0-20	NP
	12-25	Very gravelly loamy sand, very gravelly loamy coarse sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-15	10-15	50-60	35-55	15-45	5-10	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
59C: Gloucester-----	In											
	25-35	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
	35-60	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
59D: Gloucester-----	0-4	Gravelly sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	55-75	30-50	0-20	NP
	4-12	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-80	55-70	40-65	20-35	0-20	NP
	12-25	Very gravelly loamy sand, very gravelly loamy coarse sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-15	10-15	50-60	35-55	15-45	5-10	0-15	NP
	25-35	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP
	35-60	Very gravelly loamy coarse sand, very gravelly loamy sand, very gravelly loamy fine sand	GW-GM, SW-SM	A-1	0-20	10-20	50-60	35-55	15-45	5-10	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
60B: Canton-----	In											
	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-3	Gravelly fine sandy loam	SM	A-2	0-5	0-10	80-90	70-75	60-70	25-35	10-25	NP-5
	3-15	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	15-24	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	24-30	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	30-60	Very gravelly loamy sand, loamy fine sand, gravelly loamy coarse sand, gravelly loamy sand	SM	A-1, A-2	0-15	0-15	60-85	45-75	30-75	10-25	0-15	NP
Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
60B: Charlton-----	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
60C: Canton-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-3	Gravelly fine sandy loam	SM	A-2	0-5	0-10	80-90	70-75	60-70	25-35	10-25	NP-5
	3-15	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	15-24	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	24-30	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	30-60	Very gravelly loamy sand, loamy fine sand, gravelly loamy coarse sand, gravelly loamy sand	SM	A-1, A-2	0-15	0-15	60-85	45-75	30-75	10-25	0-15	NP
Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
60C: Charlton-----	In											
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
60D: Canton-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-3	Gravelly fine sandy loam	SM	A-2	0-5	0-10	80-90	70-75	60-70	25-35	10-25	NP-5
	3-15	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	15-24	Gravelly loam, very fine sandy loam, fine sandy loam	SM, ML	A-4, A-2	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	24-30	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	30-60	Very gravelly loamy sand, loamy fine sand, gravelly loamy coarse sand, gravelly loamy sand	SM	A-1, A-2	0-15	0-15	60-85	45-75	30-75	10-25	0-15	NP
Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
60D: Charlton-----	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
61B: Canton-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-3	Gravelly fine sandy loam	SM	A-2	0-5	0-10	80-90	70-75	60-70	25-35	10-25	NP-5
	3-15	Gravelly loam, very fine sandy loam, fine sandy loam	SM, ML	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	15-24	Gravelly loam, very fine sandy loam, fine sandy loam	SM, ML	A-4, A-2	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	24-30	Gravelly loam, very fine sandy loam, fine sandy loam	SM, ML	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	30-60	Very gravelly loamy sand, loamy fine sand, gravelly loamy coarse sand, gravelly loamy sand	SM	A-1, A-2	0-15	0-15	60-85	45-75	30-75	10-25	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
61B: Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
61C: Canton-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-3	Gravelly fine sandy loam	SM	A-2	0-5	0-10	80-90	70-75	60-70	25-35	10-25	NP-5
	3-15	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	15-24	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	24-30	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
61C: Canton-----	30-60	Very gravelly loamy sand, loamy fine sand, gravelly loamy coarse sand, gravelly loamy sand	SM	A-1, A-2	0-15	0-15	60-85	45-75	30-75	10-25	0-15	NP
61C: Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
62C: Canton-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-3	Gravelly fine sandy loam	SM	A-2	0-5	0-10	80-90	70-75	60-70	25-35	10-25	NP-5
	3-15	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	15-24	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
62C: Canton-----	24-30	Gravelly loam, very fine sandy loam, fine sandy loam	SM, ML	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	30-60	Very gravelly loamy sand, loamy fine sand, gravelly loamy coarse sand, gravelly loamy sand	SM	A-1, A-2	0-15	0-15	60-85	45-75	30-75	10-25	0-15	NP
Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
62D: Canton-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-3	Gravelly fine sandy loam	SM	A-2	0-5	0-10	80-90	70-75	60-70	25-35	10-25	NP-5
	3-15	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
62D: Canton-----	15-24	Gravelly loam, very fine sandy loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	24-30	Gravelly loam, very fine sandy loam, fine sandy loam	SM, ML	A-2, A-4	0-5	0-10	80-90	70-80	60-70	30-50	10-25	NP-5
	30-60	Very gravelly loamy sand, loamy fine sand, gravelly loamy coarse sand, gravelly loamy sand	SM	A-1, A-2	0-15	0-15	60-85	45-75	30-75	10-25	0-15	NP
Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
63B: Cheshire-----	0-8	Fine sandy loam	SM	A-2, A-4	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
63B: Cheshire-----	16-26	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5
63C: Cheshire-----	0-8	Fine sandy loam	SM	A-4, A-2	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	SM, ML	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	SM, ML	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5
63D: Cheshire-----	0-8	Fine sandy loam	SM	A-2, A-4	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	SM, ML	A-4, A-2	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-4, A-2	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5

Table 23.-Engineering Properties-Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
64B: Cheshire-----	0-8	Fine sandy loam	SM	A-2, A-4	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5
64C: Cheshire-----	0-8	Fine sandy loam	SM	A-2, A-4	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	SM, ML	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-4, A-2	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5
65C: Cheshire-----	0-8	Fine sandy loam	SM	A-2, A-4	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
65D: Cheshire-----	0-8	Fine sandy loam	SM	A-4, A-2	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	SM, ML	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	SM, ML	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5
66B: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5
	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP
66C: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
66C: Narragansett----	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP
67B: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5
	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP
67C: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
67C: Narragansett----	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP
68C: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5
	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP
68D: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
68D: Narragansett----	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP
69B: Yalesville-----	0-8	Fine sandy loam	SM, ML	A-4, A-2	0	0-5	90-95	80-95	75-95	30-55	15-25	NP-5
	8-14	Fine sandy loam, loam, sandy loam, gravelly fine sandy loam	SM, ML	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5
	14-25	Loam, fine sandy loam, sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5
	25-36	Gravelly sandy loam, loam, very gravelly fine sandy loam	GM, ML, SM	A-1, A-2, A-4	0-10	0-10	50-95	40-90	25-90	15-60	15-25	NP-5
	36-80	Unweathered bedrock			---	---	---	---	---	---	---	---
69C: Yalesville-----	0-8	Fine sandy loam	SM, ML	A-4, A-2	0	0-5	90-95	80-95	75-95	30-55	15-25	NP-5
	8-14	Fine sandy loam, loam, sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
				Pct	Pct					Pct		
69C: Yalesville-----	In											
	14-25	Loam, fine sandy loam, sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5
	25-36	Gravelly sandy loam, loam, very gravelly fine sandy loam	GM, ML, SM	A-1, A-2, A-4	0-10	0-10	50-95	40-90	25-90	15-60	15-25	NP-5
	36-80	Unweathered bedrock			---	---	---	---	---	---	---	---
70C: Branford-----	0-8	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-85	15-25	NP-5
	8-18	Loam, silt loam, very fine sandy loam, gravelly fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	35-85	15-25	NP-5
	18-24	Gravelly loam, gravelly fine sandy loam, loam, silt loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	75-100	55-100	55-100	25-85	15-25	NP-5
	24-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-3, A-1, A-2	0	0-10	75-90	35-85	10-75	0-30	0-15	NP
Holyoke-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	70-90	50-75	15-25	NP-5
	3-8	Silt loam, loam, fine sandy loam, very fine sandy loam, gravelly silt loam	GM, SM	A-4	0-5	0-10	65-100	55-90	55-80	35-45	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
					Pct	Pct					Pct	
70C: Holyoke-----	In											
	8-18	Gravelly silt loam, fine sandy loam, very fine sandy loam, loam	SM	A-4	0-10	0-15	70-100	60-90	55-85	35-45	15-25	NP-5
	18-80	Unweathered bedrock			---	---	---	---	---	---	---	---
71C: Brookfield-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	70-90	30-40	15-25	NP-5
	3-13	Gravelly fine sandy loam, sandy loam	ML, SM	A-2, A-4	0-5	0-5	70-100	55-90	40-90	20-50	15-25	NP-5
	13-27	Gravelly fine sandy loam, sandy loam	ML, SM	A-4, A-2	0-5	0-5	70-100	55-90	40-90	20-50	15-25	NP-5
	27-60	Gravelly sandy loam, gravelly fine sandy loam, fine sandy loam	SM	A-4, A-2	0-15	0-15	60-100	45-90	35-90	20-50	15-25	NP-5
Brimfield-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	70-90	30-40	15-25	NP-5
	3-6	Fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-5	0-5	95-100	80-90	65-90	30-55	15-25	NP-5
	6-17	Gravelly fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-10	0-10	75-100	60-90	45-90	25-55	15-25	NP-5
	17-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
71E: Brookfield-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	70-90	30-40	15-25	NP-5
	3-13	Gravelly fine sandy loam, sandy loam	ML, SM	A-4, A-2	0-5	0-5	70-100	55-90	40-90	20-50	15-25	NP-5
	13-27	Gravelly fine sandy loam, sandy loam	ML, SM	A-4, A-2	0-5	0-5	70-100	55-90	40-90	20-50	15-25	NP-5
	27-60	Gravelly sandy loam, gravelly fine sandy loam, fine sandy loam	SM	A-4, A-2	0-15	0-15	60-100	45-90	35-90	20-50	15-25	NP-5
Brimfield-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	70-90	30-40	15-25	NP-5
	3-6	Fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-5	0-5	95-100	80-90	65-90	30-55	15-25	NP-5
	6-17	Gravelly fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-10	0-10	75-100	60-90	45-90	25-55	15-25	NP-5
	17-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
73C: Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
73C: Charlton-----	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5
	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10
	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10
	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
73E: Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
73E: Charlton-----	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-4, A-2	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-4, A-2	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5
	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10
	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10
	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
74C: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5
	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
74C: Hollis-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-3	0-10	65-85	55-75	45-75	20-40	15-25	NP-5
	6-9	Channery fine sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0-5	0-10	65-100	55-90	50-90	25-50	15-25	NP-5
	9-15	Gravelly fine sandy loam, fine sandy loam	SM	A-4, A-2	0-10	0-20	70-100	65-90	55-90	25-50	15-25	NP-5
	15-80				---	---	---	---	---	---	---	---
75C: Hollis-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-3	0-10	65-85	55-75	45-75	20-40	15-25	NP-5
	6-9	Channery fine sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0-5	0-10	65-100	55-90	50-90	25-50	15-25	NP-5
	9-15	Gravelly fine sandy loam, fine sandy loam	SM	A-2, A-4	0-10	0-20	70-100	65-90	55-90	25-50	15-25	NP-5
	15-80				---	---	---	---	---	---	---	---
Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5
	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
75C: Chatfield-----	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-4, A-2	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10
	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
75E: Hollis-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-3	0-10	65-85	55-75	45-75	20-40	15-25	NP-5
	6-9	Channery fine sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0-5	0-10	65-100	55-90	50-90	25-50	15-25	NP-5
	9-15	Gravelly fine sandy loam, fine sandy loam	SM	A-2, A-4	0-10	0-20	70-100	65-90	55-90	25-50	15-25	NP-5
	15-80				---	---	---	---	---	---	---	---
Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5
	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10
	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
75E: Chatfield-----	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
76E: Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
Hollis-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-3	0-10	65-85	55-75	45-75	20-40	15-25	NP-5
	6-9	Channery fine sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0-5	0-10	65-100	55-90	50-90	25-50	15-25	NP-5
	9-15	Gravelly fine sandy loam, fine sandy loam	SM	A-2, A-4	0-10	0-20	70-100	65-90	55-90	25-50	15-25	NP-5
	15-80				---	---	---	---	---	---	---	---
76F: Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
Hollis-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-3	0-10	65-85	55-75	45-75	20-40	15-25	NP-5
	6-9	Channery fine sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0-5	0-10	65-100	55-90	50-90	25-50	15-25	NP-5
	9-15	Gravelly fine sandy loam, fine sandy loam	SM	A-2, A-4	0-10	0-20	70-100	65-90	55-90	25-50	15-25	NP-5
	15-80				---	---	---	---	---	---	---	---

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
77C: Cheshire-----	0-8	Fine sandy loam	SM	A-2, A-4	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-4, A-2	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5
Holyoke-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	70-90	50-75	15-25	NP-5
	3-8	Silt loam, loam, fine sandy loam, very fine sandy loam, gravelly silt loam	GM, SM	A-4	0-5	0-10	65-100	55-90	55-80	35-45	15-25	NP-5
	8-18	Gravelly silt loam, fine sandy loam, very fine sandy loam, loam	SM	A-4	0-10	0-15	70-100	60-90	55-85	35-45	15-25	NP-5
	18-80	Unweathered bedrock			---	---	---	---	---	---	---	---
77D: Cheshire-----	0-8	Fine sandy loam	SM	A-2, A-4	0-5	0-10	90-100	80-90	70-85	30-45	15-25	NP-5
	8-16	Fine sandy loam, silt loam, gravelly sandy loam	SM, ML	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5
	16-26	Fine sandy loam, silt loam, gravelly sandy loam	ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	40-90	20-75	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
77D: Cheshire-----	26-65	Gravelly sandy loam, fine sandy loam, gravelly fine sandy loam	SM	A-2	0-5	0-10	65-100	55-90	45-85	25-35	15-25	NP-5
Holyoke-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	70-90	50-75	15-25	NP-5
	3-8	Silt loam, loam, fine sandy loam, very fine sandy loam, gravelly silt loam	GM, SM	A-4	0-5	0-10	65-100	55-90	55-80	35-45	15-25	NP-5
	8-18	Gravelly silt loam, fine sandy loam, very fine sandy loam, loam	SM	A-4	0-10	0-15	70-100	60-90	55-85	35-45	15-25	NP-5
	18-80	Unweathered bedrock			---	---	---	---	---	---	---	---
78C: Holyoke-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	70-90	50-75	15-25	NP-5
	3-8	Silt loam, loam, fine sandy loam, very fine sandy loam, gravelly silt loam	SM, GM	A-4	0-5	0-10	65-100	55-90	55-80	35-45	15-25	NP-5
	8-18	Gravelly silt loam, fine sandy loam, very fine sandy loam, loam	SM	A-4	0-10	0-15	70-100	60-90	55-85	35-45	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
78C: Holyoke-----	18-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
78E: Holyoke-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	70-90	50-75	15-25	NP-5
	3-8	Silt loam, loam, fine sandy loam, very fine sandy loam, gravelly silt loam	GM, SM	A-4	0-5	0-10	65-100	55-90	55-80	35-45	15-25	NP-5
	8-18	Gravelly silt loam, fine sandy loam, very fine sandy loam, loam	SM	A-4	0-10	0-15	70-100	60-90	55-85	35-45	15-25	NP-5
	18-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
79E: Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
Holyoke-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Silt loam	ML	A-4	0-5	0-5	90-100	80-90	70-90	50-75	15-25	NP-5
	3-8	Silt loam, loam, fine sandy loam, very fine sandy loam, gravelly silt loam	GM, SM	A-4	0-5	0-10	65-100	55-90	55-80	35-45	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
79E: Holyoke-----	8-18	Gravelly silt loam, fine sandy loam, very fine sandy loam, loam	SM	A-4	0-10	0-15	70-100	60-90	55-85	35-45	15-25	NP-5
	18-80	Unweathered bedrock			---	---	---	---	---	---	---	---
80B: Bernardston-----	0-8	Silt loam	ML	A-4	0	0-5	90-95	80-90	75-90	50-75	20-25	NP-5
	8-14	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-75	20-25	NP-10
	14-24	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10
	24-26	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10
	26-60	Channery silt loam, channery very fine sandy loam, channery loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	65-95	55-90	50-90	35-70	20-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
80C: Bernardston-----	0-8	Silt loam	ML	A-4	0	0-5	90-95	80-90	75-90	50-75	20-25	NP-5
	8-14	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-75	20-25	NP-10
	14-24	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10
	24-26	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10
	26-60	Channery silt loam, channery very fine sandy loam, channery loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	65-95	55-90	50-90	35-70	20-25	NP-10
81C: Bernardston-----	0-8	Silt loam	ML	A-4	0	0-5	90-95	80-90	75-90	50-75	20-25	NP-5
	8-14	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-75	20-25	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
81C: Bernardston-----	In				Pct	Pct						
	14-24	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10
	24-26	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10
	26-60	Channery silt loam, channery very fine sandy loam, channery loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	65-95	55-90	50-90	35-70	20-25	NP-10
81D: Bernardston-----	0-8	Silt loam	ML	A-4	0	0-5	90-95	80-90	75-90	50-75	20-25	NP-5
	8-14	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-75	20-25	NP-10
	14-24	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
81D: Bernardston-----	In											
	24-26	Channery silt loam, channery very fine sandy loam, silt loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-5	75-95	65-90	60-90	40-70	20-25	NP-10
	26-60	Channery silt loam, channery very fine sandy loam, channery loam, very fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	65-95	55-90	50-90	35-70	20-25	NP-10
82B: Broadbrook-----	0-8	Silt loam	ML	A-4	0-5	0-5	90-100	80-100	80-100	55-90	15-25	NP-5
	8-14	Silt loam, very fine sandy loam, gravelly loam	ML, CL-ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	14-25	Silt loam, very fine sandy loam, gravelly loam	SC-SM, ML, CL-ML	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	25-65	Gravelly fine sandy loam, sandy loam, gravelly sandy loam	SC-SM, SM	A-4, A-2	0-5	0-5	75-100	60-95	45-90	25-50	15-25	NP-10
82C: Broadbrook-----	0-8	Silt loam	ML	A-4	0-5	0-5	90-100	80-100	80-100	55-90	15-25	NP-5
	8-14	Silt loam, very fine sandy loam, gravelly loam	ML, CL-ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	14-25	Silt loam, very fine sandy loam, gravelly loam	CL-ML, ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
82C: Broadbrook-----	25-65	Gravelly fine sandy loam, sandy loam, gravelly sandy loam	SC-SM, SM	A-2, A-4	0-5	0-5	75-100	60-95	45-90	25-50	15-25	NP-10
82D: Broadbrook-----	0-8	Silt loam	ML	A-4	0-5	0-5	90-100	80-100	80-100	55-90	15-25	NP-5
	8-14	Silt loam, very fine sandy loam, gravelly loam	ML, CL-ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	14-25	Silt loam, very fine sandy loam, gravelly loam	CL-ML, ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	25-65	Gravelly fine sandy loam, sandy loam, gravelly sandy loam	SC-SM, SM	A-2, A-4	0-5	0-5	75-100	60-95	45-90	25-50	15-25	NP-10
83B: Broadbrook-----	0-8	Silt loam	ML	A-4	0-5	0-5	90-100	80-100	80-100	55-90	15-25	NP-5
	8-14	Silt loam, very fine sandy loam, gravelly loam	ML, CL-ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	14-25	Silt loam, very fine sandy loam, gravelly loam	CL-ML, ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	25-65	Gravelly fine sandy loam, sandy loam, gravelly sandy loam	SC-SM, SM	A-2, A-4	0-5	0-5	75-100	60-95	45-90	25-50	15-25	NP-10
83C: Broadbrook-----	0-8	Silt loam	ML	A-4	0-5	0-5	90-100	80-100	80-100	55-90	15-25	NP-5
	8-14	Silt loam, very fine sandy loam, gravelly loam	ML, CL-ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
83C: Broadbrook-----	14-25	Silt loam, very fine sandy loam, gravelly loam	CL-ML, ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	25-65	Gravelly fine sandy loam, sandy loam, gravelly sandy loam	SC-SM, SM	A-2, A-4	0-5	0-5	75-100	60-95	45-90	25-50	15-25	NP-10
84B: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Montauk-----	0-4	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-95	75-95	35-45	15-25	NP-5
	4-14	Fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	14-25	Sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	25-39	Gravelly loamy coarse sand, loamy sand, gravelly sandy loam, sandy loam	SM, SC-SM	A-1, A-2, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
84B: Montauk-----	39-60	Gravelly sandy loam, sandy loam, gravelly loamy coarse sand, loamy sand	SM, SC-SM	A-1, A-2, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
84C: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Montauk-----	0-4	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-95	75-95	35-45	15-25	NP-5
	4-14	Fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	14-25	Sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	25-39	Gravelly loamy coarse sand, loamy sand, gravelly sandy loam, sandy loam	SM, SC-SM	A-4, A-1, A-2	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
	39-60	Gravelly sandy loam, sandy loam, gravelly loamy coarse sand, loamy sand	SM, SC-SM	A-4, A-1, A-2	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
84D: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Montauk-----	0-4	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-95	75-95	35-45	15-25	NP-5
	4-14	Fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	14-25	Sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-2, A-4	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	25-39	Gravelly loamy coarse sand, loamy sand, gravelly sandy loam, sandy loam	SM, SC-SM	A-4, A-1, A-2	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
	39-60	Gravelly sandy loam, sandy loam, gravelly loamy coarse sand, loamy sand	SM, SC-SM	A-4, A-1, A-2	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
85B: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
				Pct	Pct					Pct		
85B: Paxton-----	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Montauk-----	0-4	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-95	75-95	35-45	15-25	NP-5
	4-14	Fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	14-25	Sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	25-39	Gravelly loamy coarse sand, loamy sand, gravelly sandy loam, sandy loam	SM, SC-SM	A-4, A-1, A-2	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
	39-60	Gravelly sandy loam, sandy loam, gravelly loamy coarse sand, loamy sand	SM, SC-SM	A-1, A-2, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
85C: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
85C: Paxton-----	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
85C: Montauk-----	0-4	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-95	75-95	35-45	15-25	NP-5
	4-14	Fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	14-25	Sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-2, A-4	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	25-39	Gravelly loamy coarse sand, loamy sand, gravelly sandy loam, sandy loam	SM, SC-SM	A-4, A-1, A-2	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
	39-60	Gravelly sandy loam, sandy loam, gravelly loamy coarse sand, loamy sand	SM, SC-SM	A-2, A-1, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
86C: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
86C: Montauk-----	0-4	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-95	75-95	35-45	15-25	NP-5
	4-14	Fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	14-25	Sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	25-39	Gravelly loamy coarse sand, loamy sand, gravelly sandy loam, sandy loam	SM, SC-SM	A-1, A-2, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
	39-60	Gravelly sandy loam, sandy loam, gravelly loamy coarse sand, loamy sand	SM, SC-SM	A-1, A-2, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
86D: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Montauk-----	0-4	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-95	75-95	35-45	15-25	NP-5
	4-14	Fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
86D: Montauk-----	14-25	Sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-20	75-100	65-95	50-95	25-60	15-25	NP-5
	25-39	Gravelly loamy coarse sand, loamy sand, gravelly sandy loam, sandy loam	SM, SC-SM	A-1, A-2, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
	39-60	Gravelly sandy loam, sandy loam, gravelly loamy coarse sand, loamy sand	SM, SC-SM	A-1, A-2, A-4	0-10	0-10	75-95	60-90	30-75	15-50	15-25	NP
87B: Wethersfield---	0-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
87C: Wethersfield---	0-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
87D: Wethersfield----	0-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
88B: Wethersfield----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
88C: Wethersfield----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
88C: Wethersfield----	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
89C: Wethersfield----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
89D: Wethersfield----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
90B: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SC-SM, ML, SM, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	CL-ML, SM, SC-SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
90C: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SC-SM, ML, SM, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	CL-ML, SM, SC-SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
90D: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SC-SM, ML, SM, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	CL-ML, SM, SC-SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
91B: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SC-SM, ML, SM, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	CL-ML, SM, SC-SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
91C: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SC-SM, ML, SM, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	CL-ML, SM, SC-SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
91D: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SC-SM, ML, SM, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	ML, CL-ML, SC-SM, SM	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	ML, CL-ML, SM, SC-SM	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
92B: Nellis-----	0-8	Fine sandy loam	SM	A-4	0	0-5	85-95	80-95	75-95	35-50	15-25	NP-5
	8-14	Fine sandy loam, silt loam, gravelly fine sandy loam, very fine sandy loam, loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	14-25	Fine sandy loam, loam, silt loam, very fine sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	25-27	Loam, silt loam, gravelly fine sandy loam, very fine sandy loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-75	15-25	NP-10
	27-60	Sandy loam, gravelly loam, very gravelly fine sandy loam, loam	CL-ML, GC-GM, GM, ML	A-1, A-2, A-4	0	0-10	50-95	40-95	35-90	20-70	15-25	NP-10
92C: Nellis-----	0-8	Fine sandy loam	SM	A-4	0	0-5	85-95	80-95	75-95	35-50	15-25	NP-5
	8-14	Fine sandy loam, silt loam, gravelly fine sandy loam, very fine sandy loam, loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	14-25	Fine sandy loam, loam, silt loam, very fine sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
92C: Nellis-----	25-27	Loam, silt loam, gravelly fine sandy loam, very fine sandy loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-75	15-25	NP-10
	27-60	Sandy loam, gravelly loam, very gravelly fine sandy loam, loam	CL-ML, GC-GM, GM, ML	A-1, A-2, A-4	0	0-10	50-95	40-95	35-90	20-70	15-25	NP-10
92D: Nellis-----	0-8	Fine sandy loam	SM	A-4	0	0-5	85-95	80-95	75-95	35-50	15-25	NP-5
	8-14	Fine sandy loam, silt loam, gravelly fine sandy loam, very fine sandy loam, loam	ML, SM, CL- ML, SC-SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	14-25	Fine sandy loam, loam, silt loam, very fine sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	25-27	Loam, silt loam, gravelly fine sandy loam, very fine sandy loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-75	15-25	NP-10
	27-60	Sandy loam, gravelly loam, very gravelly fine sandy loam, loam	CL-ML, GC-GM, GM, ML	A-4, A-1, A-2	0	0-10	50-95	40-95	35-90	20-70	15-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
94C: Nellis-----	0-8	Fine sandy loam	SM	A-4	0	0-5	85-95	80-95	75-95	35-50	15-25	NP-5
	8-14	Fine sandy loam, silt loam, gravelly fine sandy loam, very fine sandy loam, loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
94C: Nellis-----	14-25	Fine sandy loam, loam, silt loam, very fine sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	25-27	Loam, silt loam, gravelly fine sandy loam, very fine sandy loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-75	15-25	NP-10
	27-60	Sandy loam, gravelly loam, very gravelly fine sandy loam, loam	CL-ML, GC-GM, GM, ML	A-1, A-2, A-4	0	0-10	50-95	40-95	35-90	20-70	15-25	NP-10
94E: Farmington-----	0-3	Fine sandy loam	SC-SM, SM	A-4	0-5	0-5	90-100	80-90	75-90	35-45	15-25	NP-5
	3-8	Fine sandy loam, loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0-5	0-10	65-100	55-90	50-90	25-70	15-25	NP-10
	8-17	Fine sandy loam, loam, silt loam, gravelly fine sandy loam, very fine sandy loam	ML, SC-SM, SM, CL-ML	A-2, A-4	0-10	0-10	65-100	60-90	50-90	30-80	15-25	NP-10
	17-80	Bedrock			---	---	---	---	---	---	---	---

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
94E: Nellis-----	0-8	Fine sandy loam	SM	A-4	0	0-5	85-95	80-95	75-95	35-50	15-25	NP-5
	8-14	Fine sandy loam, silt loam, gravelly fine sandy loam, very fine sandy loam, loam	ML, SM, CL-ML, SC-SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	14-25	Fine sandy loam, loam, silt loam, very fine sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-80	15-25	NP-10
	25-27	Loam, silt loam, gravelly fine sandy loam, very fine sandy loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-5	65-90	55-90	50-90	20-75	15-25	NP-10
	27-60	Sandy loam, gravelly loam, very gravelly fine sandy loam, loam	CL-ML, GC-GM, GM, ML	A-1, A-2, A-4	0	0-10	50-95	40-95	35-90	20-70	15-25	NP-10
95C: Farmington-----	0-3	Fine sandy loam	SM, SC-SM	A-4	0-5	0-5	90-100	80-90	75-90	35-45	15-25	NP-5
	3-8	Fine sandy loam, loam, very fine sandy loam, silt loam, gravelly fine sandy loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0-5	0-10	65-100	55-90	50-90	25-70	15-25	NP-10
	8-17	Fine sandy loam, loam, silt loam, gravelly fine sandy loam, very fine sandy loam	CL-ML, SC-SM, ML, SM	A-2, A-4	0-10	0-10	65-100	60-90	50-90	30-80	15-25	NP-10
	17-80	Bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
95E: Farmington-----	0-3	Fine sandy loam	SM, SC-SM	A-4	0-5	0-5	90-100	80-90	75-90	35-45	15-25	NP-5
	3-8	Fine sandy loam, loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SC-SM, CL-ML, SM	A-2, A-4	0-5	0-10	65-100	55-90	50-90	25-70	15-25	NP-10
	8-17	Fine sandy loam, loam, silt loam, gravelly fine sandy loam, very fine sandy loam	CL-ML, SC-SM, ML, SM	A-4, A-2	0-10	0-10	65-100	60-90	50-90	30-80	15-25	NP-10
	17-80	Bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
96: Ipswich-----	0-16	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	16-23	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	23-64	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	64-80	Muck	PT	A-8	0	0	100	100	100	100	---	---
97: Pawcatuck-----	0-12	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	12-40	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	40-46	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	46-50	Very fine sandy loam, silt loam, sandy loam	SM, ML	A-2, A-4	0	0	85-100	75-100	60-95	30-80	0-20	NP-5
	50-60	Loamy sand, loamy fine sand, gravelly sand	SM	A-1, A-2, A-4	0	0	70-100	65-100	40-75	10-40	0-15	NP
98: Westbrook-----	0-10	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	10-40	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	40-48	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	48-64	Silt loam, silty clay loam, sandy loam	ML	A-4	0	0	100	100	70-100	50-100	20-35	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
98: Westbrook-----	64-99	Silt loam, silty clay loam, sandy loam	ML	A-4	0	0	100	100	70-100	50-100	20-35	NP-5
99: Westbrook, low salt-----	0-10	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	10-40	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	40-48	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	48-64	Silt loam, silty clay loam, sandy loam	ML	A-4	0	0	100	100	70-100	50-100	20-35	NP-5
	64-99	Silt loam, silty clay loam, sandy loam	ML	A-4	0	0	100	100	70-100	50-100	20-35	NP-5
100: Suncook-----	0-7	Loamy fine sand	SM	A-2	0	0	90-100	80-100	70-95	20-50	0-15	NP
	7-15	Stratified coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0	0	85-100	75-100	10-95	1-50	0-15	NP
	15-22	Stratified coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0	0	85-100	75-100	10-95	1-50	0-15	NP
	22-32	Stratified coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0	0	85-100	75-100	10-95	1-50	0-15	NP
	32-42	Stratified coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0	0	85-100	75-100	10-95	1-50	0-15	NP
	42-65	Stratified gravelly coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0	0-10	70-100	50-100	10-95	1-50	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
101: Occum-----	0-10	Fine sandy loam	SM, ML	A-2, A-4	0	0	80-100	75-100	65-100	30-55	10-20	NP-5
	10-17	Fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0	0	85-100	75-100	50-100	25-60	10-20	NP-5
	17-28	Sandy loam, fine sandy loam	SM	A-4	0	0	85-100	75-100	65-100	40-55	10-20	NP-5
	28-32	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0-15	65-100	30-100	5-95	0-30	0-15	NP
	32-42	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-3, A-1, A-2	0	0-15	65-100	30-100	5-95	0-30	0-15	NP
	42-65	Stratified very gravelly coarse sand to loamy fine sand	SP, SP-SM, SM	A-1, A-2, A-3	0	0-15	65-100	30-100	5-95	0-30	0-15	NP
102: Pootatuck-----	0-4	Fine sandy loam	SM	A-2, A-4	0	0	95-100	75-100	70-90	30-45	0-20	NP-5
	4-16	Fine sandy loam, sandy loam	ML, SM	A-2, A-4	0	0	95-100	75-100	60-95	30-55	0-20	NP-5
	16-21	Fine sandy loam, sandy loam	SM, ML	A-2, A-4	0	0	95-100	75-100	60-95	30-55	0-20	NP-5
	21-29	Sandy loam, fine sandy loam	SM, ML	A-2, A-4	0	0	95-100	75-100	60-95	30-55	0-20	NP-5
	29-35	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0	75-100	45-100	25-95	5-35	0-20	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
				Pct	Pct					Pct		
102: Pootatuck-----	In											
	35-40	Stratified very gravelly coarse sand to loamy fine sand	SP-SM, SM	A-1, A-2, A-3	0	0	75-100	45-100	25-75	5-25	0-20	NP
	40-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0	75-100	45-100	25-75	5-25	0-20	NP
103: Rippowam-----	0-5	Fine sandy loam	SM, ML	A-4, A-2	0	0	95-100	75-100	60-95	30-55	0-15	NP-5
	5-12	Fine sandy loam, sandy loam	SM, ML	A-4, A-2	0	0	95-100	75-100	50-95	25-55	0-15	NP-5
	12-19	Fine sandy loam, sandy loam	SM, ML	A-2, A-4	0	0	95-100	75-100	50-95	25-55	0-15	NP-5
	19-24	Sandy loam, fine sandy loam, loamy fine sand, gravelly loamy fine sand	SM, ML	A-2, A-4	0	0	75-100	45-100	40-95	10-60	0-15	NP-5
	24-27	Sandy loam, fine sandy loam, loamy fine sand, gravelly loamy fine sand	SM, ML	A-2, A-4	0	0	75-100	45-100	40-95	10-60	0-15	NP-5
	27-31	Loamy sand, fine sandy loam, sandy loam, gravelly loamy fine sand	SP-SM, SP, SM	A-3, A-1, A-2	0	0-20	75-100	50-100	5-95	0-30	0-15	NP
	31-65	Stratified very gravelly coarse sand to loamy fine sand	SP-SM, SM, SP	A-1, A-2, A-3	0	0-20	75-100	50-100	5-95	0-30	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
104: Bash-----	0-11	Silt loam	ML	A-4	0	0	95-100	80-100	75-100	55-95	15-25	NP-5
	11-21	Silt loam, gravelly loam, fine sandy loam	ML, SM, CL- ML, SC-SM	A-2, A-4	0	0	75-100	55-100	50-100	20-95	15-25	NP-10
	21-28	Silt loam, gravelly loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0	0	75-100	55-100	50-100	20-95	15-25	NP-10
	28-60	Silt loam, gravelly loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0	0	75-100	55-100	50-100	20-95	10-25	NP-10
105: Hadley-----	0-12	Silt loam	ML	A-4	0	0	100	100	95-100	70-90	15-25	NP-5
	12-29	Stratified very fine sand to silt loam	ML, SM	A-4	0	0	100	100	95-100	45-80	10-20	NP-5
	29-40	Stratified very fine sand to silt loam	ML, SM	A-4	0	0	100	100	95-100	45-80	10-20	NP-5
	40-45	Stratified sand to silt loam	ML, SM	A-2, A-4	0	0	100	100	55-100	15-80	10-20	NP-5
	45-60	Stratified sand to silt loam	ML, SM	A-4, A-2	0	0	100	100	55-100	15-80	10-20	NP-5
106: Winooski-----	0-12	Silt loam	ML	A-4	0	0	100	95-100	90-100	60-95	15-25	NP-5
	12-18	Silt loam, very fine sandy loam, loamy very fine sand	ML, SM	A-4	0	0	100	95-100	90-100	40-95	15-25	NP-5
	18-36	Silt loam, very fine sandy loam, loamy very fine sand	SM, ML	A-4	0	0	100	95-100	90-100	40-95	15-25	NP-5
	36-52	Very fine sandy loam, silt loam, loamy very fine sand	SM, ML	A-4	0	0	100	95-100	90-100	40-95	15-25	NP-5
	52-65	Silt loam, very fine sandy loam, loamy very fine sand	SM, ML	A-4	0	0	100	95-100	90-100	40-95	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
					Pct	Pct						
107: Limerick-----	0-8	Silt loam	ML	A-4	0	0	100	100	90-100	60-90	0-15	NP
	8-20	Silt loam, very fine sandy loam	ML	A-4	0	0	100	100	100	60-90	0-15	NP
	20-36	Silt loam, very fine sandy loam	ML	A-4	0	0	100	100	100	60-90	0-15	NP
	36-54	Silt loam, very fine sandy loam	ML	A-4	0	0	100	100	100	60-90	0-15	NP
	54-65	Silt loam, very fine sandy loam	ML	A-4	0	0	100	100	90-100	55-85	0-15	NP
Lim-----	0-6	Very fine sandy loam	SM, ML	A-4	0	0	100	90-100	85-100	45-65	10-20	NP-5
	6-11	Very fine sandy loam, silt loam, loam	ML	A-4	0	0	100	90-100	85-100	55-95	10-25	NP-10
	11-15	Very fine sandy loam, silt loam, loam	ML	A-4	0	0	100	90-100	85-100	55-95	10-25	NP-10
	15-22	Silt loam, fine sandy loam, sandy loam	ML	A-4	0	0	100	90-100	85-100	55-95	10-25	NP-10
	22-29	Fine sandy loam, silt loam, sandy loam	SM	A-2, A-4	0	0	90-100	75-100	50-95	20-50	10-20	NP-5
	29-42	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0	75-100	35-100	20-80	5-35	0-15	NP
	42-50	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0	75-100	35-100	20-80	5-35	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
107: Lim-----	50-57	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0	75-100	35-100	20-80	5-35	0-15	NP
	57-65	Stratified very gravelly coarse sand to loamy sand	SM, SP-SM	A-1, A-2, A-3	0	0	75-100	35-100	20-80	5-35	0-15	NP
108: Saco-----	0-12	Silt loam	ML	A-4	0	0	100	90-100	80-100	60-90	15-25	NP-5
	12-32	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	50-90	15-25	NP-5
	32-48	Silt loam, very fine sandy loam	ML	A-4	0	0	100	90-100	80-100	50-90	15-25	NP-5
	48-60	Stratified very gravelly coarse sand to loamy fine sand	SM, SP-SM, SW-SM	A-1, A-2, A-3	0	0-10	70-100	50-100	20-95	5-25	15-20	NP-5
109: Fluvaquents, Frequently Flooded-----	0-4	Silt loam	ML	A-4	0	0	85-100	80-100	70-100	50-85	10-20	NP
	4-14	Very gravelly coarse sand, fine sand, silt loam, gravelly sandy loam	GP-GM, SP-SM, ML, SM	A-3, A-2, A- 1, A-4	0	0-15	50-100	30-100	15-95	5-75	10-20	NP
	14-21	Very gravelly coarse sand, gravelly sandy loam, silt loam, very fine sand	GP-GM, SM, SP-SM, ML	A-1, A-2, A- 3, A-4	0	0-15	50-100	30-100	15-95	5-75	10-20	NP
	21-38	Very gravelly loamy sand, gravelly sandy loam, silt loam	GP-GM, ML, SP-SM, SM	A-2, A-1, A-4	0	0-15	50-100	30-100	20-100	10-85	10-20	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
109: Fluvaquents, Frequently Flooded-----	38-45	Fine sandy loam, silt loam, very gravelly loamy sand, gravelly sandy loam	GP-GM, ML, SP-SM, SM	A-1, A-2, A-4	0	0-15	50-100	30-100	20-100	10-85	10-20	NP
	45-55	Very gravelly coarse sand, gravelly sandy loam, silt loam, sand	GP-GM, SM, SP-SM, ML	A-1, A-2, A-3, A-4	0	0-15	50-100	30-100	15-95	5-75	10-20	NP
	55-60	Very gravelly loamy sand, fine sandy loam, silt loam, gravelly sandy loam	GP-GM, ML, SP-SM, SM	A-1, A-2, A-4	0	0-15	50-100	30-100	20-100	10-85	10-20	NP
Udifuvents, Frequently Flooded-----	0-2	Fine sandy loam	ML, SM	A-2, A-4	0	0	85-100	80-100	70-100	25-55	10-20	NP
	2-4	Very gravelly coarse sand, silt loam, loamy fine sand	ML, SM, SP-SM, GP-GM	A-2, A-4, A-3, A-1	0	0-15	50-100	30-100	15-95	5-75	10-20	NP
	4-12	Fine sandy loam, very gravelly loamy sand, silt loam	SP-SM, ML, SM, GP-GM	A-2, A-4, A-1	0	0-15	50-100	30-100	20-100	10-85	10-20	NP
	12-18	Silt loam, fine sandy loam, very gravelly loamy sand	SP-SM, ML, SM, GP-GM	A-1, A-2, A-4	0	0-15	50-100	30-100	20-100	10-85	10-20	NP
	18-35	Silt loam, very gravelly coarse sand, loamy sand	SP-SM, ML, SM, GP-GM	A-1, A-3, A-2, A-4	0	0-15	50-100	30-100	15-95	5-75	10-20	NP
	35-38	Very gravelly loamy sand, very gravelly coarse sand, silt loam	SP-SM, ML, SM, GP-GM	A-1, A-3, A-2, A-4	0	0-15	50-100	30-100	15-95	5-75	10-20	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
109: Udifulvents, Frequently Flooded-----	38-60	Silt loam, loamy sand, very gravelly coarse sand	SP-SM, ML, SM, GP-GM	A-1, A-3, A- 2, A-4	0	0-15	50-100	30-100	15-95	5-75	10-20	NP
221A: Ninigret-----	0-8	Fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	70-100	60-100	25-55	0-35	NP-5
	8-16	Fine sandy loam, very fine sandy loam, silt loam	ML, SM	A-2, A-4	0	0	95-100	75-100	70-100	30-80	0-25	NP-5
	16-26	Fine sandy loam, very fine sandy loam, silt loam	ML, SM	A-2, A-4	0	0	95-100	75-100	70-100	30-80	0-25	NP-5
	26-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0-10	0-15	70-100	35-100	5-95	1-30	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
224A: Deerfield-----	0-8	Loamy fine sand	SM	A-2	0	0	100	85-100	70-90	20-30	0-20	NP
	8-16	Loamy sand, fine sand, loamy fine sand	SM	A-2	0	0	100	85-100	55-90	15-25	0-20	NP
	16-28	Loamy sand, fine sand, loamy fine sand	SM	A-2	0	0	100	85-100	55-90	15-25	0-20	NP
	28-34	Fine sand, sand, loamy sand	SM	A-2	0	0	100	85-100	70-80	15-25	0-15	NP
	34-60	Fine sand, loamy sand, coarse sand	SM, SP, SP-SM	A-1, A-2	0	0	100	80-100	20-85	2-30	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
225B: Brancroft-----	0-6	Silt loam	ML	A-4	0	0	100	95-100	90-100	70-95	25-35	NP-5
	6-17	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	17-22	Silty clay loam, silt loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
	22-32	Silt loam, silty clay loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	32-43	Silty clay loam, silt loam	CL, CL-ML	A-6, A-4	0	0	100	95-100	90-100	85-95	25-40	5-15
	43-66	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	90-100	85-95	25-40	5-15
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
226B: Berlin-----	0-6	Silt loam	ML	A-4	0	0	95-100	85-100	80-100	60-95	25-35	NP-5
	6-12	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	95-100	85-100	80-100	75-95	25-40	5-15
	12-20	Silty clay loam, silt loam	CL, CL-ML	A-4, A-6	0	0	95-100	85-100	80-100	75-95	25-40	5-15
	20-34	Silty clay loam, silt loam, silty clay	CL, CL-ML	A-4, A-6, A- 7, A-5	0	0	95-100	85-100	85-100	80-95	25-45	5-20
	34-48	Silty clay loam, silt loam, silty clay	CL, CL-ML	A-6, A-7, A- 4, A-5	0	0	95-100	85-100	85-100	80-95	25-45	5-20
	48-65	Silty clay loam, silt loam, silty clay	CL-ML, CL	A-4, A-6, A- 7, A-5	0	0	95-100	85-100	85-100	80-95	25-45	5-20
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
228B: Elmridge-----	0-6	Fine sandy loam	SM, ML	A-4	0	0	100	90-100	85-95	35-55	15-20	NP
	6-10	Fine sandy loam, sandy loam, loam	CL-ML, SM, ML	A-4	0	0	100	90-100	65-100	45-65	15-20	NP-5
	10-18	Fine sandy loam, sandy loam, loam	SM, CL-ML, ML	A-4	0	0	100	90-100	65-90	45-65	15-20	NP-5
	18-25	Sandy loam, fine sandy loam, loam	SM, CL-ML, ML	A-4	0	0	100	90-100	65-90	45-65	15-20	NP-5
	25-65	Silty clay, silty clay loam, clay	CL, CH	A-7	0	0	100	95-100	95-100	85-95	40-55	15-30
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
229B: Agawam-----	0-8	Fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5
	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
229C: Agawam-----	0-8	Fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
229C: Agawam-----	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
230B: Branford-----	0-8	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-85	15-25	NP-5
	8-18	Loam, silt loam, very fine sandy loam, gravelly fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	35-85	15-25	NP-5
	18-24	Gravelly loam, gravelly fine sandy loam, loam, silt loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	75-100	55-100	55-100	25-85	15-25	NP-5
	24-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-3, A-1, A-2	0	0-10	75-90	35-85	10-75	0-30	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
230C: Branford-----	0-8	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-85	15-25	NP-5
	8-18	Loam, silt loam, very fine sandy loam, gravelly fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	75-100	35-85	15-25	NP-5
	18-24	Gravelly loam, gravelly fine sandy loam, loam, silt loam, very fine sandy loam	ML, SM	A-4, A-2	0	0	75-100	55-100	55-100	25-85	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
230C: Branford-----	24-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0-10	75-90	35-85	10-75	0-30	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
232B: Haven-----	0-7	Silt loam	ML	A-4	0	0	90-100	75-100	70-100	50-100	15-25	NP-5
	7-14	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	14-20	Silt loam, very fine sandy loam	ML, SM	A-4	0	0	95-100	80-100	70-100	40-100	15-25	NP-5
	20-24	Fine sandy loam, very fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	75-100	70-95	30-65	15-25	NP-5
	24-60	Stratified very gravelly sand to gravelly fine sand	SM, SW-SM, SW	A-1, A-2, A-3	0	0-15	75-90	30-75	10-70	0-15	10-20	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
234B: Merrimac-----	0-9	Sandy loam	ML, SM	A-2, A-4	0	0	90-100	70-90	45-85	25-60	0-20	NP
	9-16	Sandy loam, fine sandy loam	SM	A-2, A-4	0	0	90-100	75-90	45-90	25-50	0-25	NP
	16-24	Gravelly sandy loam, sandy loam	SM	A-2, A-4	0	0	80-100	55-90	35-70	15-50	0-25	NP
	24-60	Stratified very gravelly coarse sand to gravelly sand	SP, SP-SM	A-1	0	10-20	65-85	35-60	35-60	1-15	0-25	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
237C: Manchester-----	0-9	Gravelly sandy loam	SM	A-2, A-4	0	0	75-85	50-70	35-70	20-45	0-15	NP
	9-18	Gravelly loamy sand, very gravelly loamy sand, very gravelly sand	SM, SP-SM	A-2, A-3, A-1	0	0-10	60-85	35-75	20-60	5-25	0-15	NP
	18-65	Stratified extremely gravelly coarse sand to very gravelly loamy sand	GM, SM, SW-SM	A-1	0	0-15	45-60	20-50	10-40	0-20	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
238A: Hinckley-----	0-8	Gravelly sandy loam	SM	A-2, A-4	0-5	1-15	65-85	60-75	45-70	25-50	0-20	NP
	8-20	Very gravelly loamy sand, gravelly loamy sand, loamy fine sand, very gravelly loamy coarse sand	SM, GM	A-1, A-2	0-10	1-15	50-100	40-90	25-90	15-25	0-20	NP
	20-27	Very gravelly sand, gravelly sand	SM, GM	A-1	1-10	5-20	50-90	40-80	25-50	5-15	0-20	NP
	27-42	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
	42-60	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
238C: Hinckley-----	0-8	Gravelly sandy loam	SM	A-2, A-4	0-5	1-15	65-85	60-75	45-70	25-50	0-20	NP
	8-20	Very gravelly loamy sand, gravelly loamy sand, loamy fine sand, very gravelly loamy coarse sand	SM, GM	A-1, A-2	0-10	1-15	50-100	40-90	25-90	15-25	0-20	NP
	20-27	Very gravelly sand, gravelly sand	SM, GM	A-1	1-10	5-20	50-90	40-80	25-50	5-15	0-20	NP
	27-42	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
	42-60	Stratified cobbly coarse sand to extremely gravelly sand	GM, GP-GM, SM, SP-SM	A-1	1-10	15-35	30-95	20-85	15-50	5-15	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
240B: Ludlow-----	0-8	Silt loam	ML	A-4	0	0	85-100	75-90	70-90	50-80	15-25	NP-5
	8-20	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	20-26	Silt loam, gravelly loam, fine sandy loam	SM, ML	A-4, A-2	0	0	70-100	65-90	60-90	30-75	15-25	NP-5
	26-65	Gravelly loam, gravelly silt loam, silt loam, fine sandy loam	ML, SM	A-4, A-2	0-5	0-15	70-100	65-90	60-90	30-75	15-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
243B: Rainbow-----	0-6	Silt loam	ML	A-4	0	0-5	90-100	80-100	70-100	50-90	15-25	NP-5
	6-18	Silt loam, very fine sandy loam, gravelly loam	SM, ML	A-4, A-2	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	18-26	Silt loam, very fine sandy loam, gravelly loam	ML, SM	A-2, A-4	0-5	0-10	85-100	75-100	70-100	30-90	15-25	NP-5
	26-65	Gravelly fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-5	0-10	75-100	60-90	45-80	20-60	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
245B: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	30-43	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
245C: Woodbridge-----	0-7	Fine sandy loam	SM	A-4	0-5	0-5	90-100	80-90	70-90	35-45	10-25	NP-5
	7-18	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	18-26	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	26-30	Fine sandy loam, gravelly fine sandy loam, loam, sandy loam	ML, SM	A-4, A-2	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	30-43	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
	43-65	Gravelly fine sandy loam, fine sandy loam, loam, sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-100	65-90	45-90	20-50	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
248B: Georgia-----	0-8	Silt loam	ML	A-4	0	0-5	90-100	80-90	75-90	60-80	20-30	NP-5
	8-14	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	14-24	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
	24-60	Gravelly fine sandy loam, gravelly loam, loam, silt loam	ML, SM	A-2, A-4	0-5	0-5	70-100	60-90	55-90	30-85	20-30	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
250B: Sutton-----	0-6	Fine sandy loam	SC-SM, SM	A-4, A-2	0	0-5	90-100	80-90	70-90	30-45	15-25	NP-5
	6-12	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, CL-ML, SC-SM, SM	A-2, A-4	0	0-5	60-100	55-90	40-90	20-55	15-25	NP-5
	12-24	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	SM, CL-ML, SC-SM, ML	A-4, A-2	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	24-28	Fine sandy loam, gravelly fine sandy loam, sandy loam, loam	ML, SM, CL-ML, SC-SM	A-4, A-2	0-5	0-5	65-100	55-90	40-90	20-55	15-25	NP-5
	28-36	Gravelly fine sandy loam, gravelly sandy loam, sandy loam	SC-SM, SM	A-2	0-5	0-10	65-100	60-90	55-85	30-40	15-20	NP-5
	36-65	Gravelly sandy loam, gravelly fine sandy loam, sandy loam	SM, SC-SM	A-4, A-2	0-5	0-10	65-100	60-90	45-85	25-50	15-20	NP-5
	Urban land-----	0-6	Material			---	---	---	---	---	---	---
253B: Wapping-----	0-11	Very fine sandy loam	ML, SM	A-4	0	0	95-100	75-100	70-100	40-75	15-25	NP-5
	11-16	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	16-20	Very fine sandy loam, loam, silt loam	ML, SM	A-4	0	0	95-100	75-100	70-95	40-80	15-25	NP-5
	20-28	Gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-85	55-75	40-70	20-35	15-25	NP-5
	28-36	Gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM	A-1, A-2	0-5	0-5	70-85	55-75	35-65	15-20	0-20	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
250B: Urban land-----	36-80	Gravelly loamy sand, very gravelly loamy sand, gravelly sandy loam, gravelly fine sandy loam	SM, SW-SM	A-1, A-2	0-10	0-15	60-85	40-75	30-65	10-20	0-20	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
255B: Watchaug-----	0-8	Fine sandy loam	SM	A-4	0	0-10	90-100	80-90	75-85	35-45	15-25	NP
	8-18	Fine sandy loam, silt loam, gravelly loam	SM, ML	A-2, A-4	0	0-10	65-100	55-90	45-90	25-70	15-25	NP-5
	18-24	Fine sandy loam, silt loam, gravelly loam	ML, SM	A-2, A-4	0-10	0-15	70-100	60-90	55-90	25-70	15-25	NP-5
	24-65	Gravelly sandy loam, gravelly fine sandy loam, fine sandy loam	SM	A-4, A-2	0-10	0-15	70-100	65-90	45-80	30-40	15-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
260B: Charlton-----	0-4	Fine sandy loam	SM	A-4, A-2	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
260B: Charlton-----	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-4, A-2	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
260C: Charlton-----	0-4	Fine sandy loam	SM	A-4, A-2	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
260D: Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
266B: Narragansett----	0-6	Silt loam	ML	A-4	1-5	1-5	90-100	80-100	75-100	55-95	15-25	NP
	6-15	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	60-100	35-95	15-25	NP-5
	15-24	Silt loam, very fine sandy loam, gravelly loam, gravelly silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	24-28	Gravelly silt loam, very fine sandy loam, gravelly loam, silt loam	ML	A-4	1-5	1-5	85-100	65-100	65-100	35-95	15-25	NP-5
	28-60	Very gravelly loamy coarse sand, gravelly loamy sand, gravelly sand	SM, SP-SM	A-1, A-2	1-15	1-15	70-95	45-85	30-85	10-35	0-15	NP
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
269B: Yalesville-----	0-8	Fine sandy loam	SM, ML	A-4, A-2	0	0-5	90-95	80-95	75-95	30-55	15-25	NP-5
	8-14	Fine sandy loam, loam, sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5
	14-25	Loam, fine sandy loam, sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5
	25-36	Gravelly sandy loam, loam, very gravelly fine sandy loam	GM, ML, SM	A-1, A-2, A-4	0-10	0-10	50-95	40-90	25-90	15-60	15-25	NP-5
	36-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
269C: Yalesville-----	0-8	Fine sandy loam	SM, ML	A-4, A-2	0	0-5	90-95	80-95	75-95	30-55	15-25	NP-5
	8-14	Fine sandy loam, loam, sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5
	14-25	Loam, fine sandy loam, sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4, A-1	0-5	0-5	65-95	55-95	35-95	20-65	15-25	NP-5
	25-36	Gravelly sandy loam, loam, very gravelly fine sandy loam	GM, ML, SM	A-1, A-2, A-4	0-10	0-10	50-95	40-90	25-90	15-60	15-25	NP-5
	36-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
273C: Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
273C: Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5
	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10
	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10
	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
273E: Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
Charlton-----	0-4	Fine sandy loam	SM	A-2, A-4	0-5	0-5	95-100	80-90	80-85	30-45	15-25	NP-5
	4-7	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	7-19	Fine sandy loam, gravelly fine sandy loam, sandy loam	SM	A-4, A-2	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	19-27	Gravelly fine sandy loam, fine sandy loam, sandy loam	SM	A-2, A-4	0-5	0-10	75-100	60-90	55-80	25-45	15-25	NP-5
	27-65	Gravelly fine sandy loam, gravelly sandy loam	SM	A-2, A-4	0-5	0-10	75-85	60-70	50-70	20-40	10-25	NP-5
Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
273E: Chatfield-----	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10
	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10
	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
275C: Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-4, A-2	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5
	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10
	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-2, A-4	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10
	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
275E: Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
Chatfield-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	NP-5
	1-6	Gravelly fine sandy loam	SM	A-2, A-4	0-5	0-5	70-85	55-75	45-75	15-40	10-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
275E: Chatfield-----	6-15	Loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0	0-10	75-100	60-90	45-90	15-60	10-25	NP-10
	15-29	Loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	SM	A-4, A-2	0	0-10	70-100	60-90	40-90	15-50	10-25	NP-10
	29-80	Unweathered bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
282B: Broadbrook-----	0-8	Silt loam	ML	A-4	0-5	0-5	90-100	80-100	80-100	55-90	15-25	NP-5
	8-14	Silt loam, very fine sandy loam, gravelly loam	ML, CL-ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	14-25	Silt loam, very fine sandy loam, gravelly loam	CL-ML, ML, SC-SM	A-4	0-5	0-5	85-100	75-100	65-100	35-90	15-25	NP-10
	25-65	Gravelly fine sandy loam, sandy loam, gravelly sandy loam	SC-SM, SM	A-2, A-4	0-5	0-5	75-100	60-95	45-90	25-50	15-25	NP-10
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
284B: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
284B: Paxton-----	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
284C: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
284D: Paxton-----	0-8	Fine sandy loam	SM	A-4	0-5	0-10	90-95	85-90	75-90	35-50	10-25	NP-5
	8-15	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	15-26	Fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
	26-65	Gravelly fine sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SM	A-2, A-4	0-10	0-20	75-90	65-90	55-85	25-60	10-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
287B: Wethersfield----	0-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
287C: Wethersfield----	0-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
287D: Wethersfield----	0-3	Loam	ML	A-4	0-5	0-5	90-100	80-90	75-90	50-70	15-25	NP-5
	3-13	Loam, gravelly loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5
	13-27	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	85-100	65-90	60-90	25-70	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
287D: Wethersfield----	27-65	Gravelly loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	0-15	80-90	65-85	55-85	25-65	15-25	NP-5
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
290B: Stockbridge-----	0-10	Loam	ML, SM	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SM, CL-ML, SC-SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	SM, SC-SM, ML, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	ML, CL-ML, SM, SC-SM	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	ML, CL-ML, SM, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
290C: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	SC-SM, ML, SM, CL-ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	ML, CL-ML, SM, SC-SM	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	SM, ML, SC- SM, CL-ML	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
					Pct	Pct					Pct	
290C: Stockbridge-----	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
290D: Stockbridge-----	0-10	Loam	SM, ML	A-4	0	0-5	85-90	75-90	60-90	45-65	20-30	NP-5
	10-20	Loam, silt loam, gravelly loam	CL-ML, SM, ML, SC-SM	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	20-28	Loam, silt loam, gravelly loam	CL-ML, SC-SM, SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-80	20-30	NP-10
	28-42	Gravelly loam, loam, silt loam	CL-ML, SM, SC-SM, ML	A-4, A-2	0	0-5	60-90	55-90	45-90	30-75	20-30	NP-10
	42-48	Gravelly loam, silt loam, very gravelly fine sandy loam	CL-ML, SM, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
	48-65	Gravelly loam, silt loam, very gravelly fine sandy loam	SM, CL-ML, ML, SC-SM	A-2, A-4	0	0-10	45-90	40-90	35-90	20-75	20-30	NP-10
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
301: Beaches-----	0-65	Gravelly sand	GP, SP	A-1, A-3	---	0-10	45-100	35-100	25-85	0-5	0-14	NP
Udipsamments----	0-38	Sand	SW-SM, SM	A-2	0	0	95-100	75-100	50-70	10-20	0-15	NP
	38-50	Coarse sand, fine sand, gravelly sand	SW-SM, SM	A-1, A-2, A-3	0	0	75-100	50-100	15-95	1-20	0-15	NP
	50-65	Sand, fine sand, coarse sand, gravelly sand	SW-SM, SM	A-1, A-2, A-3	0	0	75-100	50-100	15-95	1-20	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
302: Dumps-----	0-65	Variable			---	---	---	---	---	---	---	---
303: Pits, quarries--	0-1	Unweathered bedrock			---	---	---	---	---	---	---	---
304: Udorthents-----	0-5	Loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	90-100	80-100	70-100	45-75	15-25	NP-10
	5-21	Gravelly loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GC-GM, GM	A-3, A-2, A-1, A-4	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
	21-80	Very gravelly sandy loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GM, GC-GM	A-1, A-2, A-3, A-4	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
305: Udorthents-----	0-5	Loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	90-100	80-100	70-100	45-75	15-25	NP-10
	5-21	Gravelly loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GC-GM, GM	A-3, A-2, A-1, A-4	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
	21-80	Very gravelly sandy loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GM, GC-GM	A-4, A-1, A-2, A-3	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
Pits-----	0-65	Very gravelly sand	GW	A-1	0-5	0-20	25-65	20-50	10-30	0-10	0-15	NP

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
306: Udorthents-----	0-5	Loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	90-100	80-100	70-100	45-75	15-25	NP-10
	5-21	Gravelly loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GC-GM, GM	A-3, A-2, A- 1, A-4	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
	21-80	Very gravelly sandy loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GM, GC-GM	A-4, A-1, A- 2, A-3	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
307: Urban land-----	0-6	Material			---	---	---	---	---	---	---	---
308: Udorthents-----	0-5	Loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	90-100	80-100	70-100	45-75	15-25	NP-10
	5-21	Gravelly loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GC-GM, GM	A-2, A-1, A- 4, A-3	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
	21-80	Very gravelly sandy loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GM, GC-GM	A-2, A-3, A- 4, A-1	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
309: Udorthents-----	0-5	Loam	CL-ML, ML, SC-SM, SM	A-4	0	0-10	90-100	80-100	70-100	45-75	15-25	NP-10
	5-21	Gravelly loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GC-GM, GM	A-2, A-1, A- 4, A-3	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
309: Udorthents-----	21-80	Very gravelly sandy loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GM, GC-GM	A-2, A-3, A-4, A-1	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
310: Udorthents, Periodically Flooded-----	0-5	Loam	SM, SC-SM, ML, CL-ML	A-4	0	0-10	90-100	80-100	70-100	45-75	15-25	NP-10
	5-21	Gravelly loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GC-GM, GM	A-3, A-2, A-1, A-4	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
	21-80	Very gravelly sandy loam, extremely gravelly coarse sand, silty clay loam	SM, SC-SM, ML, CL-ML, GM, GC-GM	A-2, A-3, A-4, A-1	0-20	0-25	45-100	30-100	10-100	5-95	15-30	NP-10
401C: Macomber-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Very channery loam	GM	A-2, A-4, A-1	0	10-20	50-60	40-55	35-50	20-40	15-25	NP-5
	2-10	Very channery loam, very channery silt loam, channery loam, channery silt loam	GM	A-2, A-4, A-1	0	10-20	40-70	30-60	25-60	15-45	15-25	NP-5
	10-21	Very channery loam, very channery silt loam, channery loam, channery silt loam	GM	A-2, A-4, A-1	0	10-20	40-70	30-60	25-60	15-45	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
401C: Macomber-----	21-30	Very channery loam, very channery silt loam, extremely channery silt loam, extremely channery loam	GM	A-2, A-1	0	10-20	35-60	25-50	20-45	15-35	15-25	NP-5
Taconic-----	30-80	Bedrock			---	---	---	---	---	---	---	---
	0-1	Moderately decomposed plant material	OL		0	0	100	100	100	100	---	---
	1-4	Very gravelly loam	GM, GC-GM	A-4, A-1, A-2	10-15	10-15	55-65	45-60	35-55	20-40	20-25	NP-5
	4-11	Very gravelly loam, very gravelly silt loam, gravelly silt loam	GM, GC-GM	A-2, A-4, A-1	10-15	10-15	40-75	35-65	25-65	15-50	20-25	NP-5
	11-80	Bedrock			---	---	---	---	---	---	---	---
402D: Macomber-----	0-1	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Very channery loam	GM	A-2, A-4, A-1	0	10-20	50-60	40-55	35-50	20-40	15-25	NP-5
	2-10	Very channery loam, very channery silt loam, channery loam, channery silt loam	GM	A-2, A-4, A-1	0	10-20	40-70	30-60	25-60	15-45	15-25	NP-5
	10-21	Very channery loam, very channery silt loam, channery loam, channery silt loam	GM	A-2, A-4, A-1	0	10-20	40-70	30-60	25-60	15-45	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
402D: Macomber-----	21-30	Very channery loam, very channery silt loam, extremely channery silt loam, extremely channery loam	GM	A-2, A-1	0	10-20	35-60	25-50	20-45	15-35	15-25	NP-5
	30-80	Bedrock			---	---	---	---	---	---	---	---
Taconic-----	0-1	Moderately decomposed plant material	OL		0	0	100	100	100	100	---	---
	1-4	Very gravelly loam	GM, GC-GM	A-2, A-4, A-1	10-15	10-15	55-65	45-60	35-55	20-40	20-25	NP-5
	4-11	Very gravelly loam, very gravelly silt loam, gravelly silt loam	GC-GM, GM	A-2, A-4, A-1	10-15	10-15	40-75	35-65	25-65	15-50	20-25	NP-5
	11-80	Bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
403C: Taconic-----	0-1	Moderately decomposed plant material	OL		0	0	100	100	100	100	---	---
	1-4	Very gravelly loam	GM, GC-GM	A-4, A-1, A-2	10-15	10-15	55-65	45-60	35-55	20-40	20-25	NP-5
	4-11	Very gravelly loam, very gravelly silt loam, gravelly silt loam	GM, GC-GM	A-2, A-4, A-1	10-15	10-15	40-75	35-65	25-65	15-50	20-25	NP-5
	11-80	Bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
403E: Taconic-----	0-1	Moderately decomposed plant material	OL		0	0	100	100	100	100	---	---
	1-4	Very gravelly loam	GM, GC-GM	A-2, A-4, A-1	10-15	10-15	55-65	45-60	35-55	20-40	20-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
403E: Taconic-----	4-11	Very gravelly loam, very gravelly silt loam, gravelly silt loam	GC-GM, GM	A-2, A-4, A-1	10-15	10-15	40-75	35-65	25-65	15-50	20-25	NP-5
	11-80	Bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
403F: Taconic-----	0-1	Moderately decomposed plant material	OL		0	0	100	100	100	100	---	---
	1-4	Very gravelly loam	GM, GC-GM	A-2, A-1, A-4	10-15	10-15	55-65	45-60	35-55	20-40	20-25	NP-5
	4-11	Very gravelly loam, very gravelly silt loam, gravelly silt loam	GM, GC-GM	A-4, A-1, A-2	10-15	10-15	40-75	35-65	25-65	15-50	20-25	NP-5
	11-80	Bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
405C: Dummerston-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Gravelly loam	SM	A-2, A-4	0-5	0-10	65-85	60-75	45-75	25-50	15-25	NP-5
	2-3	Gravelly silt loam, gravelly loam	ML, SM	A-4, A-2	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	3-4	Gravelly loam, gravelly silt loam	SM, ML	A-2, A-4	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	4-6	Gravelly loam, gravelly silt loam	ML, SM	A-2, A-4	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	6-11	Gravelly loam, gravelly silt loam	SM, ML	A-4, A-2	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	11-22	Gravelly loam, gravelly silt loam	SM, ML	A-4, A-2	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
405C: Dummerston-----	22-27	Gravelly loam, gravelly silt loam, very gravelly loam, very gravelly silt loam	ML, SM	A-4, A-2	0-10	0-20	65-85	55-75	45-75	25-60	15-25	NP-5
	27-40	Very gravelly loam, gravelly fine sandy loam, gravelly loam, very stony fine sandy loam	SM	A-4	0-30	0-15	80-85	70-75	65-70	35-50	15-25	NP-5
	40-64	Very stony loam, gravelly loam, very gravelly fine sandy loam, very stony fine sandy loam	SM	A-4	0-30	0-10	80-85	70-75	65-70	35-50	15-25	NP-5
405E: Dummerston-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Gravelly loam	SM	A-4, A-2	0-5	0-10	65-85	60-75	45-75	25-50	15-25	NP-5
	2-3	Gravelly loam, gravelly silt loam	ML, SM	A-4, A-2	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	3-4	Gravelly loam, gravelly silt loam	SM, ML	A-2, A-4	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	4-6	Gravelly loam, gravelly silt loam	ML, SM	A-4, A-2	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	6-11	Gravelly loam, gravelly silt loam	SM, ML	A-2, A-4	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5
	11-22	Gravelly loam, gravelly silt loam	SM, ML	A-2, A-4	0-5	0-10	65-85	60-75	45-75	25-60	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
405E: Dummerston-----	In											
	22-27	Gravelly loam, gravelly silt loam, very gravelly loam, very gravelly silt loam	ML, SM	A-2, A-4	0-10	0-20	65-85	55-75	45-75	25-60	15-25	NP-5
	27-40	Very gravelly loam, gravelly fine sandy loam, gravelly loam, very stony fine sandy loam	SM	A-4	0-30	0-15	80-85	70-75	65-70	35-50	15-25	NP-5
	40-64	Very stony loam, gravelly loam, very gravelly fine sandy loam, very stony fine sandy loam	SM	A-4	0-30	0-10	80-85	70-75	65-70	35-50	15-25	NP-5
407C: Lanesboro-----	0-3	Highly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	3-6	Loam	ML, SM	A-4	0-5	0-5	90-100	80-100	70-90	40-60	15-25	NP-5
	6-8	Loam, silt loam, channery loam, channery silt loam	ML, SM	A-2, A-4	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5
	8-16	Channery loam, silt loam, loam, channery very fine sandy loam	ML, SM	A-4, A-2	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5
	16-22	Channery loam, silt loam, loam, channery very fine sandy loam	ML, SM	A-4, A-2	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
407C: Lanesboro-----	22-30	Channery loam, silt loam, loam, channery very fine sandy loam	ML, SM	A-2, A-4	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5
	30-60	Very channery loam, channery loam, channery very fine sandy loam, very channery very fine sandy loam	ML, SM	A-4, A-2	5-15	10-30	65-95	55-90	50-85	25-60	15-25	NP-5
407E: Lanesboro-----	0-3	Highly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	3-6	Loam	ML, SM	A-4	0-5	0-5	90-100	80-100	70-90	40-60	15-25	NP-5
	6-8	Loam, silt loam, channery loam, channery silt loam	ML, SM	A-2, A-4	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5
	8-16	Channery loam, silt loam, loam, channery very fine sandy loam	ML, SM	A-4, A-2	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5
	16-22	Channery loam, silt loam, loam, channery very fine sandy loam	ML, SM	A-2, A-4	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5
	22-30	Channery loam, silt loam, loam, channery very fine sandy loam	ML, SM	A-4, A-2	0-15	0-30	65-90	55-85	50-80	25-65	15-25	NP-5
	30-60	Very channery loam, channery loam, channery very fine sandy loam, very channery very fine sandy loam	ML, SM	A-2, A-4	5-15	10-30	65-95	55-90	50-85	25-60	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
408C: Fullam-----	In											
	0-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-4	Silt loam	ML	A-4	0	0-5	90-100	80-100	70-95	50-80	15-25	NP-5
	4-10	Silt loam, loam, gravelly loam, gravelly silt loam	ML, SM	A-4	0-10	0-10	75-90	65-90	55-90	35-75	15-25	NP-5
	10-20	Gravelly loam, gravelly silt loam	SM, ML	A-4	0-10	0-10	75-85	65-75	55-75	35-60	15-25	NP-5
	20-49	Very channery loam, channery loam, channery fine sandy loam, very channery fine sandy loam	ML, SM	A-4	5-15	10-30	80-95	75-90	65-85	35-60	15-25	NP-5
	49-60	Very channery fine sandy loam, very channery loam, channery fine sandy loam, channery loam	ML, SM	A-4	5-15	10-30	80-95	75-90	65-85	35-60	15-25	NP-5
409B: Brayton-----	0-3	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	3-6	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	6-7	Mucky silt loam	ML, SM	A-4	0	0-5	90-95	80-90	70-90	45-70	15-25	NP-5
	7-9	Silt loam, loam, gravelly loam	ML, SM	A-4	0-10	0-15	70-95	60-90	50-90	35-65	15-25	NP-5
	9-13	Gravelly loam, loam, fine sandy loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-10	0-15	70-95	60-90	60-90	30-60	15-25	NP-5
	13-18	Gravelly fine sandy loam, gravelly loam, fine sandy loam, loam	ML, SM	A-2, A-4	0-10	0-15	70-95	60-90	60-90	30-60	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
409B: Brayton-----	18-23	Gravelly fine sandy loam, gravelly loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-15	70-95	60-90	60-90	30-60	15-25	NP-5
	23-60	Gravelly fine sandy loam, gravelly loam, fine sandy loam, loam	ML, SM	A-2, A-4	0-10	0-15	70-95	60-90	60-90	30-60	15-25	NP-5
412B: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-2, A-4	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-2, A-4	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
	24-60	Gravelly sandy loam, gravelly fine sandy loam, loam, fine sandy loam, sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
412C: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-2, A-4	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-2, A-4	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
	24-60	Gravelly fine sandy loam, sandy loam, fine sandy loam, gravelly sandy loam, loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
412D: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-4, A-2	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-4, A-2	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-4, A-2	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
412D: Bice-----	24-60	Gravelly sandy loam, gravelly fine sandy loam, loam, fine sandy loam, sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
413C: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-2, A-4	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-2, A-4	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
	24-60	Gravelly sandy loam, sandy loam, gravelly fine sandy loam, loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
Millsite-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-5	Fine sandy loam	ML, SM	A-4	0	0-5	90-100	80-100	75-100	40-55	15-25	NP-5
	5-13	Stony fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4, A-2	0-25	0-15	85-100	75-90	70-85	30-60	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
413C: Millsite-----	13-24	Fine sandy loam, gravelly loam, stony sandy loam	SM, ML, CL-ML, SC-SM	A-2, A-4	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-10
	24-31	Sandy loam, gravelly loam, stony fine sandy loam	ML, SM	A-2, A-4	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-5
	31-80	Bedrock			---	---	---	---	---	---	---	---
413E: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-2, A-4	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-2, A-4	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-4, A-2	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
	24-60	Sandy loam, fine sandy loam, loam, gravelly fine sandy loam, gravelly sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
Millsite-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-5	Fine sandy loam	ML, SM	A-4	0	0-5	90-100	80-100	75-100	40-55	15-25	NP-5
	5-13	Stony fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-25	0-15	85-100	75-90	70-85	30-60	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
413E: Millsite-----	13-24	Fine sandy loam, gravelly loam, stony sandy loam	SM, ML, CL-ML, SC-SM	A-4, A-2	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-10
	24-31	Sandy loam, gravelly loam, stony fine sandy loam	ML, SM	A-2, A-4	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-5
	31-80	Bedrock			---	---	---	---	---	---	---	---
414: Fredon, cold----	0-8	Silt loam	ML, SM	A-4	0	0	85-100	75-100	70-100	45-95	15-25	NP-5
	8-17	Silt loam, loam, very fine sandy loam, fine sandy loam	SM, ML	A-4	0	0-5	90-100	80-100	75-100	35-85	15-25	NP-5
	17-24	Fine sandy loam, very fine sandy loam, silt loam, loam	ML, SM	A-4	0	0-5	90-100	80-100	75-100	35-85	15-25	NP-5
	24-29	Stratified gravelly sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-90	60-85	30-80	5-25	15-20	NP-5
	29-48	Stratified gravelly sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-90	60-85	30-80	5-25	15-20	NP-5
	48-60	Stratified gravelly sand to loamy fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-90	60-85	30-80	5-25	15-20	NP-5
415C: Millsite-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-5	Fine sandy loam	ML, SM	A-4	0	0-5	90-100	80-100	75-100	40-55	15-25	NP-5
	5-13	Stony fine sandy loam, gravelly sandy loam, loam	ML, SM	A-2, A-4	0-25	0-15	85-100	75-90	70-85	30-60	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
					Pct	Pct						
415C: Millsite-----	13-24	Fine sandy loam, gravelly loam, stony sandy loam	SM, ML, CL-ML, SC-SM	A-4, A-2	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-10
	24-31	Sandy loam, gravelly loam, stony fine sandy loam	ML, SM	A-4, A-2	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-5
	31-80	Bedrock			---	---	---	---	---	---	---	---
Westminster----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-5	Fine sandy loam	SM	A-4	0-5	0-5	90-95	85-90	80-90	40-50	15-25	NP-5
	5-12	Fine sandy loam, gravelly fine sandy loam, loam	ML, SM	A-2, A-4	0-10	0-10	80-90	70-90	65-90	30-60	15-25	NP-5
	12-16	Fine sandy loam, gravelly fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-10	80-90	70-90	65-90	30-60	15-25	NP-5
	16-80	Bedrock			---	---	---	---	---	---	---	---
Rock outcrop----	---	---	---	---	---	---	---	---	---	---	---	---
415E: Millsite-----	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-5	Fine sandy loam	ML, SM	A-4	0	0-5	90-100	80-100	75-100	40-55	15-25	NP-5
	5-13	Stony fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4, A-2	0-25	0-15	85-100	75-90	70-85	30-60	15-25	NP-5
	13-24	Fine sandy loam, gravelly loam, stony sandy loam	SM, ML, CL-ML, SC-SM	A-2, A-4	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-10
	24-31	Sandy loam, gravelly loam, stony fine sandy loam	ML, SM	A-2, A-4	0-25	0-15	80-95	75-90	55-90	30-55	15-25	NP-5
	31-80	Bedrock			---	---	---	---	---	---	---	---

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
				Pct	Pct					Pct		
416F: Westminster-----	In											
	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-5	Fine sandy loam	SM	A-4	0-5	0-5	90-95	85-90	80-90	40-50	15-25	NP-5
	5-12	Fine sandy loam, gravelly fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-10	80-90	70-90	65-90	30-60	15-25	NP-5
	12-16	Fine sandy loam, gravelly fine sandy loam, loam	ML, SM	A-4, A-2	0-10	0-10	80-90	70-90	65-90	30-60	15-25	NP-5
	16-80	Bedrock			---	---	---	---	---	---	---	
417B: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-4, A-2	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-2, A-4	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
	24-60	Sandy loam, loam, gravelly fine sandy loam, gravelly sandy loam, fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
417C: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-4, A-2	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-4, A-2	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
	24-60	Sandy loam, fine sandy loam, loam, gravelly fine sandy loam, gravelly sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
417D: Bice-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-7	Fine sandy loam	ML, SM	A-4, A-2	0	0-5	90-95	80-90	75-90	30-50	20-25	NP-5
	7-16	Fine sandy loam, gravelly sandy loam, gravelly coarse sandy loam, silt loam, sandy loam	CL-ML, ML, SM, SC-SM	A-2, A-4	0	0-10	75-95	70-90	40-90	25-70	20-25	NP-10
	16-24	Gravelly fine sandy loam, gravelly sandy loam, sandy loam, loam, fine sandy loam	SC-SM, CL-ML, ML, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
417D: Bice-----	In 24-60	Gravelly fine sandy loam, sandy loam, fine sandy loam, loam, gravelly sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0	0-10	75-95	70-90	50-90	30-60	20-25	NP-10
418C: Schroon-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	3-9	Fine sandy loam	SM	A-4	0	0-5	90-100	80-95	65-95	35-50	15-25	NP-5
	9-14	Fine sandy loam, sandy loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
	14-23	Fine sandy loam, loam, sandy loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
	23-30	Sandy loam, fine sandy loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
	30-60	Sandy loam, fine sandy loam, coarse sandy loam, gravelly sandy loam	SM	A-4, A-2	0	0-5	65-100	55-95	35-90	25-45	10-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
420A: Schroon-----	In											
	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	3-9	Fine sandy loam	SM	A-4	0	0-5	90-100	80-95	65-95	35-50	15-25	NP-5
	9-14	Fine sandy loam, sandy loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
	14-23	Fine sandy loam, loam, sandy loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
	23-30	Sandy loam, fine sandy loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
30-60	Sandy loam, fine sandy loam, coarse sandy loam, gravelly sandy loam	SM	A-2, A-4	0	0-5	65-100	55-95	35-90	25-45	10-25	NP-10	
420B: Schroon-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	3-9	Fine sandy loam	SM	A-4	0	0-5	90-100	80-95	65-95	35-50	15-25	NP-5
	9-14	Fine sandy loam, sandy loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
					Pct	Pct					Pct	
420B: Schroon-----	In											
	14-23	Fine sandy loam, loam, sandy loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
	23-30	Sandy loam, fine sandy loam, loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0	0-5	65-95	55-95	35-90	20-55	15-25	NP-5
	30-60	Sandy loam, fine sandy loam, coarse sandy loam, gravelly sandy loam	SM	A-2, A-4	0	0-5	65-100	55-95	35-90	25-45	10-25	NP-10
421A: Ninigret, cold--	0-8	Fine sandy loam	ML, SM	A-4, A-2	0	0	90-100	70-100	60-100	25-55	0-35	NP-5
	8-16	Fine sandy loam, very fine sandy loam, silt loam	ML, SM	A-2, A-4	0	0	95-100	75-100	70-100	30-80	0-25	NP-5
	16-26	Fine sandy loam, very fine sandy loam, silt loam	ML, SM	A-2, A-4	0	0	95-100	75-100	70-100	30-80	0-25	NP-5
	26-65	Stratified very gravelly coarse sand to loamy fine sand	SM, SP	A-1, A-2, A-3	0-10	0-15	70-100	35-100	5-95	1-30	0-15	NP
423A: Sudbury, cold---	0-1	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-5	Sandy loam	ML, SM	A-2, A-4	0	0-10	90-100	80-100	50-95	30-65	0-15	NP
	5-17	Gravelly sandy loam, fine sandy loam, sandy loam	SM	A-2	0	0-20	70-100	60-100	35-95	25-35	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
423A: Sudbury, cold---	In											
	17-25	Sandy loam, fine sandy loam, gravelly sandy loam	SM	A-2	0	0-20	70-100	60-100	35-95	25-35	0-15	NP
	25-60	Stratified gravel to sand	SM, SP-SM, GM, GP-GM	A-2, A-3, A-1	0-20	0-40	40-100	20-90	15-75	5-20	0-15	NP
424B: Shelburne-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Fine sandy loam	SM	A-4	0-5	0-5	95-100	85-95	80-90	35-50	15-25	NP-5
	2-7	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	7-21	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	21-27	Bouldery fine sandy loam, gravelly fine sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-25	0-10	80-95	70-90	50-90	30-55	15-25	NP-5
	27-32	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	32-43	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
424B: Shelburne-----	In											
	43-55	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	55-80	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
424C: Shelburne-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Fine sandy loam	SM	A-4	0-5	0-5	95-100	85-95	80-90	35-50	15-25	NP-5
	2-7	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	7-21	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	21-27	Bouldery fine sandy loam, gravelly fine sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-25	0-10	80-95	70-90	50-90	30-55	15-25	NP-5
	27-32	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
424C: Shelburne-----	In											
	55-80	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	32-43	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
424D: Shelburne-----	43-55	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Fine sandy loam	SM	A-4	0-5	0-5	95-100	85-95	80-90	35-50	15-25	NP-5
	2-7	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	7-21	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	21-27	Bouldery fine sandy loam, gravelly fine sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-25	0-10	80-95	70-90	50-90	30-55	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
424D: Shelburne-----	27-32	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	32-43	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	43-55	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	55-80	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
425B: Shelburne-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Fine sandy loam	SM	A-4	0-5	0-5	95-100	85-95	80-90	35-50	15-25	NP-5
	2-7	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	7-21	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
425B: Shelburne-----	21-27	Bouldery fine sandy loam, gravelly fine sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-4, A-2	0-25	0-10	80-95	70-90	50-90	30-55	15-25	NP-5
	27-32	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	32-43	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	43-55	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	55-80	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
425C: Shelburne-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Fine sandy loam	SM	A-4	0-5	0-5	95-100	85-95	80-90	35-50	15-25	NP-5
	2-7	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
425C: Shelburne-----	In				Pct	Pct						
	7-21	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	21-27	Bouldery fine sandy loam, gravelly fine sandy loam, gravelly sandy loam, fine sandy loam, loam	ML, SM	A-2, A-4	0-25	0-10	80-95	70-90	50-90	30-55	15-25	NP-5
	27-32	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	32-43	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	43-55	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	55-80	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
426D: Shelburne-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Fine sandy loam	SM	A-4	0-5	0-5	95-100	85-95	80-90	35-50	15-25	NP-5
	2-7	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	7-21	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	80-95	70-90	50-90	35-55	15-25	NP-5
	21-27	Bouldery fine sandy loam, gravelly fine sandy loam, gravelly sandy loam, fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-25	0-10	80-95	70-90	50-90	30-55	15-25	NP-5
	27-32	Gravelly fine sandy loam, fine sandy loam, gravelly sandy loam, loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	32-43	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
	43-55	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
426D: Shelburne-----	55-80	Fine sandy loam, loam, gravelly sandy loam, gravelly fine sandy loam	ML, SM	A-4	0-10	0-10	90-100	70-90	50-90	35-55	15-25	NP-10
427B: Ashfield-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	3-7	Fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-50	15-25	NP-5
	7-12	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-70	15-25	NP-5
	12-18	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	CL-ML, SC-SM, ML, SM	A-4, A-2	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	18-24	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	SM, CL-ML, SC-SM, ML	A-2, A-4	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	24-29	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM, SC-SM, CL-ML	A-2, A-4	0-5	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	29-44	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
427B: Ashfield-----	44-58	Sandy loam, fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	58-80	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
427C: Ashfield-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	3-7	Fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-50	15-25	NP-5
	7-12	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-70	15-25	NP-5
	12-18	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	CL-ML, SC-SM, ML, SM	A-4, A-2	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	18-24	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	SM, CL-ML, SC-SM, ML	A-4, A-2	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	24-29	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM, SC- SM, CL-ML	A-2, A-4	0-5	0-10	85-95	70-90	60-85	30-70	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
				Pct	Pct					Pct		
427C: Ashfield-----	In											
	29-44	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	44-58	Sandy loam, fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	58-80	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
428A: Ashfield-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	3-7	Fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-50	15-25	NP-5
	7-12	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-70	15-25	NP-5
	12-18	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	CL-ML, SC-SM, ML, SM	A-4, A-2	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	18-24	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	SM, CL-ML, SC-SM, ML	A-4, A-2	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
428A: Ashfield-----	24-29	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM, SC-SM, CL-ML	A-4, A-2	0-5	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	29-44	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	44-58	Sandy loam, fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	58-80	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
428B: Ashfield-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	3-7	Fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-50	15-25	NP-5
	7-12	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-70	15-25	NP-5
	12-18	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	CL-ML, SC-SM, ML, SM	A-2, A-4	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
428B: Ashfield-----	18-24	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	SM, CL-ML, SC-SM, ML	A-2, A-4	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	24-29	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM, SC-SM, CL-ML	A-4, A-2	0-5	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	29-44	Fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	44-58	Sandy loam, fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	58-80	Fine sandy loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
428C: Ashfield-----	0-1	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	1-2	Moderately decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-3	Highly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	3-7	Fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-50	15-25	NP-5
	7-12	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM	A-4	0	0-5	90-95	80-90	75-90	35-70	15-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
428C: Ashfield-----	12-18	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	CL-ML, SC-SM, ML, SM	A-4, A-2	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	18-24	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	SM, CL-ML, SC-SM, ML	A-4, A-2	0	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	24-29	Fine sandy loam, loam, silt loam, gravelly fine sandy loam	ML, SM, SC-SM, CL-ML	A-4, A-2	0-5	0-10	85-95	70-90	60-85	30-70	15-25	NP-5
	29-44	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	44-58	Sandy loam, fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-2, A-4	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
	58-80	Fine sandy loam, gravelly loam, gravelly sandy loam, gravelly fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2	0-5	0-10	85-95	70-90	40-85	20-70	15-25	NP-5
429A: Agawam, cold----	0-8	Fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
429A: Agawam, cold----	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP
429B: Agawam, cold----	0-8	Fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5
	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP
429C: Agawam, cold----	0-8	Fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	80-100	70-100	30-60	0-25	NP-5
	8-14	Fine sandy loam, very fine sandy loam, loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-60	0-25	NP-5
	14-24	Fine sandy loam, very fine sandy loam	ML, SM	A-2, A-4	0	0	90-100	75-100	65-100	30-65	0-20	NP-5
	24-60	Stratified very gravelly coarse sand to fine sand	SM, SP-SM	A-1, A-2, A-3	0	0-10	70-100	25-100	5-95	1-25	0-15	NP
433: Moosilauke-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-6	Loam	ML	A-4	0	0-5	90-100	80-100	70-100	50-75	0-15	NP
	6-16	Fine sandy loam, sandy loam, gravelly sandy loam	SM, ML	A-2, A-4	0	0-9	80-100	65-100	45-100	25-55	0-15	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
433: Moosilauke-----	16-24	Gravelly fine sandy loam, fine sandy loam, sandy loam, gravelly sandy loam	SM, ML	A-2, A-4	0	0-9	80-100	65-100	45-100	25-55	0-15	NP
	24-39	Loamy fine sand, very gravelly coarse sand	SW-SM, SM	A-3, A-1, A-2	0	0-17	70-100	40-100	15-100	5-35	0-15	NP
	39-65	Fine sand, very gravelly coarse sand	SW-SM, SM	A-1, A-2, A-3	0	0-17	70-100	40-100	15-100	5-35	0-15	NP
434A: Merrimac, cold--	0-9	Sandy loam	ML, SM	A-4, A-2	0	0	90-100	70-90	45-85	25-60	0-20	NP
	9-16	Sandy loam, fine sandy loam	SM	A-2, A-4	0	0	90-100	75-90	45-90	25-50	0-25	NP
	16-24	Gravelly sandy loam, sandy loam	SM	A-2, A-4	0	0	80-100	55-90	35-70	15-50	0-25	NP
	24-60	Stratified very gravelly coarse sand to gravelly sand	SP, SP-SM	A-1	0	10-20	65-85	35-60	35-60	1-15	0-25	NP
434B: Merrimac, cold--	0-9	Sandy loam	ML, SM	A-2, A-4	0	0	90-100	70-90	45-85	25-60	0-20	NP
	9-16	Sandy loam, fine sandy loam	SM	A-2, A-4	0	0	90-100	75-90	45-90	25-50	0-25	NP
	16-24	Gravelly sandy loam, sandy loam	SM	A-2, A-4	0	0	80-100	55-90	35-70	15-50	0-25	NP
	24-60	Stratified very gravelly coarse sand to gravelly sand	SP, SP-SM	A-1	0	10-20	65-85	35-60	35-60	1-15	0-25	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
434C: Merrimac, cold--	0-9	Sandy loam	ML, SM	A-4, A-2	0	0	90-100	70-90	45-85	25-60	0-20	NP
	9-16	Sandy loam, fine sandy loam	SM	A-2, A-4	0	0	90-100	75-90	45-90	25-50	0-25	NP
	16-24	Gravelly sandy loam, sandy loam	SM	A-2, A-4	0	0	80-100	55-90	35-70	15-50	0-25	NP
	24-60	Stratified very gravelly coarse sand to gravelly sand	SP, SP-SM	A-1	0	10-20	65-85	35-60	35-60	1-15	0-25	NP
435: Scarboro-----	0-12	Muck	PT	A-8	0	0	100	100	100	100	---	---
	12-17	Loamy sand, sandy loam, fine sandy loam, loamy fine sand	SM	A-2	0	0	90-100	85-100	70-90	20-30	0-20	NP
	17-31	Stratified sand to loamy fine sand	SP-SM, SM	A-2	0	0	85-100	75-100	45-95	5-30	0-15	NP
	31-72	Stratified very gravelly coarse sand to loamy fine sand	SP-SM, SM	A-1, A-2, A-3	0	0-15	70-100	40-100	20-95	5-30	0-15	NP
436: Halsey-----	0-1	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Silt loam	ML, CL-ML	A-4	0	0-5	90-100	80-100	75-100	50-85	20-25	NP-5
	8-16	Silt loam, loam, very fine sandy loam, gravelly loam	SM, CL-ML, ML, SC-SM	A-4	0	0-5	80-100	75-100	70-100	45-85	20-25	NP-10
	16-28	Fine sandy loam, very fine sandy loam, gravelly fine sandy loam, gravelly very fine sandy loam	ML, SM	A-4	0	0-5	90-100	75-100	65-100	35-70	20-25	NP-5

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
436: Halsey-----	28-38	Loamy sand, gravelly loamy sand, very gravelly loamy sand, sand	SW-SM, SM	A-1, A-2, A-3	0	0-15	55-100	40-85	25-65	5-25	15-20	NP-5
	38-60	Sand, loamy sand, gravelly loamy sand, very gravelly loamy sand	SM, SW-SM	A-1, A-2, A-3	0	0-15	55-90	40-85	25-65	5-25	15-20	NP-5
437: Wonsqueak-----	0-2	Mucky peat	PT	A-8	0	0	100	100	100	100	---	---
	2-11	Muck	PT	A-8	0	0	100	100	100	100	---	---
	11-22	Muck	PT	A-8	0	0	100	100	100	100	---	---
	22-25	Mucky silt loam, silt loam, loam, fine sandy loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0	0-10	85-100	75-100	70-95	30-75	15-25	NP-5
	25-45	Gravelly fine sandy loam, fine sandy loam, loam, silt loam	ML, SM	A-2, A-4	0	0-10	85-100	75-100	70-95	30-75	15-25	NP-5
	45-60	Fine sandy loam, gravelly fine sandy loam, loam, silt loam	ML, SM	A-2, A-4	0	0-10	85-100	75-100	70-95	30-75	15-25	NP-5
438: Bucksport-----	0-9	Muck	PT	A-8	0	0	100	100	100	100	---	---
	9-33	Muck	PT	A-8	0	0	100	100	100	100	---	---
	33-50	Muck	PT	A-8	0	0	100	100	100	100	---	---
	50-59	Muck	PT	A-8	0	0	100	100	100	100	---	---
	59-63	Sandy loam, fine sandy loam, gravelly sand, loam	ML, SM, SP-SM	A-1, A-2, A-4	0	0-10	75-100	65-100	45-90	10-70	10-25	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
440A: Boscawen-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-2	Gravelly sandy loam	SM	A-2, A-4	0	0-9	85-90	75-80	50-70	30-45	15-20	NP
	2-9	Very gravelly sandy loam, very gravelly loamy sand, gravelly sandy loam, gravelly loamy sand	SM, SW-SM	A-1, A-2, A-4	0	0-17	70-80	40-70	25-65	10-40	15-20	NP
	9-16	Very gravelly loamy sand	SM	A-1	0	0-17	65-100	30-50	20-35	10-15	15-20	NP
	16-29	Stratified fine sand to extremely gravelly coarse sand	SM, SW, SP, SW-SM	A-2, A-1, A-3	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	29-34	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-3, A-2, A-1	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	34-40	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-2, A-3, A-1	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	40-44	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-1, A-2, A-3	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	44-67	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-3, A-1, A-2	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
440C: Boscawen-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-2	Gravelly sandy loam	SM	A-4, A-2	0	0-9	85-90	75-80	50-70	30-45	15-20	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
440C: Boscawen-----	2-9	Very gravelly sandy loam, very gravelly loamy sand, gravelly sandy loam, gravelly loamy sand	SM, SW-SM	A-1, A-2, A-4	0	0-17	70-80	40-70	25-65	10-40	15-20	NP
	9-16	Very gravelly loamy sand	SM	A-1	0	0-17	65-100	30-50	20-35	10-15	15-20	NP
	16-29	Stratified fine sand to extremely gravelly coarse sand	SM, SW, SP, SW-SM	A-3, A-1, A-2	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	29-34	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-2, A-3, A-1	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	34-40	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-1, A-2, A-3	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	40-44	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-2, A-3, A-1	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	44-67	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-1, A-2, A-3	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
440E: Boscawen-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-2	Gravelly sandy loam	SM	A-2, A-4	0	0-9	85-90	75-80	50-70	30-45	15-20	NP
	2-9	Very gravelly sandy loam, very gravelly loamy sand, gravelly sandy loam, gravelly loamy sand	SM, SW-SM	A-1, A-2, A-4	0	0-17	70-80	40-70	25-65	10-40	15-20	NP

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
440E: Boscawen-----	9-16	Very gravelly loamy sand	SM	A-1	0	0-17	65-100	30-50	20-35	10-15	15-20	NP
	16-29	Stratified fine sand to extremely gravelly coarse sand	SM, SW, SP, SW-SM	A-2, A-1, A-3	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	29-34	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-1, A-2, A-3	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	34-40	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-3, A-1, A-2	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	40-44	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-3, A-1, A-2	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
	44-67	Stratified fine sand to extremely gravelly coarse sand	SM, SP, SW, SW-SM	A-2, A-3, A-1	0	0-17	65-100	15-90	5-90	0-20	15-20	NP
442: Brayton-----	0-2	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	2-10	Loam	SM	A-4	0-2	0-4	95-100	80-90	45-65	25-40	20-25	NP
	10-17	Very fine sandy loam, gravelly sandy loam, silt loam, fine sandy loam, loam	CL-ML, SC-SM, ML, SM	A-2, A-1-b, A-4	0-2	0-5	80-100	55-90	35-85	20-70	20-25	2-6
	17-22	Fine sandy loam, loam, silt loam, gravelly sandy loam, very fine sandy loam	CL-ML, ML, SC-SM, SM	A-1-b, A-4, A-2	0-2	0-5	80-100	55-90	35-85	20-70	20-25	2-6

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
442: Brayton-----	22-27	Gravelly sandy loam, fine sandy loam, silt loam, very fine sandy loam, loam, sandy loam	CL-ML, ML, SC-SM, SM	A-1-b, A-2, A-4	0-2	0-5	80-100	55-90	35-85	20-70	20-25	2-6
	27-42	Gravelly sandy loam, gravelly fine sandy loam, gravelly loam, fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-1-b, A-2, A-4	0-2	0-4	80-100	60-95	40-90	20-65	20-25	2-7
	42-65	Gravelly sandy loam, gravelly fine sandy loam, gravelly loam, fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-1-b, A-2, A-4	0-2	0-4	80-100	60-95	40-90	20-65	20-25	2-7
443: Brayton-----	0-2	Moderately decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	2-10	Loam	SM	A-4	0-2	0-4	95-100	80-90	45-65	25-40	20-25	NP
	10-17	Very fine sandy loam, gravelly sandy loam, silt loam, loam, fine sandy loam	CL-ML, SC-SM, ML, SM	A-2, A-1-b, A-4	0-2	0-5	80-100	55-90	35-85	20-70	20-25	2-6
	17-22	Loam, silt loam, fine sandy loam, gravelly sandy loam, very fine sandy loam	CL-ML, ML, SC-SM, SM	A-1-b, A-2, A-4	0-2	0-5	80-100	55-90	35-85	20-70	20-25	2-6

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
					Pct	Pct						
443: Brayton-----	22-27	Gravelly sandy loam, loam, sandy loam, fine sandy loam, silt loam, very fine sandy loam	CL-ML, ML, SC-SM, SM	A-4, A-2, A-1-b	0-2	0-5	80-100	55-90	35-85	20-70	20-25	2-6
	27-42	Gravelly sandy loam, gravelly fine sandy loam, gravelly loam, fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-4, A-2, A-1-b	0-2	0-4	80-100	60-95	40-90	20-65	20-25	2-7
	42-65	Gravelly sandy loam, gravelly fine sandy loam, gravelly loam, fine sandy loam, loam	CL-ML, ML, SC-SM, SM	A-1-b, A-2, A-4	0-2	0-4	80-100	60-95	40-90	20-65	20-25	2-7
Loonmeadow-----	0-2	Slightly decomposed plant material	OL	A-8	0	0	100	100	100	100	---	---
	2-9	Mucky fine sandy loam	SM	A-4	0-5	0-5	90-100	80-100	75-95	40-50	15-25	NP-5
	9-18	Sandy loam, fine sandy loam, gravelly sandy loam	SC-SM, SM	A-4	0-10	0-10	85-95	85-95	65-85	35-45	15-25	NP-5
	18-35	Gravelly sandy loam, gravelly fine sandy loam, sandy loam	SC-SM, SM	A-2, A-4	0-10	0-10	75-90	65-90	55-85	30-40	10-20	NP-5
	35-80	Gravelly sandy loam, gravelly fine sandy loam, sandy loam	SC-SM, SM	A-2, A-4	0-10	0-10	75-90	65-90	55-85	30-40	10-20	NP-5

Table 23.-Engineering Properties-Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
448B: Hogansburg-----	0-12	Loam	ML	A-4	0	0	85-100	80-95	70-90	50-70	25-35	5-10
	12-20	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-2	0-5	65-100	55-90	50-90	30-80	25-35	5-10
	20-29	Gravelly fine sandy loam, loam, silt loam	ML, SM	A-2, A-4	0-2	0-5	65-100	55-90	50-90	25-75	20-30	NP-10
	29-43	Gravelly fine sandy loam, loam, silt loam	SM, ML	A-4, A-2	0-2	0-5	65-100	55-90	50-90	25-75	20-30	NP-10
	43-50	Gravelly very fine sandy loam, very gravelly fine sandy loam, loam	ML, SM	A-4, A-2	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
	50-70	Loam, fine sandy loam, very gravelly fine sandy loam, gravelly very fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
	70-84	Loam, gravelly very fine sandy loam, very gravelly fine sandy loam	SM, ML	A-2, A-4	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
449B: Hogansburg-----	0-12	Loam	ML	A-4	0	0	85-100	80-95	70-90	50-70	25-35	5-10
	12-20	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-2	0-5	65-100	55-90	50-90	30-80	25-35	5-10
	20-29	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-2	0-5	65-100	55-90	50-90	25-75	20-30	NP-10
	29-43	Loam, gravelly fine sandy loam, silt loam	ML, SM	A-2, A-4	0-2	0-5	65-100	55-90	50-90	25-75	20-30	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth In	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
449B: Hogansburg-----					Pct	Pct						
	43-50	Gravelly very fine sandy loam, very gravelly fine sandy loam, loam	ML, SM	A-2, A-4	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
	50-70	Loam, fine sandy loam, very gravelly fine sandy loam, gravelly very fine sandy loam	ML, SM	A-2, A-4	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
	70-84	Loam, gravelly very fine sandy loam, very gravelly fine sandy loam	SM, ML	A-2, A-4	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
449C: Hogansburg-----	0-12	Loam	ML	A-4	0	0	85-100	80-95	70-90	50-70	25-35	5-10
	12-20	Loam, gravelly fine sandy loam, silt loam	ML, SM	A-4, A-2	0-2	0-5	65-100	55-90	50-90	30-80	25-35	5-10
	20-29	Loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-2	0-5	65-100	55-90	50-90	25-75	20-30	NP-10
	29-43	Loam, gravelly fine sandy loam, silt loam	ML, SM	A-2, A-4	0-2	0-5	65-100	55-90	50-90	25-75	20-30	NP-10
	43-50	Gravelly very fine sandy loam, very gravelly fine sandy loam, loam	ML, SM	A-2, A-4	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches Pct	inches Pct						
449C: Hogansburg-----	In											
	50-70	Loam, fine sandy loam, very gravelly fine sandy loam, gravelly very fine sandy loam	ML, SM	A-4, A-2	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
	70-84	Loam, gravelly very fine sandy loam, very gravelly fine sandy loam	SM, ML	A-2, A-4	0-5	0-5	60-100	50-90	40-90	20-70	20-30	NP-10
450B: Pyrities-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Loam	ML	A-4	0-4	0-4	90-100	80-90	75-90	45-75	25-35	NP-10
	8-13	Loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	13-26	Loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	26-45	Loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	45-65	Fine sandy loam, gravelly loam	CL-ML, SM, ML, SC-SM, SC	A-2, A-4	0-7	0-7	70-100	45-95	40-90	25-60	20-30	NP-10
450C: Pyrities-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Loam	ML	A-4	0-4	0-4	90-100	80-90	75-90	45-75	25-35	NP-10
	8-13	Loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	13-26	Loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
450C: Pyrities-----	26-45	Loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	45-65	Fine sandy loam, gravelly loam	CL-ML, SM, ML, SC-SM, SC	A-2, A-4	0-7	0-7	70-100	45-95	40-90	25-60	20-30	NP-10
450D: Pyrities-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Loam	ML	A-4	0-4	0-4	90-100	80-90	75-90	45-75	25-35	NP-10
	8-13	Loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	13-26	Loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	26-45	Loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	45-65	Fine sandy loam, gravelly loam	CL-ML, SM, ML, SC-SM, SC	A-2, A-4	0-7	0-7	70-100	45-95	40-90	25-60	20-30	NP-10
451B: Pyrities-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Loam	ML	A-4	0-4	0-4	90-100	80-90	75-90	45-75	25-35	NP-10
	8-13	Loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	13-26	Loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	26-45	Loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	45-65	Fine sandy loam, gravelly loam	CL-ML, SM, ML, SC-SM, SC	A-2, A-4	0-7	0-7	70-100	45-95	40-90	25-60	20-30	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
451C: Pyrities-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Loam	ML	A-4	0-4	0-4	90-100	80-90	75-90	45-75	25-35	NP-10
	8-13	Loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	13-26	Loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	26-45	Loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	45-65	Fine sandy loam, gravelly loam	CL-ML, SM, ML, SC-SM, SC	A-2, A-4	0-7	0-7	70-100	45-95	40-90	25-60	20-30	NP-10
451D: Pyrities-----	0-1	Slightly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	1-8	Loam	ML	A-4	0-4	0-4	90-100	80-90	75-90	45-75	25-35	NP-10
	8-13	Loam, gravelly fine sandy loam	SM, ML	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	13-26	Loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	26-45	Loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	45-65	Fine sandy loam, gravelly loam	CL-ML, SM, ML, SC-SM, SC	A-2, A-4	0-7	0-7	70-100	45-95	40-90	25-60	20-30	NP-10
457: Mudgepond-----	0-11	Silt loam	ML	A-4	0-5	0	90-95	80-90	75-85	60-75	20-30	NP-5
	11-16	Loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
457: Mudgepond-----	16-26	Fine sandy loam, loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	26-35	Gravelly fine sandy loam, very fine sandy loam, silt loam, loam	ML, SM	A-4, A-2	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	35-65	Gravelly fine sandy loam, very fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-85	25-65	20-25	NP-10
458: Mudgepond-----	0-11	Silt loam	ML	A-4	0-5	0	90-95	80-90	75-85	60-75	20-30	NP-5
	11-16	Loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	16-26	Fine sandy loam, loam, very fine sandy loam, silt loam, gravelly fine sandy loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	26-35	Gravelly fine sandy loam, very fine sandy loam, silt loam, loam	ML, SM	A-4, A-2	0-5	5-10	65-90	60-90	55-90	25-75	20-30	NP-5
	35-65	Gravelly fine sandy loam, very fine sandy loam, sandy loam, loam	ML, SM	A-2, A-4	0-5	5-10	65-90	60-90	55-85	25-65	20-25	NP-10

Table 23.—Engineering Properties—Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct					Pct	
458: Alden-----	0-4	Mucky silt loam	ML	A-4	0-5	0-5	90-100	80-100	75-100	60-95	25-35	NP-5
	4-13	Silt loam, very fine sandy loam	ML	A-4	0-5	0-5	90-100	80-100	75-100	50-95	25-35	NP-5
	13-23	Silt loam, very fine sandy loam	ML	A-4	0-5	0-5	90-100	80-100	75-100	50-95	25-35	NP-5
	23-29	Silt loam, very fine sandy loam	ML, SM	A-4	0-5	0-5	90-100	80-100	75-100	45-95	25-35	NP-5
	29-43	Gravelly loam, silt loam, loam	CL, SC	A-6	0-5	0-5	60-100	55-90	50-90	40-80	25-35	10-15
	43-60	Loam, gravelly loam, silt loam	CL, SC	A-6	0-5	0-15	70-100	60-90	50-90	40-80	25-35	10-15
501: Ondawa-----	0-1	Slightly decomposed plant material			0	0	0	0	0	0	---	---
	1-2	Highly decomposed plant material	PT	A-8	0	0	100	100	100	100	---	---
	2-14	Loam, silt loam	ML	A-4	0-4	0-4	90-100	80-90	75-90	45-75	25-35	NP-10
	14-30	Loam, gravelly fine sandy loam	SM, ML	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	30-33	Loam, gravelly fine sandy loam	ML, SM	A-4, A-2	0-4	0-4	70-100	55-90	50-85	30-75	25-35	NP-10
	33-60	Fine sandy loam, gravelly loam	SM, CL-ML, ML, SC-SM, SC	A-2, A-4	0-7	0-7	70-100	45-95	40-90	25-60	20-30	NP-10
503: Rumney-----	0-7	Silt loam	ML	A-4	0	0	95-100	75-100	70-100	50-90	19-32	NP-10
	7-22	Sandy loam, fine sandy loam, loam	CL, CL-ML, ML, SC, SC- SM, SM	A-4, A-2	0	0	95-100	75-100	60-95	30-80	18-32	NP-10
	22-38	Sandy loam, fine sandy loam, loam	SM, SC-SM, CL-ML, CL, SC, ML	A-2, A-4	0	0	95-100	75-100	60-95	30-80	18-32	NP-10
	38-42	Fine sandy loam, sandy loam, silt loam, loam	CL, SC, SC- SM, ML, CL- ML, SM	A-2, A-4	0	0	95-100	75-100	60-100	30-95	18-32	NP-10

Table 23.--Engineering Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit Pct	Plas- ticity index
			Unified	AASHTO	>10	3-10	4	10	40	200		
					inches	inches						
	In				Pct	Pct						
503: Rumney-----	42-44	Loamy fine sand, fine sandy loam, extremely gravelly coarse sand	CL, CL-ML, ML, SC, SC-SM, SM, SW-SM	A-4, A-3, A-2, A-1	0	0-9	80-100	25-100	10-85	5-55	17-27	NP-10
	44-65	Extremely gravelly coarse sand, loamy fine sand, gravelly fine sandy loam	CL-ML, SC, CL, SC-SM, SM, SW-SM, ML	A-1, A-2, A-4	0	0-9	80-100	25-100	10-50	5-50	17-27	NP-10
508: Medomak-----	0-7	Mucky silt loam	ML	A-4	0	0	95-100	90-100	80-100	60-90	20-35	NP-10
	7-24	Silt loam, fine sandy loam, very fine sandy loam	ML	A-4	0	0	95-100	90-100	85-95	60-80	20-35	NP-10
	24-33	Mucky silt loam, silt loam	ML	A-4	0	0	95-100	90-100	80-100	60-90	20-35	NP-10
	33-46	Very gravelly coarse sand, silt loam, very fine sandy loam	ML, SM, SC-SM	A-4, A-2, A-1	0	0-10	70-100	35-100	20-100	10-95	20-35	NP-10
	46-79	Very fine sandy loam, loamy very fine sand, very gravelly coarse sand	SP-SM, ML, SM, SW-SM	A-3, A-1, A-2	0	0-10	70-100	35-100	20-100	5-95	15-25	NP-5

Table 24.--Physical Properties of the Soils

(Entries under "Erosion factors--T" apply to the entire profile. Absence of an entry indicates that data were not estimated.)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
2: Ridgebury-----	0-5	55-72	25-35	3-10	1.30-1.50	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	4.0-7.0	.15	.20	3
	5-14	47-70	28-45	2-8	1.50-1.80	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.0-1.0	.28	.43	
	14-21	47-70	28-45	2-8	1.60-1.80	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.0-1.0	.28	.43	
	21-60	47-70	28-45	2-8	1.80-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.28	.43	
3: Ridgebury-----	0-5	55-72	25-35	3-10	1.30-1.50	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	4.0-7.0	.15	.20	3
	5-14	47-70	28-45	2-8	1.50-1.80	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.0-1.0	.28	.43	
	14-21	47-70	28-45	2-8	1.60-1.80	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.0-1.0	.28	.43	
	21-60	47-70	28-45	2-8	1.80-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.28	.43	
Leicester-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	5
	1-7	55-70	27-35	3-10	1.00-1.25	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	3.0-8.0	.10	.24	
	7-10	45-67	30-45	3-10	1.35-1.60	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.32	.49	
	10-18	45-67	30-45	3-10	1.35-1.60	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-1.0	.37	.55	
	18-24	45-67	30-45	3-10	1.35-1.60	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.37	.55	
	24-43	53-68	30-40	2-7	1.45-1.70	0.6-20	4.00-141.00	0.07-0.14	0.0-2.9	0.0-0.5	.28	.43	
	43-65	53-68	30-40	2-7	1.45-1.70	0.6-20	4.00-141.00	0.05-0.14	0.0-2.9	0.0-0.5	.24	.43	
Whitman-----	0-1	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	20-95	---	---	2
	1-9	53-70	25-35	5-12	0.65-1.30	0.6-6	4.00-42.00	0.11-0.13	0.0-2.9	4.0-8.0	.15	.20	
	9-16	43-85	30-45	2-12	1.30-1.65	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-2.0	.28	.43	
	16-22	47-68	30-45	2-8	1.85-2.10	0.0015-0.2	0.01-1.40	0.03-0.07	0.0-2.9	0.0-0.5	.37	.55	
	22-60	47-68	30-45	2-8	1.85-2.10	0.0015-0.2	0.01-1.40	0.03-0.07	0.0-2.9	0.0-0.5	.37	.55	
4: Leicester-----	0-7	55-70	27-35	3-10	1.00-1.25	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	3.0-8.0	.17	.24	5
	7-10	45-67	30-45	3-10	1.35-1.60	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.32	.49	
	10-18	45-67	30-45	3-10	1.35-1.60	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-1.0	.37	.55	
	18-24	45-67	30-45	3-10	1.35-1.60	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.37	.55	
	24-43	53-68	30-40	2-7	1.45-1.70	0.6-20	4.00-141.00	0.07-0.14	0.0-2.9	0.0-0.5	.28	.43	
	43-65	53-68	30-40	2-7	1.45-1.70	0.6-20	4.00-141.00	0.05-0.14	0.0-2.9	0.0-0.5	.24	.43	
5: Wilbraham-----	0-4	10-47	50-75	3-15	1.00-1.30	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	3.0-8.0	.28	.37	3
	4-8	10-62	35-75	3-15	1.40-1.65	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-1.0	.49	.64	
	8-20	10-62	35-75	3-15	1.40-1.70	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-0.5	.49	.64	
	20-65	10-62	35-75	3-15	1.80-2.10	0.0015-0.2	0.01-1.40	0.08-0.12	0.0-2.9	0.0-0.5	.43	.64	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
6: Wilbraham-----	0-4	10-47	50-75	3-15	1.00-1.30	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	3.0-8.0	.28	.37	3
	4-8	10-62	35-75	3-15	1.40-1.65	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-1.0	.49	.64	
	8-20	10-62	35-75	3-15	1.40-1.70	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-0.5	.49	.64	
	20-65	10-62	35-75	3-15	1.80-2.10	0.0015-0.2	0.01-1.40	0.08-0.12	0.0-2.9	0.0-0.5	.43	.64	
Menlo-----	0-5	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.45	0.0-20.0	20-50	---	---	3
	5-16	10-43	50-75	7-15	0.65-1.20	0.6-2	4.00-14.00	0.14-0.20	0.0-2.9	8.0-11	.10	.15	
	16-22	10-67	30-75	3-15	1.30-1.55	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	1.0-2.0	.37	.49	
	22-27	10-67	30-75	3-15	1.40-1.65	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.5-1.0	.43	.55	
	27-40	10-67	30-75	3-15	1.80-2.10	0.0015-0.2	0.01-1.40	0.08-0.12	0.0-2.9	0.0-0.5	.43	.64	
	40-60	10-67	30-75	3-15	1.80-2.10	0.0015-0.2	0.01-1.40	0.08-0.12	0.0-2.9	0.0-0.5	.43	.64	
7: Mudgepond-----	0-11	28-30	50-60	10-22	1.10-1.40	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	3.0-8.0	.20	.28	5
	11-16	28-73	20-55	7-17	1.20-1.50	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	16-26	28-75	20-55	5-17	1.30-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	26-35	28-75	20-55	5-17	1.40-1.60	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	35-65	38-72	25-45	3-17	1.50-1.80	0.6-2	4.00-14.00	0.07-0.17	0.0-2.9	0.0-1.0	.32	.49	
8: Mudgepond-----	0-11	28-30	50-60	10-22	1.10-1.40	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	3.0-8.0	.20	.28	5
	11-16	28-73	20-55	7-17	1.20-1.50	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	16-26	28-75	20-55	5-17	1.30-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	26-35	28-75	20-55	5-17	1.40-1.60	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	35-65	38-72	25-45	3-17	1.50-1.80	0.6-2	4.00-14.00	0.07-0.17	0.0-2.9	0.0-1.0	.32	.49	
Alden-----	0-4	9-33	52-65	15-26	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-5.9	10-15	.10	.15	5
	4-13	9-54	28-65	18-26	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-5.9	3.0-8.0	.28	.24	
	13-23	9-54	28-65	18-26	1.20-1.50	0.6-2	4.00-14.00	0.13-0.21	0.0-5.9	0.5-1.0	.28	.24	
	23-29	9-54	28-65	18-26	1.20-1.50	0.6-2	4.00-14.00	0.13-0.21	0.0-5.9	0.5-1.0	.28	.24	
	29-43	14-42	40-60	18-26	1.40-1.65	0.2-0.6	1.40-4.00	0.10-0.19	0.0-2.9	0.0-0.5	.32	.49	
	43-60	14-42	40-60	18-26	1.40-1.65	0.2-0.6	1.40-4.00	0.10-0.19	0.0-2.9	0.0-0.5	.32	.49	
9: Scitico-----	0-8	4-40	50-70	10-26	1.05-1.25	0.6-2	4.00-14.00	0.19-0.21	0.0-5.9	2.0-7.0	.28	.32	3
	8-11	3-20	42-65	20-55	1.40-1.70	0.06-0.6	0.42-4.00	0.15-0.21	0.0-9.0	0.0-0.5	.37	.43	
	11-18	10-20	50-65	20-55	1.40-1.70	0.06-0.6	0.42-4.00	0.15-0.21	0.0-9.0	0.0-0.5	.43	.49	
	18-30	5-15	35-55	35-60	1.50-1.75	0.0015-0.2	0.01-1.40	0.14-0.20	3.0-9.0	0.0-0.5	.32	.37	
	30-38	5-15	35-55	35-60	1.50-1.80	0.0015-0.06	0.01-0.42	0.14-0.20	3.0-9.0	0.0-0.5	.32	.37	
	38-52	5-15	35-55	35-60	1.55-1.80	0.0015-0.06	0.01-0.42	0.14-0.20	3.0-9.0	0.0-0.5	.32	.37	
	52-65	5-15	35-55	35-60	1.55-1.80	0.0015-0.06	0.01-0.42	0.14-0.20	3.0-9.0	0.0-0.5	.32	.37	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
9: Shaker-----	0-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	2-6	58-72	20-40	2-8	1.30-1.50	2-6	14.00-42.00	0.15-0.17	0.0-2.9	2.0-7.0	.24	.28	
	6-20	47-63	35-45	2-8	1.35-1.55	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-1.5	.43	.43	
	20-30	47-63	35-45	2-8	1.35-1.55	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.43	
	30-65	5-15	25-60	35-60	1.55-1.80	0.0015-0.2	0.01-1.41	0.14-0.21	3.0-9.0	0.0-0.5	.32	.32	
Maybid-----	0-9	4-25	55-70	20-26	1.00-1.30	0.6-2	4.00-14.00	0.19-0.21	0.0-6.0	3.0-10	.28	.28	3
	9-18	5-15	30-58	27-55	1.20-1.40	0.0015-0.2	0.01-1.40	0.14-0.21	3.0-9.0	0.5-2.0	.37	.37	
	18-26	5-15	30-58	27-55	1.25-1.45	0.0015-0.2	0.01-1.40	0.14-0.21	3.0-9.0	0.5-1.0	.37	.37	
	26-36	10-20	25-55	35-55	1.30-1.50	0.0015-0.2	0.01-1.40	0.14-0.21	3.0-9.0	0.0-0.5	.37	.37	
	36-60	10-20	25-55	35-55	1.55-1.80	0.0015-0.2	0.01-1.40	0.14-0.21	3.0-9.0	0.0-0.5	.37	.37	
10: Raynham-----	0-10	5-45	50-80	5-15	1.20-1.40	0.2-2	1.40-14.00	0.19-0.21	0.0-2.9	3.0-9.0	.28	.28	5
	10-16	20-65	30-65	5-15	1.20-1.40	0.2-2	1.40-14.00	0.15-0.21	0.0-2.9	0.5-2.0	.64	.64	
	16-26	20-65	30-65	5-15	1.25-1.45	0.2-2	1.40-14.00	0.15-0.21	0.0-2.9	0.5-1.0	.64	.64	
	26-34	20-65	30-65	5-15	1.25-1.50	0.2-2	1.40-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
	34-47	20-65	30-65	5-15	1.30-1.60	0.06-0.2	0.40-1.40	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
	47-60	20-65	30-65	5-15	1.30-1.60	0.06-0.2	0.40-1.40	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
12: Raypol-----	0-8	13-46	51-75	3-12	1.00-1.25	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-8.0	.32	.37	3
	8-12	13-65	32-75	3-12	1.35-1.55	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.5	.49	.55	
	12-20	13-65	32-75	3-12	1.35-1.55	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.0-0.5	.49	.55	
	20-26	13-65	32-75	3-12	1.35-1.55	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.0-0.5	.55	.64	
	26-29	13-65	32-75	3-12	1.35-1.55	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.0-0.5	.55	.64	
	29-52	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.01-0.11	0.0-2.9	0.0-0.5	.17	.43	
	52-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.01-0.11	0.0-2.9	0.0-0.5	.17	.43	
13: Walpole-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-7	46-70	28-48	2-6	1.30-1.55	2-6	14.00-42.00	0.09-0.13	0.0-2.9	0.5-2.0	.32	.37	
	7-21	46-70	28-48	2-6	1.30-1.55	2-6	14.00-42.00	0.08-0.15	0.0-2.9	0.0-1.0	.32	.43	
	21-25	52-72	28-48	0-2	1.40-1.65	6-100	42.00-703.00	0.08-0.15	0.0-2.9	0.0-0.5	.10	.15	
	25-41	52-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.10	.15	
	41-65	52-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.10	.15	
14: Fredon-----	0-8	5-45	50-80	5-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	3
	8-17	20-65	30-65	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-2.0	.43	.49	
	17-24	20-65	30-65	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-1.0	.43	.55	
	24-29	75-93	5-18	2-7	1.35-1.55	2-20	14.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
	29-48	75-93	5-18	2-7	1.35-1.60	2-20	14.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
	48-60	75-93	5-18	2-7	1.40-1.60	2-20	14.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
15: Scarboro-----	0-12	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	---	---	3
	12-17	74-83	10-25	1-5	1.25-1.45	2-20	14.00-141.00	0.08-0.11	0.0-2.9	3.0-15	.05	.05	
	17-31	75-93	5-25	0-2	1.35-1.55	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-1.0	.24	.28	
	31-72	75-93	5-25	0-2	1.35-1.55	6-100	42.00-703.00	0.02-0.08	0.0-2.9	0.0-0.5	.24	.28	
16: Halsey-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	55-70	---	---	3
	1-8	18-42	52-65	6-17	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	3.0-8.0	.28	.32	
	8-16	18-54	40-65	6-17	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-1.5	.43	.55	
	16-28	50-58	30-45	5-12	1.25-1.50	2-6	14.00-42.00	0.10-0.17	0.0-2.9	0.5-1.5	.43	.49	
	28-38	80-95	3-12	2-8	1.40-1.60	6-20	42.00-141.00	0.02-0.07	0.0-2.9	0.0-0.5	.15	.24	
	38-60	80-95	3-12	2-8	1.40-1.60	6-20	42.00-141.00	0.02-0.07	0.0-2.9	0.0-0.5	.15	.24	
17: Timakwa-----	0-10	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	2
	10-21	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	21-24	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	24-37	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	37-47	80-96	5-15	0-5	1.40-1.75	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-2.0	.15	.17	
	47-60	80-96	5-15	0-5	1.40-1.75	6-100	42.00-703.00	0.02-0.12	0.0-2.9	0.0-2.0	.15	.17	
Natchaug-----	0-2	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	2
	2-4	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	4-6	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	6-11	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	11-18	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	18-24	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.35-0.45	0.0-20.0	55-75	---	---	
	24-33	4-73	20-70	7-18	1.45-1.75	0.2-2	1.40-14.00	0.08-0.18	0.0-2.9	0.0-2.0	.37	.43	
	33-36	4-73	20-70	7-18	1.45-1.75	0.2-2	1.40-14.00	0.08-0.18	0.0-2.9	0.0-2.0	.37	.43	
	36-80	4-73	20-70	7-18	1.45-1.75	0.2-2	1.40-14.00	0.08-0.18	0.0-2.9	0.0-1.0	.37	.43	
18: Catden-----	0-2	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	3
	2-18	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	18-47	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	47-49	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	49-61	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
Freetown-----	0-4	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	3
	4-10	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	10-22	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	22-35	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	35-41	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	41-55	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	55-71	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	
	71-91	0-0	0-0	0-0	0.30-0.55	0.6-6	4.00-42.00	0.35-0.45	0.0-20.0	55-75	---	---	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											In	Pct	Pct
20A: Ellington-----	0-8	18-45	52-70	3-12	1.20-1.40	0.6-6	4.00-42.00	0.18-0.21	0.0-2.9	2.0-5.0	.43	.49	3
	8-18	23-62	35-65	3-12	1.30-1.60	0.6-6	4.00-42.00	0.12-0.21	0.0-2.9	0.5-1.5	.37	.49	
	18-26	23-62	35-65	3-12	1.35-1.60	0.6-6	4.00-42.00	0.12-0.21	0.0-2.9	0.0-0.5	.43	.49	
	26-65	73-98	2-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.20	.24	
21A: Ninigret-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.32	.37	3
	8-16	28-70	27-60	3-12	1.35-1.60	0.6-6	4.00-42.00	0.13-0.20	0.0-2.9	0.5-1.5	.43	.49	
	16-26	28-70	27-60	3-12	1.35-1.60	0.6-6	4.00-42.00	0.13-0.20	0.0-2.9	0.0-0.5	.49	.55	
	26-65	73-100	0-25	0-2	1.45-1.70	6-100	42.00-703.00	0.01-0.11	0.0-2.9	0.0-0.5	.15	.17	
Tisbury-----	0-8	9-46	51-79	3-12	1.00-1.30	0.6-2	4.00-14.00	0.18-0.21	0.0-2.9	2.0-6.0	.43	.49	3
	8-18	9-69	28-79	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.5-1.5	.55	.64	
	18-26	9-69	28-79	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.0-0.5	.55	.64	
	26-60	77-100	0-20	0-3	1.40-1.65	6-100	42.00-703.00	0.01-0.08	0.0-2.9	0.0-0.5	.17	.20	
22A: Hero-----	0-9	37-58	35-48	7-15	1.10-1.40	0.6-6	4.00-42.00	0.11-0.14	0.0-2.9	2.0-5.0	.15	.28	3
	9-18	25-70	25-60	5-15	1.30-1.55	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	0.5-2.0	.32	.43	
	18-24	25-70	25-60	5-15	1.30-1.55	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	0.5-1.5	.32	.49	
	24-27	25-70	25-60	5-15	1.30-1.55	0.6-6	4.00-42.00	0.07-0.18	0.0-2.9	0.0-0.5	.28	.43	
	27-60	80-90	5-10	3-10	1.40-1.70	6-100	42.00-703.00	0.01-0.08	0.0-2.9	0.0-0.5	.05	.15	
22B: Hero-----	0-9	37-58	35-48	7-15	1.10-1.40	0.6-6	4.00-42.00	0.11-0.14	0.0-2.9	2.0-5.0	.15	.28	3
	9-18	25-70	25-60	5-15	1.30-1.55	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	0.5-2.0	.32	.43	
	18-24	25-70	25-60	5-15	1.30-1.55	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	0.5-1.5	.32	.49	
	24-27	25-70	25-60	5-15	1.30-1.55	0.6-6	4.00-42.00	0.07-0.18	0.0-2.9	0.0-0.5	.28	.43	
	27-60	80-90	5-10	3-10	1.40-1.70	6-100	42.00-703.00	0.01-0.08	0.0-2.9	0.0-0.5	.05	.15	
23A: Sudbury-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	5
	1-5	46-68	30-48	2-6	1.10-1.40	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.24	.28	
	5-17	63-73	20-35	2-7	1.15-1.45	2-6	14.00-42.00	0.07-0.15	0.0-2.9	0.5-2.0	.24	.32	
	17-25	63-73	20-35	2-7	1.15-1.45	2-6	14.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.28	.32	
	25-60	79-98	2-8	0-3	1.30-1.45	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.15	.28	
24A: Deerfield-----	0-8	73-83	10-20	2-7	1.45-1.65	6-20	42.00-141.00	0.09-0.11	0.0-2.9	1.0-4.0	.15	.15	2
	8-16	81-94	5-12	1-7	1.45-1.65	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.15	.20	
	16-28	81-94	5-12	1-7	1.40-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.0-0.5	.17	.20	
	28-34	80-100	0-15	0-5	1.40-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.20	.24	
	34-60	80-100	0-15	0-5	1.40-1.60	6-100	42.00-703.00	0.03-0.08	0.0-2.9	0.0-0.0	.15	.15	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
25A:													
Brancroft-----	0-6	2-32	50-74	18-24	1.20-1.40	0.2-2	1.40-14.00	0.19-0.21	0.0-6.0	3.0-6.0	.28	.28	5
	6-17	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	1.0-2.0	.43	.43	
	17-22	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	0.5-1.0	.49	.49	
	22-32	10-15	50-72	18-35	1.40-1.65	0.0015-0.2	0.01-1.40	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
	32-43	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
	43-66	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
25B:													
Brancroft-----	0-6	2-32	50-74	18-24	1.20-1.40	0.2-2	1.40-14.00	0.19-0.21	0.0-6.0	3.0-6.0	.28	.28	5
	6-17	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	1.0-2.0	.43	.43	
	17-22	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	0.5-1.0	.49	.49	
	22-32	10-15	50-72	18-35	1.40-1.65	0.0015-0.2	0.01-1.40	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
	32-43	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
	43-66	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
25C:													
Brancroft-----	0-6	2-32	50-74	18-24	1.20-1.40	0.2-2	1.40-14.00	0.19-0.21	0.0-6.0	3.0-6.0	.28	.28	5
	6-17	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	1.0-2.0	.43	.43	
	17-22	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	0.5-1.0	.49	.49	
	22-32	10-15	50-72	18-35	1.40-1.65	0.0015-0.2	0.01-1.40	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
	32-43	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
	43-66	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49	
26A:													
Berlin-----	0-6	6-32	50-70	18-24	1.20-1.40	0.2-2	1.40-14.00	0.17-0.21	0.0-6.0	3.0-6.0	.28	.37	5
	6-12	12-15	50-70	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.17-0.21	0.0-6.0	1.0-2.0	.43	.49	
	12-20	12-15	50-70	18-35	1.25-1.45	0.06-0.6	0.42-4.00	0.17-0.21	0.0-6.0	0.5-1.0	.43	.49	
	20-34	5-17	50-65	18-45	1.30-1.50	0.0015-0.2	0.01-1.40	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49	
	34-48	5-17	50-65	18-45	1.50-1.65	0.0015-0.06	0.01-0.42	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49	
	48-65	5-17	50-65	18-45	1.50-1.65	0.0015-0.06	0.01-0.42	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49	
26B:													
Berlin-----	0-6	6-32	50-70	18-24	1.20-1.40	0.2-2	1.40-14.00	0.17-0.21	0.0-6.0	3.0-6.0	.28	.37	5
	6-12	12-15	50-70	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.17-0.21	0.0-6.0	1.0-2.0	.43	.49	
	12-20	12-15	50-70	18-35	1.25-1.45	0.06-0.6	0.42-4.00	0.17-0.21	0.0-6.0	0.5-1.0	.43	.49	
	20-34	5-17	50-65	18-45	1.30-1.50	0.0015-0.2	0.01-1.40	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49	
	34-48	5-17	50-65	18-45	1.50-1.65	0.0015-0.06	0.01-0.42	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49	
	48-65	5-17	50-65	18-45	1.50-1.65	0.0015-0.06	0.01-0.42	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49	
27A:													
Belgrade-----	0-8	21-35	50-74	5-15	1.20-1.40	0.6-2	4.00-14.00	0.19-0.21	0.0-2.9	2.0-5.0	.43	.43	5
	8-16	15-60	35-70	5-15	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	1.0-2.0	.55	.55	
	16-27	15-60	35-70	5-15	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.5-1.0	.64	.64	
	27-45	15-60	35-65	5-20	1.25-1.50	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
	45-60	15-60	38-65	2-20	1.25-1.50	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors							
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in
28A:																		
Elmridge-----	0-6	54-68	30-38	2-8	1.30-1.50	2-6	14.00-42.00	0.13-0.15	0.0-2.9	2.0-6.0	.24	.28	3					
	6-10	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-1.5	.37	.37						
	10-18	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.0-1.0	.43	.43						
	18-25	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.43						
	25-65	10-20	20-55	35-60	1.55-1.80	0.0015-0.06	0.01-0.42	0.14-0.20	3.0-9.0	0.0-0.5	.32	.32						
28B:																		
Elmridge-----	0-6	54-68	30-38	2-8	1.30-1.50	2-6	14.00-42.00	0.13-0.15	0.0-2.9	2.0-6.0	.24	.28	3					
	6-10	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-1.5	.37	.37						
	10-18	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.0-1.0	.43	.43						
	18-25	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.43						
	25-65	10-20	20-55	35-60	1.55-1.80	0.0015-0.06	0.01-0.42	0.14-0.20	3.0-9.0	0.0-0.5	.32	.32						
29A:																		
Agawam-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3					
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43						
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55						
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17						
29B:																		
Agawam-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3					
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43						
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55						
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17						
29C:																		
Agawam-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3					
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43						
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55						
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17						
30A:																		
Branford-----	0-8	23-46	51-65	3-12	1.20-1.40	0.6-6	4.00-42.00	0.17-0.21	0.0-2.9	2.0-5.0	.43	.49	3					
	8-18	23-69	28-65	3-12	1.25-1.45	0.6-6	4.00-42.00	0.11-0.21	0.0-2.9	0.5-1.5	.49	.55						
	18-24	23-69	28-65	3-12	1.30-1.50	0.6-6	4.00-42.00	0.09-0.21	0.0-2.9	0.0-0.5	.49	.64						
	24-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.20						
30B:																		
Branford-----	0-8	23-46	51-65	3-12	1.20-1.40	0.6-6	4.00-42.00	0.17-0.21	0.0-2.9	2.0-5.0	.43	.49	3					
	8-18	23-69	28-65	3-12	1.25-1.45	0.6-6	4.00-42.00	0.11-0.21	0.0-2.9	0.5-1.5	.49	.55						
	18-24	23-69	28-65	3-12	1.30-1.50	0.6-6	4.00-42.00	0.09-0.21	0.0-2.9	0.0-0.5	.49	.64						
	24-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.20						

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
30C:													
Branford -----	0-8	23-46	51-65	3-12	1.20-1.40	0.6-6	4.00-42.00	0.17-0.21	0.0-2.9	2.0-5.0	.43	.49	3
	8-18	23-69	28-65	3-12	1.25-1.45	0.6-6	4.00-42.00	0.11-0.21	0.0-2.9	0.5-1.5	.49	.55	
	18-24	23-69	28-65	3-12	1.30-1.50	0.6-6	4.00-42.00	0.09-0.21	0.0-2.9	0.0-0.5	.49	.64	
	24-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.20	
31A:													
Copake -----	0-6	55-65	18-35	10-17	1.40-1.60	2-6	14.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.15	.20	3
	6-13	43-72	20-40	8-17	1.40-1.60	0.6-6	4.00-42.00	0.10-0.18	0.0-2.9	0.5-2.0	.24	.32	
	13-21	28-72	20-55	8-17	1.25-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.0-0.5	.28	.43	
	21-31	28-72	20-55	8-17	1.25-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.0-0.5	.28	.43	
	31-56	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	56-65	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	65-75	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	75-80	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
31B:													
Copake -----	0-6	55-65	18-35	10-17	1.40-1.60	2-6	14.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.15	.20	3
	6-13	43-72	20-40	8-17	1.40-1.60	0.6-6	4.00-42.00	0.10-0.18	0.0-2.9	0.5-2.0	.24	.32	
	13-21	28-72	20-55	8-17	1.25-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.0-0.5	.28	.43	
	21-31	28-72	20-55	8-17	1.25-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.0-0.5	.28	.43	
	31-56	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	56-65	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	65-75	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	75-80	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
31C:													
Copake -----	0-6	55-65	18-35	10-17	1.40-1.60	2-6	14.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.15	.20	3
	6-13	43-72	20-40	8-17	1.40-1.60	0.6-6	4.00-42.00	0.10-0.18	0.0-2.9	0.5-2.0	.24	.32	
	13-21	28-72	20-55	8-17	1.25-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.0-0.5	.28	.43	
	21-31	28-72	20-55	8-17	1.25-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.0-0.5	.28	.43	
	31-56	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	56-65	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	65-75	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
	75-80	79-91	1-20	1-8	1.50-1.70	6-100	42.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.10	.20	
32A:													
Haven -----	0-7	2-44	51-80	5-18	1.10-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.32	.43	3
	7-14	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-2.0	.49	.64	
	14-20	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.0	.49	.64	
	20-24	54-70	25-28	5-18	1.25-1.50	0.6-2	4.00-14.00	0.13-0.17	0.0-2.9	0.0-0.5	.37	.43	
	24-60	92-100	0-5	0-3	1.40-1.65	20-100	141.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.15	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
32A:													
Enfield-----	0-3	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.08-0.40	---	50-80	---	---	3
	3-4	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.08-0.40	---	50-80	---	---	
	4-12	8-51	51-80	3-12	1.20-1.40	0.6-2	4.00-14.00	0.18-0.21	0.0-2.9	2.0-6.0	.43	.49	
	12-20	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.5-2.0	.55	.64	
	20-26	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.5-2.0	.55	.64	
	26-30	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.0-0.5	.64	.64	
	30-37	85-94	3-28	0-6	1.30-1.60	0.6-2	4.00-14.00	0.01-0.11	0.0-2.9	0.0-0.5	.10	.15	
	37-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.01-0.08	0.0-2.9	0.0-0.5	.17	.20	
32B:													
Haven-----	0-7	2-44	51-80	5-18	1.10-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.32	.43	3
	7-14	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-2.0	.49	.64	
	14-20	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.0	.49	.64	
	20-24	54-70	25-28	5-18	1.25-1.50	0.6-2	4.00-14.00	0.13-0.17	0.0-2.9	0.0-0.5	.37	.43	
	24-60	92-100	0-5	0-3	1.40-1.65	20-100	141.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.15	
Enfield-----	0-3	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.08-0.40	---	50-80	---	---	3
	3-4	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.08-0.40	---	50-80	---	---	
	4-12	8-51	51-80	3-12	1.20-1.40	0.6-2	4.00-14.00	0.18-0.21	0.0-2.9	2.0-6.0	.43	.49	
	12-20	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.5-2.0	.55	.64	
	20-26	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.5-2.0	.55	.64	
	26-30	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.0-0.5	.64	.64	
	30-37	85-94	3-28	0-6	1.30-1.60	0.6-2	4.00-14.00	0.01-0.11	0.0-2.9	0.0-0.5	.10	.15	
	37-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.01-0.08	0.0-2.9	0.0-0.5	.17	.20	
32C:													
Haven-----	0-7	2-44	51-80	5-18	1.10-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.32	.43	3
	7-14	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-2.0	.49	.64	
	14-20	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.0	.49	.64	
	20-24	54-70	25-28	5-18	1.25-1.50	0.6-2	4.00-14.00	0.13-0.17	0.0-2.9	0.0-0.5	.37	.43	
	24-60	92-100	0-5	0-3	1.40-1.65	20-100	141.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.15	
Enfield-----	0-3	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.08-0.40	---	50-80	---	---	3
	3-4	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.08-0.40	---	50-80	---	---	
	4-12	8-51	51-80	3-12	1.20-1.40	0.6-2	4.00-14.00	0.18-0.21	0.0-2.9	2.0-6.0	.43	.49	
	12-20	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.5-2.0	.55	.64	
	20-26	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.5-2.0	.55	.64	
	26-30	8-75	22-80	3-12	1.30-1.60	0.6-2	4.00-14.00	0.14-0.21	0.0-2.9	0.0-0.5	.64	.64	
	30-37	85-94	3-28	0-6	1.30-1.60	0.6-2	4.00-14.00	0.01-0.11	0.0-2.9	0.0-0.5	.10	.15	
	37-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.01-0.08	0.0-2.9	0.0-0.5	.17	.20	
33A:													
Hartford-----	0-8	53-70	27-40	3-7	1.25-1.50	2-6	14.00-42.00	0.10-0.12	0.0-2.9	2.0-5.0	.20	.28	3
	8-20	48-69	30-48	1-4	1.30-1.50	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.5	.24	.37	
	20-26	48-84	15-48	1-4	1.30-1.50	2-6	14.00-42.00	0.04-0.12	0.0-2.9	0.0-0.5	.28	.43	
	26-65	72-97	3-27	0-1	1.40-1.60	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.28	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
33B:													
Hartford-----	0-8	53-70	27-40	3-7	1.25-1.50	2-6	14.00-42.00	0.10-0.12	0.0-2.9	2.0-5.0	.20	.28	3
	8-20	48-69	30-48	1-4	1.30-1.50	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.5	.24	.37	
	20-26	48-84	15-48	1-4	1.30-1.50	2-6	14.00-42.00	0.04-0.12	0.0-2.9	0.0-0.5	.28	.43	
	26-65	72-97	3-27	0-1	1.40-1.60	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.28	
34A:													
Merrimac-----	0-9	45-70	27-48	3-7	1.10-1.20	2-6	14.00-42.00	0.10-0.12	0.0-2.9	1.0-5.0	.24	.28	3
	9-16	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.28	.37	
	16-24	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.0	.24	.32	
	24-60	88-100	0-9	0-3	1.30-1.50	6-100	42.00-703.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15	
34B:													
Merrimac-----	0-9	45-70	27-48	3-7	1.10-1.20	2-6	14.00-42.00	0.10-0.12	0.0-2.9	1.0-5.0	.24	.28	3
	9-16	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.28	.37	
	16-24	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.0	.24	.32	
	24-60	88-100	0-9	0-3	1.30-1.50	6-100	42.00-703.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15	
34C:													
Merrimac-----	0-9	45-70	27-48	3-7	1.10-1.20	2-6	14.00-42.00	0.10-0.12	0.0-2.9	1.0-5.0	.24	.28	3
	9-16	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.28	.37	
	16-24	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.0	.24	.32	
	24-60	88-100	0-9	0-3	1.30-1.50	6-100	42.00-703.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15	
35A:													
Penwood-----	0-8	74-83	14-25	1-3	1.45-1.60	6-100	42.00-703.00	0.05-0.08	0.0-2.9	2.0-4.0	.17	.20	3
	8-18	72-85	12-28	0-3	1.45-1.60	6-100	42.00-703.00	0.05-0.11	0.0-2.9	0.5-1.0	.24	.28	
	18-30	88-98	2-10	0-2	1.45-1.60	6-100	42.00-703.00	0.03-0.07	0.0-2.9	0.0-0.5	.15	.17	
	30-60	88-98	2-10	0-2	1.45-1.70	6-100	42.00-703.00	0.03-0.07	0.0-2.9	0.0-0.5	.15	.17	
35B:													
Penwood-----	0-8	74-83	14-25	1-3	1.45-1.60	6-100	42.00-703.00	0.05-0.08	0.0-2.9	2.0-4.0	.17	.20	3
	8-18	72-85	12-28	0-3	1.45-1.60	6-100	42.00-703.00	0.05-0.11	0.0-2.9	0.5-1.0	.24	.28	
	18-30	88-98	2-10	0-2	1.45-1.60	6-100	42.00-703.00	0.03-0.07	0.0-2.9	0.0-0.5	.15	.17	
	30-60	88-98	2-10	0-2	1.45-1.70	6-100	42.00-703.00	0.03-0.07	0.0-2.9	0.0-0.5	.15	.17	
36A:													
Windsor-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	2
	1-3	64-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.08	0.0-2.9	1.0-4.0	.17	.17	
	3-9	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10	
	9-21	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10	
	21-25	77-95	5-20	0-3	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.15	
	25-65	78-95	5-20	0-2	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.17	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors								
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct
36B:																			
Windsor-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	2						
	1-3	64-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.08	0.0-2.9	1.0-4.0	.17	.17							
	3-9	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10							
	9-21	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10							
	21-25	77-95	5-20	0-3	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.15							
	25-65	78-95	5-20	0-2	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.17							
36C:																			
Windsor-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	2						
	1-3	64-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.08	0.0-2.9	1.0-4.0	.17	.17							
	3-9	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10							
	9-21	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10							
	21-25	77-95	5-20	0-3	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.15							
	25-65	78-95	5-20	0-2	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.17							
37A:																			
Manchester-----	0-9	53-69	28-40	3-7	1.25-1.50	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-5.0	.15	.24	2						
	9-18	78-98	1-18	1-4	1.25-1.50	6-100	42.00-703.00	0.02-0.07	0.0-2.9	0.0-1.0	.15	.20							
	18-65	72-99	1-27	0-1	1.35-1.60	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.24							
37C:																			
Manchester-----	0-9	53-69	28-40	3-7	1.25-1.50	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-5.0	.15	.24	2						
	9-18	78-98	1-18	1-4	1.25-1.50	6-100	42.00-703.00	0.02-0.07	0.0-2.9	0.0-1.0	.15	.20							
	18-65	72-99	1-27	0-1	1.35-1.60	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.24							
37E:																			
Manchester-----	0-9	53-69	28-40	3-7	1.25-1.50	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-5.0	.15	.24	2						
	9-18	78-98	1-18	1-4	1.25-1.50	6-100	42.00-703.00	0.02-0.07	0.0-2.9	0.0-1.0	.15	.20							
	18-65	72-99	1-27	0-1	1.35-1.60	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.24							
38A:																			
Hinckley-----	0-8	54-69	27-38	4-8	0.90-1.10	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-7.0	.15	.28	2						
	8-20	75-83	12-24	1-5	1.20-1.40	6-20	42.00-141.00	0.03-0.10	0.0-2.9	0.5-1.5	.10	.17							
	20-27	87-93	2-12	1-5	1.20-1.40	6-20	42.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.05	.15							
	27-42	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28							
	42-60	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28							
38C:																			
Hinckley-----	0-8	54-69	27-38	4-8	0.90-1.10	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-7.0	.15	.28	2						
	8-20	75-83	12-24	1-5	1.20-1.40	6-20	42.00-141.00	0.03-0.10	0.0-2.9	0.5-1.5	.10	.17							
	20-27	87-93	2-12	1-5	1.20-1.40	6-20	42.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.05	.15							
	27-42	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28							
	42-60	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28							

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											In	Pct	Pct
38E: Hinckley-----	0-8	54-69	27-38	4-8	0.90-1.10	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-7.0	.15	.28	2
	8-20	75-83	12-24	1-5	1.20-1.40	6-20	42.00-141.00	0.03-0.10	0.0-2.9	0.5-1.5	.10	.17	
	20-27	87-93	2-12	1-5	1.20-1.40	6-20	42.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.05	.15	
	27-42	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28	
	42-60	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28	
39A: Groton-----	0-8	54-68	30-38	2-8	1.30-1.50	2-6	14.00-42.00	0.07-0.11	0.0-2.9	1.0-4.0	.10	.17	2
	8-18	50-68	30-42	2-8	1.30-1.50	2-6	14.00-42.00	0.04-0.12	0.0-2.9	0.5-1.0	.17	.37	
	18-24	53-83	15-42	2-5	1.30-1.60	2-20	14.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.10	.24	
	24-30	55-83	15-40	2-5	1.30-1.60	6-20	42.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.10	.24	
	30-52	78-95	5-20	0-2	1.40-1.70	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.20	
	52-72	78-95	5-20	0-2	1.40-1.70	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.20	
39C: Groton-----	0-8	54-68	30-38	2-8	1.30-1.50	2-6	14.00-42.00	0.07-0.11	0.0-2.9	1.0-4.0	.10	.17	2
	8-18	50-68	30-42	2-8	1.30-1.50	2-6	14.00-42.00	0.04-0.12	0.0-2.9	0.5-1.0	.17	.37	
	18-24	53-83	15-42	2-5	1.30-1.60	2-20	14.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.10	.24	
	24-30	55-83	15-40	2-5	1.30-1.60	6-20	42.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.10	.24	
	30-52	78-95	5-20	0-2	1.40-1.70	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.20	
	52-72	78-95	5-20	0-2	1.40-1.70	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.20	
39E: Groton-----	0-8	54-68	30-38	2-8	1.30-1.50	2-6	14.00-42.00	0.07-0.11	0.0-2.9	1.0-4.0	.10	.17	2
	8-18	50-68	30-42	2-8	1.30-1.50	2-6	14.00-42.00	0.04-0.12	0.0-2.9	0.5-1.0	.17	.37	
	18-24	53-83	15-42	2-5	1.30-1.60	2-20	14.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.10	.24	
	24-30	55-83	15-40	2-5	1.30-1.60	6-20	42.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.10	.24	
	30-52	78-95	5-20	0-2	1.40-1.70	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.20	
	52-72	78-95	5-20	0-2	1.40-1.70	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.20	
40A: Ludlow-----	0-8	20-44	51-65	5-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-8.0	.28	.37	3
	8-20	25-60	35-60	5-15	1.20-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.5-2.0	.32	.43	
	20-26	25-60	35-60	5-15	1.25-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.0-1.0	.37	.49	
	26-65	25-60	35-60	5-15	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.16	0.0-2.9	0.0-0.5	.37	.55	
40B: Ludlow-----	0-8	20-44	51-65	5-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-8.0	.28	.37	3
	8-20	25-60	35-60	5-15	1.20-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.5-2.0	.32	.43	
	20-26	25-60	35-60	5-15	1.25-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.0-1.0	.37	.49	
	26-65	25-60	35-60	5-15	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.16	0.0-2.9	0.0-0.5	.37	.55	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
41B: Ludlow-----	0-8	20-44	51-65	5-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-8.0	.20	.37	3
	8-20	25-60	35-60	5-15	1.20-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.5-2.0	.32	.43	
	20-26	25-60	35-60	5-15	1.25-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.0-1.0	.37	.49	
	26-65	25-60	35-60	5-15	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.16	0.0-2.9	0.0-0.5	.37	.55	
42C: Ludlow-----	0-8	20-44	51-65	5-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-8.0	.17	.37	3
	8-20	25-60	35-60	5-15	1.20-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.5-2.0	.32	.43	
	20-26	25-60	35-60	5-15	1.25-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.0-1.0	.37	.49	
	26-65	25-60	35-60	5-15	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.16	0.0-2.9	0.0-0.5	.37	.55	
43A: Rainbow-----	0-6	13-44	51-75	5-12	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-6.0	.28	.37	3
	6-18	13-71	27-75	2-12	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-2.0	.43	.49	
	18-26	13-71	27-75	2-12	1.25-1.50	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-0.5	.49	.55	
	26-65	40-71	27-48	2-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.49	
43B: Rainbow-----	0-6	13-44	51-75	5-12	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-6.0	.28	.37	3
	6-18	13-71	27-75	2-12	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-2.0	.43	.49	
	18-26	13-71	27-75	2-12	1.25-1.50	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-0.5	.49	.55	
	26-65	40-71	27-48	2-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.49	
44B: Rainbow-----	0-6	13-44	51-75	5-12	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-6.0	.28	.37	3
	6-18	13-71	27-75	2-12	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-2.0	.43	.49	
	18-26	13-71	27-75	2-12	1.25-1.50	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-0.5	.49	.55	
	26-65	40-71	27-48	2-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.49	
45A: Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	3
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37	
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37	
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37	
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
45B: Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	3
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37	
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37	
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37	
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
45C:													
Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	3
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37	
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37	
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37	
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
46B:													
Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.15	.24	3
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37	
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37	
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37	
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
46C:													
Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.15	.24	3
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37	
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37	
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37	
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
47C:													
Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.10	.24	3
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37	
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37	
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37	
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43	
48B:													
Georgia-----	0-8	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-6.0	.32	.43	5
	8-14	17-55	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-2.0	.37	.49	
	14-24	17-55	40-65	5-18	1.30-1.50	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55	
	24-60	17-55	40-65	5-18	1.35-1.65	0.2-0.6	1.40-4.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55	
Amenia-----	0-9	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	2.0-6.0	.32	.43	5
	9-16	30-42	40-65	5-18	1.20-1.40	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.5-2.0	.43	.55	
	16-25	30-42	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.55	
	25-60	30-42	40-65	5-18	1.30-1.65	0.2-0.6	1.40-4.00	0.11-0.20	0.0-2.9	0.0-0.5	.43	.55	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors							
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in
48C:																		
Georgia-----	0-8	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-6.0	.32	.43	5					
	8-14	17-55	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-2.0	.37	.49						
	14-24	17-55	40-65	5-18	1.30-1.50	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55						
	24-60	17-55	40-65	5-18	1.35-1.65	0.2-0.6	1.40-4.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55						
Amenia-----	0-9	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	2.0-6.0	.32	.43	5					
	9-16	30-42	40-65	5-18	1.20-1.40	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.5-2.0	.43	.55						
	16-25	30-42	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.55						
	25-60	30-42	40-65	5-18	1.30-1.65	0.2-0.6	1.40-4.00	0.11-0.20	0.0-2.9	0.0-0.5	.43	.55						
49B:																		
Georgia-----	0-8	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-6.0	.32	.43	5					
	8-14	17-55	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-2.0	.37	.49						
	14-24	17-55	40-65	5-18	1.30-1.50	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55						
	24-60	17-55	40-65	5-18	1.35-1.65	0.2-0.6	1.40-4.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55						
Amenia-----	0-9	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	2.0-6.0	.24	.43	5					
	9-16	30-42	40-65	5-18	1.20-1.40	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.5-2.0	.43	.55						
	16-25	30-42	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.55						
	25-60	30-42	40-65	5-18	1.30-1.65	0.2-0.6	1.40-4.00	0.11-0.20	0.0-2.9	0.0-0.5	.43	.55						
49C:																		
Georgia-----	0-8	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-6.0	.32	.43	5					
	8-14	17-55	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-2.0	.37	.49						
	14-24	17-55	40-65	5-18	1.30-1.50	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55						
	24-60	17-55	40-65	5-18	1.35-1.65	0.2-0.6	1.40-4.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55						
Amenia-----	0-9	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	2.0-6.0	.24	.43	5					
	9-16	30-42	40-65	5-18	1.20-1.40	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.5-2.0	.43	.55						
	16-25	30-42	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.55						
	25-60	30-42	40-65	5-18	1.30-1.65	0.2-0.6	1.40-4.00	0.11-0.20	0.0-2.9	0.0-0.5	.43	.55						
50A:																		
Sutton-----	0-6	56-68	20-40	4-12	1.30-1.50	2-6	14.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.15	.20	5					
	6-12	43-68	28-45	4-12	1.35-1.55	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	1.0-3.0	.24	.37						
	12-24	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-2.0	.24	.37						
	24-28	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.0-1.0	.28	.37						
	28-36	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37						
	36-65	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37						

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
50B: Sutton-----	0-6	56-68	20-40	4-12	1.30-1.50	2-6	14.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.15	.20	5
	6-12	43-68	28-45	4-12	1.35-1.55	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	1.0-3.0	.24	.37	
	12-24	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-2.0	.24	.37	
	24-28	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.0-1.0	.28	.37	
	28-36	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37	
	36-65	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37	
51B: Sutton-----	0-6	56-68	20-40	4-12	1.30-1.50	2-6	14.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.15	.20	5
	6-12	43-68	28-45	4-12	1.35-1.55	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	1.0-3.0	.24	.37	
	12-24	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-2.0	.24	.37	
	24-28	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.0-1.0	.28	.37	
	28-36	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37	
	36-65	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37	
52C: Sutton-----	0-6	56-68	20-40	4-12	1.30-1.50	2-6	14.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.15	.20	5
	6-12	43-68	28-45	4-12	1.35-1.55	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	1.0-3.0	.24	.37	
	12-24	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-2.0	.24	.37	
	24-28	43-68	28-45	4-12	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.0-1.0	.28	.37	
	28-36	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37	
	36-65	58-70	20-40	2-10	1.45-1.65	2-6	14.00-42.00	0.07-0.14	0.0-2.9	0.0-0.5	.24	.37	
53A: Wapping-----	0-11	43-66	30-45	4-12	1.20-1.40	0.6-2	4.00-14.00	0.13-0.17	0.0-2.9	2.0-6.0	.32	.43	5
	11-16	33-66	30-55	2-12	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.5	.49	.55	
	16-20	35-68	30-55	2-10	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.0-1.0	.55	.64	
	20-28	62-74	18-36	1-8	1.45-1.60	2-6	14.00-42.00	0.07-0.13	0.0-2.9	0.0-0.5	.20	.37	
	28-36	79-84	10-20	1-8	1.45-1.65	2-20	14.00-141.00	0.04-0.13	0.0-2.9	0.0-0.5	.15	.24	
	36-80	79-84	10-20	1-6	1.45-1.65	2-20	14.00-141.00	0.03-0.13	0.0-2.9	0.0-0.5	.15	.24	
53B: Wapping-----	0-11	43-66	30-45	4-12	1.20-1.40	0.6-2	4.00-14.00	0.13-0.17	0.0-2.9	2.0-6.0	.32	.43	5
	11-16	33-66	30-55	2-12	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.5	.49	.55	
	16-20	35-68	30-55	2-10	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.0-1.0	.55	.64	
	20-28	62-74	18-36	1-8	1.45-1.60	2-6	14.00-42.00	0.07-0.13	0.0-2.9	0.0-0.5	.20	.37	
	28-36	79-84	10-20	1-8	1.45-1.65	2-20	14.00-141.00	0.04-0.13	0.0-2.9	0.0-0.5	.15	.24	
	36-80	79-84	10-20	1-6	1.45-1.65	2-20	14.00-141.00	0.03-0.13	0.0-2.9	0.0-0.5	.15	.24	
54B: Wapping-----	0-11	43-66	30-45	4-12	1.20-1.40	0.6-2	4.00-14.00	0.13-0.17	0.0-2.9	2.0-6.0	.32	.43	5
	11-16	33-66	30-55	2-12	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.5	.49	.55	
	16-20	35-68	30-55	2-10	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.0-1.0	.55	.64	
	20-28	62-74	18-36	1-8	1.45-1.60	2-6	14.00-42.00	0.07-0.13	0.0-2.9	0.0-0.5	.20	.37	
	28-36	79-84	10-20	1-8	1.45-1.65	2-20	14.00-141.00	0.04-0.13	0.0-2.9	0.0-0.5	.15	.24	
	36-80	79-84	10-20	1-6	1.45-1.65	2-20	14.00-141.00	0.03-0.13	0.0-2.9	0.0-0.5	.15	.24	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
55A: Watchaug-----	0-8	56-65	20-40	4-15	1.30-1.50	0.6-2	4.00-14.00	0.12-0.15	0.0-2.9	2.0-7.0	.17	.24	5
	8-18	30-66	30-55	4-15	1.40-1.65	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	1.0-2.0	.28	.43	
	18-24	30-66	30-55	4-15	1.40-1.65	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-1.0	.32	.49	
	24-65	60-70	15-40	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	
55B: Watchaug-----	0-8	56-65	20-40	4-15	1.30-1.50	0.6-2	4.00-14.00	0.12-0.15	0.0-2.9	2.0-7.0	.17	.24	5
	8-18	30-66	30-55	4-15	1.40-1.65	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	1.0-2.0	.28	.43	
	18-24	30-66	30-55	4-15	1.40-1.65	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-1.0	.32	.49	
	24-65	60-70	15-40	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	
56B: Watchaug-----	0-8	56-65	20-40	4-15	1.30-1.50	0.6-2	4.00-14.00	0.12-0.15	0.0-2.9	2.0-7.0	.17	.24	5
	8-18	30-66	30-55	4-15	1.40-1.65	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	1.0-2.0	.28	.43	
	18-24	30-66	30-55	4-15	1.40-1.65	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-1.0	.32	.49	
	24-65	60-70	15-40	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	
57B: Gloucester-----	0-4	54-69	30-38	1-8	1.00-1.30	6-20	42.00-141.00	0.09-0.13	0.0-2.9	2.0-5.0	.17	.24	2
	4-12	54-69	30-38	1-8	1.20-1.50	6-20	42.00-141.00	0.07-0.13	0.0-2.9	0.5-1.5	.15	.28	
	12-25	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	25-35	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	35-60	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
57C: Gloucester-----	0-4	54-69	30-38	1-8	1.00-1.30	6-20	42.00-141.00	0.09-0.13	0.0-2.9	2.0-5.0	.17	.24	2
	4-12	54-69	30-38	1-8	1.20-1.50	6-20	42.00-141.00	0.07-0.13	0.0-2.9	0.5-1.5	.15	.28	
	12-25	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	25-35	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	35-60	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
57D: Gloucester-----	0-4	54-69	30-38	1-8	1.00-1.30	6-20	42.00-141.00	0.09-0.13	0.0-2.9	2.0-5.0	.17	.24	2
	4-12	54-69	30-38	1-8	1.20-1.50	6-20	42.00-141.00	0.07-0.13	0.0-2.9	0.5-1.5	.15	.28	
	12-25	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	25-35	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	35-60	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
58B: Gloucester-----	0-4	54-69	30-38	1-8	1.00-1.30	6-20	42.00-141.00	0.09-0.13	0.0-2.9	2.0-5.0	.17	.24	2
	4-12	54-69	30-38	1-8	1.20-1.50	6-20	42.00-141.00	0.07-0.13	0.0-2.9	0.5-1.5	.15	.28	
	12-25	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	25-35	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	35-60	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
58C: Gloucester-----	0-4	54-69	30-38	1-8	1.00-1.30	6-20	42.00-141.00	0.09-0.13	0.0-2.9	2.0-5.0	.17	.24	2
	4-12	54-69	30-38	1-8	1.20-1.50	6-20	42.00-141.00	0.07-0.13	0.0-2.9	0.5-1.5	.15	.28	
	12-25	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	25-35	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	35-60	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
59C: Gloucester-----	0-4	54-69	30-38	1-8	1.00-1.30	6-20	42.00-141.00	0.09-0.13	0.0-2.9	2.0-5.0	.17	.24	2
	4-12	54-69	30-38	1-8	1.20-1.50	6-20	42.00-141.00	0.07-0.13	0.0-2.9	0.5-1.5	.15	.28	
	12-25	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	25-35	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	35-60	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
59D: Gloucester-----	0-4	54-69	30-38	1-8	1.00-1.30	6-20	42.00-141.00	0.09-0.13	0.0-2.9	2.0-5.0	.17	.24	2
	4-12	54-69	30-38	1-8	1.20-1.50	6-20	42.00-141.00	0.07-0.13	0.0-2.9	0.5-1.5	.15	.28	
	12-25	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	25-35	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
	35-60	85-87	8-15	0-5	1.50-1.75	6-20	42.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.05	.17	
60B: Canton-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-3	54-70	29-38	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.05	.24	
	3-15	43-70	29-45	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	15-24	43-70	29-45	1-8	1.25-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	24-30	43-70	29-45	1-8	1.30-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.32	.55	
	30-60	77-85	15-18	0-5	1.35-1.60	6-20	42.00-141.00	0.03-0.09	0.0-2.9	0.0-0.5	.17	.32	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
60C: Canton-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-3	54-70	29-38	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.05	.24	
	3-15	43-70	29-45	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	15-24	43-70	29-45	1-8	1.25-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	24-30	43-70	29-45	1-8	1.30-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.32	.55	
	30-60	77-85	15-18	0-5	1.35-1.60	6-20	42.00-141.00	0.03-0.09	0.0-2.9	0.0-0.5	.17	.32	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permeability	Saturated hydraulic conductivity	Available water capacity	Linear extensibility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
60C:													
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
60D:													
Canton-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-3	54-70	29-38	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.05	.24	
	3-15	43-70	29-45	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	15-24	43-70	29-45	1-8	1.25-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	24-30	43-70	29-45	1-8	1.30-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.32	.55	
	30-60	77-85	15-18	0-5	1.35-1.60	6-20	42.00-141.00	0.03-0.09	0.0-2.9	0.0-0.5	.17	.32	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
61B:													
Canton-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-3	54-70	29-38	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.05	.24	
	3-15	43-70	29-45	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	15-24	43-70	29-45	1-8	1.25-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	24-30	43-70	29-45	1-8	1.30-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.32	.55	
	30-60	77-85	15-18	0-5	1.35-1.60	6-20	42.00-141.00	0.03-0.09	0.0-2.9	0.0-0.5	.17	.32	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
61C:													
Canton-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-3	54-70	29-38	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.05	.24	
	3-15	43-70	29-45	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	15-24	43-70	29-45	1-8	1.25-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	24-30	43-70	29-45	1-8	1.30-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.32	.55	
	30-60	77-85	15-18	0-5	1.35-1.60	6-20	42.00-141.00	0.03-0.09	0.0-2.9	0.0-0.5	.17	.32	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
62C:													
Canton-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-3	54-70	29-38	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.05	.24	
	3-15	43-70	29-45	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	15-24	43-70	29-45	1-8	1.25-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	24-30	43-70	29-45	1-8	1.30-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.32	.55	
	30-60	77-85	15-18	0-5	1.35-1.60	6-20	42.00-141.00	0.03-0.09	0.0-2.9	0.0-0.5	.17	.32	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
62D:													
Canton-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	3
	1-3	54-70	29-38	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.13	0.0-2.9	2.0-6.0	.05	.24	
	3-15	43-70	29-45	1-8	1.25-1.45	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	15-24	43-70	29-45	1-8	1.25-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.5-1.0	.32	.55	
	24-30	43-70	29-45	1-8	1.30-1.50	2-6	14.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.32	.55	
	30-60	77-85	15-18	0-5	1.35-1.60	6-20	42.00-141.00	0.03-0.09	0.0-2.9	0.0-0.5	.17	.32	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
63B:													
Cheshire-----	0-8	53-71	25-32	4-15	1.25-1.50	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.24	5
	8-16	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.5	.32	.43	
	16-26	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.0	.32	.43	
	26-65	68-75	10-30	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	
63C:													
Cheshire-----	0-8	53-71	25-32	4-15	1.25-1.50	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.24	5
	8-16	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.5	.32	.43	
	16-26	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.0	.32	.43	
	26-65	68-75	10-30	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	
63D:													
Cheshire-----	0-8	53-71	25-32	4-15	1.25-1.50	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.24	5
	8-16	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.5	.32	.43	
	16-26	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.0	.32	.43	
	26-65	68-75	10-30	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
67C:													
Narragansett-----	0-6	10-45	51-80	4-10	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.28	.43	3
	6-15	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	1.0-2.0	.49	.55	
	15-24	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.5-1.0	.49	.55	
	24-28	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-1.0	.49	.55	
	28-60	73-92	8-25	0-2	1.40-1.70	2-20	14.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.20	.32	
68C:													
Narragansett-----	0-6	10-45	51-80	4-10	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.24	.43	3
	6-15	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	1.0-2.0	.49	.55	
	15-24	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.5-1.0	.49	.55	
	24-28	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-1.0	.49	.55	
	28-60	73-92	8-25	0-2	1.40-1.70	2-20	14.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.20	.32	
68D:													
Narragansett-----	0-6	10-45	51-80	4-10	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.24	.43	3
	6-15	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	1.0-2.0	.49	.55	
	15-24	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.5-1.0	.49	.55	
	24-28	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-1.0	.49	.55	
	28-60	73-92	8-25	0-2	1.40-1.70	2-20	14.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.20	.32	
69B:													
Yalesville-----	0-8	53-70	25-35	5-12	1.25-1.50	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-5.0	.24	.28	2
	8-14	37-70	25-48	5-15	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-2.0	.24	.37	
	14-25	37-70	25-48	5-15	1.40-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-1.5	.24	.37	
	25-36	37-70	25-48	5-15	1.40-1.60	2-6	14.00-42.00	0.05-0.16	0.0-2.9	0.0-1.0	.24	.37	
	36-80	---	---	---	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	
69C:													
Yalesville-----	0-8	53-70	25-35	5-12	1.25-1.50	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-5.0	.24	.28	2
	8-14	37-70	25-48	5-15	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-2.0	.24	.37	
	14-25	37-70	25-48	5-15	1.40-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-1.5	.24	.37	
	25-36	37-70	25-48	5-15	1.40-1.60	2-6	14.00-42.00	0.05-0.16	0.0-2.9	0.0-1.0	.24	.37	
	36-80	---	---	---	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	
70C:													
Branford-----	0-8	23-46	51-65	3-12	1.20-1.40	0.6-6	4.00-42.00	0.17-0.21	0.0-2.9	2.0-5.0	.43	.49	3
	8-18	23-69	28-65	3-12	1.25-1.45	0.6-6	4.00-42.00	0.11-0.21	0.0-2.9	0.5-1.5	.49	.55	
	18-24	23-69	28-65	3-12	1.30-1.50	0.6-6	4.00-42.00	0.09-0.21	0.0-2.9	0.0-0.5	.49	.64	
	24-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.20	
Holyoke-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1
	1-3	18-47	50-70	3-12	1.10-1.25	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	2.0-5.0	.32	.43	
	3-8	23-67	30-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	1.0-3.0	.20	.37	
	8-18	32-68	20-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	0.5-1.0	.28	.49	
	18-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
71C:													
Brookfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	5
	1-3	57-70	20-40	3-10	1.00-1.25	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-5.0	.17	.24	
	3-13	55-72	25-35	3-10	1.35-1.60	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	1.0-3.0	.20	.32	
	13-27	55-72	25-35	3-10	1.35-1.60	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.24	.37	
	27-60	55-72	25-35	3-10	1.40-1.65	0.6-6	4.00-42.00	0.05-0.14	0.0-2.9	0.0-0.5	.24	.43	
Brimfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	1
	1-3	57-70	20-40	3-10	1.10-1.35	0.6-6	4.23-42.34	0.12-0.14	0.0-2.9	2.0-5.0	.17	.24	
	3-6	50-72	25-40	3-10	1.30-1.55	0.6-6	4.23-42.34	0.10-0.17	0.0-2.9	1.0-3.0	.28	.37	
	6-17	50-72	25-40	3-10	1.30-1.55	0.6-6	4.23-42.34	0.08-0.17	0.0-2.9	0.5-1.0	.28	.43	
	17-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
71E:													
Brookfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	5
	1-3	57-70	20-40	3-10	1.00-1.25	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-5.0	.17	.24	
	3-13	55-72	25-35	3-10	1.35-1.60	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	1.0-3.0	.20	.32	
	13-27	55-72	25-35	3-10	1.35-1.60	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.24	.37	
	27-60	55-72	25-35	3-10	1.40-1.65	0.6-6	4.00-42.00	0.05-0.14	0.0-2.9	0.0-0.5	.24	.43	
Brimfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	1
	1-3	57-70	20-40	3-10	1.10-1.35	0.6-6	4.23-42.34	0.12-0.14	0.0-2.9	2.0-5.0	.17	.24	
	3-6	50-72	25-40	3-10	1.30-1.55	0.6-6	4.23-42.34	0.10-0.17	0.0-2.9	1.0-3.0	.28	.37	
	6-17	50-72	25-40	3-10	1.30-1.55	0.6-6	4.23-42.34	0.08-0.17	0.0-2.9	0.5-1.0	.28	.43	
	17-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
73C:													
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	2
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15	
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28	
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28	
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
73E:													
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37	
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43	
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43	
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43	
Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	2
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15	
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28	
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28	
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
74C:													
Narragansett-----	0-6	10-45	51-80	4-10	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.37	.43	3
	6-15	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	1.0-2.0	.49	.55	
	15-24	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.5-1.0	.49	.55	
	24-28	10-73	23-80	4-10	1.40-1.60	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.0-1.0	.49	.55	
	28-60	73-92	8-25	0-2	1.40-1.70	2-20	14.00-141.00	0.02-0.10	0.0-2.9	0.0-0.5	.20	.32	
Hollis-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	1
	1-6	54-70	27-36	3-10	1.10-1.40	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	2.0-6.0	.10	.20	
	6-9	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.5-2.0	.15	.37	
	9-15	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	15-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
75C:													
Hollis-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	1
	1-6	54-70	27-36	3-10	1.10-1.40	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	2.0-6.0	.10	.20	
	6-9	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.5-2.0	.15	.37	
	9-15	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	15-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	2
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15	
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28	
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28	
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
75E:													
Hollis-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	1
	1-6	54-70	27-36	3-10	1.10-1.40	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	2.0-6.0	.10	.20	
	6-9	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.5-2.0	.15	.37	
	9-15	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	15-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											In	Pct	Pct
75E: Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	2
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15	
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28	
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28	
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
76E: Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
Hollis-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	1
	1-6	54-70	27-36	3-10	1.10-1.40	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	2.0-6.0	.10	.20	
	6-9	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.5-2.0	.15	.37	
	9-15	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	15-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
76F: Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
Hollis-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	1
	1-6	54-70	27-36	3-10	1.10-1.40	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	2.0-6.0	.10	.20	
	6-9	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.5-2.0	.15	.37	
	9-15	53-69	30-39	1-8	1.30-1.55	0.6-6	4.00-42.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	15-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
77C: Cheshire-----	0-8	53-71	25-32	4-15	1.25-1.50	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.24	5
	8-16	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.5	.32	.43	
	16-26	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.0	.32	.43	
	26-65	68-75	10-30	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	
Holyoke-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1
	1-3	18-47	50-70	3-12	1.10-1.25	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	2.0-5.0	.32	.43	
	3-8	23-67	30-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	1.0-3.0	.20	.37	
	8-18	32-68	20-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	0.5-1.0	.28	.49	
	18-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
77D: Cheshire-----	0-8	53-71	25-32	4-15	1.25-1.50	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.24	5
	8-16	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.5	.32	.43	
	16-26	20-71	25-65	4-15	1.40-1.65	0.6-6	4.00-42.00	0.07-0.20	0.0-2.9	0.5-1.0	.32	.43	
	26-65	68-75	10-30	2-15	1.45-1.70	0.6-6	4.00-42.00	0.07-0.15	0.0-2.9	0.0-0.5	.20	.37	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors			
											Kw	Kf	T	
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct				
77D:														
Holyoke-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1	
	1-3	18-47	50-70	3-12	1.10-1.25	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	2.0-5.0	.32	.43		
	3-8	23-67	30-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	1.0-3.0	.20	.37		
	8-18	32-68	20-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	0.5-1.0	.28	.49		
	18-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---		
78C:														
Holyoke-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1	
	1-3	18-47	50-70	3-12	1.10-1.25	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	2.0-5.0	.32	.43		
	3-8	23-67	30-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	1.0-3.0	.20	.37		
	8-18	32-68	20-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	0.5-1.0	.28	.49		
	18-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---		
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1	
78E:														
Holyoke-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1	
	1-3	18-47	50-70	3-12	1.10-1.25	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	2.0-5.0	.32	.43		
	3-8	23-67	30-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	1.0-3.0	.20	.37		
	8-18	32-68	20-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	0.5-1.0	.28	.49		
	18-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---		
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1	
79E:														
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1	
Holyoke-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1	
	1-3	18-47	50-70	3-12	1.10-1.25	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	2.0-5.0	.32	.43		
	3-8	23-67	30-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	1.0-3.0	.20	.37		
	8-18	32-68	20-65	3-12	1.30-1.55	0.6-2	4.00-14.00	0.08-0.20	0.0-2.9	0.5-1.0	.28	.49		
	18-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---		
80B:														
Bernardston-----	0-8	20-43	50-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.16-0.19	0.0-2.9	2.0-5.0	.32	.43	3	
	8-14	20-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-2.0	.43	.64		
	14-24	30-58	35-55	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-1.0	.43	.64		
	24-26	30-58	35-55	7-15	1.30-1.55	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-0.5	.43	.64		
	26-60	30-58	35-55	7-15	1.75-1.90	0.0015-0.2	0.01-1.40	0.07-0.16	0.0-2.9	0.0-0.5	.43	.64		
80C:														
Bernardston-----	0-8	20-43	50-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.16-0.19	0.0-2.9	2.0-5.0	.32	.43	3	
	8-14	20-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-2.0	.43	.64		
	14-24	30-58	35-55	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-1.0	.43	.64		
	24-26	30-58	35-55	7-15	1.30-1.55	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-0.5	.43	.64		
	26-60	30-58	35-55	7-15	1.75-1.90	0.0015-0.2	0.01-1.40	0.07-0.16	0.0-2.9	0.0-0.5	.43	.64		

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
81C: Bernardston-----	0-8	20-43	50-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.16-0.19	0.0-2.9	2.0-5.0	.32	.43	3
	8-14	20-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-2.0	.43	.64	
	14-24	30-58	35-55	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-1.0	.43	.64	
	24-26	30-58	35-55	7-15	1.30-1.55	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-0.5	.43	.64	
	26-60	30-58	35-55	7-15	1.75-1.90	0.0015-0.2	0.01-1.40	0.07-0.16	0.0-2.9	0.0-0.5	.43	.64	
81D: Bernardston-----	0-8	20-43	50-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.16-0.19	0.0-2.9	2.0-5.0	.32	.43	3
	8-14	20-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-2.0	.43	.64	
	14-24	30-58	35-55	7-15	1.25-1.50	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-1.0	.43	.64	
	24-26	30-58	35-55	7-15	1.30-1.55	0.6-2	4.00-14.00	0.11-0.19	0.0-2.9	0.0-0.5	.43	.64	
	26-60	30-58	35-55	7-15	1.75-1.90	0.0015-0.2	0.01-1.40	0.07-0.16	0.0-2.9	0.0-0.5	.43	.64	
82B: Broadbrook-----	0-8	15-45	50-70	5-15	1.15-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	3
	8-14	15-73	22-70	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	1.0-2.0	.43	.55	
	14-25	15-73	22-70	5-15	1.35-1.60	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-1.0	.49	.64	
	25-65	53-70	28-35	2-12	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.55	
82C: Broadbrook-----	0-8	15-45	50-70	5-15	1.15-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	3
	8-14	15-73	22-70	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	1.0-2.0	.43	.55	
	14-25	15-73	22-70	5-15	1.35-1.60	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-1.0	.49	.64	
	25-65	53-70	28-35	2-12	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.55	
82D: Broadbrook-----	0-8	15-45	50-70	5-15	1.15-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	3
	8-14	15-73	22-70	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	1.0-2.0	.43	.55	
	14-25	15-73	22-70	5-15	1.35-1.60	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-1.0	.49	.64	
	25-65	53-70	28-35	2-12	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.55	
83B: Broadbrook-----	0-8	15-45	50-70	5-15	1.15-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	3
	8-14	15-73	22-70	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	1.0-2.0	.43	.55	
	14-25	15-73	22-70	5-15	1.35-1.60	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-1.0	.49	.64	
	25-65	53-70	28-35	2-12	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.55	
83C: Broadbrook-----	0-8	15-45	50-70	5-15	1.15-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	3
	8-14	15-73	22-70	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	1.0-2.0	.43	.55	
	14-25	15-73	22-70	5-15	1.35-1.60	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-1.0	.49	.64	
	25-65	53-70	28-35	2-12	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.55	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors														
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct	Kw	Kf	T		
84B:																									
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.20	.28	3												
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43													
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43													
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55													
Montauk-----	0-4	54-67	15-40	6-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	3												
	4-14	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-1.5	.24	.37													
	14-25	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37													
	25-39	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													
	39-60	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													
84C:																									
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.20	.28	3												
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43													
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43													
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55													
Montauk-----	0-4	54-67	15-40	6-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	3												
	4-14	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-1.5	.24	.37													
	14-25	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37													
	25-39	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													
	39-60	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													
84D:																									
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.20	.28	3												
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43													
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43													
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55													
Montauk-----	0-4	54-67	15-40	6-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	3												
	4-14	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-1.5	.24	.37													
	14-25	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37													
	25-39	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													
	39-60	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													
85B:																									
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.17	.28	3												
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43													
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43													
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55													
Montauk-----	0-4	54-67	15-40	6-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	3												
	4-14	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-1.5	.24	.37													
	14-25	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37													
	25-39	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													
	39-60	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32													

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors						
											In	Pct	Pct	Pct	Kw	Kf	T
85C:																	
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.17	.28	3				
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43					
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43					
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55					
Montauk-----	0-4	54-67	15-40	6-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	3				
	4-14	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-1.5	.24	.37					
	14-25	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37					
	25-39	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32					
	39-60	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32					
86C:																	
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.15	.28	3				
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43					
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43					
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55					
Montauk-----	0-4	54-67	15-40	6-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	3				
	4-14	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-1.5	.24	.37					
	14-25	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37					
	25-39	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32					
	39-60	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32					
86D:																	
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.15	.28	3				
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43					
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43					
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55					
Montauk-----	0-4	54-67	15-40	6-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	3				
	4-14	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-1.5	.24	.37					
	14-25	42-69	30-40	1-18	1.70-1.90	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37					
	25-39	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32					
	39-60	50-83	16-27	1-12	1.85-2.00	0.0015-0.2	0.01-1.40	0.02-0.08	0.0-2.9	0.0-0.5	.20	.32					
87B:																	
Wethersfield-----	0-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.28	.43	3				
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43					
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43					
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49					

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors			
											Kw	Kf	T	
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct				
87C:														
Wethersfield-----	0-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.28	.43	3	
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43		
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43		
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49		
87D:														
Wethersfield-----	0-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.28	.43	3	
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43		
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43		
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49		
88B:														
Wethersfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3	
	1-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.17	.43		
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43		
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43		
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49		
88C:														
Wethersfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3	
	1-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.17	.43		
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43		
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43		
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49		
89C:														
Wethersfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3	
	1-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.17	.43		
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43		
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43		
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49		
89D:														
Wethersfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3	
	1-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.17	.43		
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43		
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43		
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49		
90B:														
Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.24	.32	3	
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55		
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55		
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55		
	42-48	35-55	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49		
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49		

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
90C: Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.24	.32	3
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55	
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
90D: Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.24	.32	3
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55	
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
91B: Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.17	.32	3
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55	
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
91C: Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.17	.32	3
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55	
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
91D: Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.17	.32	3
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55	
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
92B: Nellis-----	0-8	52-60	30-35	5-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-6.0	.17	.24	3
	8-14	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.43	
	14-25	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.49	
	25-27	22-65	30-60	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-0.5	.32	.43	
	27-60	32-62	35-50	3-18	1.70-1.80	0.6-6	4.00-42.00	0.07-0.19	0.0-2.9	0.0-1.0	.28	.49	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
92C:													
Nellis-----	0-8	52-60	30-35	5-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-6.0	.17	.24	3
	8-14	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.43	
	14-25	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.49	
	25-27	22-65	30-60	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-0.5	.32	.43	
	27-60	32-62	35-50	3-18	1.70-1.80	0.6-6	4.00-42.00	0.07-0.19	0.0-2.9	0.0-1.0	.28	.49	
92D:													
Nellis-----	0-8	52-60	30-35	5-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-6.0	.17	.24	3
	8-14	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.43	
	14-25	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.49	
	25-27	22-65	30-60	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-0.5	.32	.43	
	27-60	32-62	35-50	3-18	1.70-1.80	0.6-6	4.00-42.00	0.07-0.19	0.0-2.9	0.0-1.0	.28	.49	
93C:													
Nellis-----	0-8	52-60	30-35	5-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-6.0	.10	.24	3
	8-14	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.43	
	14-25	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.49	
	25-27	22-65	30-60	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-0.5	.32	.43	
	27-60	32-62	35-50	3-18	1.70-1.80	0.6-6	4.00-42.00	0.07-0.19	0.0-2.9	0.0-1.0	.28	.49	
94C:													
Farmington-----	0-3	60-74	10-35	5-16	1.30-1.50	0.6-6	4.00-42.00	0.11-0.14	0.0-2.9	2.0-5.0	.17	.24	1
	3-8	29-60	35-55	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.5-1.5	.28	.43	
	8-17	24-55	35-60	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.28	.49	
	17-80	---	---	---	---	0.0000-0.6	0.00-4.00	---	---	---	---	---	
Nellis-----	0-8	52-60	30-35	5-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-6.0	.17	.24	3
	8-14	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.43	
	14-25	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.49	
	25-27	22-65	30-60	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-0.5	.32	.43	
	27-60	32-62	35-50	3-18	1.70-1.80	0.6-6	4.00-42.00	0.07-0.19	0.0-2.9	0.0-1.0	.28	.49	
94E:													
Farmington-----	0-3	60-74	10-35	5-16	1.30-1.50	0.6-6	4.00-42.00	0.11-0.14	0.0-2.9	2.0-5.0	.17	.24	1
	3-8	29-60	35-55	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.5-1.5	.28	.43	
	8-17	24-55	35-60	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.28	.49	
	17-80	---	---	---	---	0.0000-0.6	0.00-4.00	---	---	---	---	---	
Nellis-----	0-8	52-60	30-35	5-18	1.30-1.60	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-6.0	.17	.24	3
	8-14	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.43	
	14-25	17-65	30-65	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.32	.49	
	25-27	22-65	30-60	5-18	1.40-1.70	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-0.5	.32	.43	
	27-60	32-62	35-50	3-18	1.70-1.80	0.6-6	4.00-42.00	0.07-0.19	0.0-2.9	0.0-1.0	.28	.49	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors							
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in
95C:																		
Farmington-----	0-3	60-74	10-35	5-16	1.30-1.50	0.6-6	4.00-42.00	0.11-0.14	0.0-2.9	2.0-5.0	.17	.24	1					
	3-8	29-60	35-55	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.5-1.5	.28	.43						
	8-17	24-55	35-60	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.28	.49						
	17-80	---	---	---	---	0.0000-0.6	0.00-4.00	---	---	---	---	---						
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1					
95E:																		
Farmington-----	0-3	60-74	10-35	5-16	1.30-1.50	0.6-6	4.00-42.00	0.11-0.14	0.0-2.9	2.0-5.0	.17	.24	1					
	3-8	29-60	35-55	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.5-1.5	.28	.43						
	8-17	24-55	35-60	5-16	1.30-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-1.0	.28	.49						
	17-80	---	---	---	---	0.0000-0.6	0.00-4.00	---	---	---	---	---						
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1					
96:																		
Ipswich-----	0-16	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.07-0.35	0.0-20.0	50-80	---	---	3					
	16-23	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.07-0.35	0.0-20.0	50-80	---	---						
	23-64	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.07-0.35	0.0-20.0	50-80	---	---						
	64-80	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.07-0.35	0.0-20.0	55-75	---	---						
97:																		
Pawcatuck-----	0-12	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.05-0.09	0.0-20.0	20-90	---	---	2					
	12-40	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.05-0.09	0.0-20.0	20-80	---	---						
	40-46	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.05-0.09	0.0-20.0	20-50	---	---						
	46-50	25-69	30-65	1-10	1.40-1.65	0.6-20	4.00-141.00	0.02-0.20	0.0-2.9	1.0-15	.20	.28						
	50-60	70-95	5-25	0-2	1.45-1.70	20-100	141.00-703.00	0.01-0.11	0.0-2.9	0.0-2.0	.20	.28						
98:																		
Westbrook-----	0-10	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.05-0.09	0.0-20.0	20-90	---	---	2					
	10-40	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.05-0.09	0.0-20.0	20-90	---	---						
	40-48	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.05-0.09	0.0-20.0	15-40	---	---						
	48-64	2-60	40-80	2-35	1.25-1.50	0.0015-2	0.01-14.00	0.02-0.07	0.0-6.0	8.0-15	.20	.20						
	64-99	2-60	40-80	2-35	1.25-1.50	0.0015-2	0.01-14.00	0.02-0.07	0.0-6.0	5.0-15	.24	.24						
99:																		
Westbrook, low salt--	0-10	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.07-0.27	0.0-20.0	20-90	---	---	2					
	10-40	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.07-0.27	0.0-20.0	20-90	---	---						
	40-48	0-0	0-0	0-0	0.30-0.55	0.6-20	4.00-141.00	0.07-0.27	0.0-20.0	15-40	---	---						
	48-64	2-60	40-80	2-35	1.25-1.50	0.0015-2	0.01-14.00	0.11-0.21	0.0-6.0	8.0-15	.20	.20						
	64-99	2-60	40-80	2-35	1.25-1.50	0.0015-2	0.01-14.00	0.11-0.21	0.0-6.0	5.0-15	.24	.24						

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
100: Suncook-----	0-7	75-100	15-22	1-3	1.10-1.30	6-100	42.00-703.00	0.08-0.11	0.0-2.9	2.0-5.0	.28	.28	5
	7-15	75-100	0-22	0-3	1.20-1.50	6-100	42.00-703.00	0.03-0.11	0.0-2.9	0.0-1.0	.28	.28	
	15-22	75-100	0-22	0-3	1.20-1.50	6-100	42.00-703.00	0.03-0.11	0.0-2.9	0.0-0.5	.28	.28	
	22-32	75-100	0-22	0-3	1.20-1.50	6-100	42.00-703.00	0.03-0.11	0.0-2.9	0.0-0.5	.28	.28	
	32-42	75-100	0-22	0-3	1.20-1.50	6-100	42.00-703.00	0.03-0.11	0.0-2.9	0.0-0.5	.28	.28	
	42-65	75-100	0-22	0-3	1.20-1.50	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.28	.32	
101: Occum-----	0-10	53-68	20-45	2-12	1.05-1.40	0.6-6	4.00-42.00	0.12-0.15	0.0-2.9	2.0-6.0	.28	.28	3
	10-17	43-70	28-45	2-12	1.20-1.50	0.6-6	4.00-42.00	0.10-0.18	0.0-2.9	0.5-1.5	.32	.37	
	17-28	50-62	30-48	2-8	1.20-1.50	0.6-6	4.00-42.00	0.10-0.15	0.0-2.9	0.0-0.5	.43	.49	
	28-32	77-100	0-18	0-5	1.30-1.60	6-100	42.00-703.00	0.01-0.11	0.0-2.9	0.0-0.5	.15	.17	
	32-42	77-100	0-18	0-5	1.30-1.60	6-100	42.00-703.00	0.01-0.11	0.0-2.9	0.0-0.5	.15	.17	
	42-65	77-100	0-18	0-5	1.30-1.60	6-100	42.00-703.00	0.01-0.11	0.0-2.9	0.0-0.5	.15	.17	
102: Pootatuck-----	0-4	54-69	25-40	2-6	1.10-1.35	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.24	.28	3
	4-16	49-69	30-45	1-6	1.20-1.45	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.37	.43	
	16-21	49-69	30-48	1-6	1.20-1.45	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.37	.43	
	21-29	49-69	30-48	1-6	1.20-1.45	0.6-6	4.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.37	.43	
	29-35	73-99	1-25	0-2	1.25-1.50	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.20	.24	
	35-40	88-100	0-10	0-2	1.25-1.50	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.20	.24	
	40-65	88-100	0-10	0-2	1.25-1.50	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.20	.24	
103: Rippowam-----	0-5	49-68	30-45	2-6	1.10-1.35	0.6-6	4.00-42.00	0.11-0.21	0.0-2.9	3.0-8.0	.15	.20	3
	5-12	49-69	30-45	1-6	1.20-1.45	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	1.0-3.0	.28	.37	
	12-19	49-69	30-45	1-6	1.20-1.45	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	0.5-1.0	.28	.37	
	19-24	49-84	15-45	1-6	1.20-1.45	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	0.0-1.0	.28	.37	
	24-27	49-84	15-45	1-6	1.20-1.45	0.6-6	4.00-42.00	0.09-0.18	0.0-2.9	0.0-0.5	.28	.37	
	27-31	73-100	0-25	0-2	1.25-1.50	6-100	42.00-703.00	0.01-0.10	0.0-2.9	0.0-0.5	.15	.17	
	31-65	73-100	0-25	0-2	1.25-1.50	6-100	42.00-703.00	0.01-0.10	0.0-2.9	0.0-0.5	.15	.17	
104: Bash-----	0-11	13-44	51-70	5-17	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	1.0-5.0	.43	.49	5
	11-21	13-70	25-70	5-17	1.20-1.55	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-2.0	.37	.43	
	21-28	13-70	25-70	5-17	1.25-1.55	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-2.0	.37	.49	
	28-60	13-69	30-70	1-17	1.25-1.55	0.2-2	1.40-14.00	0.08-0.20	0.0-2.9	0.0-0.5	.43	.55	
105: Hadley-----	0-12	15-46	52-70	5-15	1.20-1.40	0.6-2	4.00-14.00	0.19-0.21	0.0-2.9	2.0-5.0	.43	.43	5
	12-29	28-88	10-60	2-12	1.25-1.50	0.6-6	4.00-42.00	0.16-0.20	0.0-2.9	0.0-2.0	.55	.55	
	29-40	28-88	10-60	2-12	1.25-1.50	0.6-6	4.00-42.00	0.16-0.20	0.0-2.9	0.0-2.0	.55	.55	
	40-45	28-88	10-60	2-12	1.25-1.50	0.6-20	4.00-141.00	0.04-0.20	0.0-2.9	0.0-2.0	.37	.37	
	45-60	28-88	10-60	2-12	1.25-1.50	0.6-20	4.00-141.00	0.04-0.20	0.0-2.9	0.0-2.0	.37	.37	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
106: Winooski-----	0-12	2-44	51-80	5-18	1.15-1.35	0.6-2	4.00-14.00	0.19-0.21	0.0-2.9	2.0-5.0	.43	.64	5
	12-18	10-84	14-80	2-10	1.20-1.50	0.6-6	4.00-42.00	0.10-0.21	0.0-2.9	0.0-0.5	.43	.64	
	18-36	10-84	14-80	2-10	1.20-1.50	0.6-6	4.00-42.00	0.10-0.21	0.0-2.9	0.0-0.5	.43	.64	
	36-52	10-84	14-80	2-10	1.20-1.50	0.6-6	4.00-42.00	0.10-0.21	0.0-2.9	0.0-0.5	.43	.64	
	52-65	10-84	14-80	2-10	1.20-1.50	0.6-6	4.00-42.00	0.10-0.21	0.0-2.9	0.0-0.5	.43	.64	
107: Limerick-----	0-8	15-56	50-75	7-10	1.20-1.40	0.6-2	4.00-14.00	0.19-0.21	0.0-2.9	2.0-5.0	.49	.49	5
	8-20	15-58	40-75	2-10	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
	20-36	15-58	40-75	2-10	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
	36-54	15-58	40-75	2-10	1.20-1.40	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
	54-65	17-59	40-75	1-8	1.20-1.50	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.64	.64	
Lim-----	0-6	52-67	30-36	3-12	1.20-1.40	0.6-2	4.00-14.00	0.15-0.17	0.0-2.9	2.0-8.0	.28	.32	4
	6-11	5-55	42-80	3-15	1.20-1.50	0.6-2	4.00-14.00	0.15-0.19	0.0-2.9	0.0-0.5	.64	.64	
	11-15	5-55	42-80	3-15	1.20-1.50	0.6-2	4.00-14.00	0.15-0.19	0.0-2.9	0.0-0.5	.64	.64	
	15-22	5-55	42-80	3-15	1.20-1.50	0.6-2	4.00-14.00	0.15-0.19	0.0-2.9	0.0-0.5	.64	.64	
	22-29	37-77	20-51	3-12	1.20-1.50	0.6-2	4.00-14.00	0.09-0.15	0.0-2.9	0.0-0.5	.24	.37	
	29-42	75-98	2-22	0-3	1.30-1.60	6-100	42.00-703.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
	42-50	75-98	2-22	0-3	1.30-1.60	6-100	42.00-703.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
	50-57	75-98	2-22	0-3	1.30-1.60	6-100	42.00-703.00	0.02-0.08	0.0-2.9	0.0-0.5	.20	.24	
	57-65	75-98	2-22	0-3	1.30-1.60	6-100	42.00-703.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
108: Saco-----	0-12	13-41	52-70	7-17	1.20-1.40	0.6-2	4.00-14.00	0.19-0.21	0.0-2.9	3.0-8.0	.28	.32	4
	12-32	13-66	30-70	4-17	1.25-1.50	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-2.0	.49	.49	
	32-48	13-66	30-70	4-17	1.25-1.50	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.0-0.5	.49	.49	
	48-60	80-98	1-12	1-8	1.30-1.60	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.10	.15	
109: Fluvaquents, Frequently Flooded--	0-4	20-43	50-65	7-15	1.00-1.70	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.32	.37	2
	4-14	30-92	5-55	3-15	1.10-1.70	0.6-100	4.00-703.00	0.01-0.21	0.0-2.9	0.5-1.5	.28	.32	
	14-21	30-92	5-55	3-15	1.20-2.00	0.6-100	4.00-703.00	0.01-0.21	0.0-2.9	0.0-1.0	.28	.37	
	21-38	20-80	15-65	5-15	1.00-1.70	0.6-100	4.00-703.00	0.03-0.21	0.0-2.9	1.0-4.0	.24	.37	
	38-45	20-80	15-65	5-15	1.00-1.70	0.6-100	4.00-703.00	0.03-0.21	0.0-2.9	1.0-3.0	.28	.37	
	45-55	30-92	5-55	3-15	1.20-2.00	0.6-100	4.00-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.28	.37	
	55-60	20-80	15-65	5-15	1.00-1.70	0.6-100	4.00-703.00	0.03-0.21	0.0-2.9	1.0-3.0	.28	.37	
Udifulvents, Frequently Flooded--	0-2	55-75	18-30	7-15	1.20-1.45	4-36	28.00-254.00	0.11-0.15	0.0-2.9	2.0-5.0	.15	.15	2
	2-4	30-92	5-55	3-15	1.30-1.60	0.6-100	4.00-703.00	0.01-0.21	0.0-2.9	0.0-1.0	.28	.37	
	4-12	20-80	15-65	5-15	1.25-1.60	0.6-100	4.00-703.00	0.03-0.21	0.0-2.9	1.0-4.0	.24	.37	
	12-18	20-80	15-65	5-15	1.30-1.60	0.6-100	4.00-703.00	0.03-0.21	0.0-2.9	0.5-2.0	.32	.43	
	18-35	30-92	5-55	3-15	1.40-1.70	0.6-100	4.00-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.28	.37	
	35-38	30-92	5-55	3-15	1.40-1.70	0.6-100	4.00-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.28	.37	
	38-60	30-92	5-55	3-15	1.40-1.70	0.6-100	4.00-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.28	.37	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors							
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in
221A:																		
Ninigret-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.32	.37	3					
	8-16	28-70	27-60	3-12	1.35-1.60	0.6-6	4.00-42.00	0.13-0.20	0.0-2.9	0.5-1.5	.43	.49						
	16-26	28-70	27-60	3-12	1.35-1.60	0.6-6	4.00-42.00	0.13-0.20	0.0-2.9	0.0-0.5	.49	.55						
	26-65	73-100	0-25	0-2	1.45-1.70	6-100	42.00-703.00	0.01-0.11	0.0-2.9	0.0-0.5	.15	.17						
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					
224A:																		
Deerfield-----	0-8	73-83	10-20	2-7	1.45-1.65	6-20	42.00-141.00	0.09-0.11	0.0-2.9	1.0-4.0	.15	.15	2					
	8-16	81-94	5-12	1-7	1.45-1.65	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.15	.20						
	16-28	81-94	5-12	1-7	1.40-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.0-0.5	.17	.20						
	28-34	80-100	0-15	0-5	1.40-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.20	.24						
	34-60	80-100	0-15	0-5	1.40-1.60	6-100	42.00-703.00	0.03-0.08	0.0-2.9	0.0-0.0	.15	.15						
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					
225B:																		
Brancroft-----	0-6	2-32	50-74	18-24	1.20-1.40	0.2-2	1.40-14.00	0.19-0.21	0.0-6.0	3.0-6.0	.28	.28	5					
	6-17	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	1.0-2.0	.43	.43						
	17-22	10-15	50-72	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.19-0.21	0.0-6.0	0.5-1.0	.49	.49						
	22-32	10-15	50-72	18-35	1.40-1.65	0.0015-0.2	0.01-1.40	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49						
	32-43	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49						
	43-66	10-15	50-72	18-35	1.50-1.65	0.0015-0.06	0.01-0.42	0.19-0.21	0.0-6.0	0.0-0.5	.49	.49						
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					
226B:																		
Berlin-----	0-6	6-32	50-70	18-24	1.20-1.40	0.2-2	1.40-14.00	0.17-0.21	0.0-6.0	3.0-6.0	.28	.37	5					
	6-12	12-15	50-70	18-35	1.20-1.40	0.06-0.6	0.42-4.00	0.17-0.21	0.0-6.0	1.0-2.0	.43	.49						
	12-20	12-15	50-70	18-35	1.25-1.45	0.06-0.6	0.42-4.00	0.17-0.21	0.0-6.0	0.5-1.0	.43	.49						
	20-34	5-17	50-65	18-45	1.30-1.50	0.0015-0.2	0.01-1.40	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49						
	34-48	5-17	50-65	18-45	1.50-1.65	0.0015-0.06	0.01-0.42	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49						
	48-65	5-17	50-65	18-45	1.50-1.65	0.0015-0.06	0.01-0.42	0.15-0.21	0.0-6.0	0.0-0.5	.43	.49						
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					
228B:																		
Elmridge-----	0-6	54-68	30-38	2-8	1.30-1.50	2-6	14.00-42.00	0.13-0.15	0.0-2.9	2.0-6.0	.24	.28	3					
	6-10	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-1.5	.37	.37						
	10-18	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.0-1.0	.43	.43						
	18-25	47-68	30-45	2-10	1.35-1.60	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.0-0.5	.43	.43						
	25-65	10-20	20-55	35-60	1.55-1.80	0.0015-0.06	0.01-0.42	0.14-0.20	3.0-9.0	0.0-0.5	.32	.32						
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
229B:													
Agawam-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43	
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55	
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
229C:													
Agawam-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43	
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55	
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
230B:													
Branford-----	0-8	23-46	51-65	3-12	1.20-1.40	0.6-6	4.00-42.00	0.17-0.21	0.0-2.9	2.0-5.0	.43	.49	3
	8-18	23-69	28-65	3-12	1.25-1.45	0.6-6	4.00-42.00	0.11-0.21	0.0-2.9	0.5-1.5	.49	.55	
	18-24	23-69	28-65	3-12	1.30-1.50	0.6-6	4.00-42.00	0.09-0.21	0.0-2.9	0.0-0.5	.49	.64	
	24-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.20	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
230C:													
Branford-----	0-8	23-46	51-65	3-12	1.20-1.40	0.6-6	4.00-42.00	0.17-0.21	0.0-2.9	2.0-5.0	.43	.49	3
	8-18	23-69	28-65	3-12	1.25-1.45	0.6-6	4.00-42.00	0.11-0.21	0.0-2.9	0.5-1.5	.49	.55	
	18-24	23-69	28-65	3-12	1.30-1.50	0.6-6	4.00-42.00	0.09-0.21	0.0-2.9	0.0-0.5	.49	.64	
	24-65	73-100	0-25	0-2	1.40-1.65	6-100	42.00-703.00	0.02-0.10	0.0-2.9	0.0-0.5	.15	.20	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
232B:													
Haven-----	0-7	2-44	51-80	5-18	1.10-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	2.0-6.0	.32	.43	3
	7-14	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-2.0	.49	.64	
	14-20	2-44	25-80	5-18	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-2.9	0.5-1.0	.49	.64	
	20-24	54-70	25-28	5-18	1.25-1.50	0.6-2	4.00-14.00	0.13-0.17	0.0-2.9	0.0-0.5	.37	.43	
	24-60	92-100	0-5	0-3	1.40-1.65	20-100	141.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.15	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
234B:													
Merrimac-----	0-9	45-70	27-48	3-7	1.10-1.20	2-6	14.00-42.00	0.10-0.12	0.0-2.9	1.0-5.0	.24	.28	3
	9-16	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.28	.37	
	16-24	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.0	.24	.32	
	24-60	88-100	0-9	0-3	1.30-1.50	6-100	42.00-703.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors			
											In	Pct	Pct	Pct
235B:														
Penwood-----	0-8	74-83	14-25	1-3	1.45-1.60	6-100	42.00-703.00	0.05-0.08	0.0-2.9	2.0-4.0	.17	.20	3	
	8-18	72-85	12-28	0-3	1.45-1.60	6-100	42.00-703.00	0.05-0.11	0.0-2.9	0.5-1.0	.24	.28		
	18-30	88-98	2-10	0-2	1.45-1.60	6-100	42.00-703.00	0.03-0.07	0.0-2.9	0.0-0.5	.15	.17		
	30-60	88-98	2-10	0-2	1.45-1.70	6-100	42.00-703.00	0.03-0.07	0.0-2.9	0.0-0.5	.15	.17		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
236B:														
Windsor-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	45-95	---	---	2	
	1-3	64-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.08	0.0-2.9	1.0-4.0	.17	.17		
	3-9	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10		
	9-21	74-86	10-25	1-4	1.45-1.60	6-20	42.00-141.00	0.05-0.11	0.0-2.9	0.5-1.0	.10	.10		
	21-25	77-95	5-20	0-3	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.15		
	25-65	78-95	5-20	0-2	1.45-1.60	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-0.5	.15	.17		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
237A:														
Manchester-----	0-9	53-69	28-40	3-7	1.25-1.50	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-5.0	.15	.24	2	
	9-18	78-98	1-18	1-4	1.25-1.50	6-100	42.00-703.00	0.02-0.07	0.0-2.9	0.0-1.0	.15	.20		
	18-65	72-99	1-27	0-1	1.35-1.60	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.24		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
237C:														
Manchester-----	0-9	53-69	28-40	3-7	1.25-1.50	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-5.0	.15	.24	2	
	9-18	78-98	1-18	1-4	1.25-1.50	6-100	42.00-703.00	0.02-0.07	0.0-2.9	0.0-1.0	.15	.20		
	18-65	72-99	1-27	0-1	1.35-1.60	6-100	42.00-703.00	0.01-0.06	0.0-2.9	0.0-0.5	.10	.24		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
238A:														
Hinckley-----	0-8	54-69	27-38	4-8	0.90-1.10	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-7.0	.15	.28	2	
	8-20	75-83	12-24	1-5	1.20-1.40	6-20	42.00-141.00	0.03-0.10	0.0-2.9	0.5-1.5	.10	.17		
	20-27	87-93	2-12	1-5	1.20-1.40	6-20	42.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.05	.15		
	27-42	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28		
	42-60	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
238C:														
Hinckley-----	0-8	54-69	27-38	4-8	0.90-1.10	6-20	42.00-141.00	0.07-0.11	0.0-2.9	2.0-7.0	.15	.28	2	
	8-20	75-83	12-24	1-5	1.20-1.40	6-20	42.00-141.00	0.03-0.10	0.0-2.9	0.5-1.5	.10	.17		
	20-27	87-93	2-12	1-5	1.20-1.40	6-20	42.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.05	.15		
	27-42	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28		
	42-60	88-93	4-12	0-3	1.30-1.50	20-100	141.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.28		

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors			
											In	Pct	Pct	Pct
238C: Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	---	--
240B: Ludlow-----	0-8	20-44	51-65	5-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-8.0	.28	.37	3	
	8-20	25-60	35-60	5-15	1.20-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.5-2.0	.32	.43		
	20-26	25-60	35-60	5-15	1.25-1.45	0.6-2	4.00-14.00	0.10-0.20	0.0-2.9	0.0-1.0	.37	.49		
	26-65	25-60	35-60	5-15	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.16	0.0-2.9	0.0-0.5	.37	.55		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
243B: Rainbow-----	0-6	13-44	51-75	5-12	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-6.0	.28	.37	3	
	6-18	13-71	27-75	2-12	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-2.0	.43	.49		
	18-26	13-71	27-75	2-12	1.25-1.50	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-0.5	.49	.55		
	26-65	40-71	27-48	2-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.49		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
245B: Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	3	
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37		
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37		
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37		
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43		
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
245C: Woodbridge-----	0-7	57-68	20-40	3-12	1.00-1.25	0.6-2	4.00-14.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	3	
	7-18	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	1.0-3.0	.24	.37		
	18-26	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.0	.24	.37		
	26-30	48-70	27-40	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.24	.37		
	30-43	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43		
	43-65	48-70	27-40	3-12	1.70-2.00	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.43		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
248B: Georgia-----	0-8	25-31	51-70	5-18	1.20-1.40	0.6-2	4.00-14.00	0.17-0.20	0.0-2.9	2.0-6.0	.32	.43	5	
	8-14	17-55	40-65	5-18	1.25-1.45	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.5-2.0	.37	.49		
	14-24	17-55	40-65	5-18	1.30-1.50	0.6-2	4.00-14.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55		
	24-60	17-55	40-65	5-18	1.35-1.65	0.2-0.6	1.40-4.00	0.09-0.20	0.0-2.9	0.0-0.5	.37	.55		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth In	Sand Pct	Silt Pct	Clay Pct	Moist bulk density g/cc	Permea- bility In/hr	Saturated hydraulic conductivity um/sec	Available water capacity In/in	Linear extensi- bility Pct	Organic matter Pct	Erosion Factors			
											Kw	Kf	T	
269C:														
Yalesville-----	0-8	53-70	25-35	5-12	1.25-1.50	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-5.0	.24	.28	2	
	8-14	37-70	25-48	5-15	1.35-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-2.0	.24	.37		
	14-25	37-70	25-48	5-15	1.40-1.60	0.6-6	4.00-42.00	0.07-0.17	0.0-2.9	0.5-1.5	.24	.37		
	25-36	37-70	25-48	5-15	1.40-1.60	2-6	14.00-42.00	0.05-0.16	0.0-2.9	0.0-1.0	.24	.37		
	36-80	---	---	---	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
273C:														
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5	
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37		
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43		
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43		
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43		
Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	2	
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15		
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28		
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28		
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---		
273E:														
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
Charlton-----	0-4	57-72	20-40	3-8	1.25-1.45	0.6-6	4.00-42.00	0.12-0.14	0.0-2.9	2.0-6.0	.17	.24	5	
	4-7	57-72	20-40	3-8	1.30-1.45	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.5-1.0	.24	.37		
	7-19	57-72	20-40	3-8	1.35-1.50	0.6-6	4.00-42.00	0.09-0.14	0.0-2.9	0.0-0.5	.28	.43		
	19-27	57-72	20-40	3-8	1.35-1.55	0.6-6	4.00-42.00	0.08-0.14	0.0-2.9	0.0-0.5	.28	.43		
	27-65	57-72	20-40	1-8	1.35-1.60	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.28	.43		
Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	2	
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15		
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28		
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28		
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---		
275C:														
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--	
Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	2	
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15		
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28		
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28		
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---		

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors			
											In	Pct	Pct	Pct
275E:														
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	---	--
Chatfield-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	.05	.05	.28	2
	1-6	52-83	10-30	7-18	1.25-1.45	0.6-6	4.00-42.00	0.09-0.13	0.0-2.9	2.0-6.0	.10	.15	.28	
	6-15	37-83	10-45	7-18	1.30-1.45	0.6-6	4.00-42.00	0.08-0.17	0.0-2.9	0.5-2.0	.20	.28	.28	
	15-29	50-83	10-28	7-18	1.35-1.50	0.6-6	4.00-42.00	0.08-0.13	0.0-2.9	0.0-0.5	.20	.28	.28	
	29-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	---	
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	---	1
282B:														
Broadbrook-----	0-8	15-45	50-70	5-15	1.15-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	.55	3
	8-14	15-73	22-70	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	1.0-2.0	.43	.55	.55	
	14-25	15-73	22-70	5-15	1.35-1.60	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.0-1.0	.49	.64	.55	
	25-65	53-70	28-35	2-12	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.12	0.0-2.9	0.0-0.5	.32	.55	.55	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	---	--
284B:														
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.20	.28	.43	3
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43	.43	
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43	.43	
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55	.55	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	---	--
284C:														
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.20	.28	.43	3
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43	.43	
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43	.43	
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55	.55	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	---	--
284D:														
Paxton-----	0-8	53-70	27-35	3-12	1.00-1.25	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-5.0	.20	.28	.43	3
	8-15	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.5-1.5	.28	.43	.43	
	15-26	40-72	25-48	3-12	1.35-1.60	0.6-2	4.00-14.00	0.08-0.17	0.0-2.9	0.0-0.5	.28	.43	.43	
	26-65	40-72	25-48	3-12	1.70-2.00	0.0015-0.2	0.01-1.41	0.05-0.10	0.0-2.9	0.0-0.5	.37	.55	.55	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	---	--

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
287B:													
Wethersfield-----	0-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.28	.43	3
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43	
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43	
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
287C:													
Wethersfield-----	0-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.28	.43	3
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43	
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43	
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
287D:													
Wethersfield-----	0-3	37-51	42-48	7-15	1.10-1.30	0.6-2	4.00-14.00	0.14-0.17	0.0-2.9	2.0-5.0	.28	.43	3
	3-13	37-73	22-48	5-15	1.30-1.50	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.5	.32	.43	
	13-27	37-73	22-48	5-15	1.30-1.60	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.5-1.0	.32	.43	
	27-65	37-73	22-48	5-15	1.75-2.00	0.0015-0.2	0.01-1.40	0.08-0.17	0.0-2.9	0.0-0.5	.32	.49	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
290B:													
Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.24	.32	3
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55	
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--
290C:													
Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.24	.32	3
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55	
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55	
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors							
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in
290D:																		
Stockbridge-----	0-10	22-53	40-50	7-17	1.00-1.30	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.24	.32	3					
	10-20	17-53	40-65	7-17	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.0	.37	.55						
	20-28	17-53	40-65	7-17	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55						
	28-42	22-53	40-60	7-17	1.50-1.70	0.2-0.6	1.40-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.37	.55						
	42-48	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49						
	48-65	27-62	35-55	3-17	1.60-1.80	0.2-0.6	1.40-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49						
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					
301:																		
Beaches-----	0-65	---	---	0-1	1.40-1.70	20-100	141.14-705.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15	5					
Udipsamments-----	0-38	88-100	0-9	0-3	1.40-1.70	6-100	42.00-703.00	0.03-0.06	0.0-2.9	0.0-1.0	.15	.20	5					
	38-50	87-100	0-10	0-3	1.40-1.70	6-100	42.00-703.00	0.02-0.07	0.0-2.9	0.0-0.5	.10	.15						
	50-65	87-100	0-10	0-3	1.40-1.70	6-100	42.00-703.00	0.02-0.07	0.0-2.9	0.0-0.5	.10	.15						
302:																		
Dumps-----	0-65	---	---	0-10	---	0.2-2	1.40-14.00	0.00-0.00	---	0.0-2.0	.17	.20	--					
303:																		
Pits, quarries-----	0-1	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					
304:																		
Udorthents-----	0-5	35-50	43-50	7-15	1.00-1.70	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.28	.37	5					
	5-21	10-95	2-60	3-30	1.10-1.70	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28						
	21-80	10-95	2-60	3-30	1.20-2.00	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28						
305:																		
Udorthents-----	0-5	35-50	43-50	7-15	1.00-1.70	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.28	.37	3					
	5-21	10-95	2-60	3-30	1.10-1.70	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28						
	21-80	10-95	2-60	3-30	1.20-2.00	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28						
Pits-----	0-65	---	---	0-1	1.40-1.70	6-20	42.34-141.14	0.01-0.02	0.0-2.9	0.0-0.0	.02	.10	--					
306:																		
Udorthents-----	0-5	35-50	43-50	7-15	1.00-1.70	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.28	.37	3					
	5-21	10-95	2-60	3-30	1.10-1.70	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28						
	21-80	10-95	2-60	3-30	1.20-2.00	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28						
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					
307:																		
Urban land-----	0-6	---	---	0-0	---	0.01-20	0.07-141.00	0.00-0.00	---	---	---	---	--					

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors								
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct
308:																			
Udorthents-----	0-5	35-50	43-50	7-15	1.00-1.70	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.28	.37	3						
	5-21	10-95	2-60	3-30	1.10-1.70	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28							
	21-80	10-95	2-60	3-30	1.20-2.00	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28							
309:																			
Udorthents-----	0-5	35-50	43-50	7-15	1.00-1.70	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.28	.37	5						
	5-21	10-95	2-60	3-30	1.10-1.70	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28							
	21-80	10-95	2-60	3-30	1.20-2.00	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28							
310:																			
Udorthents, Periodically Flooded	0-5	35-50	43-50	7-15	1.00-1.70	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-6.0	.28	.37	5						
	5-21	10-95	2-60	3-30	1.10-1.70	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28							
	21-80	10-95	2-60	3-30	1.20-2.00	0.0015-100	0.01-703.00	0.01-0.21	0.0-2.9	0.0-0.5	.24	.28							
401C:																			
Macomber-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	2						
	1-2	35-53	43-50	7-15	1.20-1.40	0.6-2	4.00-14.00	0.08-0.12	0.0-2.9	2.0-5.0	.15	.32							
	2-10	30-53	43-55	7-15	1.20-1.40	0.6-2	4.00-14.00	0.07-0.14	0.0-2.9	0.5-2.0	.17	.49							
	10-21	30-53	43-55	7-15	1.20-1.40	0.6-2	4.00-14.00	0.07-0.14	0.0-2.9	0.0-1.0	.17	.49							
	21-30	30-53	43-55	7-15	1.20-1.40	0.6-2	4.00-14.00	0.05-0.13	0.0-2.9	0.0-0.5	.15	.49							
	30-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---							
Taconic-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1						
	1-4	37-58	35-48	7-15	1.10-1.40	0.6-6	4.00-42.00	0.08-0.12	0.0-2.9	2.0-5.0	.10	.28							
	4-11	25-58	35-60	7-15	1.20-1.50	0.6-6	4.00-42.00	0.05-0.15	0.0-2.9	0.1-2.0	.15	.43							
	11-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---							
402D:																			
Macomber-----	0-1	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	2						
	1-2	35-53	43-50	7-15	1.20-1.40	0.6-2	4.00-14.00	0.08-0.12	0.0-2.9	2.0-5.0	.15	.32							
	2-10	30-53	43-55	7-15	1.20-1.40	0.6-2	4.00-14.00	0.07-0.14	0.0-2.9	0.5-2.0	.17	.49							
	10-21	30-53	43-55	7-15	1.20-1.40	0.6-2	4.00-14.00	0.07-0.14	0.0-2.9	0.0-1.0	.17	.49							
	21-30	30-53	43-55	7-15	1.20-1.40	0.6-2	4.00-14.00	0.05-0.13	0.0-2.9	0.0-0.5	.15	.49							
	30-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---							
Taconic-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1						
	1-4	37-58	35-48	7-15	1.10-1.40	0.6-6	4.00-42.00	0.08-0.12	0.0-2.9	2.0-5.0	.10	.28							
	4-11	25-58	35-60	7-15	1.20-1.50	0.6-6	4.00-42.00	0.05-0.15	0.0-2.9	0.1-2.0	.15	.43							
	11-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---							
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1						

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors							
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in
403C:																		
Taconic-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1					
	1-4	37-58	35-48	7-15	1.10-1.40	0.6-6	4.00-42.00	0.08-0.12	0.0-2.9	2.0-5.0	.10	.28						
	4-11	25-58	35-60	7-15	1.20-1.50	0.6-6	4.00-42.00	0.05-0.15	0.0-2.9	0.1-2.0	.15	.43						
	11-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---						
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1					
403E:																		
Taconic-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1					
	1-4	37-58	35-48	7-15	1.10-1.40	0.6-6	4.00-42.00	0.08-0.12	0.0-2.9	2.0-5.0	.10	.28						
	4-11	25-58	35-60	7-15	1.20-1.50	0.6-6	4.00-42.00	0.05-0.15	0.0-2.9	0.1-2.0	.15	.43						
	11-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---						
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1					
403F:																		
Taconic-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	1					
	1-4	37-58	35-48	7-15	1.10-1.40	0.6-6	4.00-42.00	0.08-0.12	0.0-2.9	2.0-5.0	.10	.28						
	4-11	25-58	35-60	7-15	1.20-1.50	0.6-6	4.00-42.00	0.05-0.15	0.0-2.9	0.1-2.0	.15	.43						
	11-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---						
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1					
405C:																		
Dummerston-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5					
	1-2	37-58	35-48	7-15	1.00-1.20	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-6.0	.10	.28						
	2-3	25-58	35-60	7-15	1.00-1.20	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.0	.24	.43						
	3-4	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-2.0	.24	.43						
	4-6	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.5	.24	.49						
	6-11	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.5	.24	.49						
	11-22	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.0	.28	.49						
	22-27	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.0-0.5	.28	.49						
	27-40	43-58	35-45	7-12	1.40-1.60	0.6-2	4.00-14.00	0.08-0.14	0.0-2.9	0.0-0.5	.24	.49						
	40-64	43-58	35-45	7-12	1.40-1.60	0.6-2	4.00-14.00	0.08-0.14	0.0-2.9	0.0-0.5	.24	.49						
405E:																		
Dummerston-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5					
	1-2	37-58	35-48	7-15	1.00-1.20	0.6-2	4.00-14.00	0.11-0.14	0.0-2.9	2.0-6.0	.10	.28						
	2-3	25-58	35-60	7-15	1.00-1.20	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.0	.24	.43						
	3-4	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-2.0	.24	.43						
	4-6	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.5	.24	.49						
	6-11	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.5	.24	.49						
	11-22	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.17	0.0-2.9	0.5-1.0	.28	.49						
	22-27	25-58	35-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.10-0.17	0.0-2.9	0.0-0.5	.28	.49						
	27-40	43-58	35-45	7-12	1.40-1.60	0.6-2	4.00-14.00	0.08-0.14	0.0-2.9	0.0-0.5	.24	.49						
	40-64	43-58	35-45	7-12	1.40-1.60	0.6-2	4.00-14.00	0.08-0.14	0.0-2.9	0.0-0.5	.24	.49						

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											In	Pct	Pct
407C: Lanesboro-----	0-3	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	3
	3-6	37-58	35-48	7-15	1.00-1.20	0.6-2	4.00-14.00	0.15-0.18	0.0-2.9	2.0-5.0	.17	.32	
	6-8	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.5-2.0	.28	.49	
	8-16	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.5-1.5	.28	.49	
	16-22	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.0-1.0	.28	.49	
	22-30	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.0-0.5	.28	.49	
	30-60	40-60	35-48	5-12	1.75-1.90	0.0015-0.2	0.01-1.40	0.01-0.08	0.0-2.9	0.0-0.5	.24	.55	
407E: Lanesboro-----	0-3	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	3
	3-6	37-58	35-48	7-15	1.00-1.20	0.6-2	4.00-14.00	0.15-0.18	0.0-2.9	2.0-5.0	.17	.32	
	6-8	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.5-2.0	.28	.49	
	8-16	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.5-1.5	.28	.49	
	16-22	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.0-1.0	.28	.49	
	22-30	25-58	35-60	7-15	1.25-1.50	0.6-2	4.00-14.00	0.08-0.19	0.0-2.9	0.0-0.5	.28	.49	
	30-60	40-60	35-48	5-12	1.75-1.90	0.0015-0.2	0.01-1.40	0.01-0.08	0.0-2.9	0.0-0.5	.24	.55	
408C: Fullam-----	0-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	2-4	25-43	50-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.28	.43	
	4-10	25-50	43-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.21	0.0-2.9	0.5-2.0	.32	.49	
	10-20	25-50	43-60	7-15	1.20-1.40	0.6-2	4.00-14.00	0.11-0.18	0.0-2.9	0.5-1.0	.28	.49	
	20-49	40-60	35-48	5-12	1.80-2.00	0.0015-0.2	0.01-1.40	0.08-0.14	0.0-2.9	0.0-0.5	.28	.55	
	49-60	40-60	35-48	5-12	1.80-2.00	0.0015-0.2	0.01-1.40	0.08-0.14	0.0-2.9	0.0-0.5	.28	.55	
409B: Brayton-----	0-3	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	2
	3-6	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	20-60	---	---	
	6-7	31-47	50-60	3-9	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	4.0-8.0	.20	.24	
	7-9	36-50	43-55	7-9	1.20-1.40	0.6-6	4.00-42.00	0.07-0.14	0.0-2.9	0.0-2.0	.32	.49	
	9-13	46-62	35-45	3-9	1.25-1.50	0.6-6	4.00-42.00	0.06-0.12	0.0-2.9	0.0-2.0	.32	.49	
	13-18	46-62	35-45	3-9	1.65-1.90	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.37	.55	
	18-23	46-62	35-45	3-9	1.70-1.95	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.37	.55	
	23-60	46-62	35-45	3-9	1.70-1.95	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.37	.55	
412B: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
412C: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	
412D: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	
413C: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	
Millsite-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	2
	1-5	50-55	30-45	5-15	1.20-1.40	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	
	5-13	40-65	30-45	5-15	1.30-1.45	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.5-1.5	.24	.37	
	13-24	40-65	30-45	5-15	1.30-1.50	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-1.0	.24	.37	
	24-31	40-65	30-45	5-15	1.35-1.55	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-0.5	.24	.37	
	31-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
413E: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	
Millsite-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	2
	1-5	50-55	30-45	5-15	1.20-1.40	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	
	5-13	40-65	30-45	5-15	1.30-1.45	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.5-1.5	.24	.37	
	13-24	40-65	30-45	5-15	1.30-1.50	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-1.0	.24	.37	
	24-31	40-65	30-45	5-15	1.35-1.55	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-0.5	.24	.37	
	31-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											In	Pct	Pct
414: Fredon, cold-----	0-8	5-45	50-80	5-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	2.0-5.0	.37	.43	3
	8-17	20-65	30-65	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-2.0	.43	.49	
	17-24	20-65	30-65	5-15	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-1.0	.43	.55	
	24-29	75-93	5-18	2-7	1.35-1.55	2-20	14.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
	29-48	75-93	5-18	2-7	1.35-1.60	2-20	14.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
	48-60	75-93	5-18	2-7	1.40-1.60	2-20	14.00-141.00	0.02-0.08	0.0-2.9	0.0-0.5	.15	.20	
415C: Millsite-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	2
	1-5	50-55	30-45	5-15	1.20-1.40	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	
	5-13	40-65	30-45	5-15	1.30-1.45	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.5-1.5	.24	.37	
	13-24	40-65	30-45	5-15	1.30-1.50	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-1.0	.24	.37	
	24-31	40-65	30-45	5-15	1.35-1.55	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-0.5	.24	.37	
	31-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Westminster-----	0-1	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	25-50	---	---	1
	1-2	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	20-50	---	---	
	2-5	55-63	25-40	5-12	1.30-1.50	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.28	
	5-12	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	12-16	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	16-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
415E: Millsite-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	2
	1-5	50-55	30-45	5-15	1.20-1.40	0.6-6	4.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.24	
	5-13	40-65	30-45	5-15	1.30-1.45	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.5-1.5	.24	.37	
	13-24	40-65	30-45	5-15	1.30-1.50	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-1.0	.24	.37	
	24-31	40-65	30-45	5-15	1.35-1.55	0.6-6	4.00-42.00	0.07-0.16	0.0-2.9	0.0-0.5	.24	.37	
	31-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Westminster-----	0-1	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	25-50	---	---	1
	1-2	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	20-50	---	---	
	2-5	55-63	25-40	5-12	1.30-1.50	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.28	
	5-12	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	12-16	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	16-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
416E: Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
Westminster-----	0-1	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	25-50	---	---	1
	1-2	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	20-50	---	---	
	2-5	55-63	25-40	5-12	1.30-1.50	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.28	
	5-12	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	12-16	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	16-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
416F: Rock outcrop-----	---	---	---	---	---	---	---	---	---	---	---	---	1
Westminster-----	0-1	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	25-50	---	---	1
	1-2	0-0	0-0	0-0	0.30-0.55	20-100	141.00-703.00	0.08-0.40	0.0-20.0	20-50	---	---	
	2-5	55-63	25-40	5-12	1.30-1.50	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.20	.28	
	5-12	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	12-16	43-70	25-45	5-12	1.40-1.60	2-6	14.00-42.00	0.10-0.16	0.0-2.9	0.0-1.0	.32	.43	
	16-80	---	---	---	---	0.01-20	0.07-141.00	---	---	---	---	---	
417B: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	
417C: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	
417D: Bice-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5
	1-7	50-70	25-35	5-15	1.35-1.50	0.6-6	4.00-42.00	0.13-0.15	0.0-2.9	2.0-5.0	.15	.20	
	7-16	30-63	30-55	7-15	1.35-1.60	0.6-6	4.00-42.00	0.08-0.19	0.0-2.9	0.0-2.0	.28	.43	
	16-24	40-63	30-45	7-15	1.35-1.65	0.6-6	4.00-42.00	0.08-0.16	0.0-2.9	0.0-0.5	.28	.43	
	24-60	40-63	30-45	7-15	1.45-1.70	0.6-6	4.00-42.00	0.05-0.16	0.0-2.9	0.0-0.5	.28	.43	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
424B: Shelburne-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	1-2	55-67	30-30	3-15	1.20-1.40	0.6-6	4.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.20	.24	
	2-7	50-62	35-45	3-15	1.25-1.40	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	7-21	50-62	35-45	3-15	1.30-1.50	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	21-27	50-62	35-45	3-15	1.35-1.55	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-1.0	.24	.37	
	27-32	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	32-43	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	43-55	50-62	35-45	3-15	1.40-1.65	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	55-80	50-62	35-45	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
424C: Shelburne-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	1-2	55-67	30-30	3-15	1.20-1.40	0.6-6	4.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.20	.24	
	2-7	50-62	35-45	3-15	1.25-1.40	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	7-21	50-62	35-45	3-15	1.30-1.50	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	21-27	50-62	35-45	3-15	1.35-1.55	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-1.0	.24	.37	
	27-32	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	32-43	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	43-55	50-62	35-45	3-15	1.40-1.65	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	55-80	50-62	35-45	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
424D: Shelburne-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	1-2	55-67	30-30	3-15	1.20-1.40	0.6-6	4.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.20	.24	
	2-7	50-62	35-45	3-15	1.25-1.40	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	7-21	50-62	35-45	3-15	1.30-1.50	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	21-27	50-62	35-45	3-15	1.35-1.55	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-1.0	.24	.37	
	27-32	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	32-43	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	43-55	50-62	35-45	3-15	1.40-1.65	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	55-80	50-62	35-45	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
425B: Shelburne-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	1-2	55-67	30-30	3-15	1.20-1.40	0.6-6	4.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.20	.24	
	2-7	50-62	35-45	3-15	1.25-1.40	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	7-21	50-62	35-45	3-15	1.30-1.50	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	21-27	50-62	35-45	3-15	1.35-1.55	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-1.0	.24	.37	
	27-32	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	32-43	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	43-55	50-62	35-45	3-15	1.40-1.65	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	55-80	50-62	35-45	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
425C: Shelburne-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	1-2	55-67	30-30	3-15	1.20-1.40	0.6-6	4.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.20	.24	
	2-7	50-62	35-45	3-15	1.25-1.40	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	7-21	50-62	35-45	3-15	1.30-1.50	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	21-27	50-62	35-45	3-15	1.35-1.55	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-1.0	.24	.37	
	27-32	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	32-43	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	43-55	50-62	35-45	3-15	1.40-1.65	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	55-80	50-62	35-45	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
426D: Shelburne-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	3
	1-2	55-67	30-30	3-15	1.20-1.40	0.6-6	4.00-42.00	0.12-0.15	0.0-2.9	2.0-5.0	.20	.24	
	2-7	50-62	35-45	3-15	1.25-1.40	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	7-21	50-62	35-45	3-15	1.30-1.50	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-2.0	.24	.37	
	21-27	50-62	35-45	3-15	1.35-1.55	0.6-6	4.00-42.00	0.08-0.18	0.0-2.9	0.0-1.0	.24	.37	
	27-32	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	32-43	50-62	35-45	3-15	1.35-1.55	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	43-55	50-62	35-45	3-15	1.40-1.65	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
	55-80	50-62	35-45	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.04-0.08	0.0-2.9	0.0-0.5	.32	.49	
427B: Ashfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	20-95	---	---	3
	1-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	45-95	---	---	
	2-3	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	50-95	---	---	
	3-7	55-70	25-30	5-15	1.20-1.40	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.28	
	7-12	30-67	30-55	5-15	1.25-1.40	2-6	14.00-42.00	0.11-0.19	0.0-2.9	2.0-5.0	.28	.37	
	12-18	30-67	30-55	3-15	1.30-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-2.0	.37	.55	
	18-24	30-67	30-55	3-15	1.35-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-1.0	.37	.55	
	24-29	30-77	20-55	3-15	1.35-1.55	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.0-1.0	.37	.55	
	29-44	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	44-58	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	58-80	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
427C: Ashfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	20-95	---	---	3
	1-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	45-95	---	---	
	2-3	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	50-95	---	---	
	3-7	55-70	25-30	5-15	1.20-1.40	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.28	
	7-12	30-67	30-55	5-15	1.25-1.40	2-6	14.00-42.00	0.11-0.19	0.0-2.9	2.0-5.0	.28	.37	
	12-18	30-67	30-55	3-15	1.30-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-2.0	.37	.55	
	18-24	30-67	30-55	3-15	1.35-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-1.0	.37	.55	
	24-29	30-77	20-55	3-15	1.35-1.55	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.0-1.0	.37	.55	
	29-44	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	44-58	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	58-80	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth In	Sand Pct	Silt Pct	Clay Pct	Moist bulk density g/cc	Permea- bility In/hr	Saturated hydraulic conductivity um/sec	Available water capacity In/in	Linear extensi- bility Pct	Organic matter Pct	Erosion Factors		
											Kw	Kf	T
428A: Ashfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	20-95	---	---	3
	1-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	45-95	---	---	
	2-3	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	50-95	---	---	
	3-7	55-70	25-30	5-15	1.20-1.40	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.28	
	7-12	30-67	30-55	5-15	1.25-1.40	2-6	14.00-42.00	0.11-0.19	0.0-2.9	2.0-5.0	.28	.37	
	12-18	30-67	30-55	3-15	1.30-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-2.0	.37	.55	
	18-24	30-67	30-55	3-15	1.35-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-1.0	.37	.55	
	24-29	30-77	20-55	3-15	1.35-1.55	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.0-1.0	.37	.55	
	29-44	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	44-58	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	58-80	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
428B: Ashfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	20-95	---	---	3
	1-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	45-95	---	---	
	2-3	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	50-95	---	---	
	3-7	55-70	25-30	5-15	1.20-1.40	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.28	
	7-12	30-67	30-55	5-15	1.25-1.40	2-6	14.00-42.00	0.11-0.19	0.0-2.9	2.0-5.0	.28	.37	
	12-18	30-67	30-55	3-15	1.30-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-2.0	.37	.55	
	18-24	30-67	30-55	3-15	1.35-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-1.0	.37	.55	
	24-29	30-77	20-55	3-15	1.35-1.55	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.0-1.0	.37	.55	
	29-44	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	44-58	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	58-80	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
428C: Ashfield-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	20-95	---	---	3
	1-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	45-95	---	---	
	2-3	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	---	0.0-20.0	50-95	---	---	
	3-7	55-70	25-30	5-15	1.20-1.40	2-6	14.00-42.00	0.11-0.15	0.0-2.9	2.0-5.0	.17	.28	
	7-12	30-67	30-55	5-15	1.25-1.40	2-6	14.00-42.00	0.11-0.19	0.0-2.9	2.0-5.0	.28	.37	
	12-18	30-67	30-55	3-15	1.30-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-2.0	.37	.55	
	18-24	30-67	30-55	3-15	1.35-1.50	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.5-1.0	.37	.55	
	24-29	30-77	20-55	3-15	1.35-1.55	0.6-6	4.00-42.00	0.10-0.19	0.0-2.9	0.0-1.0	.37	.55	
	29-44	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	44-58	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
	58-80	30-77	20-55	3-15	1.60-2.10	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49	
429A: Agawam, cold-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43	
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55	
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors													
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct	Kw	Kf	T	
429B:																								
Agawam, cold-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3											
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43												
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55												
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17												
429C:																								
Agawam, cold-----	0-8	53-71	25-37	4-10	1.10-1.20	2-6	14.00-42.00	0.12-0.15	0.0-2.9	1.0-5.0	.28	.32	3											
	8-14	50-69	30-40	1-10	1.20-1.40	2-6	14.00-42.00	0.11-0.18	0.0-2.9	0.5-2.0	.37	.43												
	14-24	54-69	30-40	1-6	1.30-1.40	2-6	14.00-42.00	0.11-0.17	0.0-2.9	0.0-0.5	.32	.55												
	24-60	87-100	0-12	0-1	1.30-1.50	20-100	141.00-703.00	0.01-0.07	0.0-2.9	0.0-0.5	.15	.17												
433:																								
Moosilauke-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.06-0.50	0.0-20.0	45-70	---	---	3											
	1-6	40-51	42-50	7-10	1.30-1.55	2-6	14.00-42.00	0.09-0.15	0.0-2.9	0.5-2.0	.37	.43												
	6-16	52-67	30-40	3-8	1.30-1.55	2-6	14.00-42.00	0.08-0.15	0.0-2.9	0.0-1.0	.32	.37												
	16-24	52-67	30-40	0-8	1.40-1.65	6-100	42.00-703.00	0.08-0.15	0.0-2.9	0.0-0.5	.32	.37												
	24-39	80-95	5-15	0-5	1.40-1.65	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.17	.20												
	39-65	80-95	5-15	0-5	1.40-1.65	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.0-0.5	.17	.20												
434A:																								
Merrimac, cold-----	0-9	45-70	27-48	3-7	1.10-1.20	2-6	14.00-42.00	0.10-0.12	0.0-2.9	1.0-5.0	.24	.28	3											
	9-16	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.28	.37												
	16-24	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.0	.24	.32												
	24-60	88-100	0-9	0-3	1.30-1.50	6-100	42.00-703.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15												
434B:																								
Merrimac, cold-----	0-9	45-70	27-48	3-7	1.10-1.20	2-6	14.00-42.00	0.10-0.12	0.0-2.9	1.0-5.0	.24	.28	3											
	9-16	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.28	.37												
	16-24	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.0	.24	.32												
	24-60	88-100	0-9	0-3	1.30-1.50	6-100	42.00-703.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15												
434C:																								
Merrimac, cold-----	0-9	45-70	27-48	3-7	1.10-1.20	2-6	14.00-42.00	0.10-0.12	0.0-2.9	1.0-5.0	.24	.28	3											
	9-16	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.10-0.14	0.0-2.9	0.5-1.0	.28	.37												
	16-24	48-69	30-48	1-4	1.20-1.40	2-6	14.00-42.00	0.07-0.12	0.0-2.9	0.5-1.0	.24	.32												
	24-60	88-100	0-9	0-3	1.30-1.50	6-100	42.00-703.00	0.02-0.05	0.0-2.9	0.0-0.5	.10	.15												
435:																								
Scarboro-----	0-12	0-0	0-0	0-0	0.30-0.55	2-6	14.00-42.00	0.08-0.40	0.0-20.0	50-95	---	---	3											
	12-17	74-83	10-25	1-5	1.25-1.45	2-20	14.00-141.00	0.08-0.11	0.0-2.9	3.0-15	.05	.05												
	17-31	75-93	5-25	0-2	1.35-1.55	6-100	42.00-703.00	0.04-0.08	0.0-2.9	0.0-1.0	.24	.28												
	31-72	75-93	5-25	0-2	1.35-1.55	6-100	42.00-703.00	0.02-0.08	0.0-2.9	0.0-0.5	.24	.28												

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
436: Halsey-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	55-70	---	---	3
	1-8	18-42	52-65	6-17	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-2.9	3.0-8.0	.28	.32	
	8-16	18-54	40-65	6-17	1.25-1.45	0.6-2	4.00-14.00	0.12-0.21	0.0-2.9	0.5-1.5	.43	.55	
	16-28	50-58	30-45	5-12	1.25-1.50	2-6	14.00-42.00	0.10-0.17	0.0-2.9	0.5-1.5	.43	.49	
	28-38	80-95	3-12	2-8	1.40-1.60	6-20	42.00-141.00	0.02-0.07	0.0-2.9	0.0-0.5	.15	.24	
	38-60	80-95	3-12	2-8	1.40-1.60	6-20	42.00-141.00	0.02-0.07	0.0-2.9	0.0-0.5	.15	.24	
437: Wonsqueak-----	0-2	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.08-0.40	0.0-20.0	80-95	---	---	2
	2-11	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.08-0.40	0.0-20.0	80-95	---	---	
	11-22	0-0	0-0	0-0	0.30-0.55	0.6-100	4.00-703.00	0.08-0.40	0.0-20.0	80-95	---	---	
	22-25	30-65	28-52	5-18	1.50-1.70	0.2-2	1.40-14.00	0.06-0.16	0.0-2.9	0.0-2.0	.37	.43	
	28-45	30-65	28-52	5-18	1.50-1.70	0.2-2	1.40-14.00	0.06-0.16	0.0-2.9	0.0-2.0	.37	.43	
	45-60	30-65	28-52	5-18	1.50-1.70	0.2-2	1.40-14.00	0.06-0.16	0.0-2.9	0.0-2.0	.37	.43	
438: Bucksport-----	0-9	0-0	0-0	0-0	0.30-0.55	0.2-6	1.40-42.00	0.08-0.50	0.0-20.0	80-95	---	---	3
	9-33	0-0	0-0	0-0	0.30-0.55	0.2-6	1.40-42.00	0.08-0.50	0.0-20.0	80-95	---	---	
	33-50	0-0	0-0	0-0	0.30-0.55	0.2-6	1.40-42.00	0.08-0.50	0.0-20.0	80-95	---	---	
	50-59	0-0	0-0	0-0	0.30-0.55	0.2-6	1.40-42.00	0.08-0.50	0.0-20.0	80-95	---	---	
	59-63	34-93	5-48	2-18	1.50-1.70	0.2-100	1.40-703.00	0.02-0.16	0.0-2.9	0.0-2.0	.28	.37	
440A: Boscawen-----	0-1	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.07-0.11	0.0-20.0	55-75	---	---	2
	1-2	54-68	30-40	2-6	1.20-1.40	2-20	14.00-141.00	0.03-0.10	0.0-2.9	2.0-5.0	.15	.24	
	2-9	54-83	15-40	2-6	1.20-1.40	2-20	14.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.17	.37	
	9-16	77-85	15-17	2-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.05	.24	
	16-29	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	29-34	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	34-40	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	40-44	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	44-67	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
440C: Boscawen-----	0-1	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.07-0.11	0.0-20.0	55-75	---	---	2
	1-2	54-68	30-40	2-6	1.20-1.40	2-20	14.00-141.00	0.03-0.10	0.0-2.9	2.0-5.0	.15	.24	
	2-9	54-83	15-40	2-6	1.20-1.40	2-20	14.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.17	.37	
	9-16	77-85	15-17	2-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.05	.24	
	16-29	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	29-34	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	34-40	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	40-44	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	
	44-67	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15	

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors							
											In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in
440E:																		
Boscawen -----	0-1	0-0	0-0	0-0	0.30-0.55	2-20	14.00-141.00	0.07-0.11	0.0-20.0	55-75	---	---	2					
	1-2	54-68	30-40	2-6	1.20-1.40	2-20	14.00-141.00	0.03-0.10	0.0-2.9	2.0-5.0	.15	.24						
	2-9	54-83	15-40	2-6	1.20-1.40	2-20	14.00-141.00	0.02-0.05	0.0-2.9	0.0-0.5	.17	.37						
	9-16	77-85	15-17	2-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.05	.24						
	16-29	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15						
	29-34	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15						
	34-40	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15						
	40-44	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15						
	44-67	89-98	2-5	0-6	1.30-1.50	6-100	42.00-703.00	0.01-0.04	0.0-2.9	0.0-0.5	.10	.15						
442:																		
Brayton -----	0-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	2					
	2-10	35-51	42-50	7-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	4.0-8.0	.10	.37						
	10-17	30-70	25-55	5-15	1.20-1.40	0.6-2	4.00-14.00	0.07-0.14	0.0-2.9	0.0-2.0	.28	.43						
	17-22	30-70	25-55	5-15	1.25-1.50	0.6-6	4.00-42.00	0.06-0.12	0.0-2.9	0.0-2.0	.28	.43						
	22-27	30-70	25-55	5-15	1.25-1.50	0.6-6	4.00-42.00	0.06-0.12	0.0-2.9	0.0-0.8	.28	.43						
	27-42	35-70	25-50	5-15	1.65-1.90	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49						
	42-65	35-70	25-50	5-15	1.70-1.95	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49						
443:																		
Brayton -----	0-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	2					
	2-10	35-51	42-50	7-15	1.20-1.40	0.6-2	4.00-14.00	0.17-0.21	0.0-2.9	4.0-8.0	.10	.37						
	10-17	30-70	25-55	5-15	1.20-1.40	0.6-2	4.00-14.00	0.07-0.14	0.0-2.9	0.0-2.0	.28	.43						
	17-22	30-70	25-55	5-15	1.25-1.50	0.6-6	4.00-42.00	0.06-0.12	0.0-2.9	0.0-2.0	.28	.43						
	22-27	30-70	25-55	5-15	1.25-1.50	0.6-6	4.00-42.00	0.06-0.12	0.0-2.9	0.0-0.8	.28	.43						
	27-42	35-70	25-50	5-15	1.65-1.90	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49						
	42-65	35-70	25-50	5-15	1.70-1.95	0.0015-0.2	0.01-1.40	0.06-0.12	0.0-2.9	0.0-0.5	.32	.49						
Loonmeadow -----	0-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.08-0.40	0.0-20.0	50-80	---	---	5					
	2-9	57-68	20-40	3-12	1.30-1.45	2-6	14.00-42.00	0.12-0.15	0.0-2.9	8.0-15	.05	.10						
	9-18	57-68	20-40	3-12	1.40-1.80	2-6	14.00-42.00	0.09-0.14	0.0-2.9	0.0-1.0	.28	.37						
	18-35	57-70	20-40	3-10	1.45-2.00	0.2-6	1.40-42.00	0.05-0.14	0.0-2.9	0.0-1.0	.28	.43						
	35-80	57-70	20-40	3-10	1.45-2.10	0.06-20	0.42-141.00	0.05-0.14	0.0-2.9	0.0-1.0	.28	.43						
448B:																		
Hogansburg -----	0-12	29-47	35-45	18-26	1.00-1.40	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-5.0	.24	.32	5					
	12-20	23-55	30-55	15-22	1.40-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.5	.28	.43						
	20-29	25-63	30-55	7-20	1.50-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-1.0	.28	.43						
	29-43	25-63	30-55	7-20	1.50-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-1.0	.28	.43						
	43-50	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49						
	50-70	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49						
	70-84	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49						

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth In	Sand Pct	Silt Pct	Clay Pct	Moist bulk density g/cc	Permea- bility In/hr	Saturated hydraulic conductivity um/sec	Available water capacity In/in	Linear extensi- bility Pct	Organic matter Pct	Erosion Factors			
											Kw	Kf	T	
449B:														
Hogansburg-----	0-12	29-47	35-45	18-26	1.00-1.40	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-5.0	.24	.32	5	
	12-20	23-55	30-55	15-22	1.40-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.5	.28	.43		
	20-29	25-63	30-55	7-20	1.50-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-1.0	.28	.43		
	29-43	25-63	30-55	7-20	1.50-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-1.0	.28	.43		
	43-50	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49		
	50-70	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49		
	70-84	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49		
449C:														
Hogansburg-----	0-12	29-47	35-45	18-26	1.00-1.40	0.6-2	4.00-14.00	0.14-0.18	0.0-2.9	2.0-5.0	.24	.32	5	
	12-20	23-55	30-55	15-22	1.40-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.5-1.5	.28	.43		
	20-29	25-63	30-55	7-20	1.50-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-1.0	.28	.43		
	29-43	25-63	30-55	7-20	1.50-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-1.0	.28	.43		
	43-50	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49		
	50-70	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49		
	70-84	43-65	30-40	5-17	1.70-2.10	0.0015-0.2	0.01-1.40	0.05-0.10	0.0-2.9	0.0-0.5	.32	.49		
450B:														
Pyrities-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	3	
	1-8	23-52	30-50	18-27	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	2.0-5.0	.15	.43		
	8-13	25-58	30-50	12-25	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43		
	13-26	25-58	30-50	12-25	1.50-1.70	0.06-0.6	0.42-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43		
	26-45	25-58	30-50	12-25	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43		
	45-65	40-58	30-50	12-20	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49		
450C:														
Pyrities-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	3	
	1-8	23-52	30-50	18-27	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	2.0-5.0	.15	.43		
	8-13	25-58	30-50	12-25	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43		
	13-26	25-58	30-50	12-25	1.50-1.70	0.06-0.6	0.42-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43		
	26-45	25-58	30-50	12-25	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43		
	45-65	40-58	30-50	12-20	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49		
450D:														
Pyrities-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	3	
	1-8	23-52	30-50	18-27	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	2.0-5.0	.15	.43		
	8-13	25-58	30-50	12-25	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43		
	13-26	25-58	30-50	12-25	1.50-1.70	0.06-0.6	0.42-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43		
	26-45	25-58	30-50	12-25	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43		
	45-65	40-58	30-50	12-20	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49		

Table 24.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
451B: Pyrities-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	3
	1-8	23-52	30-50	18-27	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	2.0-5.0	.15	.43	
	8-13	25-58	30-50	12-25	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43	
	13-26	25-58	30-50	12-25	1.50-1.70	0.06-0.6	0.42-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43	
	26-45	25-58	30-50	12-25	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	45-65	40-58	30-50	12-20	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
451C: Pyrities-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	3
	1-8	23-52	30-50	18-27	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	2.0-5.0	.15	.43	
	8-13	25-58	30-50	12-25	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43	
	13-26	25-58	30-50	12-25	1.50-1.70	0.06-0.6	0.42-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43	
	26-45	25-58	30-50	12-25	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	45-65	40-58	30-50	12-20	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
451D: Pyrities-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	3
	1-8	23-52	30-50	18-27	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	2.0-5.0	.15	.43	
	8-13	25-58	30-50	12-25	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43	
	13-26	25-58	30-50	12-25	1.50-1.70	0.06-0.6	0.42-4.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43	
	26-45	25-58	30-50	12-25	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	45-65	40-58	30-50	12-20	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
457: Mudgepond-----	0-11	28-30	50-60	10-22	1.10-1.40	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	3.0-8.0	.20	.28	5
	11-16	28-73	20-55	7-17	1.20-1.50	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	16-26	28-75	20-55	5-17	1.30-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	26-35	28-75	20-55	5-17	1.40-1.60	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	35-65	38-72	25-45	3-17	1.50-1.80	0.6-2	4.00-14.00	0.07-0.17	0.0-2.9	0.0-1.0	.32	.49	
458: Mudgepond-----	0-11	28-30	50-60	10-22	1.10-1.40	0.6-2	4.00-14.00	0.16-0.20	0.0-2.9	3.0-8.0	.20	.28	5
	11-16	28-73	20-55	7-17	1.20-1.50	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	16-26	28-75	20-55	5-17	1.30-1.55	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	26-35	28-75	20-55	5-17	1.40-1.60	0.6-6	4.00-42.00	0.08-0.20	0.0-2.9	0.5-2.0	.32	.49	
	35-65	38-72	25-45	3-17	1.50-1.80	0.6-2	4.00-14.00	0.07-0.17	0.0-2.9	0.0-1.0	.32	.49	
Alden-----	0-4	9-33	52-65	15-26	1.20-1.40	0.6-2	4.00-14.00	0.16-0.21	0.0-5.9	10-15	.10	.15	5
	4-13	9-54	28-65	18-26	1.20-1.40	0.6-2	4.00-14.00	0.13-0.21	0.0-5.9	3.0-8.0	.28	.24	
	13-23	9-54	28-65	18-26	1.20-1.50	0.6-2	4.00-14.00	0.13-0.21	0.0-5.9	0.5-1.0	.28	.24	
	23-29	9-54	28-65	18-26	1.20-1.50	0.6-2	4.00-14.00	0.13-0.21	0.0-5.9	0.5-1.0	.28	.24	
	29-43	14-42	40-60	18-26	1.40-1.65	0.2-0.6	1.40-4.00	0.10-0.19	0.0-2.9	0.0-0.5	.32	.49	
	43-60	14-42	40-60	18-26	1.40-1.65	0.2-0.6	1.40-4.00	0.10-0.19	0.0-2.9	0.0-0.5	.32	.49	

Table 24.—Physical Properties of the Soils—Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility	Saturated hydraulic conductivity	Available water capacity	Linear extensi- bility	Organic matter	Erosion Factors		
											Kw	Kf	T
	In	Pct	Pct	Pct	g/cc	In/hr	um/sec	In/in	Pct	Pct			
501: Ondawa-----	0-1	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	3
	1-2	0-0	0-0	0-0	0.30-0.55	6-20	42.00-141.00	0.14-0.18	0.0-20.0	55-70	---	---	
	2-14	23-52	30-50	18-27	1.10-1.30	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	2.0-5.0	.15	.43	
	14-30	25-58	30-50	12-25	1.30-1.70	0.6-2	4.00-14.00	0.10-0.18	0.0-2.9	0.0-0.5	.28	.43	
	30-33	25-58	30-50	12-25	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.28	.43	
	33-60	40-58	30-50	12-20	1.60-1.80	0.06-0.6	0.42-4.00	0.06-0.18	0.0-2.9	0.0-0.5	.32	.49	
503: Rumney-----	0-7	10-45	50-65	5-25	1.05-1.60	0.6-6	4.00-42.00	0.15-0.20	0.0-2.9	2.0-5.0	.32	.37	3
	7-22	26-67	30-49	3-25	1.20-1.60	0.6-6	4.00-42.00	0.11-0.17	0.0-2.9	0.2-1.0	.37	.43	
	22-38	26-67	30-49	3-25	1.20-1.60	0.6-6	4.00-42.00	0.11-0.17	0.0-2.9	0.2-1.0	.37	.43	
	38-42	5-67	30-70	3-25	1.20-1.60	0.6-6	4.00-42.00	0.11-0.20	0.0-2.9	2.0-5.0	.28	.37	
	42-44	52-97	2-30	1-18	1.20-1.60	6-20	42.00-141.00	0.03-0.15	0.0-2.9	0.0-0.2	.15	.20	
	44-65	52-96	2-30	2-20	1.20-1.60	6-20	42.00-141.00	0.03-0.15	0.0-2.9	0.0-0.2	.15	.20	
508: Medomak-----	0-7	15-35	50-75	8-15	1.20-1.40	0.6-2	4.00-14.00	0.19-0.21	0.0-2.9	10-20	.10	.15	4
	7-24	15-60	25-75	8-15	1.20-1.40	0.6-2	4.00-14.00	0.19-0.21	0.0-2.9	0.5-6.0	.37	.43	
	24-33	10-35	50-75	8-15	1.25-1.50	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	10-20	.10	.15	
	33-46	10-95	5-75	0-15	1.25-1.50	0.6-2	4.00-14.00	0.15-0.21	0.0-2.9	0.2-1.0	.17	.20	
	46-79	10-95	5-75	0-15	1.30-1.60	6-100	42.00-703.00	0.02-0.11	0.0-2.9	0.2-1.0	.17	.20	

Table 25.—Chemical Properties of the Soils

(Absence of an entry indicates that data were not estimated.)

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
2:						
Ridgebury-----	0-5	1.7-5.6	---	4.5-6.0	0	0
	5-14	1.0-4.3	---	4.5-6.0	0	0
	14-21	1.0-4.3	---	4.5-6.0	0	0
	21-60	1.0-4.3	---	4.5-6.0	0	0
3:						
Ridgebury-----	0-5	1.7-5.6	---	4.5-6.0	0	0
	5-14	1.0-4.3	---	4.5-6.0	0	0
	14-21	1.0-4.3	---	4.5-6.0	0	0
	21-60	1.0-4.3	---	4.5-6.0	0	0
Leicester-----	0-1	---	30-63	4.5-5.0	0	0
	1-7	---	0.5-2.4	4.5-5.5	0	0
	7-10	---	0.6-2.9	4.5-5.5	0	0
	10-18	---	0.7-4.3	4.5-5.5	0	0
	18-24	---	0.7-4.3	4.5-5.5	0	0
	24-43	1.0-3.8	---	4.5-6.0	0	0
	43-65	1.0-3.8	---	4.5-6.0	0	0
Whitman-----	0-1	---	17-99	4.5-6.0	0	0
	1-9	2.8-6.7	---	4.5-6.5	0	0
	9-16	1.0-6.5	---	4.5-6.5	0	0
	16-22	1.0-4.3	---	4.5-6.5	0	0
	22-60	1.0-4.3	---	4.5-6.5	0	0
4:						
Leicester-----	0-7	---	0.5-2.4	4.5-5.5	0	0
	7-10	---	0.6-2.9	4.5-5.5	0	0
	10-18	---	0.7-4.3	4.5-5.5	0	0
	18-24	---	0.7-4.3	4.5-5.5	0	0
	24-43	1.0-3.8	---	4.5-6.0	0	0
	43-65	1.0-3.8	---	4.5-6.0	0	0
5:						
Wilbraham-----	0-4	---	0.5-3.8	4.5-6.0	0	0
	4-8	---	0.7-6.8	4.5-6.0	0	0
	8-20	---	0.7-6.8	4.5-6.0	0	0
	20-65	---	0.7-6.8	4.5-6.0	0	0
6:						
Wilbraham-----	0-4	---	0.5-3.8	4.5-6.0	0	0
	4-8	---	0.7-6.8	4.5-6.0	0	0
	8-20	---	0.7-6.8	4.5-6.0	0	0
	20-65	---	0.7-6.8	4.5-6.0	0	0
Menlo-----	0-5	---	17-65	4.5-6.0	0	0
	5-16	3.9-8.4	---	4.5-7.3	0	0
	16-22	1.6-8.2	---	5.1-7.8	0	0
	22-27	1.6-8.1	---	5.1-7.8	0	0
	27-40	1.6-8.0	---	5.1-7.8	0	0
	40-60	1.6-8.0	---	5.1-8.4	0	0
7:						
Mudgepond-----	0-11	9.3-20	---	6.6-7.8	0	0
	11-16	6.3-15	---	6.6-7.8	0-2	0
	16-26	4.6-15	---	6.6-7.8	0-15	0
	26-35	4.6-15	---	6.6-7.8	1-15	0
	35-65	2.6-14	---	6.6-8.4	5-25	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
8:						
Mudgepond-----	0-11	9.3-20	---	6.6-7.8	0	0
	11-16	6.3-15	---	6.6-7.8	0-2	0
	16-26	4.6-15	---	6.6-7.8	0-15	0
	26-35	4.6-15	---	6.6-7.8	1-15	0
	35-65	2.6-14	---	6.6-8.4	5-25	0
Alden-----	0-4	8.4-14	---	5.1-7.3	0	0
	4-13	9.8-14	---	5.1-7.3	0	0
	13-23	9.6-14	---	5.6-7.3	0	0
	23-29	9.6-14	---	5.6-7.3	0	0
	29-43	9.1-14	---	6.1-7.3	0	0
	43-60	9.1-14	---	6.1-8.4	0-15	0
9:						
Scitico-----	0-8	3.7-9.7	---	4.5-7.3	0	0
	8-11	6.7-19	---	5.1-7.3	0	0
	11-18	6.7-19	---	5.1-7.3	0	0
	18-30	12-21	---	5.6-7.8	0	0
	30-38	12-21	---	5.6-7.8	0	0
	38-52	12-21	---	5.6-7.8	0	0
	52-65	12-21	---	5.6-7.8	0	0
Shaker-----	0-2	---	32-108	4.5-6.5	0	0
	2-6	0.8-3.1	---	5.1-7.3	0	0
	6-20	0.8-3.0	---	5.1-7.3	0	0
	20-30	0.7-2.9	---	5.1-7.3	0	0
	30-65	12-21	---	5.6-7.8	0	0
Maybid-----	0-9	7.4-9.8	---	5.1-6.0	0	0
	9-18	9.6-20	---	5.1-7.3	0	0
	18-26	9.6-19	---	5.1-7.3	0	0
	26-36	12-19	---	5.1-7.3	0	0
	36-60	12-19	---	5.1-7.3	0	0
10:						
Raynham-----	0-10	2.8-8.3	---	5.1-7.3	0	0
	10-16	2.7-8.2	---	5.1-7.3	0	0
	16-26	2.7-8.1	---	5.1-7.3	0	0
	26-34	2.6-8.0	---	5.1-7.3	0	0
	34-47	2.6-8.0	---	5.6-7.8	0-5	0
	47-60	2.6-8.0	---	5.6-7.8	0-5	0
12:						
Raypol-----	0-8	---	0.5-3.1	4.5-5.5	0	0
	8-12	---	0.6-3.6	4.5-5.5	0	0
	12-20	---	0.7-5.3	4.5-5.5	0	0
	20-26	---	0.7-5.3	4.5-5.5	0	0
	26-29	---	0.7-5.3	4.5-5.5	0	0
	29-52	---	0.0-0.7	4.5-5.5	0	0
	52-65	0.0-1.1	---	4.5-6.5	0	0
13:						
Walpole-----	0-1	---	30-162	4.5-7.3	0	0
	1-7	1.9-5.4	---	4.5-7.3	0	0
	7-21	1.2-4.9	---	4.5-7.3	0	0
	21-25	0.0-1.9	---	4.5-7.3	0	0
	25-41	0.0-1.9	---	4.5-7.3	0	0
	41-65	0.0-1.9	---	4.5-7.3	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
14: Fredon-----	0-8	2.8-8.3	---	6.1-7.3	0	0
	8-17	2.7-8.2	---	6.1-7.3	0	0
	17-24	2.7-8.1	---	6.1-7.3	0	0
	24-29	1.0-3.8	---	5.6-8.4	0	0
	29-48	1.0-3.8	---	5.6-8.4	0-5	0
	48-60	1.0-3.8	---	5.6-8.4	0-10	0
15: Scarboro-----	0-12	---	32-99	4.5-6.0	0	0
	12-17	---	1.0-6.9	4.5-6.0	0	0
	17-31	---	0.1-3.6	4.5-6.0	0	0
	31-72	0.0-1.9	---	4.5-7.3	0	0
16: Halsey-----	0-1	81-104	---	5.6-7.3	0	0
	1-8	3.3-9.4	---	5.6-7.3	0	0
	8-16	3.2-9.2	---	5.6-7.3	0	0
	16-28	2.7-6.5	---	5.6-7.3	0	0
	28-38	1.0-4.3	---	6.1-8.4	0	0
	38-60	1.0-4.3	---	6.1-8.4	0-10	0
17: Timakwa-----	0-10	---	3.4-85	1.8-6.0	0	0
	10-21	---	3.4-85	1.8-6.0	0	0
	21-24	---	3.4-85	1.8-6.0	0	0
	24-37	---	3.4-85	1.8-6.0	0	0
	37-47	0.0-6.8	---	5.1-7.3	0	0
	47-60	0.0-6.8	---	5.1-7.3	0	0
Natchaug-----	0-2	---	18-85	3.5-6.0	0	0
	2-4	---	18-85	3.5-6.0	0	0
	4-6	---	18-85	3.5-6.0	0	0
	6-11	---	18-85	3.5-6.0	0	0
	11-18	---	18-85	3.5-6.0	0	0
	18-24	---	18-85	3.5-6.0	0	0
	24-33	1.8-13	---	5.1-7.3	0	0
	33-36	1.8-13	---	5.1-7.3	0	0
	36-80	1.8-10	---	5.1-7.3	0	0
18: Catden-----	0-2	---	18-138	3.5-7.3	0	0
	2-18	---	18-138	3.5-7.3	0	0
	18-47	---	18-138	3.5-7.3	0	0
	47-49	---	18-138	3.5-7.3	0	0
	49-61	---	18-138	3.5-7.3	0	0
Freetown-----	0-4	---	3.4-39	1.8-4.4	0	0
	4-10	---	3.4-39	1.8-4.4	0	0
	10-22	---	3.4-39	1.8-4.4	0	0
	22-35	---	3.4-39	1.8-4.4	0	0
	35-41	---	3.4-39	1.8-4.4	0	0
	41-55	---	3.4-39	1.8-4.4	0	0
	55-71	---	3.4-39	1.8-4.4	0	0
	71-91	---	3.4-39	1.8-4.4	0	0
20A: Ellington-----	0-8	---	0.6-2.2	4.5-6.0	0	0
	8-18	---	0.5-2.2	4.5-6.0	0	0
	18-26	---	0.5-2.2	4.5-6.0	0	0
	26-65	---	0.0-0.5	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
21A:						
Ninigret -----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-16	---	0.6-3.6	4.5-6.0	0	0
	16-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	0.0-1.1	---	4.5-6.5	0	0
Tisbury -----	0-8	---	0.5-3.1	4.5-6.0	0	0
	8-18	---	0.6-3.6	4.5-6.0	0	0
	18-26	---	0.7-5.3	4.5-6.0	0	0
	26-60	---	0.0-1.1	4.5-6.0	0	0
22A:						
Hero -----	0-9	2.6-5.7	---	5.6-7.3	0	0
	9-18	1.9-5.5	---	5.6-7.8	0	0
	18-24	1.9-5.5	---	5.6-7.8	0	0
	24-27	1.7-5.4	---	5.6-7.8	0-5	0
	27-60	1.1-3.6	---	7.3-8.4	0-20	0
22B:						
Hero -----	0-9	2.6-5.7	---	5.6-7.3	0	0
	9-18	1.9-5.5	---	5.6-7.8	0	0
	18-24	1.9-5.5	---	5.6-7.8	0	0
	24-27	1.7-5.4	---	5.6-7.8	0-5	0
	27-60	1.1-3.6	---	7.3-8.4	0-20	0
23A:						
Sudbury -----	0-1	---	30-121	4.5-6.5	0	0
	1-5	2.2-6.2	---	4.5-6.5	0	0
	5-17	1.9-6.1	---	4.5-6.5	0	0
	17-25	1.2-5.0	---	4.5-6.5	0	0
	25-60	0.0-2.6	---	4.5-6.5	0	0
24A:						
Deerfield -----	0-8	1.9-6.4	---	4.5-6.5	0	0
	8-16	1.0-5.7	---	4.5-6.5	0	0
	16-28	0.8-5.4	---	4.5-6.5	0	0
	28-34	0.0-4.0	---	4.5-6.5	0	0
	34-60	0.0-3.1	---	4.5-6.5	0	0
25A:						
Brancroft -----	0-6	9.8-13	---	4.5-6.5	0	0
	6-17	9.7-19	---	5.1-7.3	0	0
	17-22	9.6-19	---	5.1-7.3	0	0
	22-32	9.1-18	---	5.1-7.3	0	0
	32-43	9.1-18	---	5.6-7.3	0	0
	43-66	9.1-18	---	5.6-7.3	0	0
25B:						
Brancroft -----	0-6	9.8-13	---	4.5-6.5	0	0
	6-17	9.7-19	---	5.1-7.3	0	0
	17-22	9.6-19	---	5.1-7.3	0	0
	22-32	9.1-18	---	5.1-7.3	0	0
	32-43	9.1-18	---	5.6-7.3	0	0
	43-66	9.1-18	---	5.6-7.3	0	0
25C:						
Brancroft -----	0-6	9.8-13	---	4.5-6.5	0	0
	6-17	9.7-19	---	5.1-7.3	0	0
	17-22	9.6-19	---	5.1-7.3	0	0
	22-32	9.1-18	---	5.1-7.3	0	0
	32-43	9.1-18	---	5.6-7.3	0	0
	43-66	9.1-18	---	5.6-7.3	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
26A:						
Berlin-----	0-6	6.7-9.0	---	4.5-7.3	0	0
	6-12	6.5-13	---	5.1-7.3	0	0
	12-20	6.4-12	---	5.1-7.3	0	0
	20-34	6.0-16	---	5.1-7.3	0	0
	34-48	6.0-16	---	5.6-7.3	0	0
	48-65	6.0-16	---	5.6-7.3	0	0
26B:						
Berlin-----	0-6	6.7-9.0	---	4.5-7.3	0	0
	6-12	6.5-13	---	5.1-7.3	0	0
	12-20	6.4-12	---	5.1-7.3	0	0
	20-34	6.0-16	---	5.1-7.3	0	0
	34-48	6.0-16	---	5.6-7.3	0	0
	48-65	6.0-16	---	5.6-7.3	0	0
27A:						
Belgrade-----	0-8	2.8-8.3	---	4.5-7.3	0	0
	8-16	2.7-8.2	---	4.5-7.3	0	0
	16-27	2.7-8.1	---	4.5-7.3	0	0
	27-45	2.6-11	---	5.6-7.3	0	0
	45-60	1.0-11	---	5.6-7.3	0	0
28A:						
Elmridge-----	0-6	0.8-3.1	---	4.5-7.3	0	0
	6-10	0.8-3.7	---	5.1-7.3	0	0
	10-18	0.7-3.7	---	5.1-7.3	0	0
	18-25	0.7-3.6	---	5.1-7.3	0	0
	25-65	12-21	---	5.6-7.8	0	0
28B:						
Elmridge-----	0-6	0.8-3.1	---	4.5-7.3	0	0
	6-10	0.8-3.7	---	5.1-7.3	0	0
	10-18	0.7-3.7	---	5.1-7.3	0	0
	18-25	0.7-3.6	---	5.1-7.3	0	0
	25-65	12-21	---	5.6-7.8	0	0
29A:						
Agawam-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
29B:						
Agawam-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
29C:						
Agawam-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
30A:						
Branford-----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-18	---	0.6-3.6	4.5-6.0	0	0
	18-24	---	0.7-5.3	4.5-6.0	0	0
	24-65	---	0.0-0.7	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
30B:						
Branford -----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-18	---	0.6-3.6	4.5-6.0	0	0
	18-24	---	0.7-5.3	4.5-6.0	0	0
	24-65	---	0.0-0.7	4.5-6.0	0	0
30C:						
Branford -----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-18	---	0.6-3.6	4.5-6.0	0	0
	18-24	---	0.7-5.3	4.5-6.0	0	0
	24-65	---	0.0-0.7	4.5-6.0	0	0
31A:						
Copake -----	0-6	3.7-6.4	---	5.1-7.3	0	0
	6-13	2.9-6.3	---	5.1-7.3	0	0
	13-21	2.7-6.1	---	5.1-7.3	0	0
	21-31	2.7-6.1	---	5.1-7.3	0	0
	31-56	0.4-2.9	---	6.1-8.4	0-2	0
	56-65	0.4-2.9	---	6.6-8.4	1-10	0
	65-75	0.4-2.9	---	6.6-8.4	1-20	0
	75-80	0.4-2.9	---	7.4-8.4	1-25	0
31B:						
Copake -----	0-6	3.7-6.4	---	5.1-7.3	0	0
	6-13	2.9-6.3	---	5.1-7.3	0	0
	13-21	2.7-6.1	---	5.1-7.3	0	0
	21-31	2.7-6.1	---	5.1-7.3	0	0
	31-56	0.4-2.9	---	6.1-8.4	0-2	0
	56-65	0.4-2.9	---	6.6-8.4	1-10	0
	65-75	0.4-2.9	---	6.6-8.4	1-20	0
	75-80	0.4-2.9	---	7.4-8.4	1-25	0
31C:						
Copake -----	0-6	3.7-6.4	---	5.1-7.3	0	0
	6-13	2.9-6.3	---	5.1-7.3	0	0
	13-21	2.7-6.1	---	5.1-7.3	0	0
	21-31	2.7-6.1	---	5.1-7.3	0	0
	31-56	0.4-2.9	---	6.1-8.4	0-2	0
	56-65	0.4-2.9	---	6.6-8.4	1-10	0
	65-75	0.4-2.9	---	6.6-8.4	1-20	0
	75-80	0.4-2.9	---	7.4-8.4	1-25	0
32A:						
Haven -----	0-7	---	1.0-4.9	4.5-6.0	0	0
	7-14	---	1.1-5.7	4.5-6.0	0	0
	14-20	---	1.2-5.7	4.5-6.0	0	0
	20-24	---	1.3-8.4	4.5-6.0	0	0
	24-60	---	0.0-1.1	4.5-6.0	0	0
Enfield -----	0-3	---	32-71	4.5-5.5	0	0
	3-4	---	32-71	4.5-5.5	0	0
	4-12	1.7-6.7	---	4.5-6.0	0	0
	12-20	1.6-6.5	---	4.5-6.0	0	0
	20-26	1.6-6.5	---	4.5-6.0	0	0
	26-30	1.6-6.4	---	4.5-6.0	0	0
	30-37	0.0-3.2	---	4.5-6.0	0	0
	37-65	0.0-1.1	---	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
32B:						
Haven -----	0-7	---	1.0-4.9	4.5-6.0	0	0
	7-14	---	1.1-5.7	4.5-6.0	0	0
	14-20	---	1.2-5.7	4.5-6.0	0	0
	20-24	---	1.3-8.4	4.5-6.0	0	0
	24-60	---	0.0-1.1	4.5-6.0	0	0
Enfield -----	0-3	---	32-71	4.5-5.5	0	0
	3-4	---	32-71	4.5-5.5	0	0
	4-12	1.7-6.7	---	4.5-6.0	0	0
	12-20	1.6-6.5	---	4.5-6.0	0	0
	20-26	1.6-6.5	---	4.5-6.0	0	0
	26-30	1.6-6.4	---	4.5-6.0	0	0
	30-37	0.0-3.2	---	4.5-6.0	0	0
	37-65	0.0-1.1	---	4.5-6.0	0	0
32C:						
Haven -----	0-7	---	1.0-4.9	4.5-6.0	0	0
	7-14	---	1.1-5.7	4.5-6.0	0	0
	14-20	---	1.2-5.7	4.5-6.0	0	0
	20-24	---	1.3-8.4	4.5-6.0	0	0
	24-60	---	0.0-1.1	4.5-6.0	0	0
Enfield -----	0-3	---	32-71	4.5-5.5	0	0
	3-4	---	32-71	4.5-5.5	0	0
	4-12	1.7-6.7	---	4.5-6.0	0	0
	12-20	1.6-6.5	---	4.5-6.0	0	0
	20-26	1.6-6.5	---	4.5-6.0	0	0
	26-30	1.6-6.4	---	4.5-6.0	0	0
	30-37	0.0-3.2	---	4.5-6.0	0	0
	37-65	0.0-1.1	---	4.5-6.0	0	0
33A:						
Hartford -----	0-8	---	2.0-5.7	4.5-6.0	0	0
	8-20	---	0.9-3.5	4.5-6.0	0	0
	20-26	---	0.2-4.0	4.5-6.0	0	0
	26-65	---	0.1-2.0	4.5-6.0	0	0
33B:						
Hartford -----	0-8	---	2.0-5.7	4.5-6.0	0	0
	8-20	---	0.9-3.5	4.5-6.0	0	0
	20-26	---	0.2-4.0	4.5-6.0	0	0
	26-65	---	0.1-2.0	4.5-6.0	0	0
34A:						
Merrimac -----	0-9	2.8-6.9	---	5.1-6.5	0	0
	9-16	1.1-3.6	---	5.6-6.0	0	0
	16-24	1.1-3.6	---	5.6-6.0	0	0
	24-60	0.0-2.6	---	5.6-6.0	0	0
34B:						
Merrimac -----	0-9	2.8-6.9	---	5.1-6.5	0	0
	9-16	1.1-3.6	---	5.6-6.0	0	0
	16-24	1.1-3.6	---	5.6-6.0	0	0
	24-60	0.0-2.6	---	5.6-6.0	0	0
34C:						
Merrimac -----	0-9	2.8-6.9	---	5.1-6.5	0	0
	9-16	1.1-3.6	---	5.6-6.0	0	0
	16-24	1.1-3.6	---	5.6-6.0	0	0
	24-60	0.0-2.6	---	5.6-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
35A:						
Penwood-----	0-8	---	0.8-2.4	4.5-6.0	0	0
	8-18	---	0.1-1.4	4.5-6.0	0	0
	18-30	---	0.0-0.8	4.5-6.0	0	0
	30-60	---	0.0-0.8	4.5-6.0	0	0
35B:						
Penwood-----	0-8	---	0.8-2.4	4.5-6.0	0	0
	8-18	---	0.1-1.4	4.5-6.0	0	0
	18-30	---	0.0-0.8	4.5-6.0	0	0
	30-60	---	0.0-0.8	4.5-6.0	0	0
36A:						
Windsor-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-3.1	4.5-6.0	0	0
	3-9	---	0.4-1.8	4.5-6.0	0	0
	9-21	---	0.4-1.8	4.5-6.0	0	0
	21-25	0.0-2.6	---	4.5-6.5	0	0
	25-65	0.0-1.8	---	4.5-6.5	0	0
36B:						
Windsor-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-3.1	4.5-6.0	0	0
	3-9	---	0.4-1.8	4.5-6.0	0	0
	9-21	---	0.4-1.8	4.5-6.0	0	0
	21-25	0.0-2.6	---	4.5-6.5	0	0
	25-65	0.0-1.8	---	4.5-6.5	0	0
36C:						
Windsor-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-3.1	4.5-6.0	0	0
	3-9	---	0.4-1.8	4.5-6.0	0	0
	9-21	---	0.4-1.8	4.5-6.0	0	0
	21-25	0.0-2.6	---	4.5-6.5	0	0
	25-65	0.0-1.8	---	4.5-6.5	0	0
37A:						
Manchester-----	0-9	---	1.8-5.3	4.5-6.0	0	0
	9-18	---	0.1-1.8	4.5-6.0	0	0
	18-65	---	0.0-0.4	4.5-6.0	0	0
37C:						
Manchester-----	0-9	---	1.8-5.3	4.5-6.0	0	0
	9-18	---	0.1-1.8	4.5-6.0	0	0
	18-65	---	0.0-0.4	4.5-6.0	0	0
37E:						
Manchester-----	0-9	---	1.8-5.3	4.5-6.0	0	0
	9-18	---	0.1-1.8	4.5-6.0	0	0
	18-65	---	0.0-0.4	4.5-6.0	0	0
38A:						
Hinckley-----	0-8	---	2.3-6.7	3.5-6.0	0	0
	8-20	---	0.4-2.5	3.5-6.0	0	0
	20-27	---	0.1-1.6	3.5-6.0	0	0
	27-42	---	0.0-1.1	3.5-6.0	0	0
	42-60	---	0.0-1.1	3.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
38C:						
Hinckley-----	0-8	---	2.3-6.7	3.5-6.0	0	0
	8-20	---	0.4-2.5	3.5-6.0	0	0
	20-27	---	0.1-1.6	3.5-6.0	0	0
	27-42	---	0.0-1.1	3.5-6.0	0	0
	42-60	---	0.0-1.1	3.5-6.0	0	0
38E:						
Hinckley-----	0-8	---	2.3-6.7	3.5-6.0	0	0
	8-20	---	0.4-2.5	3.5-6.0	0	0
	20-27	---	0.1-1.6	3.5-6.0	0	0
	27-42	---	0.0-1.1	3.5-6.0	0	0
	42-60	---	0.0-1.1	3.5-6.0	0	0
39A:						
Groton-----	0-8	2.0-7.4	---	5.6-7.3	0	0
	8-18	1.9-6.2	---	5.6-7.3	0	0
	18-24	1.2-3.9	---	5.6-7.8	0	0
	24-30	1.2-3.9	---	5.6-7.8	0	0
	30-52	0.0-1.9	---	6.6-8.4	0-10	0
	52-72	0.0-1.9	---	6.6-8.4	1-10	0
39C:						
Groton-----	0-8	2.0-7.4	---	5.6-7.3	0	0
	8-18	1.9-6.2	---	5.6-7.3	0	0
	18-24	1.2-3.9	---	5.6-7.8	0	0
	24-30	1.2-3.9	---	5.6-7.8	0	0
	30-52	0.0-1.9	---	6.6-8.4	0-10	0
	52-72	0.0-1.9	---	6.6-8.4	1-10	0
39E:						
Groton-----	0-8	2.0-7.4	---	5.6-7.3	0	0
	8-18	1.9-6.2	---	5.6-7.3	0	0
	18-24	1.2-3.9	---	5.6-7.8	0	0
	24-30	1.2-3.9	---	5.6-7.8	0	0
	30-52	0.0-1.9	---	6.6-8.4	0-10	0
	52-72	0.0-1.9	---	6.6-8.4	1-10	0
40A:						
Ludlow-----	0-8	---	0.8-2.6	4.5-6.0	0	0
	8-20	---	0.8-2.9	4.5-6.0	0	0
	20-26	---	0.9-3.5	4.5-6.0	0	0
	26-65	---	0.9-3.5	4.5-6.0	0	0
40B:						
Ludlow-----	0-8	---	0.8-2.6	4.5-6.0	0	0
	8-20	---	0.8-2.9	4.5-6.0	0	0
	20-26	---	0.9-3.5	4.5-6.0	0	0
	26-65	---	0.9-3.5	4.5-6.0	0	0
41B:						
Ludlow-----	0-8	---	0.8-2.6	4.5-6.0	0	0
	8-20	---	0.8-2.9	4.5-6.0	0	0
	20-26	---	0.9-3.5	4.5-6.0	0	0
	26-65	---	0.9-3.5	4.5-6.0	0	0
42C:						
Ludlow-----	0-8	---	0.8-2.6	4.5-6.0	0	0
	8-20	---	0.8-2.9	4.5-6.0	0	0
	20-26	---	0.9-3.5	4.5-6.0	0	0
	26-65	---	0.9-3.5	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
43A:						
Rainbow-----	0-6	---	1.0-3.1	4.5-6.0	0	0
	6-18	---	0.4-3.6	4.5-6.0	0	0
	18-26	---	0.5-5.3	4.5-6.0	0	0
	26-65	---	0.5-5.3	4.5-6.0	0	0
43B:						
Rainbow-----	0-6	---	1.0-3.1	4.5-6.0	0	0
	6-18	---	0.4-3.6	4.5-6.0	0	0
	18-26	---	0.5-5.3	4.5-6.0	0	0
	26-65	---	0.5-5.3	4.5-6.0	0	0
44B:						
Rainbow-----	0-6	---	1.0-3.1	4.5-6.0	0	0
	6-18	---	0.4-3.6	4.5-6.0	0	0
	18-26	---	0.5-5.3	4.5-6.0	0	0
	26-65	---	0.5-5.3	4.5-6.0	0	0
45A:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0
45B:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0
45C:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0
46B:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0
46C:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
47C:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0
48B:						
Georgia-----	0-8	1.9-6.8	---	5.1-7.3	0	0
	8-14	1.9-6.6	---	5.1-7.3	0	0
	14-24	1.7-6.4	---	5.1-7.3	0	0
	24-60	1.7-6.4	---	5.1-7.3	0-25	0
Amenia-----	0-9	2.8-9.9	---	5.6-7.8	0	0
	9-16	2.7-9.8	---	5.6-7.8	0	0
	16-25	2.6-9.6	---	5.6-7.8	0	0
	25-60	2.6-9.6	---	7.4-8.4	1-25	0
48C:						
Georgia-----	0-8	1.9-6.8	---	5.1-7.3	0	0
	8-14	1.9-6.6	---	5.1-7.3	0	0
	14-24	1.7-6.4	---	5.1-7.3	0	0
	24-60	1.7-6.4	---	5.1-7.3	0-25	0
Amenia-----	0-9	2.8-9.9	---	5.6-7.8	0	0
	9-16	2.7-9.8	---	5.6-7.8	0	0
	16-25	2.6-9.6	---	5.6-7.8	0	0
	25-60	2.6-9.6	---	7.4-8.4	1-25	0
49B:						
Georgia-----	0-8	1.9-6.8	---	5.1-7.3	0	0
	8-14	1.9-6.6	---	5.1-7.3	0	0
	14-24	1.7-6.4	---	5.1-7.3	0	0
	24-60	1.7-6.4	---	5.1-7.3	0-25	0
Amenia-----	0-9	2.8-9.9	---	5.6-7.8	0	0
	9-16	2.7-9.8	---	5.6-7.8	0	0
	16-25	2.6-9.6	---	5.6-7.8	0	0
	25-60	2.6-9.6	---	7.4-8.4	1-25	0
49C:						
Georgia-----	0-8	1.9-6.8	---	5.1-7.3	0	0
	8-14	1.9-6.6	---	5.1-7.3	0	0
	14-24	1.7-6.4	---	5.1-7.3	0	0
	24-60	1.7-6.4	---	5.1-7.3	0-25	0
Amenia-----	0-9	2.8-9.9	---	5.6-7.8	0	0
	9-16	2.7-9.8	---	5.6-7.8	0	0
	16-25	2.6-9.6	---	5.6-7.8	0	0
	25-60	2.6-9.6	---	7.4-8.4	1-25	0
50A:						
Sutton-----	0-6	---	0.8-3.1	4.5-6.0	0	0
	6-12	---	0.8-3.3	4.5-6.0	0	0
	12-24	---	0.9-3.6	4.5-6.0	0	0
	24-28	---	0.9-5.3	4.5-6.0	0	0
	28-36	---	0.5-4.3	4.5-6.0	0	0
	36-65	---	0.5-4.3	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
		meq/100 g	meq/100 g	pH	Pct	mmhos/cm
50B:						
Sutton-----	0-6	---	0.8-3.1	4.5-6.0	0	0
	6-12	---	0.8-3.3	4.5-6.0	0	0
	12-24	---	0.9-3.6	4.5-6.0	0	0
	24-28	---	0.9-5.3	4.5-6.0	0	0
	28-36	---	0.5-4.3	4.5-6.0	0	0
	36-65	---	0.5-4.3	4.5-6.0	0	0
51B:						
Sutton-----	0-6	---	0.8-3.1	4.5-6.0	0	0
	6-12	---	0.8-3.3	4.5-6.0	0	0
	12-24	---	0.9-3.6	4.5-6.0	0	0
	24-28	---	0.9-5.3	4.5-6.0	0	0
	28-36	---	0.5-4.3	4.5-6.0	0	0
	36-65	---	0.5-4.3	4.5-6.0	0	0
52C:						
Sutton-----	0-6	---	0.8-3.1	4.5-6.0	0	0
	6-12	---	0.8-3.3	4.5-6.0	0	0
	12-24	---	0.9-3.6	4.5-6.0	0	0
	24-28	---	0.9-5.3	4.5-6.0	0	0
	28-36	---	0.5-4.3	4.5-6.0	0	0
	36-65	---	0.5-4.3	4.5-6.0	0	0
53A:						
Wapping-----	0-11	2.2-6.7	---	4.5-6.0	0	0
	11-16	1.1-6.5	---	4.5-6.0	0	0
	16-20	1.0-5.4	---	4.5-6.0	0	0
	20-28	0.5-4.3	---	4.5-6.0	0	0
	28-36	0.5-4.3	---	4.5-6.0	0	0
	36-80	0.5-3.2	---	4.5-6.0	0	0
53B:						
Wapping-----	0-11	2.2-6.7	---	4.5-6.0	0	0
	11-16	1.1-6.5	---	4.5-6.0	0	0
	16-20	1.0-5.4	---	4.5-6.0	0	0
	20-28	0.5-4.3	---	4.5-6.0	0	0
	28-36	0.5-4.3	---	4.5-6.0	0	0
	36-80	0.5-3.2	---	4.5-6.0	0	0
54B:						
Wapping-----	0-11	2.2-6.7	---	4.5-6.0	0	0
	11-16	1.1-6.5	---	4.5-6.0	0	0
	16-20	1.0-5.4	---	4.5-6.0	0	0
	20-28	0.5-4.3	---	4.5-6.0	0	0
	28-36	0.5-4.3	---	4.5-6.0	0	0
	36-80	0.5-3.2	---	4.5-6.0	0	0
55A:						
Watchaug-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-18	---	0.7-2.7	4.5-6.0	0	0
	18-24	---	0.7-2.9	4.5-6.0	0	0
	24-65	0.7-5.4	---	4.5-6.5	0	0
55B:						
Watchaug-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-18	---	0.7-2.7	4.5-6.0	0	0
	18-24	---	0.7-2.9	4.5-6.0	0	0
	24-65	0.7-5.4	---	4.5-6.5	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
56B:						
Watchaug-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-18	---	0.7-2.7	4.5-6.0	0	0
	18-24	---	0.7-2.9	4.5-6.0	0	0
	24-65	0.7-5.4	---	4.5-6.5	0	0
57B:						
Gloucester-----	0-4	---	1.3-6.0	3.5-6.0	0	0
	4-12	1.1-6.5	---	4.5-6.0	0	0
	12-25	0.0-3.9	---	4.5-6.0	0	0
	25-35	0.0-3.9	---	4.5-6.0	0	0
	35-60	0.0-3.9	---	4.5-6.0	0	0
57C:						
Gloucester-----	0-4	---	1.3-6.0	3.5-6.0	0	0
	4-12	1.1-6.5	---	4.5-6.0	0	0
	12-25	0.0-3.9	---	4.5-6.0	0	0
	25-35	0.0-3.9	---	4.5-6.0	0	0
	35-60	0.0-3.9	---	4.5-6.0	0	0
57D:						
Gloucester-----	0-4	---	1.3-6.0	3.5-6.0	0	0
	4-12	1.1-6.5	---	4.5-6.0	0	0
	12-25	0.0-3.9	---	4.5-6.0	0	0
	25-35	0.0-3.9	---	4.5-6.0	0	0
	35-60	0.0-3.9	---	4.5-6.0	0	0
58B:						
Gloucester-----	0-4	---	1.3-6.0	3.5-6.0	0	0
	4-12	1.1-6.5	---	4.5-6.0	0	0
	12-25	0.0-3.9	---	4.5-6.0	0	0
	25-35	0.0-3.9	---	4.5-6.0	0	0
	35-60	0.0-3.9	---	4.5-6.0	0	0
58C:						
Gloucester-----	0-4	---	1.3-6.0	3.5-6.0	0	0
	4-12	1.1-6.5	---	4.5-6.0	0	0
	12-25	0.0-3.9	---	4.5-6.0	0	0
	25-35	0.0-3.9	---	4.5-6.0	0	0
	35-60	0.0-3.9	---	4.5-6.0	0	0
59C:						
Gloucester-----	0-4	---	1.3-6.0	3.5-6.0	0	0
	4-12	1.1-6.5	---	4.5-6.0	0	0
	12-25	0.0-3.9	---	4.5-6.0	0	0
	25-35	0.0-3.9	---	4.5-6.0	0	0
	35-60	0.0-3.9	---	4.5-6.0	0	0
59D:						
Gloucester-----	0-4	---	1.3-6.0	3.5-6.0	0	0
	4-12	1.1-6.5	---	4.5-6.0	0	0
	12-25	0.0-3.9	---	4.5-6.0	0	0
	25-35	0.0-3.9	---	4.5-6.0	0	0
	35-60	0.0-3.9	---	4.5-6.0	0	0
60B:						
Canton-----	0-1	---	16-63	3.5-5.0	0	0
	1-3	---	0.1-1.4	3.5-6.0	0	0
	3-15	---	0.2-1.5	3.5-6.0	0	0
	15-24	---	0.2-1.5	3.5-6.0	0	0
	24-30	---	0.2-1.8	3.5-6.0	0	0
	30-60	0.0-1.9	---	3.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
60C:						
Canton-----	0-1	---	16-63	3.5-5.0	0	0
	1-3	---	0.1-1.4	3.5-6.0	0	0
	3-15	---	0.2-1.5	3.5-6.0	0	0
	15-24	---	0.2-1.5	3.5-6.0	0	0
	24-30	---	0.2-1.8	3.5-6.0	0	0
	30-60	0.0-1.9	---	3.5-6.0	0	0
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
60D:						
Canton-----	0-1	---	16-63	3.5-5.0	0	0
	1-3	---	0.1-1.4	3.5-6.0	0	0
	3-15	---	0.2-1.5	3.5-6.0	0	0
	15-24	---	0.2-1.5	3.5-6.0	0	0
	24-30	---	0.2-1.8	3.5-6.0	0	0
	30-60	0.0-1.9	---	3.5-6.0	0	0
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
61B:						
Canton-----	0-1	---	16-63	3.5-5.0	0	0
	1-3	---	0.1-1.4	3.5-6.0	0	0
	3-15	---	0.2-1.5	3.5-6.0	0	0
	15-24	---	0.2-1.5	3.5-6.0	0	0
	24-30	---	0.2-1.8	3.5-6.0	0	0
	30-60	0.0-1.9	---	3.5-6.0	0	0
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
61C:						
Canton-----	0-1	---	16-63	3.5-5.0	0	0
	1-3	---	0.1-1.4	3.5-6.0	0	0
	3-15	---	0.2-1.5	3.5-6.0	0	0
	15-24	---	0.2-1.5	3.5-6.0	0	0
	24-30	---	0.2-1.8	3.5-6.0	0	0
	30-60	0.0-1.9	---	3.5-6.0	0	0
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
62C:						
Canton-----	0-1	---	16-63	3.5-5.0	0	0
	1-3	---	0.1-1.4	3.5-6.0	0	0
	3-15	---	0.2-1.5	3.5-6.0	0	0
	15-24	---	0.2-1.5	3.5-6.0	0	0
	24-30	---	0.2-1.8	3.5-6.0	0	0
	30-60	0.0-1.9	---	3.5-6.0	0	0
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
62D:						
Canton-----	0-1	---	16-63	3.5-5.0	0	0
	1-3	---	0.1-1.4	3.5-6.0	0	0
	3-15	---	0.2-1.5	3.5-6.0	0	0
	15-24	---	0.2-1.5	3.5-6.0	0	0
	24-30	---	0.2-1.8	3.5-6.0	0	0
	30-60	0.0-1.9	---	3.5-6.0	0	0
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
63B:						
Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
63C:						
Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
63D:						
Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
64B:						
Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
64C:						
Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
65C:						
Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
65D:						
Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
66B:						
Narragansett-----	0-6	---	0.8-2.5	4.5-6.0	0	0
	6-15	---	0.9-2.7	4.5-6.0	0	0
	15-24	---	0.9-2.9	4.5-6.0	0	0
	24-28	---	0.9-4.3	4.5-6.0	0	0
	28-60	---	0.0-0.7	4.5-6.0	0	0
66C:						
Narragansett-----	0-6	---	0.8-2.5	4.5-6.0	0	0
	6-15	---	0.9-2.7	4.5-6.0	0	0
	15-24	---	0.9-2.9	4.5-6.0	0	0
	24-28	---	0.9-4.3	4.5-6.0	0	0
	28-60	---	0.0-0.7	4.5-6.0	0	0
67B:						
Narragansett-----	0-6	---	0.8-2.5	4.5-6.0	0	0
	6-15	---	0.9-2.7	4.5-6.0	0	0
	15-24	---	0.9-2.9	4.5-6.0	0	0
	24-28	---	0.9-4.3	4.5-6.0	0	0
	28-60	---	0.0-0.7	4.5-6.0	0	0
67C:						
Narragansett-----	0-6	---	0.8-2.5	4.5-6.0	0	0
	6-15	---	0.9-2.7	4.5-6.0	0	0
	15-24	---	0.9-2.9	4.5-6.0	0	0
	24-28	---	0.9-4.3	4.5-6.0	0	0
	28-60	---	0.0-0.7	4.5-6.0	0	0
68C:						
Narragansett-----	0-6	---	0.8-2.5	4.5-6.0	0	0
	6-15	---	0.9-2.7	4.5-6.0	0	0
	15-24	---	0.9-2.9	4.5-6.0	0	0
	24-28	---	0.9-4.3	4.5-6.0	0	0
	28-60	---	0.0-0.7	4.5-6.0	0	0
68D:						
Narragansett-----	0-6	---	0.8-2.5	4.5-6.0	0	0
	6-15	---	0.9-2.7	4.5-6.0	0	0
	15-24	---	0.9-2.9	4.5-6.0	0	0
	24-28	---	0.9-4.3	4.5-6.0	0	0
	28-60	---	0.0-0.7	4.5-6.0	0	0
69B:						
Yalesville-----	0-8	---	1.0-3.1	4.5-6.0	0	0
	8-14	---	1.1-4.6	4.5-6.0	0	0
	14-25	---	1.2-4.6	4.5-6.0	0	0
	25-36	---	1.2-6.8	4.5-6.0	0	0
	36-80	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
69C:						
Yalesville-----	0-8	---	1.0-3.1	4.5-6.0	0	0
	8-14	---	1.1-4.6	4.5-6.0	0	0
	14-25	---	1.2-4.6	4.5-6.0	0	0
	25-36	---	1.2-6.8	4.5-6.0	0	0
	36-80	---	---	---	---	---
70C:						
Branford-----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-18	---	0.6-3.6	4.5-6.0	0	0
	18-24	---	0.7-5.3	4.5-6.0	0	0
	24-65	---	0.0-0.7	4.5-6.0	0	0
Holyoke-----	0-1	---	17-89	3.5-6.0	0	0
	1-3	3.1-11	---	3.5-6.0	0	0
	3-8	3.0-11	---	3.5-6.0	0	0
	8-18	2.9-10	---	3.5-6.0	0	0
	18-80	---	---	---	0	0
71C:						
Brookfield-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-2.5	4.5-6.0	0	0
	3-13	---	0.6-2.7	4.5-6.0	0	0
	13-27	---	0.7-2.9	4.5-6.0	0	0
	27-60	---	0.7-4.3	4.5-6.0	0	0
Brimfield-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-2.5	4.5-6.0	0	0
	3-6	---	0.6-2.7	4.5-6.0	0	0
	6-17	---	0.7-2.9	4.5-6.0	0	0
	17-80	---	---	---	0	0
Rock Outcrop-----	---	---	---	---	---	---
71E:						
Brookfield-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-2.5	4.5-6.0	0	0
	3-13	---	0.6-2.7	4.5-6.0	0	0
	13-27	---	0.7-2.9	4.5-6.0	0	0
	27-60	---	0.7-4.3	4.5-6.0	0	0
Brimfield-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-2.5	4.5-6.0	0	0
	3-6	---	0.6-2.7	4.5-6.0	0	0
	6-17	---	0.7-2.9	4.5-6.0	0	0
	17-80	---	---	---	0	0
Rock Outcrop-----	---	---	---	---	---	---
73C:						
Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
Chatfield-----	0-1	---	32-63	4.5-5.0	0	0
	1-6	6.6-16	---	4.5-6.0	0	0
	6-15	6.3-16	---	4.5-6.0	0	0
	15-29	5.5-15	---	4.5-6.0	0	0
	29-80	---	---	---	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
Holyoke-----	0-1	---	17-89	3.5-6.0	0	0
	1-3	3.1-11	---	3.5-6.0	0	0
	3-8	3.0-11	---	3.5-6.0	0	0
	8-18	2.9-10	---	3.5-6.0	0	0
	18-80	---	---	---	0	0
80B: Bernardston-----	0-8	---	1.5-3.9	4.5-6.0	0	0
	8-14	---	1.6-6.8	4.5-6.0	0	0
	14-24	---	1.8-6.8	4.5-6.0	0	0
	24-26	---	1.9-6.8	4.5-6.0	0	0
	26-60	---	1.9-6.8	4.5-6.0	0	0
80C: Bernardston-----	0-8	---	1.5-3.9	4.5-6.0	0	0
	8-14	---	1.6-6.8	4.5-6.0	0	0
	14-24	---	1.8-6.8	4.5-6.0	0	0
	24-26	---	1.9-6.8	4.5-6.0	0	0
	26-60	---	1.9-6.8	4.5-6.0	0	0
81C: Bernardston-----	0-8	---	1.5-3.9	4.5-6.0	0	0
	8-14	---	1.6-6.8	4.5-6.0	0	0
	14-24	---	1.8-6.8	4.5-6.0	0	0
	24-26	---	1.9-6.8	4.5-6.0	0	0
	26-60	---	1.9-6.8	4.5-6.0	0	0
81D: Bernardston-----	0-8	---	1.5-3.9	4.5-6.0	0	0
	8-14	---	1.6-6.8	4.5-6.0	0	0
	14-24	---	1.8-6.8	4.5-6.0	0	0
	24-26	---	1.9-6.8	4.5-6.0	0	0
	26-60	---	1.9-6.8	4.5-6.0	0	0
82B: Broadbrook-----	0-8	---	1.0-3.9	4.5-6.0	0	0
	8-14	---	1.1-4.3	4.5-6.0	0	0
	14-25	---	1.2-6.8	4.5-6.0	0	0
	25-65	---	0.5-5.3	4.5-6.0	0	0
82C: Broadbrook-----	0-8	---	1.0-3.9	4.5-6.0	0	0
	8-14	---	1.1-4.3	4.5-6.0	0	0
	14-25	---	1.2-6.8	4.5-6.0	0	0
	25-65	---	0.5-5.3	4.5-6.0	0	0
82D: Broadbrook-----	0-8	---	1.0-3.9	4.5-6.0	0	0
	8-14	---	1.1-4.3	4.5-6.0	0	0
	14-25	---	1.2-6.8	4.5-6.0	0	0
	25-65	---	0.5-5.3	4.5-6.0	0	0
83B: Broadbrook-----	0-8	---	1.0-3.9	4.5-6.0	0	0
	8-14	---	1.1-4.3	4.5-6.0	0	0
	14-25	---	1.2-6.8	4.5-6.0	0	0
	25-65	---	0.5-5.3	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
83C:						
Broadbrook -----	0-8	---	1.0-3.9	4.5-6.0	0	0
	8-14	---	1.1-4.3	4.5-6.0	0	0
	14-25	---	1.2-6.8	4.5-6.0	0	0
	25-65	---	0.5-5.3	4.5-6.0	0	0
84B:						
Paxton -----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Montauk -----	0-4	1.2-3.6	---	4.5-6.0	0	0
	4-14	---	0.2-2.7	4.5-6.0	0	0
	14-25	---	0.2-2.7	4.5-6.0	0	0
	25-39	---	0.2-1.8	4.5-6.0	0	0
	39-60	---	0.2-1.8	4.5-6.0	0	0
84C:						
Paxton -----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Montauk -----	0-4	1.2-3.6	---	4.5-6.0	0	0
	4-14	---	0.2-2.7	4.5-6.0	0	0
	14-25	---	0.2-2.7	4.5-6.0	0	0
	25-39	---	0.2-1.8	4.5-6.0	0	0
	39-60	---	0.2-1.8	4.5-6.0	0	0
84D:						
Paxton -----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Montauk -----	0-4	1.2-3.6	---	4.5-6.0	0	0
	4-14	---	0.2-2.7	4.5-6.0	0	0
	14-25	---	0.2-2.7	4.5-6.0	0	0
	25-39	---	0.2-1.8	4.5-6.0	0	0
	39-60	---	0.2-1.8	4.5-6.0	0	0
85B:						
Paxton -----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Montauk -----	0-4	1.2-3.6	---	4.5-6.0	0	0
	4-14	---	0.2-2.7	4.5-6.0	0	0
	14-25	---	0.2-2.7	4.5-6.0	0	0
	25-39	---	0.2-1.8	4.5-6.0	0	0
	39-60	---	0.2-1.8	4.5-6.0	0	0
85C:						
Paxton -----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
		meq/100 g	meq/100 g	pH	Pct	mmhos/cm
Montauk-----	0-4	1.2-3.6	---	4.5-6.0	0	0
	4-14	---	0.2-2.7	4.5-6.0	0	0
	14-25	---	0.2-2.7	4.5-6.0	0	0
	25-39	---	0.2-1.8	4.5-6.0	0	0
	39-60	---	0.2-1.8	4.5-6.0	0	0
86C:						
Paxton-----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Montauk-----	0-4	1.2-3.6	---	4.5-6.0	0	0
	4-14	---	0.2-2.7	4.5-6.0	0	0
	14-25	---	0.2-2.7	4.5-6.0	0	0
	25-39	---	0.2-1.8	4.5-6.0	0	0
	39-60	---	0.2-1.8	4.5-6.0	0	0
86D:						
Paxton-----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Montauk-----	0-4	1.2-3.6	---	4.5-6.0	0	0
	4-14	---	0.2-2.7	4.5-6.0	0	0
	14-25	---	0.2-2.7	4.5-6.0	0	0
	25-39	---	0.2-1.8	4.5-6.0	0	0
	39-60	---	0.2-1.8	4.5-6.0	0	0
87B:						
Wethersfield-----	0-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
87C:						
Wethersfield-----	0-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
87D:						
Wethersfield-----	0-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
88B:						
Wethersfield-----	0-1	---	32-89	4.5-6.0	0	0
	1-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
88C:						
Wethersfield-----	0-1	---	32-89	4.5-6.0	0	0
	1-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
89C:						
Wethersfield-----	0-1	---	32-89	4.5-6.0	0	0
	1-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
89D:						
Wethersfield-----	0-1	---	32-89	4.5-6.0	0	0
	1-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
90B:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
90C:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
90D:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
91B:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
91C:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
91D:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
95E:						
Farmington-----	0-3	2.8-8.8	---	5.1-7.3	0	0
	3-8	2.7-8.7	---	5.6-7.8	0	0
	8-17	2.6-8.6	---	5.6-7.8	0-5	0
	17-80	---	---	---	---	---
Rock Outcrop-----	---	---	---	---	---	---
96:						
Ipswich-----	0-16	73-118	---	5.1-7.8	0	8.0- 16.0
	16-23	73-118	---	5.1-7.8	0	8.0- 16.0
	23-64	73-118	---	5.1-7.8	0	16.0- 60.0
	64-80	108-160	---	5.1-7.8	0	16.0- 60.0
97:						
Pawcatuck-----	0-12	38-130	---	5.1-7.8	0	8.0- 32.0
	12-40	38-118	---	5.1-7.8	0	8.0- 32.0
	40-46	38-83	---	5.1-7.8	0	16.0- 32.0
	46-50	2.5-18	---	5.1-7.8	0	16.0- 32.0
	50-60	0.0-4.3	---	5.1-7.8	0	16.0- 32.0
98:						
Westbrook-----	0-10	38-130	---	5.1-7.8	0	16.0- 60.0
	10-40	38-130	---	5.1-7.8	0	16.0- 40.0
	40-48	32-71	---	5.1-7.8	0	16.0- 40.0
	48-64	6.8-34	---	5.1-7.8	0	16.0- 32.0
	64-99	5.8-34	---	5.1-7.8	0	16.0- 32.0
99:						
Westbrook, low salt--	0-10	38-130	---	5.1-7.8	0	2.0- 16.0
	10-40	38-130	---	5.1-7.8	0	2.0- 16.0
	40-48	32-71	---	5.1-7.8	0	2.0- 16.0
	48-64	6.8-34	---	5.1-7.8	0	2.0- 16.0
	64-99	5.8-34	---	5.1-7.8	0	2.0- 16.0
100:						
Suncook-----	0-7	1.1-3.1	---	4.5-6.5	0	0
	7-15	0.0-2.7	---	4.5-6.5	0	0
	15-22	0.0-2.6	---	4.5-6.5	0	0
	22-32	0.0-2.6	---	4.5-6.5	0	0
	32-42	0.0-2.6	---	4.5-6.5	0	0
	42-65	0.0-2.6	---	4.5-6.5	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
101: Occum-----	0-10	2.1-11	---	4.5-6.5	0	0
	10-17	2.0-11	---	4.5-6.5	0	0
	17-28	1.8-7.1	---	4.5-6.5	0	0
	28-32	0.0-4.6	---	4.5-6.5	0	0
	32-42	0.0-4.6	---	4.5-6.5	0	0
	42-65	0.0-4.6	---	4.5-6.5	0	0
102: Pootatuck-----	0-4	1.1-3.4	---	4.5-6.5	0	0
	4-16	0.6-3.3	---	4.5-6.5	0	0
	16-21	0.6-3.3	---	4.5-6.5	0	0
	21-29	0.6-3.3	---	4.5-6.5	0	0
	29-35	0.0-1.1	---	4.5-6.5	0	0
	35-40	0.0-1.1	---	4.5-6.5	0	0
	40-65	0.0-1.1	---	4.5-6.5	0	0
103: Rippowam-----	0-5	2.2-6.1	---	4.5-7.3	0	0
	5-12	1.1-5.9	---	4.5-7.3	0	0
	12-19	1.1-5.6	---	4.5-7.3	0	0
	19-24	1.0-5.6	---	4.5-7.3	0	0
	24-27	1.0-5.5	---	5.6-7.3	0	0
	27-31	0.0-2.0	---	5.6-7.3	0	0
	31-65	0.0-2.0	---	5.6-7.3	0	0
104: Bash-----	0-11	1.9-6.4	---	4.5-6.0	0	0
	11-21	1.9-6.3	---	4.5-6.0	0	0
	21-28	1.9-6.3	---	4.5-6.0	0	0
	28-60	---	0.2-4.0	4.5-6.0	0	0
105: Hadley-----	0-12	4.9-14	---	4.5-7.3	0	0
	12-29	1.8-11	---	4.5-7.3	0	0
	29-40	1.8-11	---	4.5-7.3	0	0
	40-45	1.8-11	---	5.6-7.8	0	0
	45-60	1.8-11	---	5.6-7.8	0	0
106: Winooski-----	0-12	2.8-9.9	---	5.6-6.5	0	0
	12-18	1.0-5.4	---	5.6-7.3	0	0
	18-36	1.0-5.4	---	5.6-7.3	0	0
	36-52	1.0-5.4	---	5.6-7.3	0	0
	52-65	1.0-5.4	---	5.6-7.3	0	0
107: Limerick-----	0-8	3.8-5.5	---	5.1-7.3	0	0
	8-20	1.0-5.4	---	5.6-7.3	0	0
	20-36	1.0-5.4	---	5.6-7.3	0	0
	36-54	1.0-5.4	---	5.6-7.3	0	0
	54-65	0.5-4.3	---	5.6-7.3	0	0
Lim-----	0-6	3.1-11	---	5.1-7.3	0	0
	6-11	2.6-12	---	5.1-7.3	0	0
	11-15	2.6-12	---	5.1-7.3	0	0
	15-22	2.6-12	---	5.1-7.3	0	0
	22-29	2.6-10	---	5.1-7.3	0	0
	29-42	0.0-2.9	---	5.1-7.3	0	0
	42-50	0.0-2.9	---	5.1-7.3	0	0
	50-57	0.0-2.9	---	5.1-7.3	0	0
	57-65	0.0-2.9	---	5.1-7.3	0	0

Table 25.--Chemical Properties of the Soils--Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
108:						
Saco-----	0-12	3.9-9.4	---	5.1-7.3	0	0
	12-32	2.1-9.2	---	5.1-7.3	0	0
	32-48	2.1-9.0	---	5.6-7.3	0	0
	48-60	0.5-4.3	---	5.6-7.3	0	0
109:						
Fluvaquents, Frequently Flooded--	0-4	5.0-15	---	4.5-7.8	0	0
	4-14	0.0-10	---	4.5-7.8	0	0
	14-21	0.0-10	---	4.5-7.8	0	0
	21-38	5.0-15	---	4.5-7.8	0	0
	38-45	5.0-15	---	4.5-7.8	0	0
	45-55	0.0-10	---	4.5-7.8	0	0
	55-60	5.0-15	---	4.5-7.8	0	0
Udifulvents, Frequently Flooded--	0-2	5.0-15	---	4.5-7.3	0	0
	2-4	0.0-5.0	---	4.5-7.3	0	0
	4-12	0.0-10	---	4.5-7.3	0	0
	12-18	0.0-10	---	4.5-7.3	0	0
	18-35	0.0-5.0	---	4.5-7.3	0	0
	35-38	0.0-5.0	---	4.5-7.3	0	0
	38-60	0.0-5.0	---	4.5-7.3	0	0
221A:						
Ninigret-----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-16	---	0.6-3.6	4.5-6.0	0	0
	16-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	0.0-1.1	---	4.5-6.5	0	0
Urban Land-----	0-6	---	---	---	---	---
224A:						
Deerfield-----	0-8	1.9-6.4	---	4.5-6.5	0	0
	8-16	1.0-5.7	---	4.5-6.5	0	0
	16-28	0.8-5.4	---	4.5-6.5	0	0
	28-34	0.0-4.0	---	4.5-6.5	0	0
	34-60	0.0-3.1	---	4.5-6.5	0	0
Urban Land-----	0-6	---	---	---	---	---
225B:						
Brancroft-----	0-6	9.8-13	---	4.5-6.5	0	0
	6-17	9.7-19	---	5.1-7.3	0	0
	17-22	9.6-19	---	5.1-7.3	0	0
	22-32	9.1-18	---	5.1-7.3	0	0
	32-43	9.1-18	---	5.6-7.3	0	0
	43-66	9.1-18	---	5.6-7.3	0	0
Urban Land-----	0-6	---	---	---	---	---
226B:						
Berlin-----	0-6	6.7-9.0	---	4.5-7.3	0	0
	6-12	6.5-13	---	5.1-7.3	0	0
	12-20	6.4-12	---	5.1-7.3	0	0
	20-34	6.0-16	---	5.1-7.3	0	0
	34-48	6.0-16	---	5.6-7.3	0	0
	48-65	6.0-16	---	5.6-7.3	0	0
Urban Land-----	0-6	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
228B:						
Elmridge-----	0-6	0.8-3.1	---	4.5-7.3	0	0
	6-10	0.8-3.7	---	5.1-7.3	0	0
	10-18	0.7-3.7	---	5.1-7.3	0	0
	18-25	0.7-3.6	---	5.1-7.3	0	0
	25-65	12-21	---	5.6-7.8	0	0
Urban Land-----	0-6	---	---	---	---	---
229B:						
Agawam-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
Urban Land-----	0-6	---	---	---	---	---
229C:						
Agawam-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
Urban Land-----	0-6	---	---	---	---	---
230B:						
Branford-----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-18	---	0.6-3.6	4.5-6.0	0	0
	18-24	---	0.7-5.3	4.5-6.0	0	0
	24-65	---	0.0-0.7	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
230C:						
Branford-----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-18	---	0.6-3.6	4.5-6.0	0	0
	18-24	---	0.7-5.3	4.5-6.0	0	0
	24-65	---	0.0-0.7	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
232B:						
Haven-----	0-7	---	1.0-4.9	4.5-6.0	0	0
	7-14	---	1.1-5.7	4.5-6.0	0	0
	14-20	---	1.2-5.7	4.5-6.0	0	0
	20-24	---	1.3-8.4	4.5-6.0	0	0
	24-60	---	0.0-1.1	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
234B:						
Merrimac-----	0-9	2.8-6.9	---	5.1-6.5	0	0
	9-16	1.1-3.6	---	5.6-6.0	0	0
	16-24	1.1-3.6	---	5.6-6.0	0	0
	24-60	0.0-2.6	---	5.6-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
235B:						
Penwood-----	0-8	---	0.8-2.4	4.5-6.0	0	0
	8-18	---	0.1-1.4	4.5-6.0	0	0
	18-30	---	0.0-0.8	4.5-6.0	0	0
	30-60	---	0.0-0.8	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
236B:						
Windsor-----	0-1	---	30-99	4.5-6.0	0	0
	1-3	---	0.6-3.1	4.5-6.0	0	0
	3-9	---	0.4-1.8	4.5-6.0	0	0
	9-21	---	0.4-1.8	4.5-6.0	0	0
	21-25	0.0-2.6	---	4.5-6.5	0	0
	25-65	0.0-1.8	---	4.5-6.5	0	0
Urban Land-----	0-6	---	---	---	---	---
237A:						
Manchester-----	0-9	---	1.8-5.3	4.5-6.0	0	0
	9-18	---	0.1-1.8	4.5-6.0	0	0
	18-65	---	0.0-0.4	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
237C:						
Manchester-----	0-9	---	1.8-5.3	4.5-6.0	0	0
	9-18	---	0.1-1.8	4.5-6.0	0	0
	18-65	---	0.0-0.4	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
238A:						
Hinckley-----	0-8	---	2.3-6.7	3.5-6.0	0	0
	8-20	---	0.4-2.5	3.5-6.0	0	0
	20-27	---	0.1-1.6	3.5-6.0	0	0
	27-42	---	0.0-1.1	3.5-6.0	0	0
	42-60	---	0.0-1.1	3.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
238C:						
Hinckley-----	0-8	---	2.3-6.7	3.5-6.0	0	0
	8-20	---	0.4-2.5	3.5-6.0	0	0
	20-27	---	0.1-1.6	3.5-6.0	0	0
	27-42	---	0.0-1.1	3.5-6.0	0	0
	42-60	---	0.0-1.1	3.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
240B:						
Ludlow-----	0-8	---	0.8-2.6	4.5-6.0	0	0
	8-20	---	0.8-2.9	4.5-6.0	0	0
	20-26	---	0.9-3.5	4.5-6.0	0	0
	26-65	---	0.9-3.5	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
243B:						
Rainbow-----	0-6	---	1.0-3.1	4.5-6.0	0	0
	6-18	---	0.4-3.6	4.5-6.0	0	0
	18-26	---	0.5-5.3	4.5-6.0	0	0
	26-65	---	0.5-5.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
245B:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
245C:						
Woodbridge-----	0-7	---	0.5-3.1	4.5-6.0	0	0
	7-18	---	0.6-3.3	4.5-6.0	0	0
	18-26	---	0.7-3.6	4.5-6.0	0	0
	26-30	---	0.7-5.3	4.5-6.0	0	0
	30-43	---	0.7-5.3	4.5-6.0	0	0
	43-65	---	0.7-5.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
248B:						
Georgia-----	0-8	1.9-6.8	---	5.1-7.3	0	0
	8-14	1.9-6.6	---	5.1-7.3	0	0
	14-24	1.7-6.4	---	5.1-7.3	0	0
	24-60	1.7-6.4	---	5.1-7.3	0-25	0
Urban Land-----	0-6	---	---	---	---	---
250B:						
Sutton-----	0-6	---	0.8-3.1	4.5-6.0	0	0
	6-12	---	0.8-3.3	4.5-6.0	0	0
	12-24	---	0.9-3.6	4.5-6.0	0	0
	24-28	---	0.9-5.3	4.5-6.0	0	0
	28-36	---	0.5-4.3	4.5-6.0	0	0
	36-65	---	0.5-4.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
253B:						
Wapping-----	0-11	2.2-6.7	---	4.5-6.0	0	0
	11-16	1.1-6.5	---	4.5-6.0	0	0
	16-20	1.0-5.4	---	4.5-6.0	0	0
	20-28	0.5-4.3	---	4.5-6.0	0	0
	28-36	0.5-4.3	---	4.5-6.0	0	0
	36-80	0.5-3.2	---	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
255B:						
Watchaug-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-18	---	0.7-2.7	4.5-6.0	0	0
	18-24	---	0.7-2.9	4.5-6.0	0	0
	24-65	0.7-5.4	---	4.5-6.5	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
Urban Land-----	0-6	---	---	---	---	---
260B: Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
260C: Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
260D: Charlton-----	0-4	1.7-4.5	---	4.5-6.0	0	0
	4-7	1.6-4.3	---	4.5-6.0	0	0
	7-19	1.6-4.3	---	4.5-6.0	0	0
	19-27	1.6-4.3	---	4.5-6.0	0	0
	27-65	0.5-4.3	---	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
263B: Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
263C: Cheshire-----	0-8	---	0.6-2.6	4.5-6.0	0	0
	8-16	---	0.7-2.9	4.5-6.0	0	0
	16-26	---	0.7-2.9	4.5-6.0	0	0
	26-65	---	0.3-3.5	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
266B: Narragansett-----	0-6	---	0.8-2.5	4.5-6.0	0	0
	6-15	---	0.9-2.7	4.5-6.0	0	0
	15-24	---	0.9-2.9	4.5-6.0	0	0
	24-28	---	0.9-4.3	4.5-6.0	0	0
	28-60	---	0.0-0.7	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
269B: Yalesville-----	0-8	---	1.0-3.1	4.5-6.0	0	0
	8-14	---	1.1-4.6	4.5-6.0	0	0
	14-25	---	1.2-4.6	4.5-6.0	0	0
	25-36	---	1.2-6.8	4.5-6.0	0	0
	36-80	---	---	---	---	---
Urban Land-----	0-6	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
282B:						
Broadbrook-----	0-8	---	1.0-3.9	4.5-6.0	0	0
	8-14	---	1.1-4.3	4.5-6.0	0	0
	14-25	---	1.2-6.8	4.5-6.0	0	0
	25-65	---	0.5-5.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
284B:						
Paxton-----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
284C:						
Paxton-----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
284D:						
Paxton-----	0-8	1.7-6.6	---	4.5-6.5	0	0
	8-15	---	0.6-3.6	4.5-6.0	0	0
	15-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	---	0.7-5.3	4.5-6.0	0	0
Urban Land-----	0-6	---	---	---	---	---
287B:						
Wethersfield-----	0-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
Urban Land-----	0-6	---	---	---	---	---
287C:						
Wethersfield-----	0-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
Urban Land-----	0-6	---	---	---	---	---
287D:						
Wethersfield-----	0-3	---	1.5-3.9	4.5-6.0	0	0
	3-13	---	1.2-4.6	4.5-6.0	0	0
	13-27	---	1.2-4.6	4.5-6.0	0	0
	27-65	2.6-8.0	---	5.1-7.8	0	0
Urban Land-----	0-6	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
290B:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
Urban Land-----	0-6	---	---	---	---	---
290C:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
Urban Land-----	0-6	---	---	---	---	---
290D:						
Stockbridge-----	0-10	2.6-6.4	---	5.1-7.3	0	0
	10-20	2.6-6.2	---	5.6-7.3	0	0
	20-28	2.4-6.1	---	5.6-7.3	0	0
	28-42	2.4-6.1	---	5.6-7.3	0	0
	42-48	1.1-6.1	---	5.6-8.4	1-10	0
	48-65	1.1-6.1	---	5.6-8.4	1-10	0
Urban Land-----	0-6	---	---	---	---	---
301:						
Beaches-----	0-65	---	---	6.1-7.8	---	8.0- 32.0
Udipsamments-----	0-38	0.0-3.0	---	5.6-7.3	0	0
	38-50	0.0-2.0	---	5.6-7.3	0	0
	50-65	0.0-2.0	---	5.6-7.3	0	0
302:						
Dumps-----	0-65	---	---	---	---	---
303:						
Pits, Quarries-----	0-1	---	---	---	---	---
304:						
Udorthents-----	0-5	5.0-15	---	4.5-7.8	0	0
	5-21	0.0-10	---	4.5-7.8	0	0
	21-80	0.0-10	---	4.5-7.8	0	0
305:						
Udorthents-----	0-5	5.0-15	---	4.5-7.8	0	0
	5-21	0.0-10	---	4.5-7.8	0	0
	21-80	0.0-10	---	4.5-7.8	0	0
Pits-----	0-65	---	---	4.5-7.8	---	---
306:						
Udorthents-----	0-5	5.0-15	---	4.5-7.8	0	0
	5-21	0.0-10	---	4.5-7.8	0	0
	21-80	0.0-10	---	4.5-7.8	0	0
Urban Land-----	0-6	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
403F:						
Taconic-----	0-1	---	32-71	4.5-5.5	0	0
	1-4	---	1.5-3.9	4.5-5.5	0	0
	4-11	---	1.6-6.0	4.5-5.5	0	0
	11-80	---	---	---	0	0
Rock Outcrop-----	---	---	---	---	---	---
405C:						
Dummerston-----	0-1	---	32-89	4.5-6.0	0	0
	1-2	---	1.4-3.9	4.5-6.0	0	0
	2-3	---	3.7-13	4.5-6.0	0	0
	3-4	---	1.6-4.6	4.5-6.0	0	0
	4-6	---	1.7-4.6	4.5-6.0	0	0
	6-11	---	1.7-4.6	4.5-6.0	0	0
	11-22	---	1.8-4.6	4.5-6.0	0	0
	22-27	---	1.9-6.8	4.5-6.0	0	0
	27-40	---	1.9-5.3	4.5-6.0	0	0
	40-64	---	1.9-5.3	4.5-6.0	0	0
405E:						
Dummerston-----	0-1	---	32-89	4.5-6.0	0	0
	1-2	---	1.4-3.9	4.5-6.0	0	0
	2-3	---	3.7-13	4.5-6.0	0	0
	3-4	---	1.6-4.6	4.5-6.0	0	0
	4-6	---	1.7-4.6	4.5-6.0	0	0
	6-11	---	1.7-4.6	4.5-6.0	0	0
	11-22	---	1.8-4.6	4.5-6.0	0	0
	22-27	---	1.9-6.8	4.5-6.0	0	0
	27-40	---	1.9-5.3	4.5-6.0	0	0
	40-64	---	1.9-5.3	4.5-6.0	0	0
407C:						
Lanesboro-----	0-3	---	17-73	4.5-6.0	0	0
	3-6	---	1.5-3.9	4.5-6.0	0	0
	6-8	---	1.6-4.6	4.5-6.0	0	0
	8-16	---	1.7-4.6	4.5-6.0	0	0
	16-22	---	1.8-6.8	4.5-6.0	0	0
	22-30	---	1.9-6.8	4.5-6.0	0	0
	30-60	---	1.3-5.3	4.5-6.0	0	0
407E:						
Lanesboro-----	0-3	---	17-73	4.5-6.0	0	0
	3-6	---	1.5-3.9	4.5-6.0	0	0
	6-8	---	1.6-4.6	4.5-6.0	0	0
	8-16	---	1.7-4.6	4.5-6.0	0	0
	16-22	---	1.8-6.8	4.5-6.0	0	0
	22-30	---	1.9-6.8	4.5-6.0	0	0
	30-60	---	1.3-5.3	4.5-6.0	0	0
408C:						
Fullam-----	0-2	---	32-89	4.5-6.0	0	0
	2-4	---	1.5-3.9	4.5-6.0	0	0
	4-10	---	1.6-4.6	4.5-6.0	0	0
	10-20	---	1.8-4.6	4.5-6.0	0	0
	20-49	---	1.3-5.3	4.5-6.0	0	0
	49-60	---	1.3-5.3	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
409B:						
Brayton-----	0-3	---	17-89	3.5-6.0	0	0
	3-6	---	9.2-73	3.5-6.0	0	0
	6-7	---	0.5-2.0	3.5-6.0	0	0
	7-9	3.6-4.9	---	5.1-6.5	0	0
	9-13	1.6-4.9	---	5.1-6.5	0	0
	13-18	1.6-4.8	---	5.6-7.3	0	0
	18-23	1.6-4.8	---	5.6-7.3	0	0
	23-60	1.6-4.8	---	5.6-7.3	0	0
412B:						
Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
412C:						
Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
412D:						
Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
413C:						
Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
Millsite-----	0-1	---	32-71	4.5-5.5	0	0
	1-5	---	1.0-3.9	4.5-6.5	0	0
	5-13	---	1.2-4.6	4.5-6.5	0	0
	13-24	2.6-8.1	---	4.5-6.5	0	0
	24-31	2.6-8.0	---	4.5-6.5	0	0
	31-80	---	---	---	---	0
413E:						
Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
Millsite-----	0-1	---	32-71	4.5-5.5	0	0
	1-5	---	1.0-3.9	4.5-6.5	0	0
	5-13	---	1.2-4.6	4.5-6.5	0	0
	13-24	2.6-8.1	---	4.5-6.5	0	0
	24-31	2.6-8.0	---	4.5-6.5	0	0
	31-80	---	---	---	---	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
414:						
Fredon, cold-----	0-8	2.8-8.3	---	6.1-7.3	0	0
	8-17	2.7-8.2	---	6.1-7.3	0	0
	17-24	2.7-8.1	---	6.1-7.3	0	0
	24-29	1.0-3.8	---	5.6-8.4	0	0
	29-48	1.0-3.8	---	5.6-8.4	0-5	0
	48-60	1.0-3.8	---	5.6-8.4	0-10	0
415C:						
Millsite-----	0-1	---	32-71	4.5-5.5	0	0
	1-5	---	1.0-3.9	4.5-6.5	0	0
	5-13	---	1.2-4.6	4.5-6.5	0	0
	13-24	2.6-8.1	---	4.5-6.5	0	0
	24-31	2.6-8.0	---	4.5-6.5	0	0
	31-80	---	---	---	---	0
Westminster-----	0-1	---	11-65	3.5-6.0	0	0
	1-2	---	9.2-41	3.5-5.0	0	0
	2-5	---	1.0-3.1	3.5-5.5	0	0
	5-12	---	1.2-5.3	3.5-5.5	0	0
	12-16	---	1.2-5.3	3.5-5.5	0	0
	16-80	---	---	---	---	---
Rock Outcrop-----	---	---	---	---	---	---
415E:						
Millsite-----	0-1	---	32-71	4.5-5.5	0	0
	1-5	---	1.0-3.9	4.5-6.5	0	0
	5-13	---	1.2-4.6	4.5-6.5	0	0
	13-24	2.6-8.1	---	4.5-6.5	0	0
	24-31	2.6-8.0	---	4.5-6.5	0	0
	31-80	---	---	---	---	0
Westminster-----	0-1	---	11-52	3.5-5.5	0	0
	1-2	---	9.2-41	3.5-5.0	0	0
	2-5	---	1.0-3.1	3.5-5.5	0	0
	5-12	---	1.2-5.3	3.5-5.5	0	0
	12-16	---	1.2-5.3	3.5-5.5	0	0
	16-80	---	---	---	---	---
Rock Outcrop-----	---	---	---	---	---	---
416E:						
Rock Outcrop-----	---	---	---	---	---	---
Westminster-----	0-1	---	11-52	3.5-5.5	0	0
	1-2	---	9.2-52	3.5-5.5	0	0
	2-5	---	1.0-3.1	3.5-5.5	0	0
	5-12	---	1.2-5.3	3.5-5.5	0	0
	12-16	---	1.2-5.3	3.5-5.5	0	0
	16-80	---	---	---	---	---
416F:						
Rock Outcrop-----	---	---	---	---	---	---
Westminster-----	0-1	---	11-52	3.5-5.5	0	0
	1-2	---	9.2-52	3.5-5.5	0	0
	2-5	---	1.0-3.1	3.5-5.5	0	0
	5-12	---	1.2-5.3	3.5-5.5	0	0
	12-16	---	1.2-5.3	3.5-5.5	0	0
	16-80	---	---	---	---	---

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
417B: Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
417C: Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
417D: Bice-----	0-1	---	32-89	4.5-6.0	0	0
	1-7	2.8-8.3	---	4.5-6.0	0	0
	7-16	3.6-8.2	---	4.5-6.0	0	0
	16-24	3.6-8.0	---	4.5-6.0	0	0
	24-60	3.6-8.0	---	4.5-6.0	0	0
418C: Schroon-----	0-1	---	17-71	3.5-5.5	0	0
	1-2	---	17-71	3.5-5.5	0	0
	2-3	---	9.2-59	3.5-5.5	0	0
	3-9	6.6-14	---	3.5-6.0	0	0
	9-14	6.4-13	---	4.5-6.0	0	0
	14-23	6.3-13	---	4.5-6.0	0	0
	23-30	6.3-13	---	4.5-6.0	0	0
	30-60	2.6-12	---	5.1-6.5	0	0
420A: Schroon-----	0-1	---	17-71	3.5-5.5	0	0
	1-2	---	17-71	3.5-5.5	0	0
	2-3	---	9.2-59	3.5-5.5	0	0
	3-9	6.6-14	---	3.5-6.0	0	0
	9-14	6.4-13	---	4.5-6.0	0	0
	14-23	6.3-13	---	4.5-6.0	0	0
	23-30	6.3-13	---	4.5-6.0	0	0
	30-60	2.6-12	---	5.1-6.5	0	0
420B: Schroon-----	0-1	---	17-71	3.5-5.5	0	0
	1-2	---	17-71	3.5-5.5	0	0
	2-3	---	9.2-59	3.5-5.5	0	0
	3-9	6.6-14	---	3.5-6.0	0	0
	9-14	6.4-13	---	4.5-6.0	0	0
	14-23	6.3-13	---	4.5-6.0	0	0
	23-30	6.3-13	---	4.5-6.0	0	0
	30-60	2.6-12	---	5.1-6.5	0	0
421A: Ninigret, cold-----	0-8	---	0.6-3.1	4.5-6.0	0	0
	8-16	---	0.6-3.6	4.5-6.0	0	0
	16-26	---	0.7-5.3	4.5-6.0	0	0
	26-65	0.0-1.1	---	4.5-6.5	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
423A: Sudbury, cold-----	0-1	---	30-121	4.5-6.5	0	0
	1-5	2.2-6.2	---	4.5-6.5	0	0
	5-17	1.9-6.1	---	4.5-6.5	0	0
	17-25	1.2-5.0	---	4.5-6.5	0	0
	25-60	0.0-2.6	---	4.5-6.5	0	0
	424B: Shelburne-----	0-1	---	32-89	4.5-6.0	0
1-2	---	0.6-3.9	4.5-6.0	0	0	
2-7	---	0.6-6.8	4.5-6.0	0	0	
7-21	---	0.6-6.8	4.5-6.0	0	0	
21-27	---	0.7-6.8	4.5-6.0	0	0	
27-32	1.6-8.0	---	5.1-6.0	0	0	
32-43	1.6-8.0	---	5.1-6.0	0	0	
43-55	1.6-8.0	---	5.1-6.0	0	0	
55-80	1.6-8.0	---	5.1-6.0	0	0	
424C: Shelburne-----	0-1	---	32-89	4.5-6.0	0	0
	1-2	---	0.6-3.9	4.5-6.0	0	0
	2-7	---	0.6-6.8	4.5-6.0	0	0
	7-21	---	0.6-6.8	4.5-6.0	0	0
	21-27	---	0.7-6.8	4.5-6.0	0	0
	27-32	1.6-8.0	---	5.1-6.0	0	0
	32-43	1.6-8.0	---	5.1-6.0	0	0
	43-55	1.6-8.0	---	5.1-6.0	0	0
	55-80	1.6-8.0	---	5.1-6.0	0	0
	424D: Shelburne-----	0-1	---	32-89	4.5-6.0	0
1-2		---	0.6-3.9	4.5-6.0	0	0
2-7		---	0.6-6.8	4.5-6.0	0	0
7-21		---	0.6-6.8	4.5-6.0	0	0
21-27		---	0.7-6.8	4.5-6.0	0	0
27-32		1.6-8.0	---	5.1-6.0	0	0
32-43		1.6-8.0	---	5.1-6.0	0	0
43-55		1.6-8.0	---	5.1-6.0	0	0
55-80		1.6-8.0	---	5.1-6.0	0	0
425B: Shelburne-----		0-1	---	32-89	4.5-6.0	0
	1-2	---	0.6-3.9	4.5-6.0	0	0
	2-7	---	0.6-6.8	4.5-6.0	0	0
	7-21	---	0.6-6.8	4.5-6.0	0	0
	21-27	---	0.7-6.8	4.5-6.0	0	0
	27-32	1.6-8.0	---	5.1-6.0	0	0
	32-43	1.6-8.0	---	5.1-6.0	0	0
	43-55	1.6-8.0	---	5.1-6.0	0	0
	55-80	1.6-8.0	---	5.1-6.0	0	0
	425C: Shelburne-----	0-1	---	32-89	4.5-6.0	0
1-2		---	0.6-3.9	4.5-6.0	0	0
2-7		---	0.6-6.8	4.5-6.0	0	0
7-21		---	0.6-6.8	4.5-6.0	0	0
21-27		---	0.7-6.8	4.5-6.0	0	0
27-32		1.6-8.0	---	5.1-6.0	0	0
32-43		1.6-8.0	---	5.1-6.0	0	0
43-55		1.6-8.0	---	5.1-6.0	0	0
55-80		1.6-8.0	---	5.1-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
426D: Shelburne-----	0-1	---	32-89	4.5-6.0	0	0
	1-2	---	0.6-3.9	4.5-6.0	0	0
	2-7	---	0.6-6.8	4.5-6.0	0	0
	7-21	---	0.6-6.8	4.5-6.0	0	0
	21-27	---	0.7-6.8	4.5-6.0	0	0
	27-32	1.6-8.0	---	5.1-6.0	0	0
	32-43	1.6-8.0	---	5.1-6.0	0	0
	43-55	1.6-8.0	---	5.1-6.0	0	0
	55-80	1.6-8.0	---	5.1-6.0	0	0
	427B: Ashfield-----	0-1	---	9.2-80	3.5-5.5	0
1-2		---	16-80	3.5-5.5	0	0
2-3		---	17-80	3.5-5.5	0	0
3-7		---	1.0-3.9	3.5-5.5	0	0
7-12		---	1.0-3.9	3.5-5.5	0	0
12-18		---	0.6-4.6	3.5-5.5	0	0
18-24		---	0.7-4.6	3.5-5.5	0	0
24-29		---	0.7-6.8	3.5-5.5	0	0
29-44		1.6-8.0	---	5.1-6.5	0	0
44-58		1.6-8.0	---	5.1-6.5	0	0
58-80		1.6-8.0	---	5.1-6.5	0	0
427C: Ashfield-----		0-1	---	9.2-80	3.5-5.5	0
	1-2	---	16-80	3.5-5.5	0	0
	2-3	---	17-80	3.5-5.5	0	0
	3-7	---	1.0-3.9	3.5-5.5	0	0
	7-12	---	1.0-3.9	3.5-5.5	0	0
	12-18	---	0.6-4.6	3.5-5.5	0	0
	18-24	---	0.7-4.6	3.5-5.5	0	0
	24-29	---	0.7-6.8	3.5-5.5	0	0
	29-44	1.6-8.0	---	5.1-6.5	0	0
	44-58	1.6-8.0	---	5.1-6.5	0	0
	58-80	1.6-8.0	---	5.1-6.5	0	0
	428A: Ashfield-----	0-1	---	9.2-80	3.5-5.5	0
1-2		---	16-80	3.5-5.5	0	0
2-3		---	17-80	3.5-5.5	0	0
3-7		---	1.0-3.9	3.5-5.5	0	0
7-12		---	1.0-3.9	3.5-5.5	0	0
12-18		---	0.6-4.6	3.5-5.5	0	0
18-24		---	0.7-4.6	3.5-5.5	0	0
24-29		---	0.7-6.8	3.5-5.5	0	0
29-44		1.6-8.0	---	5.1-6.5	0	0
44-58		1.6-8.0	---	5.1-6.5	0	0
58-80		1.6-8.0	---	5.1-6.5	0	0
428B: Ashfield-----		0-1	---	9.2-80	3.5-5.5	0
	1-2	---	16-80	3.5-5.5	0	0
	2-3	---	17-80	3.5-5.5	0	0
	3-7	---	1.0-3.9	3.5-5.5	0	0
	7-12	---	1.0-3.9	3.5-5.5	0	0
	12-18	---	0.6-4.6	3.5-5.5	0	0
	18-24	---	0.7-4.6	3.5-5.5	0	0
	24-29	---	0.7-6.8	3.5-5.5	0	0
	29-44	1.6-8.0	---	5.1-6.5	0	0
	44-58	1.6-8.0	---	5.1-6.5	0	0
	58-80	1.6-8.0	---	5.1-6.5	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
428C:						
Ashfield-----	0-1	---	9.2-80	3.5-5.5	0	0
	1-2	---	16-80	3.5-5.5	0	0
	2-3	---	17-80	3.5-5.5	0	0
	3-7	---	1.0-3.9	3.5-5.5	0	0
	7-12	---	1.0-3.9	3.5-5.5	0	0
	12-18	---	0.6-4.6	3.5-5.5	0	0
	18-24	---	0.7-4.6	3.5-5.5	0	0
	24-29	---	0.7-6.8	3.5-5.5	0	0
	29-44	1.6-8.0	---	5.1-6.5	0	0
	44-58	1.6-8.0	---	5.1-6.5	0	0
	58-80	1.6-8.0	---	5.1-6.5	0	0
429A:						
Agawam, cold-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
429B:						
Agawam, cold-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
429C:						
Agawam, cold-----	0-8	2.2-5.5	---	4.5-6.5	0	0
	8-14	0.6-5.5	---	4.5-6.5	0	0
	14-24	0.5-3.2	---	4.5-6.5	0	0
	24-60	0.0-0.6	---	4.5-6.5	0	0
433:						
Moosilauke-----	0-1	---	30-81	4.5-6.0	0	0
	1-6	---	2.0-6.0	4.5-6.0	0	0
	6-16	---	0.3-7.2	4.5-6.0	0	0
	16-24	---	0.1-5.3	4.5-6.0	0	0
	24-39	0.0-3.9	---	4.5-6.0	0	0
	39-65	0.0-3.9	---	4.5-6.0	0	0
434A:						
Merrimac, cold-----	0-9	2.8-6.9	---	5.1-6.5	0	0
	9-16	1.1-3.6	---	5.6-6.0	0	0
	16-24	1.1-3.6	---	5.6-6.0	0	0
	24-60	0.0-2.6	---	5.6-6.0	0	0
434B:						
Merrimac, cold-----	0-9	2.8-6.9	---	5.1-6.5	0	0
	9-16	1.1-3.6	---	5.6-6.0	0	0
	16-24	1.1-3.6	---	5.6-6.0	0	0
	24-60	0.0-2.6	---	5.6-6.0	0	0
434C:						
Merrimac, cold-----	0-9	2.8-6.9	---	5.1-6.5	0	0
	9-16	1.1-3.6	---	5.6-6.0	0	0
	16-24	1.1-3.6	---	5.6-6.0	0	0
	24-60	0.0-2.6	---	5.6-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
435:						
Scarboro-----	0-12	---	32-99	4.5-6.0	0	0
	12-17	---	1.0-6.9	4.5-6.0	0	0
	17-31	---	0.1-3.6	4.5-6.0	0	0
	31-72	0.0-1.9	---	4.5-7.3	0	0
436:						
Halsey-----	0-1	81-104	---	5.6-7.3	0	0
	1-8	3.3-9.4	---	5.6-7.3	0	0
	8-16	3.2-9.2	---	5.6-7.3	0	0
	16-28	2.7-6.5	---	5.6-7.3	0	0
	28-38	1.0-4.3	---	6.1-8.4	0	0
	38-60	1.0-4.3	---	6.1-8.4	0-10	0
437:						
Wonsqueak-----	0-2	---	23-121	3.5-6.5	0	0
	2-11	---	23-121	3.5-6.5	0	0
	11-22	---	43-121	4.5-6.5	0	0
	22-25	1.5-13	---	4.5-7.3	0	0
	25-45	1.5-13	---	4.5-7.3	0	0
	45-60	1.5-13	---	4.5-7.3	0	0
438:						
Bucksport-----	0-9	---	23-80	3.5-5.5	0	0
	9-33	---	43-121	4.5-6.5	0	0
	33-50	---	43-121	4.5-6.5	0	0
	50-59	133-171	---	4.5-6.5	0	0
	59-63	1.0-13	---	4.5-7.3	0	0
440A:						
Boscawen-----	0-1	---	34-68	4.5-5.5	0	0
	1-2	---	1.3-4.6	4.5-5.5	0	0
	2-9	---	0.2-1.9	4.5-5.5	0	0
	9-16	---	0.2-1.9	4.5-6.0	0	0
	16-29	---	0.0-1.9	4.5-6.0	0	0
	29-34	---	0.0-1.9	4.5-6.0	0	0
	34-40	0.0-4.7	---	4.5-6.0	0	0
	40-44	0.0-4.7	---	4.5-6.0	0	0
	44-67	0.0-4.7	---	4.5-6.0	0	0
440C:						
Boscawen-----	0-1	---	34-68	4.5-5.5	0	0
	1-2	---	1.3-4.6	4.5-5.5	0	0
	2-9	---	0.2-1.9	4.5-5.5	0	0
	9-16	---	0.2-1.9	4.5-6.0	0	0
	16-29	---	0.0-1.9	4.5-6.0	0	0
	29-34	---	0.0-1.9	4.5-6.0	0	0
	34-40	0.0-4.7	---	4.5-6.0	0	0
	40-44	0.0-4.7	---	4.5-6.0	0	0
	44-67	0.0-4.7	---	4.5-6.0	0	0
440E:						
Boscawen-----	0-1	---	34-68	4.5-5.5	0	0
	1-2	---	1.3-4.6	4.5-5.5	0	0
	2-9	---	0.2-1.9	4.5-5.5	0	0
	9-16	---	0.2-1.9	4.5-6.0	0	0
	16-29	---	0.0-1.9	4.5-6.0	0	0
	29-34	---	0.0-1.9	4.5-6.0	0	0
	34-40	0.0-4.7	---	4.5-6.0	0	0
	40-44	0.0-4.7	---	4.5-6.0	0	0
	44-67	0.0-4.7	---	4.5-6.0	0	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
442:						
Brayton-----	0-2	---	17-89	3.5-6.0	0	0
	2-10	---	1.4-3.6	3.5-6.0	0	0
	10-17	2.6-8.2	---	5.1-6.5	0	0
	17-22	2.6-8.2	---	5.1-6.5	0	0
	22-27	2.6-8.0	---	5.1-7.3	0	0
	27-42	2.6-8.0	---	5.6-7.3	0	0
	42-65	2.6-8.0	---	5.6-7.3	0	0
443:						
Brayton-----	0-2	---	17-89	3.5-6.0	0	0
	2-10	---	1.4-3.6	3.5-6.0	0	0
	10-17	2.6-8.2	---	5.1-6.5	0	0
	17-22	2.6-8.2	---	5.1-6.5	0	0
	22-27	2.6-8.0	---	5.1-7.3	0	0
	27-42	2.6-8.0	---	5.6-7.3	0	0
	42-65	2.6-8.0	---	5.6-7.3	0	0
Loonmeadow-----	0-2	65-88	---	5.1-6.0	0	0
	2-9	1.7-6.7	---	5.1-7.3	0	0
	9-18	1.6-6.5	---	5.1-7.8	0-5	0
	18-35	1.6-5.4	---	6.6-8.4	0-5	0
	35-80	1.6-5.4	---	6.6-8.4	0-5	0
448B:						
Hogansburg-----	0-12	6.6-9.7	---	5.1-7.3	0	0
	12-20	5.4-8.0	---	5.1-7.8	0	0
	20-29	2.4-7.2	---	5.1-7.8	0	0
	29-43	2.4-7.2	---	5.1-8.0	1-5	0
	43-50	1.7-6.1	---	7.4-8.4	1-10	0
	50-70	1.7-6.1	---	7.4-8.4	1-10	0
	70-84	1.7-6.1	---	7.4-8.4	10-25	0
449B:						
Hogansburg-----	0-12	6.6-9.7	---	5.1-7.3	0	0
	12-20	5.4-8.0	---	5.1-7.8	0	0
	20-29	2.4-7.2	---	5.1-7.8	0	0
	29-43	2.4-7.2	---	5.1-8.0	1-5	0
	43-50	1.7-6.1	---	7.4-8.4	1-10	0
	50-70	1.7-6.1	---	7.4-8.4	1-10	0
	70-84	1.7-6.1	---	7.4-8.4	10-25	0
449C:						
Hogansburg-----	0-12	6.6-9.7	---	5.1-7.3	0	0
	12-20	5.4-8.0	---	5.1-7.8	0	0
	20-29	2.4-7.2	---	5.1-7.8	0	0
	29-43	2.4-7.2	---	5.1-8.0	1-5	0
	43-50	1.7-6.1	---	7.4-8.4	1-10	0
	50-70	1.7-6.1	---	7.4-8.4	1-10	0
	70-84	1.7-6.1	---	7.4-8.4	10-25	0
450B:						
Pyrities-----	0-1	69-81	---	5.6-6.5	0	0
	1-8	9.8-15	---	5.6-7.8	0	0
	8-13	6.1-13	---	6.1-7.8	0	0
	13-26	6.1-13	---	6.1-7.8	0	0
	26-45	6.1-13	---	6.1-8.4	0	0
	45-65	6.1-11	---	6.1-8.4	5-35	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
450C: Pyrities-----	0-1	69-81	---	5.6-6.5	0	0
	1-8	9.8-15	---	5.6-7.8	0	0
	8-13	6.1-13	---	6.1-7.8	0	0
	13-26	6.1-13	---	6.1-7.8	0	0
	26-45	6.1-13	---	6.1-8.4	0	0
	45-65	6.1-11	---	6.1-8.4	5-35	0
450D: Pyrities-----	0-1	69-81	---	5.6-6.5	0	0
	1-8	9.8-15	---	5.6-7.8	0	0
	8-13	6.1-13	---	6.1-7.8	0	0
	13-26	6.1-13	---	6.1-7.8	0	0
	26-45	6.1-13	---	6.1-8.4	0	0
	45-65	6.1-11	---	6.1-8.4	5-35	0
451B: Pyrities-----	0-1	69-81	---	5.6-6.5	0	0
	1-8	9.8-15	---	5.6-7.8	0	0
	8-13	6.1-13	---	6.1-7.8	0	0
	13-26	6.1-13	---	6.1-7.8	0	0
	26-45	6.1-13	---	6.1-8.4	0	0
	45-65	6.1-11	---	6.1-8.4	5-35	0
451C: Pyrities-----	0-1	69-81	---	5.6-6.5	0	0
	1-8	9.8-15	---	5.6-7.8	0	0
	8-13	6.1-13	---	6.1-7.8	0	0
	13-26	6.1-13	---	6.1-7.8	0	0
	26-45	6.1-13	---	6.1-8.4	0	0
	45-65	6.1-11	---	6.1-8.4	5-35	0
451D: Pyrities-----	0-1	69-81	---	5.6-6.5	0	0
	1-8	9.8-15	---	5.6-7.8	0	0
	8-13	6.1-13	---	6.1-7.8	0	0
	13-26	6.1-13	---	6.1-7.8	0	0
	26-45	6.1-13	---	6.1-8.4	0	0
	45-65	6.1-11	---	6.1-8.4	5-35	0
457: Mudgepond-----	0-11	9.3-20	---	6.6-7.8	0	0
	11-16	6.3-15	---	6.6-7.8	0-2	0
	16-26	4.6-15	---	6.6-7.8	0-15	0
	26-35	4.6-15	---	6.6-7.8	1-15	0
	35-65	2.6-14	---	6.6-8.4	5-25	0
458: Mudgepond-----	0-11	9.3-20	---	6.6-7.8	0	0
	11-16	6.3-15	---	6.6-7.8	0-2	0
	16-26	4.6-15	---	6.6-7.8	0-15	0
	26-35	4.6-15	---	6.6-7.8	1-15	0
	35-65	2.6-14	---	6.6-8.4	5-25	0
Alden-----	0-4	8.4-14	---	5.1-7.3	0	0
	4-13	9.8-14	---	5.1-7.3	0	0
	13-23	9.6-14	---	5.6-7.3	0	0
	23-29	9.6-14	---	5.6-7.3	0	0
	29-43	9.1-14	---	6.1-7.3	0	0
	43-60	9.1-14	---	6.1-8.4	0-15	0

Table 25.—Chemical Properties of the Soils—Continued

Map symbol and soil name	Depth	Cation exchange capacity	Effective cation exchange capacity	Soil reaction	Calcium carbonate	Salinity
	In	meq/100 g	meq/100 g	pH	Pct	mmhos/cm
501: Ondawa-----	0-1	---	34-65	4.5-5.5	---	---
	1-2	---	34-65	4.5-5.5	0	0
	2-14	---	4.4-7.8	4.5-6.5	0	0
	14-30	---	3.6-12	4.5-6.5	0	0
	30-33	---	3.6-12	4.5-6.5	0	0
	33-60	---	3.6-9.5	4.5-6.5	5-35	0
503: Rumney-----	0-7	2.8-14	---	5.6-7.3	0	0
	7-22	1.6-13	---	5.6-7.3	0	0
	22-38	1.6-13	---	5.6-7.3	0	0
	38-42	1.7-14	---	5.6-7.3	0	0
	42-44	0.5-9.5	---	6.1-7.8	0	0
	44-65	1.0-10	---	6.1-7.8	0	0
508: Medomak-----	0-7	29-43	---	5.1-7.3	0	0
	7-24	7.1-14	---	5.1-7.3	0	0
	24-33	29-43	---	5.1-7.3	0	0
	33-46	0.0-13	---	5.6-7.8	0	0
	46-79	0.0-13	---	5.6-7.8	0	0

Table 26.—Water Features

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated. Where no water tables are observed within the area of observation, depths are recorded as >6'.)

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
2: Ridgebury-----	D	January	0.0-0.5	1.7-2.5	---	---	None	---	None
		February	0.0-0.5	1.7-2.5	---	---	None	---	None
		March	0.0-0.5	1.7-2.5	---	---	None	---	None
		April	0.0-0.5	1.7-2.5	---	---	None	---	None
		May	0.0-0.5	1.7-2.5	---	---	None	---	None
		November	0.0-0.5	1.7-2.5	---	---	None	---	None
		December	0.0-0.5	1.7-2.5	---	---	None	---	None
3: Ridgebury-----	D	January	0.0-0.5	1.7-2.5	---	---	None	---	None
		February	0.0-0.5	1.7-2.5	---	---	None	---	None
		March	0.0-0.5	1.7-2.5	---	---	None	---	None
		April	0.0-0.5	1.7-2.5	---	---	None	---	None
		May	0.0-0.5	1.7-2.5	---	---	None	---	None
		November	0.0-0.5	1.7-2.5	---	---	None	---	None
		December	0.0-0.5	1.7-2.5	---	---	None	---	None
Leicester-----	D	January	0.0-1.5	>6.0	---	---	None	---	None
		February	0.0-1.5	>6.0	---	---	None	---	None
		March	0.0-1.5	>6.0	---	---	None	---	None
		April	0.0-1.5	>6.0	---	---	None	---	None
		May	0.0-1.5	>6.0	---	---	None	---	None
		November	0.0-1.5	>6.0	---	---	None	---	None
		December	0.0-1.5	>6.0	---	---	None	---	None
Whitman-----	D	January	0.0-1.0	1.0-2.5	0.0-1.0	Long	Occasional	---	None
		February	0.0-1.0	1.0-2.5	0.0-1.0	Long	Occasional	---	None
		March	0.0-1.0	1.0-2.5	0.0-1.0	Long	Occasional	---	None
		April	0.0-1.0	1.0-2.5	0.0-1.0	Long	Occasional	---	None
		May	0.0-1.0	1.0-2.5	0.0-1.0	Long	Occasional	---	None
		June	0.0-1.0	1.0-2.5	0.0-1.0	Brief	Occasional	---	None
		July	0.0-1.0	1.0-2.5	---	---	None	---	None
		August	0.0-1.0	1.0-2.5	---	---	None	---	None
		September	0.0-1.0	1.0-2.5	0.0-1.0	Brief	Occasional	---	None
		October	0.0-1.0	1.0-2.5	0.0-1.0	Brief	Occasional	---	None
		November	0.0-1.0	1.0-2.5	0.0-1.0	Brief	Occasional	---	None
		December	0.0-1.0	1.0-2.5	0.0-1.0	Long	Occasional	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
4: Leicester-----	D	January	0.0-1.5	>6.0	---	---	None	---	None
		February	0.0-1.5	>6.0	---	---	None	---	None
		March	0.0-1.5	>6.0	---	---	None	---	None
		April	0.0-1.5	>6.0	---	---	None	---	None
		May	0.0-1.5	>6.0	---	---	None	---	None
		November	0.0-1.5	>6.0	---	---	None	---	None
		December	0.0-1.5	>6.0	---	---	None	---	None
5: Wilbraham-----	D	January	0.0-1.5	1.7-3.0	---	---	None	---	None
		February	0.0-1.5	1.7-3.0	---	---	None	---	None
		March	0.0-1.5	1.7-3.0	---	---	None	---	None
		April	0.0-1.5	1.7-3.0	---	---	None	---	None
		May	0.0-1.5	1.7-3.0	---	---	None	---	None
		November	0.0-1.5	1.7-3.0	---	---	None	---	None
		December	0.0-1.5	1.7-3.0	---	---	None	---	None
6: Wilbraham-----	D	January	0.0-1.5	1.7-3.0	---	---	None	---	None
		February	0.0-1.5	1.7-3.0	---	---	None	---	None
		March	0.0-1.5	1.7-3.0	---	---	None	---	None
		April	0.0-1.5	1.7-3.0	---	---	None	---	None
		May	0.0-1.5	1.7-3.0	---	---	None	---	None
		November	0.0-1.5	1.7-3.0	---	---	None	---	None
		December	0.0-1.5	1.7-3.0	---	---	None	---	None
6: Menlo-----	D	January	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		February	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		March	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		April	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		May	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		June	0.0-1.0	1.5-3.0	---	---	None	---	None
		July	0.0-1.0	1.5-3.0	---	---	None	---	None
		August	0.0-1.0	1.5-3.0	---	---	None	---	None
		September	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		October	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		November	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None
		December	0.0-1.0	1.5-3.0	0.0-1.0	Long	Frequent	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
7: Mudgepond-----	D	January	0.0-1.0	5.4-5.4	---	---	None	---	None
		February	0.0-1.0	5.4-5.4	---	---	None	---	None
		March	0.0-1.0	5.4-5.4	---	---	None	---	None
		April	0.0-1.0	5.4-5.4	---	---	None	---	None
		May	0.0-1.0	5.4-5.4	---	---	None	---	None
		June	0.0-1.0	5.4-5.4	---	---	None	---	None
		July	0.0-1.0	5.4-5.4	---	---	None	---	None
		August	0.0-1.0	5.4-5.4	---	---	None	---	None
		September	0.0-1.0	5.4-5.4	---	---	None	---	None
		October	0.0-1.0	5.4-5.4	---	---	None	---	None
		November	0.0-1.0	5.4-5.4	---	---	None	---	None
		December	0.0-1.0	5.4-5.4	---	---	None	---	None
8: Mudgepond-----	D	January	0.0-1.0	5.4-5.4	---	---	None	---	None
		February	0.0-1.0	5.4-5.4	---	---	None	---	None
		March	0.0-1.0	5.4-5.4	---	---	None	---	None
		April	0.0-1.0	5.4-5.4	---	---	None	---	None
		May	0.0-1.0	5.4-5.4	---	---	None	---	None
		June	0.0-1.0	5.4-5.4	---	---	None	---	None
		July	0.0-1.0	5.4-5.4	---	---	None	---	None
		August	0.0-1.0	5.4-5.4	---	---	None	---	None
		September	0.0-1.0	5.4-5.4	---	---	None	---	None
		October	0.0-1.0	5.4-5.4	---	---	None	---	None
		November	0.0-1.0	5.4-5.4	---	---	None	---	None
		December	0.0-1.0	5.4-5.4	---	---	None	---	None
8: Alden-----	D	January	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		February	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		March	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		April	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		May	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		June	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		December	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding			
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency		
			Ft	Ft	Ft						
9: Scitico-----	D	January	0.0-1.0	>6.0	---	---	None	---	None		
		February	0.0-1.0	>6.0	---	---	None	---	None		
		March	0.0-1.0	>6.0	---	---	None	---	None		
		April	0.0-1.0	>6.0	---	---	None	---	None		
		May	0.0-1.0	>6.0	---	---	None	---	None		
		June	0.0-1.0	>6.0	---	---	None	---	None		
		October	0.0-1.0	>6.0	---	---	None	---	None		
		November	0.0-1.0	>6.0	---	---	None	---	None		
		December	0.0-1.0	>6.0	---	---	None	---	None		
		Shaker-----	D	January	0.0-1.5	>6.0	---	---	None	---	None
				February	0.0-1.5	>6.0	---	---	None	---	None
				March	0.0-1.5	>6.0	---	---	None	---	None
April	0.0-1.5			>6.0	---	---	None	---	None		
May	0.0-1.5			>6.0	---	---	None	---	None		
June	0.0-1.5			>6.0	---	---	None	---	None		
October	0.0-1.5			>6.0	---	---	None	---	None		
November	0.0-1.5			>6.0	---	---	None	---	None		
December	0.0-1.5			>6.0	---	---	None	---	None		
9: Maybid-----	D			January	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None
				February	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None
				March	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None
		April	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None		
		May	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None		
		June	0.0-0.5	>6.0	---	---	None	---	None		
		July	0.0-0.5	>6.0	---	---	None	---	None		
		August	0.0-0.5	>6.0	---	---	None	---	None		
		September	0.5-3.0	>6.0	---	---	None	---	None		
		October	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None		
		November	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None		
		December	0.0-0.5	>6.0	0.0-0.5	Long	Occasional	---	None		
10: Raynham-----	D	January	0.0-1.0	>6.0	---	---	None	---	None		
		February	0.0-1.0	>6.0	---	---	None	---	None		
		March	0.0-1.0	>6.0	---	---	None	---	None		
		April	0.0-1.0	>6.0	---	---	None	---	None		
		May	1.0-4.0	>6.0	---	---	None	---	None		
		November	0.0-1.0	>6.0	---	---	None	---	None		
		December	0.0-1.0	>6.0	---	---	None	---	None		

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
12: Raypol-----	D	January	0.0-1.0	>6.0	---	---	None	---	None
		February	0.0-1.0	>6.0	---	---	None	---	None
		March	0.0-1.0	>6.0	---	---	None	---	None
		April	0.0-1.0	>6.0	---	---	None	---	None
		May	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	---	---	None	---	None
		December	0.0-1.0	>6.0	---	---	None	---	None
13: Walpole-----	D	January	0.0-1.0	>6.0	---	---	None	---	None
		February	0.0-1.0	>6.0	---	---	None	---	None
		March	0.0-1.0	>6.0	---	---	None	---	None
		April	0.0-1.0	>6.0	---	---	None	---	None
		May	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	---	---	None	---	None
		December	0.0-1.0	>6.0	---	---	None	---	None
14: Fredon-----	D	January	0.0-1.0	>6.0	---	---	None	---	None
		February	0.0-1.0	>6.0	---	---	None	---	None
		March	0.0-1.0	>6.0	---	---	None	---	None
		April	0.0-1.0	>6.0	---	---	None	---	None
		May	0.0-1.0	>6.0	---	---	None	---	None
		June	0.0-1.0	>6.0	---	---	None	---	None
		July	1.5-3.0	>6.0	---	---	None	---	None
		August	1.5-3.0	>6.0	---	---	None	---	None
		September	1.5-3.0	>6.0	---	---	None	---	None
		October	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	---	---	None	---	None
		December	0.0-1.0	>6.0	---	---	None	---	None
15: Scarboro-----	D	January	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		February	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		March	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		April	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		May	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		June	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		July	0.0-1.0	>6.0	---	---	None	---	None
		August	0.0-1.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		November	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		December	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
16: Halsey-----	D	January	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		February	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		March	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		April	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		May	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		June	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		July	0.0-1.0	>6.0	---	---	None	---	None
		August	0.0-1.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		November	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		December	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
17: Timakwa-----	D	January	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		February	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		March	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		April	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		May	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		June	1.0-2.0	>6.0	---	---	None	---	None
		July	1.0-2.0	>6.0	---	---	None	---	None
		August	1.0-2.0	>6.0	---	---	None	---	None
		September	1.0-2.0	>6.0	---	---	None	---	None
		October	1.0-2.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		December	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
Natchaug-----	D	January	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		February	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		March	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		April	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		May	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		June	0.0-1.0	>6.0	---	---	None	---	None
		July	0.0-1.0	>6.0	---	---	None	---	None
		August	0.0-1.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		December	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
18: Catden-----	D	January	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		February	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		March	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		April	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		May	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		June	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		July	1.0-2.0	>6.0	---	---	None	---	None
		August	1.0-2.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		October	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		November	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		December	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
18: Freetown-----	D	January	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		February	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		March	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		April	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		May	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		June	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		July	0.8-2.0	>6.0	---	---	None	---	None
		August	0.8-2.0	>6.0	---	---	None	---	None
		September	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		October	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		November	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		December	0.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
20A: Ellington-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
21A: Ninigret-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		May	2.5-5.0	>6.0	---	---	None	---	None
		September	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
21A: Tisbury-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		May	2.5-5.0	>6.0	---	---	None	---	None
		September	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
22A: Hero-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
22B: Hero-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
23A: Sudbury-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
24A: Deerfield-----	A	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
25A: Brancroft-----	C	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		October	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
25B: Brancroft-----	C	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		October	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
25C: Brancroft-----	C	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		October	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
26A: Berlin-----	C	January	1.0-2.5	>6.0	---	---	None	---	None
		February	1.0-2.5	>6.0	---	---	None	---	None
		March	1.0-2.5	>6.0	---	---	None	---	None
		April	1.0-2.5	>6.0	---	---	None	---	None
		October	1.0-2.5	>6.0	---	---	None	---	None
		November	1.0-2.5	>6.0	---	---	None	---	None
		December	1.0-2.5	>6.0	---	---	None	---	None
26B: Berlin-----	C	January	1.0-2.5	>6.0	---	---	None	---	None
		February	1.0-2.5	>6.0	---	---	None	---	None
		March	1.0-2.5	>6.0	---	---	None	---	None
		April	1.0-2.5	>6.0	---	---	None	---	None
		October	1.0-2.5	>6.0	---	---	None	---	None
		November	1.0-2.5	>6.0	---	---	None	---	None
		December	1.0-2.5	>6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
27A: Belgrade-----	B	January	1.5-3.5	>6.0	---	---	None	---	None
		February	1.5-3.5	>6.0	---	---	None	---	None
		March	1.5-3.5	>6.0	---	---	None	---	None
		April	1.5-3.5	>6.0	---	---	None	---	None
		November	1.5-3.5	>6.0	---	---	None	---	None
		December	1.5-3.5	>6.0	---	---	None	---	None
28A: Elmridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
28B: Elmridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
29A: Agawam-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
29B: Agawam-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
29C: Agawam-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
30A: Branford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
30B: Branford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
30C: Branford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features--Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding		Flooding		
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
31A: Copake-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
31B: Copake-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
31C: Copake-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
32A: Haven-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Enfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
32B: Haven-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Enfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
32C: Haven-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Enfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
33A: Hartford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
33B: Hartford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
34A: Merrimac-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
34B: Merrimac-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
34C: Merrimac-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
35A: Penwood-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
35B: Penwood-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
36A: Windsor-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
36B: Windsor-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
36C: Windsor-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
37A: Manchester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
37C: Manchester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
37E: Manchester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
38A: Hinckley-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
38C: Hinckley-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
38E: Hinckley-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
39A: Groton-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
39C: Groton-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
39E: Groton-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
40A: Ludlow-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
40B: Ludlow-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
41B: Ludlow-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
42C: Ludlow-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
43A: Rainbow-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
February		1.5-2.5	1.7-3.3	---	---	None	---	None	
March		1.5-2.5	1.7-3.3	---	---	None	---	None	
April		1.5-2.5	1.7-3.3	---	---	None	---	None	
May		1.5-2.5	1.7-3.3	---	---	None	---	None	
November		1.5-2.5	1.7-3.3	---	---	None	---	None	
December		1.5-2.5	1.7-3.3	---	---	None	---	None	
43B: Rainbow-----		C	January	1.5-2.5	1.7-3.3	---	---	None	---
February	1.5-2.5		1.7-3.3	---	---	None	---	None	
March	1.5-2.5		1.7-3.3	---	---	None	---	None	
April	1.5-2.5		1.7-3.3	---	---	None	---	None	
May	1.5-2.5		1.7-3.3	---	---	None	---	None	
November	1.5-2.5		1.7-3.3	---	---	None	---	None	
December	1.5-2.5		1.7-3.3	---	---	None	---	None	
44B: Rainbow-----	C		January	1.5-2.5	1.7-3.3	---	---	None	---
February		1.5-2.5	1.7-3.3	---	---	None	---	None	
March		1.5-2.5	1.7-3.3	---	---	None	---	None	
April		1.5-2.5	1.7-3.3	---	---	None	---	None	
May		1.5-2.5	1.7-3.3	---	---	None	---	None	
November		1.5-2.5	1.7-3.3	---	---	None	---	None	
December		1.5-2.5	1.7-3.3	---	---	None	---	None	
45A: Woodbridge-----		C	January	1.5-2.5	1.7-3.3	---	---	None	---
February	1.5-2.5		1.7-3.3	---	---	None	---	None	
March	1.5-2.5		1.7-3.3	---	---	None	---	None	
April	1.5-2.5		1.7-3.3	---	---	None	---	None	
May	1.5-2.5		1.7-3.3	---	---	None	---	None	
November	1.5-2.5		1.7-3.3	---	---	None	---	None	
December	1.5-2.5		1.7-3.3	---	---	None	---	None	
45B: Woodbridge-----	C		January	1.5-2.5	1.7-3.3	---	---	None	---
February		1.5-2.5	1.7-3.3	---	---	None	---	None	
March		1.5-2.5	1.7-3.3	---	---	None	---	None	
April		1.5-2.5	1.7-3.3	---	---	None	---	None	
May		1.5-2.5	1.7-3.3	---	---	None	---	None	
November		1.5-2.5	1.7-3.3	---	---	None	---	None	
December		1.5-2.5	1.7-3.3	---	---	None	---	None	

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
45C: Woodbridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
46B: Woodbridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
46C: Woodbridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
47C: Woodbridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
48B: Georgia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
48B: Amenia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
48C: Georgia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
Amenia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
49B: Georgia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
Amenia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
49C: Georgia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
49C: Amenia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		May	1.5-3.0	>6.0	---	---	None	---	None
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
50A: Sutton-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
50B: Sutton-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
51B: Sutton-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
52C: Sutton-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
53A: Wapping-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
53B: Wapping-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
54B: Wapping-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
55A: Watchaug-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
55B: Watchaug-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
56B: Watchaug-----		B	January	1.5-2.5	>6.0	---	---	None	---
	February		1.5-2.5	>6.0	---	---	None	---	None
	March		1.5-2.5	>6.0	---	---	None	---	None
	April		1.5-2.5	>6.0	---	---	None	---	None
	November		1.5-2.5	>6.0	---	---	None	---	None
	December		1.5-2.5	>6.0	---	---	None	---	None
57B: Gloucester-----	B		Jan-Dec	> 6.0	> 6.0	---	---	None	---
57C: Gloucester-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
57D: Gloucester-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
58B: Gloucester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
58C: Gloucester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
59C: Gloucester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
59D: Gloucester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding		Flooding		
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
60B: Canton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
60C: Canton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
60C: Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
60D: Canton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
61B: Canton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
61C: Canton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
62C: Canton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
62D: Canton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
63B: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
63C: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
63D: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
64B: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
64C: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
65C: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
65D: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
66B: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
66C: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
67B: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
67C: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
68C: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
68D: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
69B: Yalesville-----	C	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
69C: Yalesville-----	C	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
70C: Branford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Holyoke-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
71C: Brookfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Brimfield-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
71E: Brookfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Brimfield-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
73C: Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
73E: Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
74C: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Hollis-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
75C: Hollis-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
75E: Hollis-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
76E: Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Hollis-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.--Water Features--Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
76F: Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Hollis-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
77C: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Holyoke-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
77D: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Holyoke-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
78C: Holyoke-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
78E: Holyoke-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
79E: Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Holyoke-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
80B: Bernardston-----	C	February	1.5-2.0	1.7-2.5	---	---	None	---	None
		March	1.5-2.0	1.7-2.5	---	---	None	---	None
		April	1.5-2.0	1.7-2.5	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
80C: Bernardston-----	C	February	1.5-2.0	1.7-2.5	---	---	None	---	None
		March	1.5-2.0	1.7-2.5	---	---	None	---	None
		April	1.5-2.0	1.7-2.5	---	---	None	---	None
81C: Bernardston-----	C	February	1.5-2.0	1.7-2.5	---	---	None	---	None
		March	1.5-2.0	1.7-2.5	---	---	None	---	None
		April	1.5-2.0	1.7-2.5	---	---	None	---	None
81D: Bernardston-----	C	February	1.5-2.0	1.7-2.5	---	---	None	---	None
		March	1.5-2.0	1.7-2.5	---	---	None	---	None
		April	1.5-2.0	1.7-2.5	---	---	None	---	None
82B: Broadbrook-----	C	March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
82C: Broadbrook-----	C	March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
82D: Broadbrook-----	C	March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
83B: Broadbrook-----	C	March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
83C: Broadbrook-----	C	March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
84B: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Montauk-----	C	January	2.0-2.5	2.2-3.2	---	---	None	---	None
		February	2.0-2.5	2.2-3.2	---	---	None	---	None
		March	2.0-2.5	2.2-3.2	---	---	None	---	None
		April	2.0-2.5	2.2-3.2	---	---	None	---	None
		November	2.0-2.5	2.2-3.2	---	---	None	---	None
		December	2.0-2.5	2.2-3.2	---	---	None	---	None
84C: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Montauk-----	C	January	2.0-2.5	2.2-3.2	---	---	None	---	None
		February	2.0-2.5	2.2-3.2	---	---	None	---	None
		March	2.0-2.5	2.2-3.2	---	---	None	---	None
		April	2.0-2.5	2.2-3.2	---	---	None	---	None
		November	2.0-2.5	2.2-3.2	---	---	None	---	None
		December	2.0-2.5	2.2-3.2	---	---	None	---	None
84D: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
84D: Montauk-----	C	January	2.0-2.5	2.2-3.2	---	---	None	---	None
		February	2.0-2.5	2.2-3.2	---	---	None	---	None
		March	2.0-2.5	2.2-3.2	---	---	None	---	None
		April	2.0-2.5	2.2-3.2	---	---	None	---	None
		November	2.0-2.5	2.2-3.2	---	---	None	---	None
		December	2.0-2.5	2.2-3.2	---	---	None	---	None
85B: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
85B: Montauk-----	C	January	2.0-2.5	2.2-3.2	---	---	None	---	None
		February	2.0-2.5	2.2-3.2	---	---	None	---	None
		March	2.0-2.5	2.2-3.2	---	---	None	---	None
		April	2.0-2.5	2.2-3.2	---	---	None	---	None
		November	2.0-2.5	2.2-3.2	---	---	None	---	None
		December	2.0-2.5	2.2-3.2	---	---	None	---	None
85C: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Montauk-----	C	January	2.0-2.5	2.2-3.2	---	---	None	---	None
		February	2.0-2.5	2.2-3.2	---	---	None	---	None
		March	2.0-2.5	2.2-3.2	---	---	None	---	None
		April	2.0-2.5	2.2-3.2	---	---	None	---	None
		November	2.0-2.5	2.2-3.2	---	---	None	---	None
		December	2.0-2.5	2.2-3.2	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
86C: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Montauk-----		C	January	2.0-2.5	2.2-3.2	---	---	None	---
	February		2.0-2.5	2.2-3.2	---	---	None	---	None
	March		2.0-2.5	2.2-3.2	---	---	None	---	None
	April		2.0-2.5	2.2-3.2	---	---	None	---	None
	November		2.0-2.5	2.2-3.2	---	---	None	---	None
	December		2.0-2.5	2.2-3.2	---	---	None	---	None
86D: Paxton-----	C		January	1.5-2.5	1.7-3.3	---	---	None	---
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Montauk-----		C	January	2.0-2.5	2.2-3.2	---	---	None	---
	February		2.0-2.5	2.2-3.2	---	---	None	---	None
	March		2.0-2.5	2.2-3.2	---	---	None	---	None
	April		2.0-2.5	2.2-3.2	---	---	None	---	None
	November		2.0-2.5	2.2-3.2	---	---	None	---	None
	December		2.0-2.5	2.2-3.2	---	---	None	---	None
87B: Wethersfield-----	C		February	1.5-2.5	1.7-3.3	---	---	None	---
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
87C: Wethersfield-----		C	February	1.5-2.5	1.7-3.3	---	---	None	---
	March		1.5-2.5	1.7-3.3	---	---	None	---	None
	April		1.5-2.5	1.7-3.3	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
87D: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
88B: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
88C: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
89C: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
89D: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
90B: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
90C: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
90D: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
91B: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
91C: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
91D: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
92B: Nellis-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
92C: Nellis-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
92D: Nellis-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
93C: Nellis-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
94C: Farmington-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Nellis-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
94E: Farmington-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Nellis-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
95C: Farmington-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
95E: Farmington-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
96: Ipswich-----	D	Jan-Dec	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Very frequent
97: Pawcatuck-----	D	Jan-Dec	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Very
98: Westbrook-----	D	Jan-Dec	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Very
99: Westbrook, low salt-----	D	Jan-Dec	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Very
100: Suncook-----	A	January	5.0-6.0	>6.0	---	---	None	---	None
		February	5.0-6.0	>6.0	---	---	None	---	None
		March	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		April	5.0-6.0	>6.0	---	---	None	Brief	Occasional
101: Occum-----	B	January	5.0-6.0	>6.0	---	---	None	---	None
		February	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		March	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		April	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		November	5.0-6.0	>6.0	---	---	None	---	None
		December	5.0-6.0	>6.0	---	---	None	---	None
102: Pootatuck-----	B	January	1.5-2.5	>6.0	---	---	None	Brief	Frequent
		February	1.5-2.5	>6.0	---	---	None	Brief	Frequent
		March	1.5-2.5	>6.0	---	---	None	Brief	Frequent
		April	1.5-2.5	>6.0	---	---	None	Brief	Frequent
		November	1.5-2.5	>6.0	---	---	None	Brief	Frequent
		December	1.5-2.5	>6.0	---	---	None	Brief	Frequent

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
103: Rippowam-----	D	January	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		February	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		March	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		April	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		May	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		June	0.0-1.5	>6.0	---	---	None	---	None
		September	0.0-1.5	>6.0	---	---	None	---	None
		October	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		November	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		December	0.0-1.5	>6.0	---	---	None	Brief	Frequent
104: Bash-----	D	January	0.5-1.5	>6.0	---	---	None	Brief	Frequent
		February	0.5-1.5	>6.0	---	---	None	Long	Frequent
		March	0.5-1.5	>6.0	---	---	None	Long	Frequent
		April	0.5-1.5	>6.0	---	---	None	Long	Frequent
		May	0.5-1.5	>6.0	---	---	None	Long	Frequent
		June	1.5-3.5	>6.0	---	---	None	---	None
		July	3.5-6.0	>6.0	---	---	None	---	None
		August	3.5-6.0	>6.0	---	---	None	---	None
		September	1.5-3.5	>6.0	---	---	None	---	None
		October	1.5-3.5	>6.0	---	---	None	---	None
		November	1.5-3.5	>6.0	---	---	None	---	None
		December	0.5-1.5	>6.0	---	---	None	Brief	Frequent
105: Hadley-----	B	January	5.0-6.0	>6.0	---	---	None	---	None
		February	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		March	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		April	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		November	5.0-6.0	>6.0	---	---	None	---	None
		December	5.0-6.0	>6.0	---	---	None	---	None
106: Winooski-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	Brief	Frequent
		March	1.5-3.0	>6.0	---	---	None	Brief	Frequent
		April	1.5-3.0	>6.0	---	---	None	Brief	Frequent
		November	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
107: Limerick-----	D	January	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		February	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		March	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		April	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		May	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		June	0.0-1.5	>6.0	---	---	None	---	None
		September	0.0-1.5	>6.0	---	---	None	---	None
		October	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		November	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		December	0.0-1.5	>6.0	---	---	None	Brief	Frequent
Lim-----	D	January	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		February	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		March	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		April	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		May	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		June	0.0-1.5	>6.0	---	---	None	---	None
		September	0.0-1.5	>6.0	---	---	None	---	None
		October	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		November	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		December	0.0-1.5	>6.0	---	---	None	Brief	Frequent
108: Saco-----	D	January	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		February	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		March	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		April	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		May	0.0-0.5	>6.0	---	---	None	Brief	Frequent
		June	0.0-0.5	>6.0	---	---	None	---	None
		July	0.0-0.5	>6.0	---	---	None	---	None
		August	0.0-0.5	>6.0	---	---	None	---	None
		September	0.0-0.5	>6.0	---	---	None	---	None
		October	0.0-0.5	>6.0	---	---	None	Brief	Frequent
		November	0.0-0.5	>6.0	---	---	None	Brief	Frequent
		December	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
109: Fluvaquents, Frequently Flooded-----	D	January	0.0-1.0	>6.0	---	---	None	Brief	Frequent
		February	0.0-1.0	>6.0	---	---	None	Brief	Frequent
		March	0.0-1.0	>6.0	---	---	None	Brief	Frequent
		April	0.0-1.0	>6.0	---	---	None	Brief	Frequent
		May	0.0-1.0	>6.0	---	---	None	Brief	Frequent
		June	1.5-2.5	>6.0	---	---	None	Brief	Frequent
		July	1.5-2.5	>6.0	---	---	None	---	None
		August	1.5-2.5	>6.0	---	---	None	---	None
		September	1.5-2.5	>6.0	---	---	None	Brief	Frequent
		October	0.0-1.0	>6.0	---	---	None	Brief	Frequent
		November	0.0-1.0	>6.0	---	---	None	Brief	Frequent
		December	0.0-1.0	>6.0	---	---	None	Brief	Frequent
Udifluents, Frequently Flooded-----	B	January	6.0	>6.0	---	---	None	Brief	Frequent
		February	6.0	>6.0	---	---	None	Brief	Frequent
		March	6.0	>6.0	---	---	None	Brief	Frequent
		April	6.0	>6.0	---	---	None	Brief	Frequent
		May	6.0	>6.0	---	---	None	Brief	Frequent
		June	6.0	>6.0	---	---	None	Brief	Frequent
		July	6.0	>6.0	---	---	None	---	None
		August	6.0	>6.0	---	---	None	---	None
		September	6.0	>6.0	---	---	None	Brief	Frequent
		October	6.0	>6.0	---	---	None	Brief	Frequent
		November	6.0	>6.0	---	---	None	Brief	Frequent
		December	6.0	>6.0	---	---	None	Brief	Frequent
221A: Ninigret-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		May	2.5-5.0	>6.0	---	---	None	---	None
		September	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
Urban Land-----	---								

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
224A: Deerfield-----	A	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
Urban Land-----	---								
225B: Brancroft-----	C	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		October	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
Urban Land-----	---								
226B: Berlin-----	C	January	1.0-2.5	>6.0	---	---	None	---	None
		February	1.0-2.5	>6.0	---	---	None	---	None
		March	1.0-2.5	>6.0	---	---	None	---	None
		April	1.0-2.5	>6.0	---	---	None	---	None
		October	1.0-2.5	>6.0	---	---	None	---	None
		November	1.0-2.5	>6.0	---	---	None	---	None
		December	1.0-2.5	>6.0	---	---	None	---	None
Urban Land-----	---								
228B: Elmridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
229B: Agawam-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
229C: Agawam-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
230B: Branford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
230C: Branford-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
232B: Haven-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
234B: Merrimac-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
235B: Penwood-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
236B: Windsor-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
237A: Manchester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
237C: Manchester-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
238A: Hinckley-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
238C: Hinckley-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
240B: Ludlow-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								
243B: Rainbow-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		May	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft					
245B: Woodbridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
February		1.5-2.5	1.7-3.3	---	---	None	---	None	
March		1.5-2.5	1.7-3.3	---	---	None	---	None	
April		1.5-2.5	1.7-3.3	---	---	None	---	None	
May		1.5-2.5	1.7-3.3	---	---	None	---	None	
November		1.5-2.5	1.7-3.3	---	---	None	---	None	
December		1.5-2.5	1.7-3.3	---	---	None	---	None	
Urban Land-----		---							
245C: Woodbridge-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
February		1.5-2.5	1.7-3.3	---	---	None	---	None	
March		1.5-2.5	1.7-3.3	---	---	None	---	None	
April		1.5-2.5	1.7-3.3	---	---	None	---	None	
May		1.5-2.5	1.7-3.3	---	---	None	---	None	
November		1.5-2.5	1.7-3.3	---	---	None	---	None	
December		1.5-2.5	1.7-3.3	---	---	None	---	None	
Urban Land-----		---							
248B: Georgia-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
February		1.5-3.0	>6.0	---	---	None	---	None	
March		1.5-3.0	>6.0	---	---	None	---	None	
April		1.5-3.0	>6.0	---	---	None	---	None	
May		1.5-3.0	>6.0	---	---	None	---	None	
November		1.5-3.0	>6.0	---	---	None	---	None	
December		1.5-3.0	>6.0	---	---	None	---	None	
Urban Land-----		---							
248B: Urban Land-----	---								
250B: Sutton-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
February		1.5-2.5	>6.0	---	---	None	---	None	
March		1.5-2.5	>6.0	---	---	None	---	None	
April		1.5-2.5	>6.0	---	---	None	---	None	
November		1.5-2.5	>6.0	---	---	None	---	None	
December		1.5-2.5	>6.0	---	---	None	---	None	
Urban Land-----		---							

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
253B: Wapping-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
Urban Land-----	---								
255B: Watchaug-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
Urban Land-----	---								
260B: Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
260C: Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
260D: Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
263B: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
263C: Cheshire-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
266B: Narragansett-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
269B: Yalesville-----	C	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
269C: Yalesville-----	C	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
273C: Urban Land-----	---								
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
273E: Urban Land-----	---								
Charlton-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
275C: Urban Land-----	---								
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
275E: Urban Land-----	---								
Chatfield-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
282B: Broadbrook-----	C	March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								
284B: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								
284C: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								
284D: Paxton-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
287B: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								
287C: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								
287D: Wethersfield-----	C	February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
Urban Land-----	---								
290B: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
290C: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
290D: Stockbridge-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Urban Land-----	---								
301: Beaches-----	D	Jan-Dec	0.0-1.0	>6.0	---	---	None	Very brief	Very frequent

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
301: Udipsamments-----	A	Jan-Dec	4.0-6.0	>6.0	---	---	None	Extremely brief	Occasional
302: Dumps-----	---	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
303: Pits, Quarries-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
304: Udorthents-----	B	January	4.5-6.0	>6.0	---	---	None	---	None
		February	4.5-6.0	>6.0	---	---	None	---	None
		March	4.5-6.0	>6.0	---	---	None	---	None
		April	4.5-6.0	>6.0	---	---	None	---	None
		November	4.5-6.0	>6.0	---	---	None	---	None
		December	4.5-6.0	>6.0	---	---	None	---	None
305: Udorthents-----	B	January	2.0-4.5	>6.0	---	---	None	---	None
		February	2.0-4.5	>6.0	---	---	None	---	None
		March	2.0-4.5	>6.0	---	---	None	---	None
		April	2.0-4.5	>6.0	---	---	None	---	None
		November	2.0-4.5	>6.0	---	---	None	---	None
		December	2.0-4.5	>6.0	---	---	None	---	None
Pits-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
306: Udorthents-----	B	January	4.5-6.0	>6.0	---	---	None	---	None
		February	4.5-6.0	>6.0	---	---	None	---	None
		March	4.5-6.0	>6.0	---	---	None	---	None
		April	4.5-6.0	>6.0	---	---	None	---	None
		November	4.5-6.0	>6.0	---	---	None	---	None
		December	4.5-6.0	>6.0	---	---	None	---	None
Urban Land-----	---								
307: Urban Land-----	---								

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
308: Udorthents-----	B	January	2.0-4.5	>6.0	---	---	None	---	None
		February	2.0-4.5	>6.0	---	---	None	---	None
		March	2.0-4.5	>6.0	---	---	None	---	None
		April	2.0-4.5	>6.0	---	---	None	---	None
		November	2.0-4.5	>6.0	---	---	None	---	None
		December	2.0-4.5	>6.0	---	---	None	---	None
309: Udorthents-----		B	January	2.0-4.5	>6.0	---	---	None	Very brief
	February		2.0-4.5	>6.0	---	---	None	Very brief	Rare
	March		2.0-4.5	>6.0	---	---	None	Very brief	Rare
	April		2.0-4.5	>6.0	---	---	None	Very brief	Rare
	May		---	---	---	---	None	Very brief	Rare
	June		---	---	---	---	None	Very brief	Very rare
	July		---	---	---	---	None	Very brief	Very rare
	August		---	---	---	---	None	Very brief	Rare
	September		---	---	---	---	None	Very brief	Rare
	October		---	---	---	---	None	Very brief	Rare
	November		2.0-4.5	>6.0	---	---	None	Very brief	Rare
	December		2.0-4.5	>6.0	---	---	None	Very brief	Rare
310: Udorthents, Periodically Flooded-----	B	January	2.0-4.5	>6.0	---	---	None	Very brief	Occasional
		February	2.0-4.5	>6.0	---	---	None	Very brief	Occasional
		March	2.0-4.5	>6.0	---	---	None	Very brief	Frequent
		April	2.0-4.5	>6.0	---	---	None	Very brief	Frequent
		May	---	---	---	---	None	Very brief	Frequent
		June	---	---	---	---	None	Very brief	Rare
		July	---	---	---	---	None	Very brief	Very rare
		August	---	---	---	---	None	Very brief	Very rare
		September	---	---	---	---	None	Very brief	Rare
		October	---	---	---	---	None	Very brief	Rare
		November	2.0-4.5	>6.0	---	---	None	Very brief	Occasional
		December	2.0-4.5	>6.0	---	---	None	Very brief	Occasional
401C: Macomber-----	C	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Taconic-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
402D: Macomber-----	C	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Taconic-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
403C: Taconic-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
403E: Taconic-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
403F: Taconic-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
405C: Dummerston-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
405E: Dummerston-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
407C: Lanesboro-----	C	February	1.5-2.5	1.7-2.9	---	---	None	---	None
		March	1.5-2.5	1.7-2.9	---	---	None	---	None
		April	1.5-2.5	1.7-2.9	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
407E: Lanesboro-----	C	February	1.5-2.5	1.7-2.9	---	---	None	---	None
		March	1.5-2.5	1.7-2.9	---	---	None	---	None
		April	1.5-2.5	1.7-2.9	---	---	None	---	None
408C: Fullam-----	C	January	1.5-2.5	1.7-3.0	---	---	None	---	None
		February	1.5-2.5	1.7-3.0	---	---	None	---	None
		March	1.5-2.5	1.7-3.0	---	---	None	---	None
		April	1.5-2.5	1.7-3.0	---	---	None	---	None
		November	1.5-2.5	1.7-3.0	---	---	None	---	None
		December	1.5-2.5	1.7-3.0	---	---	None	---	None
409B: Brayton-----	D	January	0.0-1.0	0.8-2.2	---	---	None	---	None
		February	0.0-1.0	0.8-2.2	---	---	None	---	None
		March	0.0-1.0	0.8-2.2	---	---	None	---	None
		April	0.0-1.0	0.8-2.2	---	---	None	---	None
		May	0.0-1.0	0.8-2.2	---	---	None	---	None
		June	0.0-1.0	0.8-2.2	---	---	None	---	None
		October	0.0-1.0	0.8-2.2	---	---	None	---	None
		November	0.0-1.0	0.8-2.2	---	---	None	---	None
		December	0.0-1.0	0.8-2.2	---	---	None	---	None
412B: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
412C: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
412D: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
413C: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Millsite-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
413E: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Millsite-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
414: Fredon, cold-----	D	January	0.0-1.0	>6.0	---	---	None	---	None
		February	0.0-1.0	>6.0	---	---	None	---	None
		March	0.0-1.0	>6.0	---	---	None	---	None
		April	0.0-1.0	>6.0	---	---	None	---	None
		May	0.0-1.0	>6.0	---	---	None	---	None
		June	0.0-1.0	>6.0	---	---	None	---	None
		July	1.5-3.0	>6.0	---	---	None	---	None
		August	1.5-3.0	>6.0	---	---	None	---	None
		September	1.5-3.0	>6.0	---	---	None	---	None
		October	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	---	---	None	---	None
		December	0.0-1.0	>6.0	---	---	None	---	None
415C: Millsite-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Westminster-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
415E: Millsite-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Westminster-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
416E: Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Westminster-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
Rock Outcrop-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
416F: Westminster-----	D	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
417B: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
417C: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
417D: Bice-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
418C: Schroon-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
420A: Schroon-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None

Table 26.--Water Features--Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
420B: Schroon-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
421A: Ninigret, cold-----	B	January	1.5-2.5	>6.0	---	---	None	---	None
		February	1.5-2.5	>6.0	---	---	None	---	None
		March	1.5-2.5	>6.0	---	---	None	---	None
		April	1.5-2.5	>6.0	---	---	None	---	None
		May	2.5-5.0	>6.0	---	---	None	---	None
		September	1.5-2.5	>6.0	---	---	None	---	None
		November	1.5-2.5	>6.0	---	---	None	---	None
		December	1.5-2.5	>6.0	---	---	None	---	None
423A: Sudbury, cold-----	B	January	1.5-3.0	>6.0	---	---	None	---	None
		February	1.5-3.0	>6.0	---	---	None	---	None
		March	1.5-3.0	>6.0	---	---	None	---	None
		April	1.5-3.0	>6.0	---	---	None	---	None
		December	1.5-3.0	>6.0	---	---	None	---	None
424B: Shelburne-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
424C: Shelburne-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
424D: Shelburne-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
425B: Shelburne-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
425C: Shelburne-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
426D: Shelburne-----	C	January	1.5-2.5	1.7-3.3	---	---	None	---	None
		February	1.5-2.5	1.7-3.3	---	---	None	---	None
		March	1.5-2.5	1.7-3.3	---	---	None	---	None
		April	1.5-2.5	1.7-3.3	---	---	None	---	None
		November	1.5-2.5	1.7-3.3	---	---	None	---	None
		December	1.5-2.5	1.7-3.3	---	---	None	---	None
427B: Ashfield-----	C	January	1.0-2.0	1.1-2.2	---	---	None	---	None
		February	1.0-2.0	1.1-2.2	---	---	None	---	None
		March	1.0-2.0	1.1-2.2	---	---	None	---	None
		April	1.0-2.0	1.1-2.2	---	---	None	---	None
		May	1.0-2.0	1.1-2.2	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
427C: Ashfield-----	C	January	1.0-2.0	1.1-2.2	---	---	None	---	None
		February	1.0-2.0	1.1-2.2	---	---	None	---	None
		March	1.0-2.0	1.1-2.2	---	---	None	---	None
		April	1.0-2.0	1.1-2.2	---	---	None	---	None
		May	1.0-2.0	1.1-2.2	---	---	None	---	None
428A: Ashfield-----	C	January	1.0-2.0	1.1-2.2	---	---	None	---	None
		February	1.0-2.0	1.1-2.2	---	---	None	---	None
		March	1.0-2.0	1.1-2.2	---	---	None	---	None
		April	1.0-2.0	1.1-2.2	---	---	None	---	None
		May	1.0-2.0	1.1-2.2	---	---	None	---	None
428B: Ashfield-----	C	January	1.0-2.0	1.1-2.2	---	---	None	---	None
		February	1.0-2.0	1.1-2.2	---	---	None	---	None
		March	1.0-2.0	1.1-2.2	---	---	None	---	None
		April	1.0-2.0	1.1-2.2	---	---	None	---	None
		May	1.0-2.0	1.1-2.2	---	---	None	---	None
428C: Ashfield-----	C	January	1.0-2.0	1.1-2.2	---	---	None	---	None
		February	1.0-2.0	1.1-2.2	---	---	None	---	None
		March	1.0-2.0	1.1-2.2	---	---	None	---	None
		April	1.0-2.0	1.1-2.2	---	---	None	---	None
		May	1.0-2.0	1.1-2.2	---	---	None	---	None
429A: Agawam, cold-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
429B: Agawam, cold-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
429C: Agawam, cold-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
433: Moosilauke-----	D	January	0.0-1.0	>6.0	---	---	None	---	None
		February	0.0-1.0	>6.0	---	---	None	---	None
		March	0.0-1.0	>6.0	---	---	None	---	None
		April	0.0-1.0	>6.0	---	---	None	---	None
		May	0.0-1.0	>6.0	---	---	None	---	None
		June	0.0-1.0	>6.0	---	---	None	---	None
		August	0.0-1.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	---	---	None	---	None
		December	0.0-1.0	>6.0	---	---	None	---	None
434A: Merrimac, cold-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
434B: Merrimac, cold-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
434C: Merrimac, cold-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
435: Scarboro-----	D	January	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		February	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		March	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		April	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		May	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		June	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		July	0.0-1.0	>6.0	---	---	None	---	None
		August	0.0-1.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		November	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		December	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
436: Halsey-----	D	January	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		February	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		March	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		April	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		May	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		June	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		July	0.0-1.0	>6.0	---	---	None	---	None
		August	0.0-1.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		November	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
		December	0.0-0.5	>6.0	0.0-0.5	Brief	Occasional	---	None
437: Wonsqueak-----	D	January	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		February	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		March	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		April	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		May	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		June	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		July	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		August	0.0-3.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		October	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		November	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		December	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
438: Bucksport-----	D	January	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		February	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		March	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		April	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		May	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		June	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		July	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		August	0.0-3.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		October	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		November	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare
		December	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	Very brief	Rare

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
440A: Boscawen-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
440C: Boscawen-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
440E: Boscawen-----	A	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
442: Brayton-----	D	January	0.0-1.0	0.8-2.2	---	---	None	---	None
		February	0.0-1.0	0.8-2.2	---	---	None	---	None
		March	0.0-1.0	0.8-2.2	---	---	None	---	None
		April	0.0-1.0	0.8-2.2	---	---	None	---	None
		May	0.0-1.0	0.8-2.2	---	---	None	---	None
		June	0.0-1.0	0.8-2.2	---	---	None	---	None
		October	0.0-1.0	0.8-2.2	---	---	None	---	None
		November	0.0-1.0	0.8-2.2	---	---	None	---	None
		December	0.0-1.0	0.8-2.2	---	---	None	---	None
443: Brayton-----	D	January	0.0-1.0	0.8-2.2	---	---	None	---	None
		February	0.0-1.0	0.8-2.2	---	---	None	---	None
		March	0.0-1.0	0.8-2.2	---	---	None	---	None
		April	0.0-1.0	0.8-2.2	---	---	None	---	None
		May	0.0-1.0	0.8-2.2	---	---	None	---	None
		June	0.0-1.0	0.8-2.2	---	---	None	---	None
		October	0.0-1.0	0.8-2.2	---	---	None	---	None
		November	0.0-1.0	0.8-2.2	---	---	None	---	None
		December	0.0-1.0	0.8-2.2	---	---	None	---	None

Table 26.-Water Features-Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
443: Loonmeadow-----	D	January	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	---	None
		February	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	---	None
		March	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	---	None
		April	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	---	None
		May	0.0-1.0	>6.0	0.0-1.0	Brief	Frequent	---	None
		June	0.0-1.0	>6.0	---	---	None	---	None
		July	0.0-1.0	>6.0	---	---	None	---	None
		August	0.0-1.0	>6.0	---	---	None	---	None
		September	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	0.0-1.0	Brief	Frequent	---	None
		December	0.0-1.0	>6.0	0.0-1.0	Long	Frequent	---	None
448B: Hogansburg-----	B	January	1.5-3.0	3.0-3.5	---	---	None	---	None
		February	1.5-3.0	3.0-3.5	---	---	None	---	None
		March	1.5-3.0	3.0-3.5	---	---	None	---	None
		April	1.5-3.0	3.0-3.5	---	---	None	---	None
		May	1.5-3.0	3.0-3.5	---	---	None	---	None
		November	1.5-3.0	3.0-3.5	---	---	None	---	None
		December	1.5-3.0	3.0-3.5	---	---	None	---	None
449B: Hogansburg-----	B	January	1.5-3.0	3.0-3.5	---	---	None	---	None
		February	1.5-3.0	3.0-3.5	---	---	None	---	None
		March	1.5-3.0	3.0-3.5	---	---	None	---	None
		April	1.5-3.0	3.0-3.5	---	---	None	---	None
		May	1.5-3.0	3.0-3.5	---	---	None	---	None
		November	1.5-3.0	3.0-3.5	---	---	None	---	None
		December	1.5-3.0	3.0-3.5	---	---	None	---	None
449C: Hogansburg-----	B	January	1.5-3.0	3.0-3.5	---	---	None	---	None
		February	1.5-3.0	3.0-3.5	---	---	None	---	None
		March	1.5-3.0	3.0-3.5	---	---	None	---	None
		April	1.5-3.0	3.0-3.5	---	---	None	---	None
		May	1.5-3.0	3.0-3.5	---	---	None	---	None
		November	1.5-3.0	3.0-3.5	---	---	None	---	None
		December	1.5-3.0	3.0-3.5	---	---	None	---	None
450B: Pyrities-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
450C: Pyrities-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
450D: Pyrities-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
451B: Pyrities-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
451C: Pyrities-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
451D: Pyrities-----	B	Jan-Dec	> 6.0	> 6.0	---	---	None	---	None
457: Mudgepond-----	D	Jan-Dec	0.0-1.0	5.4-5.4	---	---	None	---	None
458: Mudgepond-----	D	Jan-Dec	0.0-1.0	5.4-5.4	---	---	None	---	None
Alden-----	D	January	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		February	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		March	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		April	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		May	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		June	0.0-1.0	>6.0	---	---	None	---	None
		October	0.0-1.0	>6.0	---	---	None	---	None
		November	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
		December	0.0-1.0	>6.0	0.0-0.5	Long	Occasional	---	None
501: Ondawa-----	B	January	5.0-6.0	>6.0	---	---	None	---	None
		February	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		March	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		April	5.0-6.0	>6.0	---	---	None	Brief	Occasional
		November	5.0-6.0	>6.0	---	---	None	---	None
		December	5.0-6.0	>6.0	---	---	None	---	None

Table 26.—Water Features—Continued

Map symbol and soil name	Hydrologic group	Month	Water table		Ponding			Flooding	
			Upper limit	Lower limit	Surface water depth	Duration	Frequency	Duration	Frequency
			Ft	Ft	Ft				
503: Rumney-----	D	January	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		February	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		March	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		April	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		May	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		June	0.0-1.5	>6.0	---	---	None	---	None
		September	0.0-1.5	>6.0	---	---	None	---	None
		October	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		November	0.0-1.5	>6.0	---	---	None	Brief	Frequent
		December	0.0-1.5	>6.0	---	---	None	Brief	Frequent
508: Medomak-----	D	January	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		February	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		March	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		April	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent
		May	0.0-0.5	>6.0	---	---	None	Brief	Frequent
		June	0.0-0.5	>6.0	---	---	None	---	None
		July	0.0-0.5	>6.0	---	---	None	---	None
		August	0.0-0.5	>6.0	---	---	None	---	None
		September	0.0-0.5	>6.0	---	---	None	---	None
		October	0.0-0.5	>6.0	---	---	None	Brief	Frequent
		November	0.0-0.5	>6.0	---	---	None	Brief	Frequent
		December	0.0-0.5	>6.0	0.0-1.5	Long	Frequent	Brief	Frequent

Table 27.—Soil Features

(See text for definitions of terms used in this table. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
2: Ridgebury-----	Dense material	20-30	---	very firm	0	0	High	High	Moderate
3: Ridgebury-----	Dense material	20-30	---	very firm	0	0	High	High	Moderate
Leicester-----	---	---	---	---	0	0	High	High	High
Whitman-----	Dense material	12-20	---	very firm	0	0	High	High	Moderate
4: Leicester-----	---	---	---	---	0	0	High	High	High
5: Wilbraham-----	Dense material	20-36	---	very firm	0	0	High	Moderate	Moderate
6: Wilbraham-----	Dense material	20-36	---	very firm	0	0	High	Moderate	Moderate
Menlo-----	Dense material	20-36	---	very firm	0	0	High	Moderate	Moderate
7: Mudgepond-----	---	---	---	---	0	0	High	High	Low
8: Mudgepond-----	---	---	---	---	0	0	High	High	Low
Alden-----	---	---	---	---	0	0	High	Moderate	Low
9: Scitico-----	---	---	---	---	0	0	High	High	Moderate
Shaker-----	---	---	---	---	0	0	High	High	Moderate
Maybid-----	---	---	---	---	0	0	High	Moderate	Moderate
10: Raynham-----	---	---	---	---	0	0	High	High	Moderate
12: Raypol-----	---	---	---	---	0	0	High	High	High
13: Walpole-----	---	---	---	---	0	0	Moderate	High	Moderate

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top	Thickness	Hardness	Initial		Total	Uncoated steel	Concrete
		In	In		In	In			
14: Fredon-----	---	---	---	---	0	0	High	High	Low
15: Scarboro-----	---	---	---	---	0	0	Moderate	High	Moderate
16: Halsey-----	---	---	---	---	0	0	High	High	Low
17: Timakwa-----	---	---	---	---	9-18	9-28	High	High	High
Natchaug-----	---	---	---	---	6-12	6-18	High	High	High
18: Catden-----	---	---	---	---	15-30	15-46	High	High	High
Freetown-----	---	---	---	---	23-46	23-68	High	High	High
20A: Ellington-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
21A: Ninigret-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Tisbury-----	---	---	---	---	0	0	High	Low	Moderate
22A: Hero-----	---	---	---	---	0	0	Moderate	Low	Low
22B: Hero-----	---	---	---	---	0	0	Moderate	Low	Low
23A: Sudbury-----	---	---	---	---	0	0	Low	Moderate	Moderate
24A: Deerfield-----	---	---	---	---	0	0	Low	Low	Moderate
25A: Brancroft-----	---	---	---	---	0	0	High	High	Moderate
25B: Brancroft-----	---	---	---	---	0	0	High	High	Moderate
25C: Brancroft-----	---	---	---	---	0	0	High	High	Moderate

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
26A: Berlin-----	---	---	---	---	0	0	High	High	Moderate
26B: Berlin-----	---	---	---	---	0	0	High	High	Moderate
27A: Belgrade-----	---	---	---	---	0	0	High	Moderate	Moderate
28A: Elmridge-----	---	---	---	---	0	0	Moderate	High	Moderate
28B: Elmridge-----	---	---	---	---	0	0	Moderate	High	Moderate
29A: Agawam-----	---	---	---	---	0	0	Moderate	Low	Moderate
29B: Agawam-----	---	---	---	---	0	0	Moderate	Low	Moderate
29C: Agawam-----	---	---	---	---	0	0	Moderate	Low	Moderate
30A: Branford-----	---	---	---	---	0	0	Moderate	Low	Moderate
30B: Branford-----	---	---	---	---	0	0	Moderate	Low	Moderate
30C: Branford-----	---	---	---	---	0	0	Moderate	Low	Moderate
31A: Copake-----	---	---	---	---	0	0	Moderate	Low	Low
31B: Copake-----	---	---	---	---	0	0	Moderate	Low	Low
31C: Copake-----	---	---	---	---	0	0	Moderate	Low	Low
32A: Haven-----	---	---	---	---	0	0	Moderate	Low	Moderate
Enfield-----	---	---	---	---	0	0	High	Low	Moderate

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
32B: Haven-----	---	---	---	---	0	0	Moderate	Low	Moderate
Enfield-----	---	---	---	---	0	0	High	Low	Moderate
32C: Haven-----	---	---	---	---	0	0	Moderate	Low	Moderate
Enfield-----	---	---	---	---	0	0	High	Low	Moderate
33A: Hartford-----	---	---	---	---	0	0	Low	Low	High
33B: Hartford-----	---	---	---	---	0	0	Low	Low	High
34A: Merrimac-----	---	---	---	---	0	0	Low	Low	Moderate
34B: Merrimac-----	---	---	---	---	0	0	Low	Low	Moderate
34C: Merrimac-----	---	---	---	---	0	0	Low	Low	Moderate
35A: Penwood-----	---	---	---	---	0	0	Low	Low	High
35B: Penwood-----	---	---	---	---	0	0	Low	Low	High
36A: Windsor-----	---	---	---	---	0	0	Low	Low	High
36B: Windsor-----	---	---	---	---	0	0	Low	Low	High
36C: Windsor-----	---	---	---	---	0	0	Low	Low	High
37A: Manchester-----	---	---	---	---	0	0	Low	Low	High
37C: Manchester-----	---	---	---	---	0	0	Low	Low	High
37E: Manchester-----	---	---	---	---	0	0	Low	Low	High

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
38A: Hinckley-----	---	---	---	---	0	0	Low	Low	High
38C: Hinckley-----	---	---	---	---	0	0	Low	Low	High
38E: Hinckley-----	---	---	---	---	0	0	Low	Low	High
39A: Groton-----	---	---	---	---	0	0	Low	Low	Low
39C: Groton-----	---	---	---	---	0	0	Low	Low	Low
39E: Groton-----	---	---	---	---	0	0	Low	Low	Low
40A: Ludlow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
40B: Ludlow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
41B: Ludlow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
42C: Ludlow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
43A: Rainbow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
43B: Rainbow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
44B: Rainbow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
45A: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
45B: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
45C: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
46B: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
46C: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
47C: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
48B: Georgia-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Amenia-----	---	---	---	---	0	0	Moderate	Moderate	Low
48C: Georgia-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Amenia-----	---	---	---	---	0	0	Moderate	Moderate	Low
49B: Georgia-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Amenia-----	---	---	---	---	0	0	Moderate	Moderate	Low
49C: Georgia-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Amenia-----	---	---	---	---	0	0	Moderate	Moderate	Low
50A: Sutton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
50B: Sutton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
51B: Sutton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
52C: Sutton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
53A: Wapping-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
53B: Wapping-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
54B: Wapping-----	---	---	---	---	0	0	Moderate	Moderate	Moderate

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top	Thickness	Hardness	Initial		Total	Uncoated steel	Concrete
55A: Watchaug-----	---	In ---	In ---	---	0	0	Moderate	Moderate	Moderate
55B: Watchaug-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
56B: Watchaug-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
57B: Gloucester-----	---	---	---	---	0	0	Low	Low	High
57C: Gloucester-----	---	---	---	---	0	0	Low	Low	High
57D: Gloucester-----	---	---	---	---	0	0	Low	Low	High
58B: Gloucester-----	---	---	---	---	0	0	Low	Low	High
58C: Gloucester-----	---	---	---	---	0	0	Low	Low	High
59C: Gloucester-----	---	---	---	---	0	0	Low	Low	High
59D: Gloucester-----	---	---	---	---	0	0	Low	Low	High
60B: Canton-----	---	---	---	---	0	0	Moderate	Moderate	High
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
60C: Canton-----	---	---	---	---	0	0	Moderate	Moderate	High
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
60D: Canton-----	---	---	---	---	0	0	Moderate	Moderate	High
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
61B: Canton-----	---	---	---	---	0	0	Moderate	Moderate	High
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
61C: Canton-----	---	---	---	---	0	0	Moderate	Moderate	High
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
62C: Canton-----	---	---	---	---	0	0	Moderate	Moderate	High
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
62D: Canton-----	---	---	---	---	0	0	Moderate	Moderate	High
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
63B: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
63C: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
63D: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
64B: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
64C: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
65C: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
65D: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
66B: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High
66C: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High
67B: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High
67C: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
68C: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High
68D: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High
69B: Yalesville-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
69C: Yalesville-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
70C: Branford-----	---	---	---	---	0	0	Moderate	Low	Moderate
Holyoke-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
71C: Brookfield-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Brimfield-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
71E: Brookfield-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Brimfield-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
73C: Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
73E: Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
74C: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High
Hollis-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top	Thickness	Hardness	Initial		Total	Uncoated steel	Concrete
		In	In		In	In			
75C:									
Hollis-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
75E:									
Hollis-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
76E:									
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
Hollis-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
76F:									
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
Hollis-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
77C:									
Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Holyoke-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
77D:									
Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Holyoke-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
78C:									
Holyoke-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
78E:									
Holyoke-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
79E:									
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
Holyoke-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Moderate

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
80B: Bernardston-----	Dense material	20-30	---	very firm	0	0	Moderate	Moderate	High
80C: Bernardston-----	Dense material	20-30	---	very firm	0	0	Moderate	Moderate	High
81C: Bernardston-----	Dense material	20-30	---	very firm	0	0	Moderate	Moderate	High
81D: Bernardston-----	Dense material	20-30	---	very firm	0	0	Moderate	Moderate	High
82B: Broadbrook-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
82C: Broadbrook-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
82D: Broadbrook-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
83B: Broadbrook-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
83C: Broadbrook-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
84B: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Montauk-----	Dense material	20-38	---	very firm	0	0	Moderate	Moderate	Moderate
84C: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Montauk-----	Dense material	20-38	---	very firm	0	0	Moderate	Moderate	Moderate
84D: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Montauk-----	Dense material	20-38	---	very firm	0	0	Moderate	Moderate	Moderate
85B: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Montauk-----	Dense material	20-38	---	very firm	0	0	Moderate	Moderate	Moderate

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top	Thickness	Hardness	Initial		Total	Uncoated steel	Concrete
		In	In		In	In			
85C: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Montauk-----	Dense material	20-38	---	very firm	0	0	Moderate	Moderate	Moderate
86C: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Montauk-----	Dense material	20-38	---	very firm	0	0	Moderate	Moderate	Moderate
86D: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Montauk-----	Dense material	20-38	---	very firm	0	0	Moderate	Moderate	Moderate
87B: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
87C: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
87D: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
88B: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
88C: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
89C: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
89D: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
90B: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
90C: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
90D: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
91B: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
91C: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
91D: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
92B: Nellis-----	---	---	---	---	0	0	Moderate	Moderate	Low
92C: Nellis-----	---	---	---	---	0	0	Moderate	Moderate	Low
92D: Nellis-----	---	---	---	---	0	0	Moderate	Moderate	Low
93C: Nellis-----	---	---	---	---	0	0	Moderate	Moderate	Low
94C: Farmington-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Low
Nellis-----	---	---	---	---	0	0	Moderate	Moderate	Low
94E: Farmington-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Low
Nellis-----	---	---	---	---	0	0	Moderate	Moderate	Low
95C: Farmington-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Low
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
95E: Farmington-----	Bedrock (lithic)	10-20	---	extremely firm	0	0	High	Low	Low
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
96: Ipswich-----	Salic	20-40	---	Noncemented	20-40	20-60	High	High	High
	Sulfuric	20-40	---	Noncemented					
97: Pawcatuck-----	Salic	0-60	---	Noncemented	12-23	12-35	High	High	High
	Sulfuric	0-60	---	Noncemented					

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top	Thickness	Hardness	Initial		Total	Uncoated steel	Concrete
		In	In		In	In			
98: Westbrook-----	Salic	0-51	---	Noncemented	12-24	12-36	High	High	High
	Sulfuric	0-51	---	Noncemented					
99: Westbrook, low salt----	Sulfuric	0-51	---	Noncemented	12-24	12-36	High	High	High
100: Suncook-----	---	---	---	---	0	0	Low	Low	High
101: Occum-----	---	---	---	---	0	0	Moderate	Low	Moderate
102: Pootatuck-----	---	---	---	---	0	0	Moderate	Low	Moderate
103: Rippowam-----	---	---	---	---	0	0	High	High	Moderate
104: Bash-----	---	---	---	---	0	0	Moderate	High	High
105: Hadley-----	---	---	---	---	0	0	High	Moderate	Moderate
106: Winooski-----	---	---	---	---	0	0	High	Moderate	Low
107: Limerick-----	---	---	---	---	0	0	High	High	Low
	Lim-----	---	---	---	0	0	High	High	Low
108: Saco-----	---	---	---	---	0	0	High	Moderate	Low
109: Fluvaquents, Frequently Flooded-----	---	---	---	---	0	0	High	High	Low
	Udifluents, Frequently Flooded-----	---	---	---	0	0	Moderate	Moderate	Moderate
221A: Ninigret-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
	Urban Land-----	---	---	---	---	---	None	---	---

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
224A: Deerfield-----	---	---	---	---	0	0	Low	Low	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
225B: Brancroft-----	---	---	---	---	0	0	High	High	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
226B: Berlin-----	---	---	---	---	0	0	High	High	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
228B: Elmridge-----	---	---	---	---	0	0	Moderate	High	Moderate
Urban Land-----	---	---	---	---	---	---	---	---	---
229B: Agawam-----	---	---	---	---	0	0	Moderate	Low	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
229C: Agawam-----	---	---	---	---	0	0	Moderate	Low	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
230B: Branford-----	---	---	---	---	0	0	Moderate	Low	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
230C: Branford-----	---	---	---	---	0	0	Moderate	Low	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
232B: Haven-----	---	---	---	---	0	0	Moderate	Low	Moderate
Urban Land-----	---	---	---	---	---	---	---	---	---
234B: Merrimac-----	---	---	---	---	0	0	Low	Low	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top	Thickness	Hardness	Initial		Total	Uncoated steel	Concrete
		In	In		In	In			
235B: Penwood-----	---	---	---	---	0	0	Low	Low	High
Urban Land-----	---	---	---	---	---	---	None	---	---
236B: Windsor-----	---	---	---	---	0	0	Low	Low	High
Urban Land-----	---	---	---	---	---	---	None	---	---
237A: Manchester-----	---	---	---	---	0	0	Low	Low	High
Urban Land-----	---	---	---	---	---	---	---	---	---
237C: Manchester-----	---	---	---	---	0	0	Low	Low	High
Urban Land-----	---	---	---	---	---	---	None	---	---
238A: Hinckley-----	---	---	---	---	0	0	Low	Low	High
Urban Land-----	---	---	---	---	---	---	None	---	---
238C: Hinckley-----	---	---	---	---	0	0	Low	Low	High
Urban Land-----	---	---	---	---	---	---	None	---	---
240B: Ludlow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
243B: Rainbow-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
245B: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
245C: Woodbridge-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
248B: Georgia-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
250B: Sutton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
253B: Wapping-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
255B: Watchaug-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
260B: Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
260C: Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
260D: Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
263B: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
263C: Cheshire-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
266B: Narragansett-----	---	---	---	---	0	0	Moderate	Low	High
Urban Land-----	---	---	---	---	---	---	None	---	---

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top	Thickness	Hardness	Initial		Total	Uncoated steel	Concrete
		In	In			In	In		
269B: Yalesville-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
269C: Yalesville-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
273C: Urban Land-----	---	---	---	---	---	---	None	---	---
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
273E: Urban Land-----	---	---	---	---	---	---	None	---	---
Charlton-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
275C: Urban Land-----	---	---	---	---	---	---	None	---	---
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
275E: Urban Land-----	---	---	---	---	---	---	None	---	---
Chatfield-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	Moderate
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
282B: Broadbrook-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
284B: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
284C: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
284D: Paxton-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
287B: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
287C: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
287D: Wethersfield-----	Dense material	20-40	---	very firm	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
290B: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
Urban Land-----	---	---	---	---	---	---	None	---	---
290C: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
Urban Land-----	---	---	---	---	---	---	None	---	---
290D: Stockbridge-----	---	---	---	---	0	0	Moderate	Moderate	Low
Urban Land-----	---	---	---	---	---	---	None	---	---
301: Beaches-----	---	---	---	---	0	0	Low	High	High
Udipsammments-----	---	---	---	---	0	0	Low	High	High
302: Dumps-----	---	---	---	---	0	0	None	---	---
303: Pits, Quarries-----	Bedrock (lithic)	0-4	---	extremely firm	0	0	None	---	---

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
304: Udorthents-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
305: Udorthents-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Pits-----	---	---	---	---	0	0	Low	---	---
306: Udorthents-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
Urban Land-----	---	---	---	---	---	---	None	---	---
307: Urban Land-----	---	---	---	---	---	---	None	---	---
308: Udorthents-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
309: Udorthents-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
310: Udorthents, Periodically Flooded--	---	---	---	---	0	0	Moderate	Moderate	Moderate
401C: Macomber-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	High
Taconic-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	Moderate	Low	High
402D: Macomber-----	Bedrock (lithic)	20-40	---	extremely firm	0	0	Moderate	Moderate	High
Taconic-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	Moderate	Low	High
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
403C: Taconic-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	Moderate	Low	High
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
403E: Taconic-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	Moderate	Low	High
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
403F: Taconic-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	Moderate	Low	High
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
405C: Dummerston-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
405E: Dummerston-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
407C: Lanesboro-----	Dense material	20-35	---	very firm	---	0	Moderate	Moderate	Moderate
407E: Lanesboro-----	Dense material	20-35	---	very firm	---	0	Moderate	Moderate	Moderate
408C: Fullam-----	Dense material	20-30	---	very firm	0	0	Moderate	Moderate	Moderate
409B: Brayton-----	Dense material	10-20	---	very firm	0	0	High	High	Moderate
412B: Bice-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
412C: Bice-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
412D: Bice-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
413C: Bice-----	---	---	---	---	---	0	Moderate	Moderate	Moderate
Millsite-----	Bedrock (lithic)	20-40	---	extremely firm	---	0	Moderate	Moderate	Moderate
413E: Bice-----	---	---	---	---	---	0	Moderate	Moderate	Moderate
Millsite-----	Bedrock (lithic)	20-40	---	extremely firm	---	0	Moderate	Moderate	Moderate
414: Fredon, cold-----	---	---	---	---	0	0	High	High	Low

Table 27.--Soil Features--Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
415C: Millsite-----	Bedrock (lithic)	20-40	---	extremely firm	---	0	Moderate	Moderate	Moderate
Westminster-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	High	Low	High
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
415E: Millsite-----	Bedrock (lithic)	20-40	---	extremely firm	---	0	Moderate	Moderate	Moderate
Westminster-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	High	Low	High
Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
416E: Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
Westminster-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	High	Low	High
416F: Rock Outcrop-----	Bedrock (lithic)	0-4	---	extremely firm	---	---	None	---	---
Westminster-----	Bedrock (lithic)	10-20	---	extremely firm	---	0	High	Low	High
417B: Bice-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
417C: Bice-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
417D: Bice-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
418C: Schroon-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
420A: Schroon-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
420B: Schroon-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
421A: Ninigret, cold-----	---	---	---	---	0	0	Moderate	Moderate	Moderate
423A: Sudbury, cold-----	---	---	---	---	0	0	Low	Moderate	Moderate

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer			Subsidence		Potential for frost action	Risk of corrosion		
	Kind	Depth to top In	Thickness In	Hardness	Initial In		Total In	Uncoated steel	Concrete
424B: Shelburne-----	Dense material	20-30	---	very firm	---	0	Moderate	Moderate	Moderate
424C: Shelburne-----	Dense material	20-30	---	very firm	---	0	Moderate	Moderate	Moderate
424D: Shelburne-----	Dense material	20-30	---	very firm	---	0	Moderate	Moderate	Moderate
425B: Shelburne-----	Dense material	20-30	---	very firm	---	0	Moderate	Moderate	Moderate
425C: Shelburne-----	Dense material	20-30	---	very firm	---	0	Moderate	Moderate	Moderate
426D: Shelburne-----	Dense material	20-30	---	very firm	---	0	Moderate	Moderate	Moderate
427B: Ashfield-----	Dense material	20-33	---	extremely firm	---	0	Moderate	Moderate	High
427C: Ashfield-----	Dense material	20-33	---	very firm	---	0	Moderate	Moderate	High
428A: Ashfield-----	Dense material	20-33	---	very firm	---	0	Moderate	Moderate	High
428B: Ashfield-----	Dense material	20-33	---	very firm	---	0	Moderate	Moderate	High
428C: Ashfield-----	Dense material	20-33	---	very firm	---	0	Moderate	Moderate	High
429A: Agawam, cold-----	---	---	---	---	0	0	Moderate	Low	Moderate
429B: Agawam, cold-----	---	---	---	---	0	0	Moderate	Low	Moderate
429C: Agawam, cold-----	---	---	---	---	0	0	Moderate	Low	Moderate
433: Moosilauke-----	---	---	---	---	0	0	Moderate	High	Moderate
434A: Merrimac, cold-----	---	---	---	---	0	0	Low	Low	Moderate

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
434B: Merrimac, cold-----	---	---	---	---	0	0	Low	Low	Moderate
434C: Merrimac, cold-----	---	---	---	---	0	0	Low	Low	Moderate
435: Scarboro-----	---	---	---	---	0	0	Moderate	High	Moderate
436: Halsey-----	---	---	---	---	0	0	High	High	Low
437: Wonsqueak-----	---	---	---	---	6-11	6-17	High	High	High
438: Bucksport-----	---	---	---	---	15-30	15-44	High	High	High
440A: Boscawen-----	---	---	---	---	0	0	Low	Low	High
440C: Boscawen-----	---	---	---	---	0	0	Low	Low	High
440E: Boscawen-----	---	---	---	---	0	0	Low	Low	High
442: Brayton-----	Dense material	20-27	---	very firm	0	0	High	High	Moderate
443: Brayton-----	Dense material	20-27	---	very firm	0	0	High	High	Moderate
Loonmeadow-----	---	---	---	---	0	0	High	High	Low
448B: Hogansburg-----	Dense material	20-43	---	very firm	0	0	Moderate	Moderate	Moderate
449B: Hogansburg-----	Dense material	20-43	---	very firm	0	0	Moderate	Moderate	Moderate
449C: Hogansburg-----	Dense material	20-43	---	very firm	0	0	Moderate	Moderate	Moderate
450B: Pyrities-----	---	---	---	---	0	0	Moderate	Moderate	Low

Table 27.—Soil Features—Continued

Map symbol and soil name	Restrictive layer				Subsidence		Potential for frost action	Risk of corrosion	
	Kind	Depth to top	Thickness	Hardness	Initial	Total		Uncoated steel	Concrete
		In	In		In	In			
450C: Pyrities-----	---	---	---	---	0	0	Moderate	Moderate	Low
450D: Pyrities-----	---	---	---	---	0	0	Moderate	Moderate	Low
451B: Pyrities-----	---	---	---	---	0	0	Moderate	Moderate	Low
451C: Pyrities-----	---	---	---	---	0	0	Moderate	Moderate	Low
451D: Pyrities-----	---	---	---	---	0	0	Moderate	Moderate	Low
457: Mudgepond-----	---	---	---	---	0	0	High	High	Low
458: Mudgepond-----	---	---	---	---	0	0	High	High	Low
Alden-----	---	---	---	---	0	0	High	Moderate	Low
501: Ondawa-----	---	---	---	---	0	0	Moderate	Low	Moderate
503: Rumney-----	---	---	---	---	0	0	High	High	Moderate
508: Medomak-----	---	---	---	---	0	0	High	Moderate	Low

Table 28.—Storm Water Runoff Systems

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
2: Ridgebury-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Restricted permeability Dense layer	1.00 1.00 0.50 0.34	Unlimited		Unlimited	
3: Ridgebury-----	40	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Restricted permeability Dense layer	1.00 1.00 0.50 0.34	Unlimited		Unlimited	
Leicester-----	35	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage Somewhat Limited Water Quantity	1.00 0.01	Unlimited Somewhat Limited Water Quantity	0.01
Whitman-----	15	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer	1.00 1.00	Unlimited		Unlimited	
4: Leicester-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage Somewhat Limited Water Quantity	1.00 0.01	Unlimited Somewhat Limited Water Quantity	0.01

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
5: Wilbraham-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Restricted permeability	1.00 1.00 0.97	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01
6: Wilbraham-----	60	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Restricted permeability	1.00 1.00 0.97	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01
Menlo-----	25	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Restricted permeability	1.00 1.00 0.90	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01
7: Mudgepond-----	85	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability Dense layer	1.00 0.86 0.03	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	
8: Mudgepond-----	45	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability Dense layer	1.00 0.86 0.03	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8: Alden-----	35	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01
9: Scitico-----	40	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability Dense layer	1.00 1.00 0.17	Unlimited		Unlimited	
Shaker-----	30	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability Dense layer	1.00 0.82 0.17	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01
Maybid-----	15	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability Dense layer	1.00 1.00 0.17	Unlimited		Unlimited	
10: Raynham-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Somewhat limited Somewhat Limited Water Quantity	0.34	Somewhat limited Somewhat Limited Water Quantity	0.34
12: Raypol-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
13: Walpole-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
14: Fredon-----	85	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.07
15: Scarboro-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
16: Halsey-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
17: Timakwa-----	45	Very limited Flooding Depth limited	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
Natchaug-----	40	Very limited Flooding Depth limited	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 0.40	Unlimited		Unlimited	
18: Catden-----	40	Very limited Flooding Depth limited	1.00 1.00	Very limited Depth to saturated zone Restricted permeability Flooding	1.00 0.50 0.40	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
18: Freetown-----	40	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	
		Depth limited	1.00	Restricted permeability Flooding	0.50 0.40				
20A: Ellington-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.02	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.02	Very limited Limited Water Quantity Subject To Seepage Slope	0.99 0.28 0.02
21A: Ninigret-----	60	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.02	Somewhat limited Ksat - Subject to seepage Too droughty Somewhat Limited Water Quantity Slope	1.00 0.75 0.12 0.02	Unlimited Subject To Seepage Somewhat Limited Water Quantity Slope	0.28 0.12 0.02
Tisbury-----	25	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage Too droughty Somewhat Limited Water Quantity Slope	1.00 0.75 0.12	Unlimited Subject To Seepage Somewhat Limited Water Quantity	0.28 0.12
22A: Hero-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Limited Water Quantity Subject To Seepage	0.99 0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
22B: Hero-----	85	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.28
23A: Sudbury-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.02	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.02	Very limited Limited Water Quantity Subject To Seepage Slope	0.99 0.28 0.02
24A: Deerfield-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited High infiltration Limited Water Quantity Subject To Seepage	1.00 0.99 0.28
25A: Brancroft-----	80	Unlimited		Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Somewhat limited Somewhat droughty Somewhat Limited Water Quantity	0.25 0.08	Unlimited Somewhat Limited Water Quantity	0.08

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
25B: Brancroft-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Restricted permeability	1.00	Somewhat droughty	0.25	Somewhat Limited Water Quantity	0.08
				Slope	0.74	Somewhat Limited Water Quantity	0.08		
25C: Brancroft-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	1.00	Somewhat droughty	0.25	Somewhat Limited Water Quantity	0.08
				Slope	1.00	Somewhat Limited Water Quantity	0.08		
26A: Berlin-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Somewhat droughty	0.25	Unlimited Somewhat Limited Water Quantity	0.08
				Restricted permeability	1.00	Somewhat Limited Water Quantity	0.08		
26B: Berlin-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Restricted permeability	1.00	Somewhat droughty	0.25	Somewhat Limited Water Quantity	0.08
				Slope	0.74	Somewhat Limited Water Quantity	0.08		
27A: Belgrade-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability	0.97	Limited Water Quantity	0.99	Slope	0.02
				Slope	0.02	Too droughty Slope	0.75 0.02		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
28A: Elmridge-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Somewhat droughty	0.25	Unlimited Somewhat Limited Water Quantity	0.12
				Restricted permeability	0.93	Somewhat Limited Water Quantity	0.12		
				Dense layer	0.17				
28B: Elmridge-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Restricted permeability	0.93	Somewhat droughty	0.25	Somewhat Limited Water Quantity	0.12
				Slope	0.74	Somewhat Limited Water Quantity	0.12		
				Dense layer	0.17				
29A: Agawam-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited Subject To Seepage	1.00
						Limited Water Quantity	0.99	Limited Water Quantity	0.99
						Too droughty	0.75		
29B: Agawam-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Subject To Seepage	1.00
						Limited Water Quantity	0.99	Limited Water Quantity	0.99
						Too droughty	0.75	Slope	0.74
						Slope	0.74		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
29C: Agawam-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00 1.00	Very limited Slope	1.00
						Limited Water Quantity Too droughty	0.99 0.75	Subject To Seepage Limited Water Quantity	1.00 0.99
30A: Branford-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Limited Water Quantity Subject To Seepage	0.99 0.28
30B: Branford-----	80	Unlimited		Somewhat limited Slope	0.26	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.26	Very limited Limited Water Quantity Subject To Seepage Slope	0.99 0.28 0.26
30C: Branford-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
31A: Copake-----	85	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
31B: Copake-----	85	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Slope	0.74
						Too droughty	0.75	Subject To Seepage	0.28
						Slope	0.74		
31C: Copake-----	85	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
						Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
32A: Haven-----	60	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited Subject To Seepage	1.00
						Limited Water Quantity	0.99	Limited Water Quantity	0.99
						Too droughty	0.75		
Enfield-----	25	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
32B: Haven-----	60	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Subject To Seepage Limited Water Quantity Slope	1.00 0.99 0.74
Enfield-----	25	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.28
32C: Haven-----	60	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Subject To Seepage Limited Water Quantity	1.00 1.00 0.99
Enfield-----	25	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
33A: Hartford-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
33B: Hartford-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Slope	0.74
						Too droughty	0.75	Subject To Seepage	0.28
						Slope	0.74		
34A: Merrimac-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
34B: Merrimac-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Slope	0.74
						Too droughty	0.75	Subject To Seepage	0.28
						Slope	0.74		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
34C: Merrimac-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope	1.00
						Limited Water Quantity	0.99	Limited Water Quantity Subject To Seepage	0.28
						Too droughty	0.75		
35A: Penwood-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
						Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
35B: Penwood-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
						Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Slope	0.74
						Slope	0.74	Subject To Seepage	0.28
36A: Windsor-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
						Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
36B: Windsor-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Too droughty	1.00 1.00	Very limited High infiltration	1.00
						Limited Water Quantity Slope	0.99 0.74	Slope Subject To Seepage	0.74 0.28
36C: Windsor-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Too droughty	1.00 1.00 1.00	Very limited High infiltration	1.00
						Limited Water Quantity	0.99	Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28
37A: Manchester-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage Too droughty	1.00 1.00	Very limited High infiltration	1.00
						Limited Water Quantity	0.99	Limited Water Quantity Subject To Seepage	0.99 0.28
37C: Manchester-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Too droughty Slope	1.00 1.00 1.00	Very limited High infiltration	1.00
						Limited Water Quantity	0.99	Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
37E: Manchester-----	80	Very limited Slope	0.88	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Too droughty	1.00 1.00 1.00	Very limited High infiltration Slope Limited Water Quantity Subject To Seepage	1.00 1.00 0.99 0.28
38A: Hinckley-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited High infiltration Subject To Seepage Limited Water Quantity	1.00 1.00 0.99
38C: Hinckley-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Too droughty Slope Limited Water Quantity	1.00 1.00 1.00 0.99	Very limited High infiltration Subject To Seepage Slope Limited Water Quantity	1.00 1.00 1.00 0.99
38E: Hinckley-----	80	Very limited Slope	0.88	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Too droughty Limited Water Quantity	1.00 1.00 1.00 0.99	Very limited High infiltration Slope Subject To Seepage Limited Water Quantity	1.00 1.00 1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
39A: Groton-----	85	Unlimited		Unlimited		Very limited Ksat - Subject to seepage Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited High infiltration Limited Water Quantity Subject To Seepage	1.00 0.99 0.28
39C: Groton-----	85	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Too droughty Slope Limited Water Quantity	1.00 1.00 1.00 0.99	Very limited High infiltration Slope Limited Water Quantity Subject To Seepage	1.00 1.00 0.99 0.28
39E: Groton-----	85	Very limited Slope	0.88	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Too droughty Limited Water Quantity	1.00 1.00 1.00 0.99	Very limited High infiltration Slope Limited Water Quantity Subject To Seepage	1.00 1.00 0.99 0.28
40A: Ludlow-----	80	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability	1.00 1.00 0.97	Very limited Limited Water Quantity Somewhat droughty	0.99 0.25	Very limited Limited Water Quantity	0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
40B: Ludlow-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Thin layer	1.00	Slope	0.74	Slope	0.74
				Restricted permeability	0.97	Somewhat droughty	0.25		
				Slope	0.74				
41B: Ludlow-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Thin layer	1.00	Slope	0.41	Slope	0.41
				Restricted permeability	0.97	Somewhat droughty	0.25		
				Slope	0.41				
42C: Ludlow-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Slope	1.00	Somewhat droughty	0.25		
				Restricted permeability	0.97				
43A: Rainbow-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Somewhat droughty	0.25	Unlimited Somewhat Limited Water Quantity	0.12
				Thin layer	1.00	Somewhat Limited Water Quantity	0.12		
				Restricted permeability	0.97				

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
43B: Rainbow-----	80	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Restricted permeability Slope	0.97 0.74	Somewhat droughty Somewhat Limited Water Quantity	0.25 0.12	Somewhat Limited Water Quantity	0.12
44B: Rainbow-----	80	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Restricted permeability Slope	0.97 0.74	Somewhat droughty Somewhat Limited Water Quantity	0.25 0.12	Somewhat Limited Water Quantity	0.12
45A: Woodbridge-----	80	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Somewhat droughty	0.25	Unlimited Somewhat Limited Water Quantity	0.12
				Restricted permeability	0.97	Somewhat Limited Water Quantity	0.12		
45B: Woodbridge-----	80	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Restricted permeability Slope	0.97 0.74	Somewhat droughty Somewhat Limited Water Quantity	0.25 0.12	Somewhat Limited Water Quantity	0.12

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
45C: Woodbridge-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope Somewhat droughty	1.00 0.25	Very limited Slope Somewhat Limited Water Quantity	1.00 0.12
				Thin layer Restricted permeability	1.00 0.97	Somewhat Limited Water Quantity	0.12		
46B: Woodbridge-----	80	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Slope Somewhat droughty	0.41 0.25	Somewhat limited Slope Somewhat Limited Water Quantity	0.41 0.12
				Restricted permeability Slope	0.97 0.41	Somewhat Limited Water Quantity	0.12		
46C: Woodbridge-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope Somewhat droughty	1.00 0.25	Very limited Slope Somewhat Limited Water Quantity	1.00 0.12
				Thin layer Restricted permeability	1.00 0.97	Somewhat Limited Water Quantity	0.12		
47C: Woodbridge-----	80	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope Somewhat droughty	1.00 0.25	Very limited Slope Somewhat Limited Water Quantity	1.00 0.12
				Slope Restricted permeability	1.00 0.97	Somewhat Limited Water Quantity	0.12		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
48B: Georgia-----	50	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Too droughty	0.75	Somewhat limited Slope	0.41
				Restricted permeability	1.00	Slope	0.41	Somewhat Limited Water Quantity	0.17
				Slope	0.41	Somewhat Limited Water Quantity	0.17		
Amenia-----	35	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Too droughty	0.75	Somewhat limited Slope	0.41
				Restricted permeability	1.00	Slope	0.41	Somewhat Limited Water Quantity	0.17
				Slope	0.41	Somewhat Limited Water Quantity	0.17		
48C: Georgia-----	50	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Too droughty	0.75	Somewhat Limited Water Quantity	0.17
				Restricted permeability	1.00	Somewhat Limited Water Quantity	0.17		
Amenia-----	35	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Too droughty	0.75	Somewhat Limited Water Quantity	0.17
				Restricted permeability	1.00	Somewhat Limited Water Quantity	0.17		
49B: Georgia-----	50	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Too droughty	0.75	Somewhat limited Slope	0.74
				Restricted permeability	1.00	Slope	0.74	Somewhat Limited Water Quantity	0.17
				Slope	0.74	Somewhat Limited Water Quantity	0.17		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
49B: Amenia-----	35	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Too droughty	0.75	Somewhat limited Slope	0.74
				Restricted permeability	1.00	Slope	0.74	Somewhat Limited Water Quantity	0.17
				Slope	0.74	Somewhat Limited Water Quantity	0.17		
49C: Georgia-----	50	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Too droughty	0.75	Somewhat Limited Water Quantity	0.17
				Restricted permeability	1.00	Somewhat Limited Water Quantity	0.17		
Amenia-----	35	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Too droughty	0.75	Somewhat Limited Water Quantity	0.17
				Restricted permeability	1.00	Somewhat Limited Water Quantity	0.17		
50A: Sutton-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability	0.34	Limited Water Quantity	0.99	Subject To Seepage	0.07
						Too droughty	0.75		
50B: Sutton-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Slope	0.74	Limited Water Quantity	0.99	Slope	0.74
				Restricted permeability	0.34	Too droughty	0.75	Subject To Seepage	0.07
						Slope	0.74		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
51B: Sutton-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.41	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.41	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.41 0.07
52C: Sutton-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Ksat - Subject to seepage Slope	1.00 1.00	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.07
53A: Wapping-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Limited Water Quantity Subject To Seepage	0.99 0.07
53B: Wapping-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.07

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
54B: Wapping-----	80	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.41	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.41	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.41 0.07
55A: Watchaug-----	80	Unlimited		Very limited Depth to saturated zone Restricted permeability	1.00 0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Limited Water Quantity	0.99
55B: Watchaug-----	80	Unlimited		Very limited Depth to saturated zone Slope Restricted permeability	1.00 0.74 0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
56B: Watchaug-----	80	Unlimited		Very limited Depth to saturated zone Restricted permeability Slope	1.00 0.74 0.41	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.41	Very limited Limited Water Quantity Slope	0.99 0.41

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
57B: Gloucester-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.28
57C: Gloucester-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28
57D: Gloucester-----	80	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28
58B: Gloucester-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Too droughty Limited Water Quantity Slope	1.00 1.00 0.99 0.74	Very limited High infiltration Limited Water Quantity Slope Subject To Seepage	1.00 0.99 0.74 0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
58C: Gloucester-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Too droughty	1.00 1.00 1.00	Very limited High infiltration Slope Limited Water Quantity Subject To Seepage	1.00 1.00 0.99 0.28
59C: Gloucester-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Too droughty Slope	1.00 1.00 1.00	Very limited High infiltration Slope Limited Water Quantity Subject To Seepage	1.00 1.00 0.99 0.28
59D: Gloucester-----	80	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Too droughty	1.00 1.00 1.00	Very limited High infiltration Slope Limited Water Quantity Subject To Seepage	1.00 1.00 0.99 0.28
60B: Canton-----	45	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
60B: Charlton-----	35	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability	0.50	Limited Water Quantity	0.99	Slope	0.74
						Too droughty Slope	0.75 0.74		
60C: Canton-----	45	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
						Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
Charlton-----	35	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
60D: Canton-----	45	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
						Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems			
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value		
60D: Charlton-----	35	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope	1.00		
				Restricted permeability	0.50	Limited Water Quantity Too droughty	0.99	Limited Water Quantity	0.99		
								Too droughty	0.75		
61B: Canton-----	45	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99		
						Limited Water Quantity Too droughty	0.99	Slope	0.74	Subject To Seepage	0.28
								Slope	0.74		
Charlton-----	35	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99		
						Restricted permeability	0.50	Limited Water Quantity Too droughty	0.99	Slope	0.74
								Slope	0.74		
61C: Canton-----	45	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope	1.00		
								Limited Water Quantity	0.99	Limited Water Quantity	0.99
								Limited Water Quantity Too droughty	0.99	Subject To Seepage	0.28
						Too droughty	0.75				

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
61C: Charlton-----	35	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
62C: Canton-----	45	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
						Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
Charlton-----	35	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
62D: Canton-----	45	Somewhat limited Slope	0.50	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
						Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
62D: Charlton-----	35	Somewhat limited Slope	0.50	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Too droughty	0.75
63B: Cheshire-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability	0.50	Limited Water Quantity	0.99	Slope	0.74
						Too droughty	0.75	Slope	0.74
63C: Cheshire-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Too droughty	0.75
63D: Cheshire-----	80	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Too droughty	0.75

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
64B: Cheshire-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability	0.50	Limited Water Quantity	0.99	Slope	0.74
						Too droughty	0.75		
						Slope	0.74		
64C: Cheshire-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
65C: Cheshire-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
65D: Cheshire-----	80	Somewhat limited Slope	0.50	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
66B: Narragansett-----	80	Unlimited		Somewhat limited Slope	0.41	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.41	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.41 0.07
66C: Narragansett-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.07
67B: Narragansett-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.07
67C: Narragansett-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.07

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
68C: Narragansett-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00 1.00	Very limited Slope	1.00
						Limited Water Quantity Too droughty	0.99 0.07	Limited Water Quantity Subject To Seepage	0.07
68D: Narragansett-----	80	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00 1.00	Very limited Slope	1.00
						Limited Water Quantity Too droughty	0.99 0.07	Limited Water Quantity Subject To Seepage	0.07
69B: Yalesville-----	75	Somewhat limited Bedrock	0.66	Very limited Thin layer Slope Restricted permeability	1.00 0.74 0.50	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.74 0.25	Very limited Limited Water Quantity Slope	0.99 0.74
69C: Yalesville-----	75	Somewhat limited Bedrock	0.66	Very limited Slope Thin layer Restricted permeability	1.00 1.00 0.50	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
70C: Branford-----	50	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00 1.00	Very limited Slope	1.00
						Limited Water Quantity Too droughty	0.99 0.99 0.75	Limited Water Quantity Subject To Seepage	0.99 0.28
Holyoke-----	30	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.83				
71C: Brookfield-----	45	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00 1.00	Very limited Slope	1.00
				Restricted permeability	0.42	Limited Water Quantity Too droughty	0.99 0.75	Limited Water Quantity	0.99
Brimfield-----	30	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.25				
Rock outcrop-----	15	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
71E: Brookfield-----	45	Very limited Slope	0.88	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.42	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
Brimfield-----	30	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Slope	0.88	Restricted permeability	0.25	Limited Water Quantity	0.99	Limited Water Quantity	0.99
Rock outcrop-----	15	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Slope	0.88			Limited Water Quantity	0.99	Limited Water Quantity	0.99
73C: Charlton-----	45	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
Chatfield-----	30	Very limited Bedrock	0.89	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.49	Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
73E: Charlton-----	45	Very limited Slope	0.88	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope	1.00
	Restricted permeability			0.50	Limited Water Quantity			0.99	
					Limited Water Quantity Too droughty			0.99 0.75	
Chatfield-----	30	Very limited Bedrock Slope	0.89 0.88	Very limited Slope	1.00	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope	1.00
	Thin layer			1.00	Limited Water Quantity			0.99	
	Restricted permeability			0.49	Too droughty			0.75	
74C: Narragansett-----	55	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope	1.00
					Limited Water Quantity Subject To Seepage			0.99 0.07	
Hollis-----	20	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer	1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope	1.00
	Slope			1.00	Limited Water Quantity			0.99	
	Restricted permeability			0.48					
75C: Hollis-----	35	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer	1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope	1.00
	Slope			1.00	Limited Water Quantity			0.99	
	Restricted permeability			0.48					

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
75C: Chatfield-----	30	Very limited Bedrock	0.89	Very limited Slope Thin layer	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.49	Too droughty	0.75		
Rock outcrop-----	15	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
75E: Hollis-----	35	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
		Slope	0.88	Restricted permeability	0.48				
Chatfield-----	30	Very limited Bedrock Slope	0.89 0.88	Very limited Slope Thin layer	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.49	Too droughty	0.75		
Rock outcrop-----	15	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
		Slope	0.88						
76E: Rock outcrop-----	55	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
		Slope	0.41						

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
76E: Hollis-----	25	Very limited Depth limited Bedrock Slope	1.00 1.00 0.41	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.48	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
76F: Rock outcrop-----	55	Very limited Slope Depth limited Bedrock	1.00 1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Hollis-----	25	Very limited Slope Depth limited Bedrock	1.00 1.00 1.00	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.48	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
77C: Cheshire-----	45	Unlimited		Very limited Slope Restricted permeability	1.00 0.50	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
Holyoke-----	35	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.83	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
77D: Cheshire-----	45	Somewhat limited Slope	0.50	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Limited Water Quantity Too droughty	0.99 0.75	Limited Water Quantity	0.99
Holyoke-----	35	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
		Slope	0.50	Restricted permeability	0.83				
78C: Holyoke-----	50	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.83				
Rock outcrop-----	25	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
78E: Holyoke-----	50	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
		Slope	0.88	Restricted permeability	0.83				

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
78E: Rock outcrop-----	25	Very limited Depth limited Bedrock Slope	 1.00 1.00 0.88	Very limited Thin layer Slope	 1.00 1.00	Very limited Slope Limited Water Quantity	 1.00 0.99	Very limited Slope Limited Water Quantity	 1.00 0.99
79E: Rock outcrop-----	55	Very limited Depth limited Bedrock Slope	 1.00 1.00 0.41	Very limited Thin layer Slope	 1.00 1.00	Very limited Slope Limited Water Quantity	 1.00 0.99	Very limited Slope Limited Water Quantity	 1.00 0.99
Holyoke-----	25	Very limited Depth limited Bedrock Slope	 1.00 1.00 0.41	Very limited Thin layer Slope Restricted permeability	 1.00 1.00 0.83	Very limited Slope Limited Water Quantity	 1.00 0.99	Very limited Slope Limited Water Quantity	 1.00 0.99
80E: Bernardston-----	80	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	 1.00 1.00 0.97 0.74	Very limited Limited Water Quantity Slope Somewhat droughty	 0.99 0.74 0.25	Very limited Limited Water Quantity Slope	 0.99 0.74
80C: Bernardston-----	80	Unlimited		Very limited Depth to saturated zone Thin layer Slope Restricted permeability	 1.00 1.00 1.00 0.97	Very limited Slope Limited Water Quantity Somewhat droughty	 1.00 0.99 0.25	Very limited Slope Limited Water Quantity	 1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
81C: Bernardston-----	80	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope Restricted permeability	1.00 0.97	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
81D: Bernardston-----	80	Somewhat limited Slope	0.12	Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope Restricted permeability	1.00 0.97	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
82B: Broadbrook-----	80	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.97 0.41	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.41 0.25	Very limited Limited Water Quantity Slope	0.99 0.41
82C: Broadbrook-----	80	Unlimited		Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.97	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
82D: Broadbrook-----	80	Somewhat limited Slope	0.12	Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer Restricted permeability	1.00 0.97	Limited Water Quantity	0.99	Limited Water Quantity	0.99
						Somewhat droughty	0.25		
83B: Broadbrook-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Thin layer Restricted permeability	1.00 0.97	Slope	0.74	Slope	0.74
				Slope	0.74	Somewhat droughty	0.25		
83C: Broadbrook-----	80	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Thin layer Restricted permeability	1.00 0.97	Somewhat droughty	0.25		
84B: Paxton-----	55	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Thin layer Restricted permeability	1.00 0.97	Slope	0.74	Slope	0.74
				Slope	0.74	Somewhat droughty	0.25		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
84B: Montauk-----	30	Unlimited		Very limited Depth to saturated zone Thin layer Dense layer Slope Restricted permeability	1.00 1.00 1.00 0.74 0.50	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.74 0.25	Very limited Limited Water Quantity Slope	0.99 0.74
84C: Paxton-----	55	Unlimited		Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.97	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
Montauk-----	30	Unlimited		Very limited Depth to saturated zone Thin layer Slope Dense layer Restricted permeability	1.00 1.00 1.00 1.00 0.50	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
84D: Paxton-----	55	Somewhat limited Slope	0.12	Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.97	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
84D: Montauk-----	30	Somewhat limited Slope	0.12	Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
85B: Paxton-----	55	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.97 0.74	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.74 0.25	Very limited Limited Water Quantity Slope	0.99 0.74
Montauk-----	30	Unlimited		Very limited Depth to saturated zone Thin layer Dense layer Slope Restricted permeability	1.00 1.00 1.00 0.74 0.50	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.74 0.25	Very limited Limited Water Quantity Slope	0.99 0.74
85C: Paxton-----	55	Unlimited		Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.97	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
85C: Montauk-----	30	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope Dense layer Restricted permeability	1.00 1.00 0.50	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
86C: Paxton-----	55	Unlimited		Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer Restricted permeability	1.00 0.97	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
Montauk-----	30	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope Dense layer Restricted permeability	1.00 1.00 0.50	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
86D: Paxton-----	55	Somewhat limited Slope	0.50	Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer Restricted permeability	1.00 0.97	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
86D: Montauk-----	30	Somewhat limited Slope	0.50	Very limited		Very limited		Very limited	
	Depth to saturated zone			1.00	Slope	1.00	Slope	1.00	
	Thin layer			1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99	
	Slope			1.00	Somewhat droughty	0.25			
	Dense layer			1.00					
	Restricted permeability	0.50							
87B: Wethersfield-----	80	Unlimited		Very limited		Very limited		Very limited	
	Depth to saturated zone			1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99	
	Thin layer			1.00	Slope	0.74	Slope	0.74	
	Restricted permeability			0.97	Somewhat droughty	0.25			
	Slope			0.74					
87C: Wethersfield-----	80	Unlimited		Very limited		Very limited		Very limited	
	Depth to saturated zone			1.00	Slope	1.00	Slope	1.00	
	Slope			1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99	
	Thin layer			1.00	Somewhat droughty	0.25			
	Restricted permeability			0.97					
87D: Wethersfield-----	80	Somewhat limited Slope	0.12	Very limited		Very limited		Very limited	
	Depth to saturated zone			1.00	Slope	1.00	Slope	1.00	
	Slope			1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99	
	Thin layer			1.00	Somewhat droughty	0.25			
	Restricted permeability			0.97					

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
88B: Wethersfield-----	80	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.89 0.74	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.74 0.25	Very limited Limited Water Quantity Slope	0.99 0.74
88C: Wethersfield-----	80	Unlimited		Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.89	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
89C: Wethersfield-----	80	Unlimited		Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.89	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
89D: Wethersfield-----	80	Somewhat limited Slope	0.50	Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.89	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
90B: Stockbridge-----	80	Unlimited		Very limited Restricted permeability Dense layer Slope	1.00 0.83 0.74	Very limited Limited Water Quantity Too droughty Slope	0.99 0.75 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
90C: Stockbridge-----	80	Unlimited		Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
90D: Stockbridge-----	80	Somewhat limited Slope	0.12	Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
91B: Stockbridge-----	80	Unlimited		Very limited Restricted permeability Dense layer Slope	1.00 0.83 0.74	Very limited Limited Water Quantity Too droughty Slope	0.99 0.75 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
91C: Stockbridge-----	80	Unlimited		Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
91D: Stockbridge-----	80	Somewhat limited Slope	0.50	Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
92B: Nellis-----	85	Unlimited		Very limited Dense layer	0.97	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability	0.86	Limited Water Quantity	0.99	Slope	0.74
				Slope	0.74	Too droughty	0.75		
						Slope	0.74		
92C: Nellis-----	85	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Dense layer	0.97	Slope	1.00	Limited Water Quantity	0.99
				Restricted permeability	0.86	Limited Water Quantity	0.99		
						Too droughty	0.75		
92D: Nellis-----	85	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Dense layer	0.97	Slope	1.00	Limited Water Quantity	0.99
				Restricted permeability	0.86	Limited Water Quantity	0.99		
						Too droughty	0.75		
93C: Nellis-----	85	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Dense layer	0.97	Slope	1.00	Limited Water Quantity	0.99
				Restricted permeability	0.86	Limited Water Quantity	0.99		
						Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value						
94C: Farmington-----	40	Very limited		Very limited		Very limited		Very limited	
		Depth limited	1.00	Thin layer	1.00	Slope	1.00	Slope	1.00
		Bedrock	1.00	Slope	1.00	Limited Water	0.99	Limited Water	0.99
				Restricted permeability	0.50	Quantity		Quantity	
Nellis-----	35	Unlimited		Very limited		Very limited		Very limited	
				Slope	1.00	Ksat - Subject to seepage	1.00	Slope	1.00
				Dense layer	0.97	Slope	1.00	Limited Water	0.99
				Restricted permeability	0.86	Limited Water	0.99	Quantity	
						Quantity		Quantity	
						Too droughty	0.75		
94E: Farmington-----	40	Very limited		Very limited		Very limited		Very limited	
		Depth limited	1.00	Thin layer	1.00	Slope	1.00	Slope	1.00
		Bedrock	1.00	Slope	1.00	Limited Water	0.99	Limited Water	0.99
		Slope	0.50	Restricted permeability	0.50	Quantity		Quantity	
Nellis-----	35	Somewhat limited		Very limited		Very limited		Very limited	
		Slope	0.50	Slope	1.00	Ksat - Subject to seepage	1.00	Slope	1.00
				Dense layer	0.97	Slope	1.00	Limited Water	0.99
				Restricted permeability	0.86	Limited Water	0.99	Quantity	
						Quantity		Quantity	
						Too droughty	0.75		
95C: Farmington-----	60	Very limited		Very limited		Very limited		Very limited	
		Depth limited	1.00	Thin layer	1.00	Slope	1.00	Slope	1.00
		Bedrock	1.00	Slope	1.00	Limited Water	0.99	Limited Water	0.99
				Restricted permeability	0.50	Quantity		Quantity	

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
95C: Rock outcrop-----	20	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
95E: Farmington-----	60	Very limited Depth limited Bedrock Slope	1.00 1.00 0.88	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.50	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Rock outcrop-----	20	Very limited Depth limited Bedrock Slope	1.00 1.00 0.88	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
96: Ipswich-----	85	Very limited Flooding Depth limited	1.00 1.00	Very limited Depth to saturated zone Flooding Thin layer	1.00 1.00 1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	
97: Pawcatuck-----	85	Very limited Flooding Depth limited	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 1.00	Somewhat limited Ksat - Subject to seepage	1.00	Somewhat limited Subject To Seepage	1.00
98: Westbrook-----	80	Very limited Flooding Depth limited	1.00 1.00	Very limited Depth to saturated zone Flooding	1.00 1.00	Unlimited		Unlimited	

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
99: Westbrook, low salt-	80	Very limited Flooding	1.00	Very limited Depth to saturated zone Flooding	1.00	Unlimited		Unlimited	
		Depth limited	1.00		1.00				
100: Suncook-----	80	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
				Depth to saturated zone	0.14	Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
101: Occum-----	80	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Depth to saturated zone	0.14	Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
102: Pootatuck-----	80	Very limited Flooding	1.00	Very limited Depth to saturated zone Flooding	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
					1.00	Limited Water Quantity	0.99	Subject To Seepage	0.28
						Too droughty	0.75		
103: Rippowam-----	80	Very limited Flooding	1.00	Very limited Depth to saturated zone Flooding	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
		Depth limited	1.00		1.00	Somewhat Limited Water Quantity	0.01	Somewhat Limited Water Quantity	0.01

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
104: Bash-----	80	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01
		Depth limited	1.00	Flooding Restricted permeability	1.00 0.97				
105: Hadley-----	80	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability Depth to saturated zone	0.63 0.05	Limited Water Quantity Too droughty	0.99 0.75		
106: Winooski-----	80	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Flooding Restricted permeability	1.00 0.82	Limited Water Quantity Too droughty	0.99 0.75		
107: Limerick-----	50	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Somewhat Limited Water Quantity	0.01
		Depth limited	1.00	Flooding Restricted permeability	1.00 0.97	Somewhat Limited Water Quantity	0.01		
Lim-----	30	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
		Depth limited	1.00	Flooding	1.00	Somewhat Limited Water Quantity	0.01	Somewhat Limited Water Quantity	0.01

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
108: Saco-----	80	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
		Depth limited	1.00	Flooding Restricted permeability	1.00 0.07				
109: Fluvaquents, Frequently Flooded-	50	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	
		Depth limited	1.00	Flooding	1.00				
Udifulvents, Frequently Flooded-	35	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty	1.00 0.98 0.75	Very limited Limited Water Quantity	0.98
221A: Ninigret-----	40	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.02	Somewhat limited Ksat - Subject to seepage Too droughty Somewhat Limited Water Quantity Slope	1.00 0.75 0.12 0.02	Unlimited Subject To Seepage Somewhat Limited Water Quantity Slope	0.28 0.12 0.02
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.02	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.02	Very limited Limited Water Quantity Slope	0.99 0.02

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
224A: Deerfield-----	40	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Too droughty	1.00	Very limited High infiltration	1.00
						Limited Water Quantity	0.99	Limited Water Quantity Subject To Seepage	0.28
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability	1.00	Very limited Too droughty	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
225B: Brancroft-----	40	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Somewhat droughty	0.25	Somewhat limited Slope	0.15
				Restricted permeability	1.00	Slope	0.15	Somewhat Limited Water Quantity	0.08
				Slope	0.15	Somewhat Limited Water Quantity	0.08		
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability	1.00	Very limited Too droughty	1.00	Very limited Limited Water Quantity	0.99
				Slope	0.15	Limited Water Quantity Slope	0.99 0.15	Slope	0.15
226B: Berlin-----	40	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Somewhat droughty	0.25	Somewhat limited Slope	0.15
				Restricted permeability	1.00	Slope	0.15	Somewhat Limited Water Quantity	0.08
				Slope	0.15	Somewhat Limited Water Quantity	0.08		

Table 28.--Storm Water Runoff Systems--Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
226B: Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
228B: Elmridge-----	40	Unlimited		Very limited Depth to saturated zone Restricted permeability Dense layer Slope	1.00 0.93 0.17 0.15	Somewhat limited Somewhat droughty Slope Somewhat Limited Water Quantity	0.25 0.15 0.12	Somewhat limited Slope Somewhat Limited Water Quantity	0.15 0.12
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
229B: Agawam-----	40	Unlimited		Somewhat limited Slope	0.15	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.15	Very limited Subject To Seepage Limited Water Quantity Slope	1.00 0.99 0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
229C: Agawam-----	40	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00 1.00	Very limited Slope	1.00
						Limited Water Quantity Too droughty	0.99 0.75	Subject To Seepage Limited Water Quantity	1.00 0.99
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
230B: Branford-----	40	Unlimited		Somewhat limited Slope	0.15	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.15	Very limited Limited Water Quantity Subject To Seepage Slope	0.99 0.28 0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
230C: Branford-----	40	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
230C: Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
232B: Haven-----	40	Unlimited		Somewhat limited Slope	0.15	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.15	Very limited Subject To Seepage Limited Water Quantity Slope	1.00 0.99 0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
234B: Merrimac-----	40	Unlimited		Somewhat limited Slope	0.15	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.15	Very limited Limited Water Quantity Subject To Seepage Slope	0.99 0.28 0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
235B: Penwood-----	40	Unlimited		Somewhat limited Slope	0.15	Very limited Ksat - Subject to seepage Too droughty	1.00	Very limited High infiltration	1.00
						Limited Water Quantity	0.99	Limited Water Quantity	0.99
						Subject To Seepage Slope	0.28	Subject To Seepage Slope	0.28
						Slope	0.15	Slope	0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00	Very limited Too droughty	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity Slope	0.99	Limited Water Quantity Slope	0.15
						Slope	0.15	Slope	0.15
236B: Windsor-----	40	Unlimited		Somewhat limited Slope	0.15	Very limited Ksat - Subject to seepage Too droughty	1.00	Very limited High infiltration	1.00
						Limited Water Quantity	0.99	Limited Water Quantity	0.99
						Subject To Seepage Slope	0.28	Subject To Seepage Slope	0.28
						Slope	0.15	Slope	0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00	Very limited Too droughty	1.00	Very limited Limited Water Quantity	0.99
						Limited Water Quantity Slope	0.99	Limited Water Quantity Slope	0.15
						Slope	0.15	Slope	0.15
237A: Manchester-----	40	Unlimited		Unlimited		Very limited Ksat - Subject to seepage Too droughty	1.00	Very limited High infiltration	1.00
						Limited Water Quantity	0.99	Limited Water Quantity	0.99
						Subject To Seepage	0.28	Subject To Seepage	0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
237A: Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability	1.00	Very limited Too droughty Limited Water Quantity	1.00 0.99	Very limited Limited Water Quantity	0.99
237C: Manchester-----	40	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Too droughty Slope Limited Water Quantity	1.00 1.00 1.00 0.99	Very limited High infiltration Slope Limited Water Quantity Subject To Seepage	1.00 1.00 0.99 0.28
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
238A: Hinckley-----	40	Unlimited		Unlimited		Very limited Ksat - Subject to seepage Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited High infiltration Subject To Seepage Limited Water Quantity	1.00 1.00 0.99
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability	1.00	Very limited Too droughty Limited Water Quantity	1.00 0.99	Very limited Limited Water Quantity	0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
238C: Hinckley-----	40	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Too droughty	1.00 1.00	Very limited High infiltration	1.00
						Slope Limited Water Quantity	1.00 0.99	Subject To Seepage Slope Limited Water Quantity	1.00 1.00 0.99
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
240B: Ludlow-----	40	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.97 0.15	Very limited Limited Water Quantity Somewhat droughty Slope	0.99 0.25 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
243B: Rainbow-----	40	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.97 0.15	Somewhat limited Somewhat droughty Slope Somewhat Limited Water Quantity	0.25 0.15 0.12	Somewhat limited Slope Somewhat Limited Water Quantity	0.15 0.12

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
243B: Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
245B: Woodbridge-----	40	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.97 0.15	Somewhat limited Somewhat droughty Slope Somewhat Limited Water Quantity	0.25 0.15 0.12	Somewhat limited Slope Somewhat Limited Water Quantity	0.15 0.12
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
245C: Woodbridge-----	40	Unlimited		Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.97	Very limited Slope Somewhat droughty Somewhat Limited Water Quantity	1.00 0.25 0.12	Very limited Slope Somewhat Limited Water Quantity	1.00 0.12
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
248B: Georgia-----	40	Unlimited		Very limited Depth to saturated zone Restricted permeability Slope	1.00 1.00 0.41	Somewhat limited Too droughty Slope Somewhat Limited Water Quantity	0.75 0.41 0.17	Somewhat limited Slope Somewhat Limited Water Quantity	0.41 0.17
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.41	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.41	Very limited Limited Water Quantity Slope	0.99 0.41
250B: Sutton-----	40	Unlimited		Very limited Depth to saturated zone Restricted permeability Slope	1.00 0.34 0.15	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.15	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.15 0.07
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
253B: Wapping-----	40	Unlimited		Very limited Depth to saturated zone Slope	1.00 0.15	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.15	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.15 0.07

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
253B: Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
255B: Watchaug-----	40	Unlimited		Very limited Depth to saturated zone Restricted permeability Slope	1.00 0.74 0.15	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.15	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.15	Very limited Limited Water Quantity Slope	0.99 0.15
260B: Charlton-----	40	Unlimited		Somewhat limited Slope Restricted permeability	0.74 0.50	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.74	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.74	Very limited Limited Water Quantity Slope	0.99 0.74

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
260C: Charlton-----	40	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	1.00	Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
260D: Charlton-----	40	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
Urban land-----	35	Very limited Depth limited Slope	1.00 0.12	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	1.00	Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
263B: Cheshire-----	40	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity	0.99
				Restricted permeability	0.50	Limited Water Quantity	0.99	Slope	0.74
						Too droughty	0.75		
						Slope	0.74		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
263B: Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.74	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
263C: Cheshire-----	40	Unlimited		Very limited Slope Restricted permeability	1.00 0.50	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
266B: Narragansett-----	40	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.07
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.74	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.74	Very limited Limited Water Quantity Slope	0.99 0.74

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value						
269B: Yalesville-----	40	Somewhat limited Bedrock	0.66	Very limited	1.00	Very limited	0.99	Very limited	0.99
	Thin layer			Limited Water Quantity		Limited Water Quantity			
	Slope			Slope		Slope			
				Restricted permeability	0.74	Somewhat droughty	0.74	0.25	0.74
Urban land-----	35	Very limited Depth limited	1.00	Very limited	1.00	Very limited	1.00	Very limited	0.99
	Restricted permeability			Too droughty		Limited Water Quantity			
	Slope			Limited Water Quantity		Slope			
					0.74		0.74		0.74
269C: Yalesville-----	40	Somewhat limited Bedrock	0.66	Very limited	1.00	Very limited	1.00	Very limited	1.00
	Slope			Slope		Limited Water Quantity			
	Thin layer			Limited Water Quantity		Limited Water Quantity			
	Restricted permeability			Somewhat droughty		Somewhat droughty			
Urban land-----	35	Very limited Depth limited	1.00	Very limited	1.00	Very limited	1.00	Very limited	1.00
	Slope			Slope		Limited Water Quantity			
	Restricted permeability			Too droughty		Limited Water Quantity			
					1.00		0.99		0.99
273C: Urban land-----	35	Very limited Depth limited	1.00	Very limited	1.00	Very limited	1.00	Very limited	1.00
	Slope			Slope		Limited Water Quantity			
	Restricted permeability			Too droughty		Limited Water Quantity			
					1.00		0.99		0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
273C: Charlton-----	25	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
Chatfield-----	15	Very limited Bedrock	0.89	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.49	Too droughty	0.75		
273E: Urban land-----	35	Very limited Depth limited Slope	1.00 0.88	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	1.00	Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
Charlton-----	25	Very limited Slope	0.88	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.50	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
						Too droughty	0.75		
Chatfield-----	15	Very limited Bedrock Slope	0.89 0.88	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.49	Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
275C: Urban land-----	45	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Chatfield-----	30	Very limited Bedrock	0.89	Very limited Slope Thin layer Restricted permeability	1.00 1.00 0.49	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
275E: Urban land-----	35	Very limited Depth limited Slope	1.00 0.88	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Chatfield-----	25	Very limited Bedrock Slope	0.89 0.88	Very limited Slope Thin layer Restricted permeability	1.00 1.00 0.49	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
Rock outcrop-----	15	Very limited Depth limited Bedrock Slope	1.00 1.00 0.88	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
282B: Broadbrook-----	40	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Thin layer	1.00	Slope	0.74	Slope	0.74
				Restricted permeability	0.97	Somewhat droughty	0.25		
				Slope	0.74				
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability	1.00	Very limited Too droughty	1.00	Very limited Limited Water Quantity	0.99
				Slope	0.74	Limited Water Quantity	0.99	Slope	0.74
						Slope	0.74		
284B: Paxton-----	40	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Thin layer	1.00	Slope	0.74	Slope	0.74
				Restricted permeability	0.97	Somewhat droughty	0.25		
				Slope	0.74				
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability	1.00	Very limited Too droughty	1.00	Very limited Limited Water Quantity	0.99
				Slope	0.74	Limited Water Quantity	0.99	Slope	0.74
						Slope	0.74		
284C: Paxton-----	40	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Thin layer	1.00	Somewhat droughty	0.25		
				Restricted permeability	0.97				

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
284C: Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
284D: Paxton-----	40	Somewhat limited Slope	0.12	Very limited Depth to saturated zone Slope Thin layer Restricted permeability	1.00 1.00 1.00 0.97	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
Urban land-----	35	Very limited Depth limited Slope	1.00 0.12	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
287B: Wethersfield-----	40	Unlimited		Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.97 0.74	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.74 0.25	Very limited Limited Water Quantity Slope	0.99 0.74
Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.74	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.74	Very limited Limited Water Quantity Slope	0.99 0.74

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
287C: Wethersfield-----	40	Unlimited		Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer Restricted permeability	1.00 0.97	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
287D: Wethersfield-----	40	Somewhat limited Slope	0.12	Very limited Depth to saturated zone Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer Restricted permeability	1.00 0.97	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
Urban land-----	35	Very limited Depth limited Slope	1.00 0.12	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
290B: Stockbridge-----	40	Unlimited		Very limited Restricted permeability Dense layer Slope	1.00 0.83 0.74	Very limited Limited Water Quantity Too droughty Slope	0.99 0.75 0.74	Very limited Limited Water Quantity Slope	0.99 0.74

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
290B: Urban land-----	35	Very limited Depth limited	1.00	Very limited Restricted permeability Slope	1.00 0.74	Very limited Too droughty Limited Water Quantity Slope	1.00 0.99 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
290C: Stockbridge-----	40	Unlimited		Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
Urban land-----	35	Very limited Depth limited	1.00	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
290D: Stockbridge-----	40	Somewhat limited Slope	0.12	Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
Urban land-----	35	Very limited Depth limited Slope	1.00 0.12	Very limited Slope Restricted permeability	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
301: Beaches-----	50	Very limited Flooding Depth limited	1.00 1.00	Very limited Depth to saturated zone Flooding Slope	1.00 1.00 0.41	Somewhat limited Ksat - Subject to seepage Slope	1.00 0.41	Somewhat limited Subject To Seepage Slope	1.00 0.41

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
301: Udipsamments-----	35	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
				Slope	1.00	Too droughty	1.00	Slope	1.00
				Depth to saturated zone	0.24	Slope	1.00	Limited Water Quantity	0.87
						Limited Water Quantity	0.87	Subject To Seepage	0.28
302: Dumps-----	95	Very limited Depth limited	1.00	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	0.95	Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99		
303: Pits, quarries-----	90	Very limited Depth limited	1.00	Very limited Thin layer	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Bedrock	1.00	Restricted permeability	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
		Slope	0.50	Slope	1.00				
304: Udorthents-----	90	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	0.53	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Depth to saturated zone	0.28	Too droughty	0.75		
305: Udorthents-----	65	Somewhat limited Slope	0.04	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.53	Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
305: Pits-----	25	Very limited Depth limited Slope	1.00 1.00	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28
306: Udorthents-----	50	Somewhat limited Slope	0.04	Very limited Slope Restricted permeability Depth to saturated zone	1.00 0.53 0.28	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
Urban land-----	35	Very limited Depth limited Slope	1.00 0.04	Very limited Restricted permeability Slope	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
307: Urban land-----	80	Very limited Depth limited Slope	1.00 0.32	Very limited Restricted permeability Slope	1.00 1.00	Very limited Slope Too droughty Limited Water Quantity	1.00 1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
308: Udorthents-----	80	Somewhat limited Slope	0.04	Very limited Depth to saturated zone Slope Restricted permeability	1.00 1.00 0.53	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
309: Udorthents-----	80	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Slope	0.04	Slope	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability Flooding	0.53 0.40	Too droughty	0.75		
310: Udorthents, Periodically Flooded-----	85	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Slope	1.00	Very limited Slope	1.00
		Slope	0.04	Depth to saturated zone	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Slope	1.00	Too droughty	0.75		
				Restricted permeability	0.53				
401C: Macomber-----	55	Very limited Bedrock	0.87	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.95	Somewhat droughty	0.25		
Taconic-----	30	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	0.16	Limited Water Quantity	0.99	Limited Water Quantity	0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
402D: Maconber-----	50	Very limited Bedrock Slope	0.87 0.12	Very limited Slope Thin layer Restricted permeability	1.00 1.00 0.95	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
Taconic-----	25	Very limited Depth limited Bedrock Slope	1.00 1.00 0.12	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.16	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Rock outcrop-----	15	Very limited Depth limited Bedrock Slope	1.00 1.00 0.12	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
403C: Taconic-----	70	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.16	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Rock outcrop-----	25	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
403E: Taconic-----	70	Very limited Depth limited Bedrock Slope	1.00 1.00 1.00	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.16	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
403E: Rock outcrop-----	20	Very limited Depth limited Bedrock Slope	1.00 1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
403F: Taconic-----	70	Very limited Slope Depth limited Bedrock	1.00 1.00 1.00	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.16	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Rock outcrop-----	20	Very limited Slope Depth limited Bedrock	1.00 1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
405C: Dummerston-----	85	Unlimited		Very limited Slope Restricted permeability	1.00 0.94	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
405E: Dummerston-----	85	Somewhat limited Slope	0.50	Very limited Slope Restricted permeability	1.00 0.94	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
407C: Lanesboro-----	85	Unlimited		Very limited Depth to saturated zone Thin layer Slope Restricted permeability	1.00 1.00 1.00 0.93	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
407E: Lanesboro-----	85	Very limited Slope	0.88	Very limited Depth to saturated zone Thin layer Slope Restricted permeability	1.00 1.00 1.00 0.93	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
408C: Fullam-----	85	Unlimited		Very limited Depth to saturated zone Thin layer Slope Restricted permeability	1.00 1.00 1.00 0.73	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99
409B: Brayton-----	85	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Slope Restricted permeability	1.00 1.00 0.41 0.02	Somewhat limited Slope	0.41	Somewhat limited Slope	0.41

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
412B: Bice-----	85	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Limited Water Quantity Slope	0.99
				Restricted permeability	0.45				
412C: Bice-----	85	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.45				
412D: Bice-----	85	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.45				
413C: Bice-----	45	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope Limited Water Quantity	1.00 0.99
				Restricted permeability	0.45				

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
413C: Millsite-----	40	Very limited Bedrock	0.84	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.37	Too droughty	0.75		
413E: Bice-----	45	Somewhat limited Slope	0.50	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
				Restricted permeability	0.45	Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Too droughty	0.75
Millsite-----	40	Very limited Bedrock Slope	0.84 0.50	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.37	Too droughty	0.75		
414: Fredon, cold-----	85	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.07
415C: Millsite-----	40	Very limited Bedrock	0.84	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Thin layer	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Restricted permeability	0.37	Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
415C: Westminster-----	40	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.01	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Rock outcrop-----	15	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
415E: Millsite-----	40	Very limited Slope Bedrock	0.88 0.84	Very limited Slope Thin layer Restricted permeability	1.00 1.00 0.37	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
Westminster-----	40	Very limited Depth limited Bedrock Slope	1.00 1.00 0.88	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.01	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Rock outcrop-----	15	Very limited Depth limited Bedrock Slope	1.00 1.00 0.88	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
416E: Rock outcrop-----	70	Very limited Depth limited Bedrock Slope	1.00 1.00 0.50	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
416E: Westminster-----	20	Very limited Depth limited Bedrock	1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
		Slope	0.50	Restricted permeability	0.01				
416F: Rock outcrop-----	70	Very limited Slope Depth limited Bedrock	1.00 1.00 1.00	Very limited Thin layer Slope	1.00 1.00	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
Westminster-----	20	Very limited Slope Depth limited Bedrock	1.00 1.00 1.00	Very limited Thin layer Slope Restricted permeability	1.00 1.00 0.01	Very limited Slope Limited Water Quantity	1.00 0.99	Very limited Slope Limited Water Quantity	1.00 0.99
417B: Bice-----	85	Unlimited		Somewhat limited Slope Restricted permeability	0.74 0.45	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
417C: Bice-----	85	Unlimited		Very limited Slope Restricted permeability	1.00 0.45	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems			
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value		
417D: Bice-----	85	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope	1.00		
	Restricted permeability			0.45	Limited Water Quantity			0.99			
					Too droughty			0.75			
418C: Schroon-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Slope	1.00		
				Slope	1.00			Limited Water Quantity	0.99		
				Restricted permeability	0.23			Limited Water Quantity	0.99	Subject To Seepage	0.07
						Too droughty	0.75				
420A: Schroon-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Limited Water Quantity	0.99		
				Restricted permeability	0.23			Limited Water Quantity	0.99	Subject To Seepage	0.07
								Too droughty	0.75		
420B: Schroon-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Slope	1.00	Very limited Limited Water Quantity	0.99		
				Slope	0.41			Limited Water Quantity	0.99	Slope	0.41
				Restricted permeability	0.23			Too droughty	0.75	Subject To Seepage	0.07
										Slope	0.41

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
421A: Ninigret, cold-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage Too droughty	1.00 0.75	Unlimited Subject To Seepage Somewhat Limited Water Quantity	0.28 0.12
423A: Sudbury, cold-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Limited Water Quantity Subject To Seepage	0.99 0.28
424B: Shelburne-----	85	Unlimited		Very limited Depth to saturated zone Thin layer Slope Restricted permeability	1.00 1.00 0.74 0.39	Very limited Limited Water Quantity Slope Somewhat droughty	0.99 0.74 0.25	Very limited Limited Water Quantity Slope	0.99 0.74
424C: Shelburne-----	85	Unlimited		Very limited Depth to saturated zone Thin layer Slope Restricted permeability	1.00 1.00 1.00 0.39	Very limited Slope Limited Water Quantity Somewhat droughty	1.00 0.99 0.25	Very limited Slope Limited Water Quantity	1.00 0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
424D: Shelburne-----	85	Somewhat limited Slope	0.12	Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope Restricted permeability	1.00 0.39	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
425B: Shelburne-----	85	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Slope Restricted permeability	0.74 0.39	Slope Somewhat droughty	0.74 0.25	Slope	0.74
425C: Shelburne-----	85	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope Restricted permeability	1.00 0.39	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99
426D: Shelburne-----	85	Somewhat limited Slope	0.50	Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope Restricted permeability	1.00 0.39	Limited Water Quantity Somewhat droughty	0.99 0.25	Limited Water Quantity	0.99

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
427B: Ashfield-----	85	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Dense layer	1.00	Somewhat droughty	0.25	Somewhat Limited Water Quantity	0.06
				Restricted permeability Slope	0.88 0.74	Somewhat Limited Water Quantity	0.06		
427C: Ashfield-----	85	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Somewhat droughty	0.25	Somewhat Limited Water Quantity	0.06
				Restricted permeability	0.14	Somewhat Limited Water Quantity	0.06		
428A: Ashfield-----	85	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Somewhat droughty	0.25	Unlimited Somewhat Limited Water Quantity	0.06
				Restricted permeability	0.14	Somewhat Limited Water Quantity	0.06		
428B: Ashfield-----	85	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Somewhat limited Slope	0.74	Somewhat limited Slope	0.74
				Slope	0.74	Somewhat droughty	0.25	Somewhat Limited Water Quantity	0.06
				Restricted permeability	0.14	Somewhat Limited Water Quantity	0.06		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
428C: Ashfield-----	85	Unlimited		Very limited Depth to saturated zone Thin layer	1.00 1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Somewhat Limited Water Quantity	0.06	Somewhat Limited Water Quantity	0.06
				Restricted permeability	0.14				
429A: Agawam, cold-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited Subject To Seepage	1.00
						Limited Water Quantity Too droughty	0.99 0.75	Limited Water Quantity	0.99
429B: Agawam, cold-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage	1.00	Very limited Subject To Seepage	1.00
						Limited Water Quantity Too droughty Slope	0.99 0.75 0.74	Limited Water Quantity Slope	0.99 0.74
429C: Agawam, cold-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited Slope	1.00
						Slope	1.00	Subject To Seepage	1.00
						Limited Water Quantity Too droughty	0.99 0.75	Limited Water Quantity	0.99
433: Moosilauke-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
434A: Merrimac, cold-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Limited Water Quantity Subject To Seepage	0.99 0.28
434B: Merrimac, cold-----	80	Unlimited		Somewhat limited Slope	0.74	Very limited Ksat - Subject to seepage Limited Water Quantity Too droughty Slope	1.00 0.99 0.75 0.74	Very limited Limited Water Quantity Slope Subject To Seepage	0.99 0.74 0.28
434C: Merrimac, cold-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage Slope Limited Water Quantity Too droughty	1.00 1.00 0.99 0.75	Very limited Slope Limited Water Quantity Subject To Seepage	1.00 0.99 0.28
435: Scarboro-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28
436: Halsey-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
437: Wonsqueak-----	85	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Unlimited		Unlimited	
		Depth limited	1.00	Flooding	0.40				
438: Bucksport-----	85	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Unlimited		Unlimited	
		Depth limited	1.00	Flooding	0.40	Somewhat Limited Water Quantity	0.01	Somewhat Limited Water Quantity	0.01
				Restricted permeability	0.22				
440A: Boscawen-----	80	Unlimited		Unlimited		Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
						Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
440C: Boscawen-----	80	Unlimited		Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
						Too droughty	1.00	Slope	1.00
						Slope	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28
440E: Boscawen-----	85	Very limited Slope	0.88	Very limited Slope	1.00	Very limited Ksat - Subject to seepage	1.00	Very limited High infiltration	1.00
						Slope	1.00	Slope	1.00
						Too droughty	1.00	Limited Water Quantity	0.99
						Limited Water Quantity	0.99	Subject To Seepage	0.28

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
442: Brayton-----	85	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.57 0.41	Somewhat limited Slope	0.41	Somewhat limited Slope	0.41
443: Brayton-----	50	Very limited Depth limited	1.00	Very limited Depth to saturated zone Thin layer Restricted permeability Slope	1.00 1.00 0.57 0.41	Somewhat limited Slope	0.41	Somewhat limited Slope	0.41
Loonmeadow-----	35	Very limited Depth limited	1.00	Very limited Depth to saturated zone Dense layer Restricted permeability	1.00 1.00 0.89	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01
448B: Hogansburg-----	85	Unlimited		Very limited Depth to saturated zone Restricted permeability Slope Dense layer	1.00 0.97 0.41 0.03	Somewhat limited Too droughty Slope Somewhat Limited Water Quantity	0.75 0.41 0.17	Somewhat limited Slope Somewhat Limited Water Quantity	0.41 0.17

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
449B: Hogansburg-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Somewhat limited Too droughty	0.75	Somewhat limited Slope	0.74
				Restricted permeability	0.97	Slope	0.74	Somewhat Limited Water Quantity	0.17
				Slope	0.74	Somewhat Limited Water Quantity	0.17		
				Dense layer	0.03				
449C: Hogansburg-----	85	Unlimited		Very limited Depth to saturated zone	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Slope	1.00	Too droughty	0.75	Somewhat Limited Water Quantity	0.17
				Restricted permeability	0.97	Somewhat Limited Water Quantity	0.17		
				Dense layer	0.03				
450B: Pyrities-----	80	Unlimited		Very limited Restricted permeability	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99
				Dense layer	0.83	Too droughty	0.75	Slope	0.74
				Slope	0.74	Slope	0.74		
450C: Pyrities-----	80	Unlimited		Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Dense layer	0.83	Too droughty	0.75		
450D: Pyrities-----	80	Somewhat limited Slope	0.12	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
				Restricted permeability	1.00	Limited Water Quantity	0.99	Limited Water Quantity	0.99
				Dense layer	0.83	Too droughty	0.75		

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems	
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
451B: Pyrities-----	80	Unlimited		Very limited Restricted permeability Dense layer Slope	1.00 0.83 0.74	Very limited Limited Water Quantity Too droughty Slope	0.99 0.75 0.74	Very limited Limited Water Quantity Slope	0.99 0.74
451C: Pyrities-----	80	Unlimited		Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
451D: Pyrities-----	80	Somewhat limited Slope	0.50	Very limited Slope Restricted permeability Dense layer	1.00 1.00 0.83	Very limited Slope Limited Water Quantity Too droughty	1.00 0.99 0.75	Very limited Slope Limited Water Quantity	1.00 0.99
457: Mudgepond-----	80	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability Dense layer	1.00 0.86 0.03	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	
458: Mudgepond-----	55	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability Dense layer	1.00 0.86 0.03	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited	
Alden-----	35	Very limited Depth limited	1.00	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Unlimited Somewhat Limited Water Quantity	0.01	Unlimited Somewhat Limited Water Quantity	0.01

Table 28.—Storm Water Runoff Systems—Continued

Map symbol and soil name	Pct. of map unit	Storm water basins		Infiltration systems		Perennial wetland systems		Intermittent wetland systems			
		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value		
501: Ondawa-----	85	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Limited Water Quantity	0.99	Very limited Limited Water Quantity	0.99		
				Restricted permeability	0.97					Too droughty	0.75
				Dense layer Depth to saturated zone	0.83 0.14						
503: Rumney-----	80	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28		
				Depth limited	1.00					Somewhat Limited Water Quantity	0.01
				Depth limited	1.00						
508: Medomak-----	85	Very limited Flooding	1.00	Very limited Depth to saturated zone	1.00	Somewhat limited Ksat - Subject to seepage	1.00	Unlimited Subject To Seepage	0.28		
				Depth limited	1.00					Somewhat limited Ksat - Subject to seepage	1.00
				Depth limited	1.00						

Table 29.—Taxonomic Classification of the Soils

(An asterisk in the first column indicates a taxadjunct to the series. See text for a description of those characteristics that are outside the range of the series.)

Soil name	Family or higher taxonomic class
Agawam-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts
*Agawam-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, frigid Typic Dystrudepts
Alden-----	Fine-loamy, mixed, active, nonacid, mesic Mollic Endoaquepts
*Alden-----	Fine-loamy, mixed, active, nonacid, frigid Mollic Endoaquepts
Amenia-----	Coarse-loamy, mixed, active, mesic Aquic Eutrudepts
Ashfield-----	Coarse-loamy, mixed, active, frigid Aquic Dystrudepts
Bash-----	Coarse-loamy, mixed, semiactive, mesic Fluvaquentic Dystrudepts
Belgrade-----	Coarse-silty, mixed, active, mesic Aquic Dystric Eutrudepts
Berlin-----	Fine-silty, mixed, semiactive, mesic Aquic Dystric Eutrudepts
Bernardston-----	Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts
Bice-----	Coarse-loamy, mixed, active, frigid Typic Dystrudepts
Boscawen-----	Sandy-skeletal, mixed, frigid Typic Udorthents
Brancroft-----	Fine-silty, mixed, active, mesic Aquic Dystric Eutrudepts
Branford-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts
*Brayton-----	Coarse-loamy, mixed, active, nonacid, frigid Typic Humaquepts
*Brayton-----	Loamy, mixed, active, nonacid, frigid, shallow Typic Endoaquepts
Brimfield-----	Loamy, mixed, active, mesic Lithic Dystrudepts
Broadbrook-----	Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts
Brookfield-----	Coarse-loamy, mixed, active, mesic Typic Dystrudepts
Bucksport-----	Euic, frigid Typic Haplosaprists
Canton-----	Coarse-loamy over sandy or sandy-skeletal, mixed, semiactive, mesic Typic Dystrudepts
Catden-----	Euic, mesic Typic Haplosaprists
Charlton-----	Coarse-loamy, mixed, active, mesic Typic Dystrudepts
Chatfield-----	Coarse-loamy, mixed, superactive, mesic Typic Dystrudepts
Cheshire-----	Coarse-loamy, mixed, semiactive, mesic Typic Dystrudepts
Copake-----	Coarse-loamy over sandy or sandy-skeletal, mixed, semiactive, mesic Dystric Eutrudepts
Deerfield-----	Mixed, mesic Aquic Udipsamments
Dummerston-----	Coarse-loamy, mixed, active, frigid Typic Dystrudepts
Ellington-----	Coarse-loamy over sandy or sandy-skeletal, mixed, subactive, mesic Aquic Dystrudepts
Elmridge-----	Coarse-loamy over clayey, mixed, semiactive, mesic Aquic Dystric Eutrudepts
Enfield-----	Coarse-silty over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts
Farmington-----	Loamy, mixed, active, mesic Lithic Eutrudepts
Fluvaquents-----	Fluvaquents
Fredon-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, nonacid, mesic Aeric Endoaquepts
*Fredon-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, nonacid, frigid Aeric Endoaquepts
Freetown-----	Dysic, mesic Typic Haplosaprists
Fullam-----	Coarse-loamy, mixed, active, frigid Aquic Dystrudepts
Georgia-----	Coarse-loamy, mixed, semiactive, mesic Aquic Dystric Eutrudepts
Gloucester-----	Sandy-skeletal, mixed, mesic Typic Dystrudepts
Groton-----	Sandy-skeletal, mixed, mesic Typic Eutrudepts
Hadley-----	Coarse-silty, mixed, superactive, nonacid, mesic Typic Udifluvents
Halsey-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, nonacid, mesic Typic Humaquepts
*Halsey-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, nonacid, frigid Typic Humaquepts
Hartford-----	Sandy, mixed, mesic Typic Dystrudepts
Haven-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts
Hero-----	Coarse-loamy over sandy or sandy-skeletal, mixed, semiactive, mesic Aquic Eutrudepts
Hinckley-----	Sandy-skeletal, mixed, mesic Typic Udorthents
Hogansburg-----	Coarse-loamy, mixed, semiactive, frigid Aquic Eutrudepts
Hollis-----	Loamy, mixed, active, mesic Lithic Dystrudepts

Table 29.--Taxonomic Classification of the Soils--Continued

Soil name	Family or higher taxonomic class
Holyoke-----	Loamy, mixed, superactive, mesic Lithic Dystrudepts
Ipswich-----	Euic, mesic Typic Sulfihemists
Lanesboro-----	Coarse-loamy, mixed, active, frigid Oxyaquic Dystrudepts
Leicester-----	Coarse-loamy, mixed, active, acid, mesic Aeric Endoaquepts
Lim-----	Coarse-loamy, mixed, superactive, nonacid, mesic Fluvaquentic Endoaquepts
Limerick-----	Coarse-silty, mixed, active, nonacid, mesic Fluvaquentic Endoaquepts
Loonmeadow-----	Coarse-loamy, mixed, active, nonacid, frigid Mollic Endoaquepts
Ludlow-----	Coarse-loamy, mixed, semiactive, mesic Aquic Dystrudepts
Macomber-----	Loamy-skeletal, mixed, active, frigid Typic Dystrudepts
Manchester-----	Sandy-skeletal, mixed, mesic Typic Udorthents
Maybid-----	Fine, mixed, semiactive, nonacid, mesic Typic Humaquepts
*Medomak-----	Coarse-loamy over sandy or sandy-skeletal, mixed, superactive, nonacid, frigid Fluvaquentic Endoaquepts
Menlo-----	Coarse-loamy, mixed, active, mesic Typic Endoaquolls
Merrimac-----	Sandy, mixed, mesic Typic Dystrudepts
*Merrimac-----	Sandy, mixed, frigid Typic Dystrudepts
Millsite-----	Coarse-loamy, mixed, active, frigid Typic Dystrudepts
Montauk-----	Coarse-loamy, mixed, subactive, mesic Oxyaquic Dystrudepts
Moosilauke-----	Sandy, mixed, frigid Aeric Endoaquepts
Mudgepond-----	Coarse-loamy, mixed, superactive, mesic Typic Endoaquolls
*Mudgepond-----	Coarse-loamy, mixed, superactive, frigid Typic Endoaquolls
Narragansett-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts
Natchaug-----	Loamy, mixed, euic, mesic Terric Haplosaprists
Nellis-----	Coarse-loamy, mixed, superactive, mesic Typic Eutrudepts
Ninigret-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Aquic Dystrudepts
*Ninigret-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, frigid Aquic Dystrudepts
Occum-----	Coarse-loamy, mixed, superactive, mesic Fluventic Dystrudepts
Ondawa-----	Coarse-loamy, mixed, active, frigid Fluventic Dystrudepts
Pawcatuck-----	Sandy or sandy-skeletal, mixed, euic, mesic Terric Sulfihemists
Paxton-----	Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts
Penwood-----	Mixed, mesic Typic Udipsamments
Pootatuck-----	Coarse-loamy, mixed, active, mesic Fluvaquentic Dystrudepts
Pyrities-----	Coarse-loamy, mixed, active, frigid Dystric Eutrudepts
Rainbow-----	Coarse-loamy, mixed, active, mesic Aquic Dystrudepts
Raynham-----	Coarse-silty, mixed, active, nonacid, mesic Aeric Epiaquepts
Raypol-----	Coarse-loamy over sandy or sandy-skeletal, mixed, active, acid, mesic Aeric Endoaquepts
*Ridgebury-----	Coarse-loamy, mixed, active, nonacid, mesic Aeric Endoaquepts
Rippowam-----	Coarse-loamy, mixed, superactive, nonacid, mesic Fluvaquentic Endoaquepts
Rumney-----	Coarse-loamy, mixed, active, nonacid, frigid Fluvaquentic Endoaquepts
Saco-----	Coarse-silty, mixed, active, nonacid, mesic Fluvaquentic Humaquepts
Scarboro-----	Sandy, mixed, mesic Histic Humaquepts
*Scarboro-----	Sandy, mixed, frigid Histic Humaquepts
Schroon-----	Coarse-loamy, mixed, superactive, frigid Aquic Dystrudepts
Scitico-----	Fine, mixed, semiactive, nonacid, mesic Typic Endoaquepts
Shaker-----	Coarse-loamy over clayey, mixed, semiactive, nonacid, mesic Aeric Epiaquepts
Shelburne-----	Coarse-loamy, mixed, active, frigid Oxyaquic Dystrudepts
Stockbridge-----	Coarse-loamy, mixed, semiactive, mesic Dystric Eutrudepts
Sudbury-----	Sandy, mixed, mesic Aquic Dystrudepts
*Sudbury-----	Sandy, mixed, frigid Aquic Dystrudepts
Suncook-----	Mixed, mesic Typic Udipsamments
Sutton-----	Coarse-loamy, mixed, active, mesic Aquic Dystrudepts
Taconic-----	Loamy-skeletal, mixed, active, frigid Lithic Dystrudepts
Timakwa-----	Sandy or sandy-skeletal, mixed, euic, mesic Terric Haplosaprists
Tisbury-----	Coarse-silty over sandy or sandy-skeletal, mixed, active, mesic Aquic Dystrudepts
Udifuvents-----	Udifuvents
Udipsamments-----	Mesic Udipsamments
Udorthents-----	Udorthents
Walpole-----	Sandy, mixed, mesic Aeric Endoaquepts
Wapping-----	Coarse-loamy, mixed, active, mesic Aquic Dystrudepts
Watchaug-----	Coarse-loamy, mixed, semiactive, mesic Aquic Dystrudepts

Table 29.--Taxonomic Classification of the Soils--Continued

Soil name	Family or higher taxonomic class
Westbrook-----	Loamy, mixed, euic, mesic Terric Sulphemists
Westminster-----	Loamy, mixed, active, frigid Lithic Dystrudepts
Wethersfield-----	Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts
Whitman-----	Loamy, mixed, active, nonacid, mesic, shallow Typic Humaquepts
Wilbraham-----	Coarse-loamy, mixed, active, mesic Aquic Dystrudepts
Windsor-----	Mixed, mesic Typic Udipsamments
Winooski-----	Coarse-silty, mixed, active, mesic Fluvaquentic Dystrudepts
Wonsqueak-----	Loamy, mixed, euic, frigid Terric Haplosaprists
Woodbridge-----	Coarse-loamy, mixed, active, mesic Aquic Dystrudepts
Yalesville-----	Coarse-loamy, mixed, active, mesic Typic Dystrudepts

Table 30.—Relationships Among Parent Material, Dominant Texture, and Drainage of the Soils

(An asterisk indicates soils with a mean annual soil temperature less than 8 degrees C. These soils are at elevations greater than 1,300 feet in Litchfield County.)

Parent material	Excessively drained	Somewhat excessively drained	Well drained	Moderately well drained	Somewhat poorly drained	Poorly drained	Very poorly drained
Soils Formed in Glacial Till with Granite, Schist, and Gneiss Rock Lithology							
Sandy		Gloucester					
Loamy			Charlton Bice*	Sutton Schroon*		Leicester	Loonmeadow
Loamy over sandy substratum			Canton				
Loamy, firm substratum			Paxton Shelburne*	Woodbridge Ashfield*		Ridgebury	Whitman
Loamy over firm, sandy substratum			Montauk				
Loamy, 10 to 20 inches over bedrock		Hollis Westminster*					
Loamy, 20 to 40 inches over bedrock			Chatfield Millsite*				
Soils Formed in Glacial Till with Limestone and Crystalline Rock Lithology							
Loamy		Stockbridge Nellis Pyrities*	Georgia Amenia Hogansburg*			Mudgepond	Alden
Loamy, 10 to 20 inches over limestone bedrock			Farmington				

Table 30.—Relationships Among Parent Material, Dominant Texture, and Drainage of the Soils

Parent material	Excessively drained	Somewhat excessively drained	Well drained	Moderately well drained	Somewhat poorly drained	Poorly drained	Very poorly drained
Soils Formed in Glacial Till with Sandstone, Shale, and Basalt Rock Lithology							
Loamy			Cheshire	Ludlow			
Loamy, firm substratum			Wethersfield	Watchaug		Wilbraham	Menlo
Loamy, 10 to 20 inches over bedrock			Holyoke				
Loamy, 20 to 40 inches over bedrock			Yalesville				
Soils Formed in Glacial Till with Micaceous Schist Rock Lithology							
Loamy			Brookfield				
Loamy, 10 to 20 inches over bedrock			Brimfield				
Soils Formed in Glacial Till with Phyllite Schist, and Slate Rock Lithology							
Loamy			Bernardston Dummerston*				
Loamy, firm substratum			Lanesboro*	Fullam*		Brayton*	
Loamy, 10 to 20 inches over bedrock		Taconic*					
Loamy, 20 to 40 inches over bedrock			Macomber*				
Soils Formed in Glacial Till with Sandstone, Shale, Basalt, and Crystalline Rock Lithology							
Loamy			Narragansett	Wapping			
Loamy, firm substratum			Broadbrook	Rainbow			

Table 30.—Relationships Among Parent Material, Dominant Texture, and Drainage of the Soils

Parent material	Excessively drained	Somewhat excessively drained	Well drained	Moderately well drained	Somewhat poorly drained	Poorly drained	Very poorly drained
Soils Formed in Glacial Outwash with Crystalline Rock Lithology							
Sandy and gravelly	Hinckley Boscawen*	Merrimac		Sudbury		Walpole Moosilauke*	
Sandy	Windsor			Deerfield			Scarboro
Loamy over sand and gravel			Agawam	Ninigret			
Silty over sand and gravel			Enfield Haven	Tisbury		Raypol	
Soils Formed in Glacial Outwash with Red Sandstone, Shale and Conglomerate Rock Lithology							
Sandy and gravelly	Manchester	Hartford					
Sandy	Penwood						
Silty over sand and gravel			Branford	Ellington			
Soils Formed in Glacial Outwash with Limestone and Crystalline Rock Lithology							
Sandy and gravelly		Groton					
Loamy over sand and gravel			Copake	Hero		Fredon	Halsey

Table 30.—Relationships Among Parent Material, Dominant Texture, and Drainage of the Soils

Parent material	Excessively drained	Somewhat excessively drained	Well drained	Moderately well drained	Somewhat poorly drained	Poorly drained	Very poorly drained
Soils Formed in Glaciolacustrine Deposits							
Silty				Belgrade		Raynham	
Loamy over clayey				Elmridge		Shaker	
Silty and clayey				Brancroft Berlin		Scitico	Maybid
Soils Formed in Alluvium							
Sandy	Suncook						
Loamy			Occum	Pootatuck		Rippowam	
Silty			Hadley Ondawa*	Winooski	Bash	Limerick Lim Rumney*	Saco Medomak*
Variable Texture			Unifluvents			Fluvaquents	
Soils Formed in Organic Material							
Organic deposits more than 51 inches deep							Catden Freetown Bucksport*
Organic deposits 16 to 51 inches deep over loamy mineral material							Natchaug Wonsqueak*

Table 30.—Relationships Among Parent Material, Dominant Texture, and Drainage of the Soils

Parent material	Excessively drained	Somewhat excessively drained	Well drained	Moderately well drained	Somewhat poorly drained	Poorly drained	Very poorly drained
Soils Formed in Organic Material							
Organic deposits 16 to 51 inches deep over sandy mineral material							Timakwa
Organic deposits 16 to 51 inches deep over sandy mineral material, in tidal areas							Pawcatuck
Organic deposits 16 to 51 inches deep over mineral material, in tidal areas							Westbrook
Organic deposits more than 51 inches deep, in tidal areas							Ipswich

About This Soil Survey Report

Welcome. This soil survey report contains information about the soils of Connecticut. Click on a subject of interest or browse the menu to view the files.

This online document consists of two main sections: the soil survey manuscript and the detailed soil maps.

If users save this document to their computers, they will only have saved the manuscript. The soil maps are not saved in the downloaded document. Complete versions of this survey (including both the manuscript and the detailed soil maps) are available on one CD from the [NRCS Connecticut](#) state office.

Soil Survey Manuscript

This contains general information about the survey area, the general and detailed soil map unit descriptions, the taxonomic unit descriptions, and the soil interpretation and soil properties tables.

Detailed Soil Maps

The detailed soil maps can be useful in planning the use and management of small areas (but not less than 3 acres in size). Each of the 409 detailed soil maps can be viewed online or saved individually to the user's computer.

Online Viewing of Soil Maps

From the Index to Map Sheets, click on any quarter-quadrangle to view the soil delineations and map unit symbols. The maps were made at a scale of 1:12,000 (one inch on the map is 1,000 feet on the ground) and are each 22 x 27 inches in size.