

SOIL SURVEY OF

# DeKalb County, Illinois



**United States Department of Agriculture**  
**Soil Conservation Service**  
In cooperation with  
**Illinois Agricultural Experiment Station**

This is a publication of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and agencies of the States, usually the Agricultural Experiment Stations. In some surveys, other Federal and local agencies also contribute. The Soil Conservation Service has leadership for the Federal part of the National Cooperative Soil Survey. In line with Department of Agriculture policies, benefits of this program are available to all who need the information, regardless of race, color, national origin, sex, religion, marital status, or age.

Major fieldwork for this soil survey was completed in the period 1967-71. Soil names and descriptions were approved in 1973. Unless otherwise indicated, statements in the publication refer to conditions in the county in 1971. This survey was made cooperatively by the Soil Conservation Service and the Illinois Agricultural Experiment Station. It is part of the technical assistance furnished to the DeKalb County Soil and Water Conservation District.

Soil maps in this survey may be copied without permission, but any enlargement of these maps could cause misunderstanding of the detail of mapping and result in erroneous interpretations. Enlarged maps do not show small areas of contrasting soils that could have been shown at a larger mapping scale.

## HOW TO USE THIS SOIL SURVEY

**T**HIS SOIL SURVEY contains information that can be applied in managing farms and woodlands; in selecting sites for roads, ponds, buildings, and other structures; and in judging the suitability of tracts of land for farming, industry, and recreation.

### Locating Soils

All the soils of DeKalb County are shown on the detailed map at the back of this publication. This map consists of many sheets made from aerial photographs. Each sheet is numbered to correspond with a number on the Index to Map Sheets.

On each sheet of the detailed map, soil areas are outlined and are identified by symbols. All areas marked with the same symbol are the same kind of soil. The soil symbol is inside the area if there is enough room; otherwise, it is outside and a pointer shows where the symbol belongs.

### Finding and Using Information

The "Guide to Mapping Units" can be used to find information. This guide lists all the soils of the county in numeric order by map symbol and gives the management group of each. It also shows the page where each soil is described and gives the tree and shrub planting group and the recreation group in which the soil has been placed.

Individual colored maps showing the relative suitability or degree of limitation of soils for many specific purposes can be developed by using the soil map and the information in the text. Translucent material can

be used as an overlay over the soil map and colored to show the soils that have the same limitation or suitability. For example, soils that have a slight limitation for a given use can be colored green, those that have a moderate limitation can be colored yellow, and those that have a severe limitation can be colored red.

*Farmers and those who work with farmers* can learn about use and management of the soils from the soil descriptions and from the discussions of the interpretive groups.

*Homeowners and others* can refer to the section "Use of the Soils for Ornamental Trees and Shrubs," where the soils of the county are grouped according to their suitability for trees and shrubs.

*Game managers, sportsmen, and others* can refer to the section "Use of the Soils for Wildlife," where general wildlife areas are described.

*Recreation specialists* will find pertinent information in the section "Recreational Uses of the Soils."

*Engineers and builders* will find, under "Engineering Uses of the Soils," tables that contain test data, estimates of soil properties, and information about soil features that affect engineering practices.


*Scientists, students, and others* can read about how the soils formed and how they are classified in the section "Formation and Classification of the Soils."

*Newcomers in DeKalb County* may be especially interested in the section "General Soil Map," where broad patterns of soils are described. They may also be interested in information about the county given at the beginning of the publication.

Cover: Typical area in the Saybrook-Drummer-Octagon association.

## Contents

	<i>Page</i>		<i>Page</i>
<b>General nature of the county</b> .....	1	<b>Muscatine series</b> .....	25
Relief, physiography, and drainage .....	1	<b>Octagon series</b> .....	25
Climate .....	1	<b>Peotone series</b> .....	26
<b>How this survey was made</b> .....	2	<b>Plano series</b> .....	26
<b>General soil map</b> .....	3	<b>Proctor series</b> .....	27
1. Saybrook-Drummer-Octagon		<b>Rush series</b> .....	28
association .....	3	<b>Sable series</b> .....	28
2. Drummer-Flanagan-Catlin		<b>St. Charles series</b> .....	29
association .....	4	<b>Sawmill series</b> .....	30
3. Drummer-Elburn-Batavia		<b>Saybrook series</b> .....	30
association .....	5	<b>Strawn series</b> .....	31
4. Sawmill-Camden-Harvard		<b>Virgil series</b> .....	31
association .....	6	<b>Will series</b> .....	32
5. Dodge-Miami association .....	8	<b>Use and management of the soils</b> .....	33
6. Muscatine-Sable-Catlin		General management of crops .....	33
association .....	8	Capability grouping .....	33
<b>Descriptions of the soils</b> .....	9	Management groups .....	34
Batavia series .....	10	Predicted yields .....	36
Bowes series .....	11	Use of the soils for ornamental trees	
Camden series .....	12	and shrubs .....	38
Catlin series .....	13	Use of the soils for wildlife .....	39
Cut and fill land .....	13	Recreational uses of the soils .....	40
Dodge series .....	14	Engineering uses of the soils .....	41
Dresden series .....	14	Engineering soil classification	
Drummer series .....	15	systems .....	60
Elburn series .....	16	Soil properties significant in	
Flanagan series .....	17	engineering .....	60
Gravel pits .....	17	Engineering interpretations of soils ..	62
Harpster series .....	17	Soil test data .....	63
Harvard series .....	18	<b>Formation and classification of the soils</b> ..	64
Herbert series .....	19	Factors of soil formation .....	64
Houghton series .....	20	Parent material .....	64
Kendall series .....	20	Climate .....	64
Knight series .....	21	Plants and animals .....	65
La Rose series .....	21	Relief and drainage .....	65
Limestone quarries .....	22	Time .....	65
Lisbon series .....	22	Classification of the soils .....	66
Lorenzo series .....	23	<b>Literature cited</b> .....	67
Miami series .....	23	<b>Glossary</b> .....	67
Millbrook series .....	24	<b>Guide to mapping units</b> .....	Following 67



Location of DeKalb County in Illinois.

# SOIL SURVEY OF DEKALB COUNTY, ILLINOIS

BY KENNETH C. HINKLEY, SOIL CONSERVATION SERVICE

SOILS SURVEYED BY R. J. HERMAN, K. C. HINKLEY, G. T. KELLER, B. G. STEWART, AND L. C. YOUNG,  
SOIL CONSERVATION SERVICE

UNITED STATES DEPARTMENT OF AGRICULTURE, SOIL CONSERVATION SERVICE, IN COOPERATION WITH THE  
ILLINOIS AGRICULTURAL EXPERIMENT STATION

**D**EKALB COUNTY is in the northeastern part of Illinois. It occupies 636 square miles. Sycamore, the county seat, is northeast of the center of the county.

## *General Nature of the County*

DeKalb County was established in 1835. Since settlement of the county, the population has been increasing to the present time. In 1970, the population of the county was 71,654 and that of DeKalb, the largest community, was about 33,000. DeKalb is the home of Northern Illinois University.

Farming has been the main enterprise in DeKalb County since its settlement. In 1969, according to the U.S. Census of Agriculture, there were 1,344 farms. The average farm was 287 acres. Slightly more than one-fourth of the farms have livestock.

Corn, soybeans, and oats are the main crops. According to the 1969 U.S. Census of Agriculture, 180,987 acres was in corn, 68,080 acres was in soybeans, and 17,751 acres was in oats. Hay was grown on 10,577 acres, including 7,675 acres of alfalfa. Vegetables were grown on 13,459 acres.

The total number of cattle in 1969 was 76,739, including 3,920 dairy cattle, and swine numbered 103,201.

The county has a well-developed transportation system. State Highways Nos. 72, 64, and 38 and U.S. Highway No. 30 cross the county from east to west. U.S. Highway No. 34 crosses the southeastern corner, and the Illinois Tollway crosses through the middle of county from east to west. State Highway No. 23 crosses from north to south. Main secondary roads are blacktop, and every farm is accessible by an all-weather road. Railroads also serve the county.

Many small industries are in DeKalb County, but most people are employed in businesses that serve farmers.

## **Relief, Physiography, and Drainage**

DeKalb County has relatively low relief. Elevation ranges from 650 feet above sea level one-half mile southwest of Sandwich to 990 feet above sea level 3

miles north of Lee. Elevation in most of the county is between 700 and 950 feet.

Most soils in the county are on uplands, which consist mainly of a glacial till plain covered by loess. The loess varies in thickness, but it is thicker in the western part of the county, especially the southwestern part, than it is in the eastern part. The major bottom-land areas are along Somonauk Creek, Indian Creek, and the Kishwaukee River.

The northern two-thirds of the county is drained by the Kishwaukee River and its tributaries. The water from these streams flows into the Rock River. The southern third of the county is drained by several smaller tributaries that flow southward into the Fox River, which in turn discharges into the Illinois River. The principal source of water is wells.

## **Climate**

DeKalb County has the continental climate typical of north-central Illinois. The annual temperature range is wide. Storm centers and associated weather fronts bring frequent changes in temperature, humidity, cloudiness, and wind direction during much of the year, but such changes are less frequent in summer.

Table 1 gives temperature and precipitation data for DeKalb County. It shows that summers are warm, but continuous warm periods are seldom prolonged. January is normally the coldest month. February often has days as cold as days in January, but cold periods in February are usually of shorter duration. The record low temperature of 25° F. below zero occurred in 1933.

Annual precipitation averages about 35 inches but has ranged from a low of about 21 inches to a high of about 50 inches. Monthly precipitation averages about 4 inches in May and June but only about 1¾ inches for the normally driest months of December to February (3).<sup>2</sup>

<sup>1</sup> Data furnished by WILLIAM L. DENMARK, climatologist for Illinois, National Weather Service, U.S. Department of Commerce, Champaign, Ill.

<sup>2</sup> Italic numbers in parentheses refer to Literature Cited, page 67.

TABLE 1.—*Temperature and precipitation data for DeKalb County, Illinois*

Month	Temperature				Precipitation	
	Average daily maximum	Average daily minimum	Record highest	Record lowest	Average total	Average snow-fall
	°F	°F	°F	°F	Inches	Inches
January	31	16	65	-24	1.8	7.2
February	33	18	73	-25	1.5	6.1
March	43	27	83	-12	2.4	5.3
April	58	38	93	13	2.9	.8
May	69	49	106	22	4.1	-----
June	80	59	107	33	4.5	-----
July	85	64	110	42	3.8	-----
August	83	62	107	35	3.9	-----
September	75	54	102	25	3.3	-----
October	63	44	92	10	2.9	-----
November	46	30	81	-11	2.3	2.1
December	34	20	67	-24	1.9	7.1
Year	58	40	110	-25	35.2	28.6

Because normal rainfall in July and August is not sufficient to meet the demands of a vigorously growing field crop, moisture must be stored in the subsoil during the previous fall and winter. Severe droughts are infrequent, but rather prolonged dry periods during part of the growing season are not unusual. Such periods usually cause reduced crop yields.

Most summer showers or thunderstorms are brief. A single thunderstorm often produces more than an inch of rain and is occasionally accompanied by hail and damaging winds. Damage to growing field crops by hail is most likely in June, July, and August. Hail-producing thunderstorms average about three per year in any one place and less than one during the

critical growing period (5). Not all hailstorms have stones of sufficient size or quantity to produce extensive crop damage.

The number of days between the average date of the last freeze (32° F. or below) in spring and the first freeze in fall has been termed the "growing season." This is approximately 157 days in DeKalb County. The term "growing season" is misleading, because different crops have different temperatures at which growth is affected. Table 2 indicates the probability of occurrence of several different freezing temperatures (6). Temperatures often vary considerably between ridges and valleys during radiation freezes, the type most common in Illinois. All freeze data are based on temperatures in a standard U.S. Weather Bureau thermometer shelter at a height of approximately 5 feet above the ground and in a representative exposure. Lower temperatures exist at times nearer the ground and in places that are subject to extreme air drainage.

### *How This Survey Was Made*

Soil scientists made this survey to learn what kinds of soil are in DeKalb County, where they are located, and how they can be used. They went into the county knowing they were likely to find many soils they had already seen and perhaps some they had not. As they worked throughout the county, they observed the steepness, length, and shape of slopes; the size and speed of streams; the kinds of native plants or crops; the kinds of rock; and many facts about the soils. They dug or bored many holes to expose soil profiles. A profile is the sequence of natural layers, or horizons, in a soil; it extends from the surface down to material that has not been changed much by leaching or by roots of plants.

The soil scientists made comparisons among the profiles they studied, and they compared these profiles with those in counties nearby and in places more distant. They classified and named the soils according to nationwide, uniform procedures. The *soil series* and *soil phase* are the categories of soil classification most used in a local survey.

TABLE 2.—*Probability of last freezing temperature in spring and first in fall*

Probability	Dates for given probability and temperature				
	32° F	28° F	24° F	20° F	16° F
<b>Last in spring:</b>					
Average date -----	May 5	April 23	April 6	March 30	March 19
25 percent chance after---	May 7	May 1	April 14	April 6	March 25
10 percent chance after---	May 17	May 9	April 22	April 14	April 2
<b>First in fall:</b>					
Average date -----	October 6	October 17	October 30	November 10	November 20
25 percent chance before---	September 30	October 14	October 24	October 30	November 10
10 percent chance before---	September 23	October 7	October 17	October 23	November 3

Soils that have profiles almost alike make up a soil series. Except for different texture in the surface layer, all the soils of one series have major horizons that are similar in thickness, arrangement, and other important characteristics. Each soil series is named for a town or other geographic feature near the place where a soil of that series was first observed and mapped. Miami and Flanagan, for example, are the names of two soil series. All the soils in the United States that have the same series name are essentially alike in those characteristics that affect their behavior in the undisturbed landscape.

Soils of one series can differ in texture of the surface soil and in slope, stoniness, or some other characteristic that affects use of the soils by man. On the basis of such differences, a soil series is divided into phases. The name of a soil phase indicates a feature that affects management. For example, Miami silt loam, 4 to 7 percent slopes, eroded, is one of three phases within the Miami series.

After a guide for classifying and naming the soils had been worked out, the soil scientists drew the boundaries of the individual soils on aerial photographs. These photographs show roads, buildings, field borders, trees, and other details that help in drawing boundaries accurately. The soil map in the back of this publication was prepared from the aerial photographs.

The areas shown on the soil map are called mapping units. On most maps detailed enough to be useful in planning the management of farms and fields, a mapping unit is nearly equivalent to a soil phase. It is not exactly equivalent, because it is not practical to show on such a map all the small, scattered bits of soil of some other kind that have been seen within an area that is dominantly of a recognized soil phase.

In most areas surveyed there are places where the soil material is so rocky, so shallow, or so severely eroded that it cannot be classified by soil series. These places are shown on the map and are described in the survey, but they are called land types and are given descriptive names. Cut and fill land is a land type in DeKalb County.

While a soil survey is in progress, samples of soils are taken, as needed, for laboratory measurements and for engineering tests. Laboratory data from the same kinds of soil in other places are assembled. Data on yields of crops under defined practices are assembled from farm records and from field or plot experiments on the same kinds of soil. Yields under defined management are estimated for all the soils.

But only part of a soil survey is done when the soils have been named, described, and delineated on the map, and the laboratory data and yield data have been assembled. The mass of detailed information then needs to be organized in such a way as to be readily useful to different groups, among them farmers, managers of woodland, and engineers.

On the basis of yield and practice tables and other data, the soil scientists set up trial groups. They test these groups by further study and by consultation with farmers, agronomists, engineers, and others; then they adjust the groups according to the results of their studies and consultation. Thus, the groups that are

finally evolved reflect up-to-date knowledge of the soils and their behavior under present methods of use and management.

## General Soil Map

The general soil map at the back of this survey shows, in color, the soil associations in DeKalb County. A soil association is a landscape that has a distinctive proportional pattern of soils. It normally consists of one or more major soils and at least one minor soil, and it is named for the major soils. The soils in one association may be in another, but in a different pattern.

A map that shows soil associations is useful to people who want a general idea of the soils in a county, who want to compare different parts of a county, or who want to know the location of large tracts that are suitable for a certain kind of farming or other land use. Such a map is not suitable for planning the management of a farm or field or for selecting the exact location of a road, building, or similar structure, because the soils in any one association ordinarily differ in slope, depth, drainage, or other characteristics that affect management.

The general soil map of DeKalb County does not join with that of La Salle County in most areas, because the composition of the soils in the two counties is different. The distribution and extent of the soils in each series vary from one survey area to the other. One soil association in La Salle County in range 3 does not join with DeKalb County, because the kinds of soil in that association only go a short distance into DeKalb County and it would have made a delineation too small to recognize at the scale used.

Also, the general soil map of DeKalb County does not join with that of Kendall County, because the composition of the soils is different. The distribution and extent of the soils in each series vary from one survey area to the other.

The soil associations in DeKalb County are described in the following pages.

### 1. Saybrook-Drummer-Octagon association

*Well-drained to poorly drained, sloping to nearly level soils that formed in silty material and the underlying loam glacial till; on uplands*

This soil association (fig. 1) consists of gently rolling, irregularly shaped ridges and drainageways on the glacial till plain and of rolling end moraines in areas of irregular topography and dominantly short slopes. Differences in elevation are commonly 10 to 20 feet. Ridges and slopes are interwoven with drainageways.

This association occupies about 48 percent of the county. It is about 40 percent Saybrook soils, 25 percent Drummer soils, 10 percent Octagon soils, and 25 percent minor soils.

Saybrook soils are gently sloping to sloping, well drained and moderately well drained soils that commonly occur on ridgetops. The surface layer typically is very dark brown and brown silt loam about 12 inches thick. The subsoil is dark yellowish-brown and dark-




Figure 1.—Pattern of soils in association 1 and their relationship to parent material.

brown silty clay loam and clay loam. It is underlain by loam till below a depth of 44 inches.

Drummer soils are nearly level and poorly drained. The surface layer typically is black silty clay loam about 18 inches thick. The subsoil is olive-gray and gray silty clay loam. The underlying material is stratified silt loam and sandy clay loam.

Octagon soils are gently sloping to sloping and well drained and moderately well drained. They commonly occur on side slopes and are more extensive in areas of end moraines. The surface layer typically is very dark grayish-brown silt loam about 7 inches thick. The subsoil is dark-brown and yellowish-brown clay loam. The underlying material is loam till below a depth of 28 inches.

Minor soils in this association are the Flanagan, Catlin, La Rose, and Harpster soils. The somewhat poorly drained Flanagan soils occupy the upper parts of drainageways, the edges of areas of gently sloping Saybrook soils, and small rises in poorly drained areas. The more sloping Octagon soils include areas of the thinner La Rose soils. Small areas of Catlin soils occur on gently sloping ridgetops. Small spots of Harpster soils are within areas of Drummer soils.

The soils in this association are well suited to all cultivated crops commonly grown in the county. The available water capacity is moderate in the La Rose soils but high in the other soils. Organic-matter is high in the Saybrook, Drummer, Flanagan, Catlin, and Harpster soils and medium in the Octagon and La Rose soils. The main concerns in managing these soils are improving drainage, controlling water erosion in the sloping areas, and maintaining tilth and fertility.

Most of this association is used for cultivated crops, but some of the steeper soils are used for pasture. All but the more sloping soils have a high potential for farming enterprises.

## 2. Drummer-Flanagan-Catlin association

*Poorly drained to well-drained, nearly level to sloping soils that formed in silty material and the underlying loam glacial till; on uplands*

This soil association (fig. 2) consists of broad, gently undulating, irregular ridges and drainageways on a large till plain. The gentle slopes in most places are long. Differences in elevation are commonly 5 to 10 feet. Some steeper areas occur, especially along small streams.




Figure 2.—Pattern of soils in association 2 and their relationship to parent material.

This association occupies about 24 percent of the county. It is about 40 percent Drummer soils, 25 percent Flanagan soils, 20 percent Catlin soils, and 15 percent minor soils.

Drummer soils are on large, somewhat flat areas and in drainageways. They are nearly level and poorly drained. The surface layer typically is black silty clay loam about 18 inches thick. The subsoil is olive-gray and gray silty clay loam. The underlying material is stratified silt loam and loam.

Flanagan soils are on the edges of gentle slopes. They are nearly level and somewhat poorly drained. The surface layer typically is black silt loam about 13 inches thick. The subsoil is mostly brown silty clay loam. The underlying material is loam till below a depth of 45 inches.

Catlin soils are on ridges. They are well drained and moderately well drained and gently sloping to sloping. The surface layer typically is very dark brown and dark-brown silt loam about 11 inches thick. The subsoil is mostly dark yellowish-brown silty clay loam. The underlying material is loam till below a depth of 65 inches.

Minor soils in this association are Saybrook and Octagon soils that are near the more sloping Catlin soils. The Saybrook and Octagon soils generally occur along small streams.

The soils in this association are well suited to all cultivated crops commonly grown in the county. All the soils have high available water capacity. The Drummer, Flanagan, Catlin, and Saybrook soils are high in organic-matter content, and the Octagon soil is medium in organic-matter content. The main concerns in managing these soils are controlling water erosion in the sloping areas, improving drainage, and maintaining tilth and fertility.

Nearly all of this association is used for cultivated crops. All the soils have a potential for farming enterprises.

### 3. Drummer-Elburn-Batavia association

*Poorly drained to well-drained, nearly level to sloping soils that formed in silty material and the underlying outwash, stratified water deposited silts, or water-worked loam drift; on uplands*

This soil association (fig. 3) consists of broad, nearly level and gently sloping or water-worked drift areas. The areas are mainly nearly level, but some of them are gently sloping and generally long. Differences in elevation are commonly less than 5 feet. Also in the association, in the vicinity of DeKalb and Malta, is an area of circular to elliptical, raised, flat-topped mounds. The mounds rise 5 to 15 feet above the adjacent gently




Figure 3.—Pattern of soils in association 3 and their relationship to parent material.

rolling areas and are commonly surrounded by low, poorly drained soils.

This association occupies about 15 percent of the county. It is about 40 percent Drummer soils, 20 percent Elburn soils, 15 percent Batavia soils, and 25 percent minor soils.

Drummer soils are nearly level and poorly drained. The surface layer typically is black silty clay loam about 18 inches thick. The subsoil is olive-gray silty clay loam. The underlying material is stratified silt loam and sandy clay loam.

Elburn soils are nearly level and somewhat poorly drained. The surface layer typically is very dark brown and very dark grayish-brown silt loam and light silty clay loam. The subsoil is dark grayish-brown and yellowish-brown silty clay loam. The underlying material is stratified loam and silt loam.

Batavia soils are on circular or elliptical mounds. They are nearly level on the tops of the mounds and gently sloping to sloping on the edges. They are well drained and moderately well drained. The surface layer typically is very dark gray silt loam about 10 inches thick. The subsoil is brown silty clay loam. The underlying material is stratified loam.

Minor soils in this association are the Plano, Virgil, Saybrook, and Harvard soils. The well drained and moderately well drained Plano soils are nearly level to gently sloping and occupy higher positions in the landscape. Nearly level, somewhat poorly drained Virgil soils are on the tops of some mounds. A few

small knobs are occupied by the well drained and moderately well drained Saybrook soils. The well drained and moderately well drained Harvard soils commonly are on the more sloping rims or edges of mounds.

The soils in this association are well suited to all crops in the county. All the soils have high available water capacity. The Drummer and Elburn soils are high in organic-matter content. The Harvard soils are medium in organic-matter content. Improving drainage and maintaining tilth and fertility are the main concerns in managing the Drummer and Elburn soils. Controlling water erosion is the main concern of management on the Harvard soils.

Most of this association is used for cultivated crops. All but the more sloping soils have a high potential for farming enterprises.

#### 4. Sawmill-Camden-Harvard association

*Poorly drained, nearly level soils that formed in clay loam and silty clay loam alluvium on bottom lands and well drained and moderately well drained, sloping to nearly level soils that formed in silty material and the underlying outwash material; on terraces and uplands*

Most of this association (fig. 4) is along the major streams and rivers. The association consists of nearly level Sawmill soils that border the streams and that commonly are between areas of the well-drained Camden and Harvard soils. Much of the area adjacent




Figure 4.—Pattern of soils in association 4 and their relationship to parent material.

to the Sawmill soils is nearly level to gently sloping, but some areas are steeper and generally have short slopes. Some gravel pits occur throughout the association.

This association occupies about 7 percent of the county. It is about 45 percent Sawmill soils, 25 percent Camden soils, 15 percent Harvard soils, and 15 percent minor soils.

Sawmill soils are nearly level and poorly drained. They are low lying and subject to flooding. The surface layer typically is black silty clay loam about 31 inches thick. The subsoil is very dark gray and dark-gray silty clay loam. The underlying material is clay loam that has strata of sand and gravel.

Camden soils are nearly level to sloping and well drained and moderately well drained. The surface layer typically is very dark grayish-brown silt loam about 5 inches thick. The subsurface layer is dark grayish-brown silt loam. The subsoil is brown and dark-brown silty clay loam and clay loam. The underlying material is stratified sandy loam and silt loam.

Harvard soils are nearly level to sloping and well drained and moderately well drained. The surface layer typically is black silt loam about 12 inches thick. The subsoil is dark yellowish-brown silty clay loam and clay loam. The underlying material is loam.

Minor soils in this association are the Millbrook, St. Charles, Kendall, Drummer, Dresden, Bowes, and Rush

soils. The somewhat poorly drained Millbrook soils occupy the upper end of drainageways or are on nearly level uplands. The well drained and moderately well drained St. Charles soils and the somewhat poorly drained Kendall soils are associated, and they occur mostly along Somonauk Creek in Sandwich Township. The poorly drained Drummer soils occupy the lower end of natural drainageways and small depressions. The well-drained Dresden, Bowes, and Rush soils occupy areas that are underlain by sand and gravel and occur mostly in the northern part of the county.

The soils of this association are suited to all cultivated crops commonly grown in the county. The Sawmill soils are high in organic-matter content and have high available water capacity. The Harvard soils have high available water capacity and are moderate in organic-matter content. The Camden soils have high available water capacity and are low in organic-matter content. The main concerns in managing these soils are improving drainage on the poorly drained and somewhat poorly drained soils, protecting soils on bottom lands from flooding, controlling water erosion on the sloping soils, and maintaining tilth and fertility.

Most of this association is used for cultivated crops, although some areas are used for pasture and some remain in native woods. Most of the gravel pits in the county occur in this association.

### 5. Dodge-Miami association

*Well drained and moderately well drained, nearly level to strongly sloping soils that formed in silty material and the underlying loam glacial till; on uplands*

This soil association (fig. 5) occupies parts of the rolling end moraines and occurs on valley sides along the major streams. Slopes are irregular and dominantly short, and steeper areas are commonly adjacent to drainageways or streams.

This association occupies about 5 percent of the county. It is about 50 percent Dodge soils, 30 percent Miami soils, and 20 percent minor soils.

Dodge soils are nearly level to sloping and well drained and moderately well drained. The surface layer typically is brown and dark-brown silt loam about 6 inches thick. The subsoil is dark yellowish-brown silty clay loam and dark-brown clay loam. The underlying material is loam till below a depth of 31 inches.

Miami soils are gently sloping to strongly sloping and well drained and moderately well drained. The surface layer typically is dark grayish-brown silt loam about 6 inches thick. The subsoil is dark yellowish-brown and brown clay loam. The underlying material

is loam till below a depth of 31 inches.

Minor soils in this association are the Herbert and Drummer soils. Small areas of the somewhat poorly drained Herbert soils occur along the edges of areas of Dodge and Miami soils and as slight rises in the poorly drained areas. Drummer soils occur in lower lying, nearly level drainageways throughout the association.

The Dodge and Miami soils in this association are suited to all crops grown in the county. Both soils have high available water capacity and are medium in organic-matter content. Controlling water erosion is the main concern in managing these soils.

Much of this association is used for cultivated crops, but some areas, especially along streams, remain in native woods.

### 6. Muscatine-Sable-Catlin association

*Poorly drained to well-drained, nearly level to sloping soils that formed in silty material and the underlying loam glacial till; on uplands*

This association (fig. 6) consists of broad, gently undulating, irregularly shaped ridges and drainageways. Differences in elevation are commonly 5 to 10 feet. Gentle slopes are commonly long, but shorter, steeper




Figure 5.—Pattern of soils in association 5 and their relationship to parent material.

slopes are along some drainageways.

This association occupies less than 1 percent of the county. It is about 50 percent Muscatine soils, 25 percent Sable soils, 20 percent Catlin soils, and 5 percent minor soils.

Muscatine soils are nearly level and somewhat poorly drained. The surface layer typically is black silt loam about 14 inches thick. The subsoil is dark grayish-brown silty clay loam. The underlying material is silt loam.

Sable soils are nearly level and poorly drained. The surface layer typically is black and very dark gray silty clay loam about 17 inches thick. The subsoil is olive-gray silty clay loam. The underlying material is silt loam.

The Catlin soils, which occupy both the gently sloping rises and associated steeper areas, are well drained and moderately well drained and gently sloping to sloping. The surface layer typically is very dark brown and dark-brown silt loam. The subsoil is dark yellowish-brown to yellowish-brown silty clay loam and dark yellowish-brown clay loam. The underlying material is loam till below a depth of 65 inches.

Minor soils in this association are the Harpster and Drummer soils. The poorly drained Harpster soils occupy the rims of slight depressions that are within

areas of Sable soils. The poorly drained Drummer soils occupy the lower lying drainageways along streams.

The soils of this association are well suited to all cultivated crops commonly grown in the county. All the soils have high available water capacity and high organic-matter content. The main concerns in managing these soils are controlling water erosion in the sloping areas, improving drainage, and maintaining tilth and fertility.

Nearly all of this association is used for cultivated crops. It has a high potential for farming enterprises.

### *Descriptions of the Soils*

This section describes the soil series and mapping units in DeKalb County. Each soil series is described in considerable detail, and then, briefly, each mapping unit in that series is described. Unless it is specifically mentioned otherwise, it is to be assumed that what is stated about the soil series holds true for the mapping units in that series. Thus, to get full information about any one mapping unit, it is necessary to read both the description of the mapping unit and the description of the soil series to which it belongs.

An important part of the description of each soil




Figure 6.—Pattern of soils in association 6 and their relationship to parent material.

series is the soil profile, that is, the sequence of layers from the surface downward to rock or other underlying material. Each series contains two descriptions of this profile. The first is brief and in terms familiar to the layman. The second, detailed and in technical terms, is for scientists, engineers, and others who need to make thorough and precise studies of soils. Unless it is otherwise stated, the colors given in the descriptions are those of a dry soil.

As mentioned in the section "How This Survey Was Made," not all mapping units are members of a soil series. Cut and fill land, for example, does not belong to a soil series, but it is listed in alphabetic order along with the soil series.

Following the name of each mapping unit is a symbol in parentheses. This symbol identifies the mapping unit on the detailed soil map. Listed at the end of each description of a mapping unit is the management group in which the mapping unit has been placed. The page for the description of each management group can be found by referring to the "Guide to Mapping Units" at the back of this survey.

In some instances the mapping units on the detailed soil map do not join with those of La Salle County, because the mapping unit in one survey area is of minor extent and is correlated with a similar mapping unit and described as an inclusion.

The acreage and proportionate extent of each mapping unit are shown in table 3. Many of the terms used in describing soils can be found in the Glossary at the back of this survey, and more detailed information about the terminology and methods of soil mapping can be obtained from the Soil Survey Manual (7).

### Batavia Series

The Batavia series consists of deep, nearly level to sloping soils that are well drained and moderately well drained. These soils are in broad outwash areas or on unique circular, raised, flat-topped mounds. They formed in loess and in the underlying stratified fine sand and silt loam.

In a representative profile the surface layer is very dark gray silt loam about 8 inches thick. The subsurface layer is dark grayish-brown silt loam about 2 inches thick. The subsoil is about 45 inches thick. The upper 32 inches of the subsoil is brown to dark-brown, yellowish-brown, and light olive-gray silty clay loam, and the lower 13 inches is light olive-gray and yellowish-brown silt loam that has a noticeable amount of sand. The underlying material is light reddish-brown and yellowish-brown stratified loam.

These soils are moderately permeable and have high available water capacity. Internal drainage is medium,

TABLE 3.—Approximate acreage and proportionate extent of the soils

Soil	Area		Soil	Area	
	Acres	Percent		Acres	Percent
Batavia silt loam, 0 to 2 percent slopes	4,310	1.1	La Rose silt loam, 7 to 12 percent slopes, eroded	2,170	0.5
Batavia silt loam, 2 to 4 percent slopes	9,610	2.4	Limestone quarries	20	( <sup>1</sup> )
Batavia silt loam, 4 to 7 percent slopes, eroded	560	.1	Lisbon silt loam	980	.2
Bowes silt loam, 0 to 2 percent slopes	595	.1	Lorenzo loam, 6 to 15 percent slopes, eroded	125	( <sup>1</sup> )
Bowes silt loam, 2 to 4 percent slopes	615	.1	Miami silt loam, 2 to 4 percent slopes	2,580	.6
Camden silt loam, 0 to 2 percent slopes	2,015	.5	Miami silt loam, 4 to 7 percent slopes, eroded	3,245	.8
Camden silt loam, 2 to 6 percent slopes	2,335	.6	Miami silt loam, 7 to 12 percent slopes, eroded	360	.1
Catlin silt loam, 0 to 2 percent slopes	6,410	1.6	Millbrook silt loam	2,430	.6
Catlin silt loam, 2 to 4 percent slopes	37,660	9.2	Muscatine silt loam	1,110	.3
Catlin silt loam, 4 to 7 percent slopes, eroded	725	.2	Octagon silt loam, 2 to 4 percent slopes	8,560	2.1
Cut and fill land	425	.1	Octagon silt loam, 4 to 7 percent slopes, eroded	12,485	3.1
Dodge silt loam, 0 to 2 percent slopes	520	.1	Peotone silty clay loam	3,090	.8
Dodge silt loam, 2 to 4 percent slopes	8,325	2.0	Plano silt loam, 0 to 2 percent slopes	2,320	.6
Dodge silt loam, 4 to 7 percent slopes, eroded	705	.2	Plano silt loam, 2 to 4 percent slopes	1,090	.3
Dresden silt loam, 2 to 4 percent slopes	400	.1	Proctor silt loam, 0 to 2 percent slopes	835	.2
Dresden silt loam, 4 to 7 percent slopes, eroded	480	.1	Proctor silt loam, 2 to 4 percent slopes	330	.1
Drummer silty clay loam	117,415	28.8	Rush silt loam, 0 to 2 percent slopes	500	.1
Elburn silt loam	16,590	4.1	Rush silt loam, 2 to 4 percent slopes	320	.1
Flanagan silt loam	48,265	11.9	Sable silty clay loam	440	.1
Gravel pits	410	.1	St. Charles silt loam, 0 to 2 percent slopes	1,135	.3
Harpster silty clay loam	5,740	1.4	St. Charles silt loam, 2 to 4 percent slopes	195	( <sup>1</sup> )
Harvard silt loam, 0 to 2 percent slopes	2,055	.5	Sawmill soils	9,615	2.4
Harvard silt loam, 2 to 4 percent slopes	4,495	1.1	Saybrook silt loam, 2 to 4 percent slopes	62,015	15.2
Harvard silt loam, 4 to 7 percent slopes, eroded	1,200	.3	Saybrook silt loam, 4 to 7 percent slopes, eroded	5,725	1.4
Herbert silt loam	3,295	.8	Strawn soils, 10 to 20 percent slopes, eroded	275	.1
Houghton muck	1,235	.3	Virgil silt loam	4,180	1.0
Kendall silt loam	700	.2	Will clay loam	610	.1
Knight silt loam	355	.1	Sewage lagoons	10	( <sup>1</sup> )
La Rose silt loam, 2 to 4 percent slopes	240	.1	Water <sup>2</sup>	510	.1
La Rose silt loam, 4 to 7 percent slopes, eroded	2,095	.5	Total	407,040	100.0

<sup>1</sup> Less than 0.05 percent.

<sup>2</sup> Includes all bodies of water.

and the soils remain stable with changes in moisture content. The water table is more than 5 feet below the surface.

These soils are well suited to intensive use for row crops, and they have few limitations for other uses.

Representative profile of Batavia silt loam, 2 to 4 percent slopes, 660 feet north and 156 feet east of the SW. corner of NW $\frac{1}{4}$  sec. 26, T. 40 N., R. 3 E.

- Ap—0 to 8 inches, very dark gray (10YR 3/1) silt loam; moderate, fine and very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A2—8 to 10 inches, dark grayish-brown (10YR 4/2) silt loam; weak, medium, platy structure; friable; neutral; abrupt, smooth boundary.
- B1—10 to 18 inches, brown to dark-brown (10YR 4/3) silty clay loam; moderate, very fine, subangular blocky structure; firm; slightly acid; clear, smooth boundary.
- B21t—18 to 23 inches, brown to dark-brown (10YR 4/3) silty clay loam; moderate, fine, subangular blocky structure; firm; medium acid; clear, smooth boundary.
- B22t—23 to 33 inches, yellowish-brown (10YR 5/4) silty clay loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; moderate, medium, subangular blocky structure; friable; slightly acid; gradual, smooth boundary.
- B23t—33 to 42 inches, mixed yellowish-brown (10YR 5/6 and 5/8) and light olive-gray (5Y 6/2) light silty clay loam; weak, coarse, prismatic structure; friable; neutral; abrupt, smooth boundary.
- IIB3—42 to 55 inches, mixed light olive-gray (5Y 6/2) and yellowish-brown (10YR 5/6 and 5/8) silt loam that has a noticeable amount of sand; weak, coarse, prismatic structure; friable; mildly alkaline; abrupt, smooth boundary.
- IIC—55 to 60 inches, light reddish-brown (5YR 6/3) and yellowish-brown (10YR 5/8) loam that has thin lenses or strata of sand; massive; friable; moderately alkaline.

The A horizon ranges from 6 to 10 inches in thickness. The B horizon ranges from 20 to 50 inches in thickness and from medium acid to mildly alkaline. The IIB horizon is 5 to 15 inches thick. It is silt loam that has a noticeable amount of sand to clay loam. The C horizon is stratified sandy loam, silt loam, or loam, but in places it is sand or gravel.

Batavia soils are associated with Virgil and Harvard soils. They are better drained than Virgil soils and have a thicker layer of silty material over the stratified outwash material than Harvard soils.

**Batavia silt loam, 0 to 2-percent slopes (105A).**—This soil is in high outwash areas or on raised circular mounds. Included with this soil in mapping were small areas of Virgil silt loam and areas that have a layer of silt more than 5 feet thick.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. It has few limitations for most other uses. Management group I-2.

**Batavia silt loam, 2 to 4 percent slopes (105B).**—This soil is in higher areas on outwash plains or on the rims of raised circular mounds. It has the profile described as representative of the series.

Included with this soil in mapping were small areas of Virgil silt loam. Also included were small areas that are calcareous at depths of less than 40 inches and areas that have a layer of silt more than 5 feet thick.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has few limitations for most other uses. Management group IIe-2.

**Batavia silt loam, 4 to 7 percent slopes, eroded (105C2).**—This soil is on rims and edges of raised circular mounds. It has a profile similar to the one described as representative of the series, but some of the brown to dark-brown, finer textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas of Harvard silt loam and small areas where none of the subsoil has been mixed into the surface layer.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses because of slope. Management group IIe-4.

## Bowes Series

The Bowes series consists of deep, nearly level to gently sloping, well-drained soils. These soils are in upland areas near the Kishwaukee River. They formed in silty material and in the underlying sand and gravel.

In a representative profile the surface layer is very dark grayish-brown silt loam about 7 inches thick. The subsoil is about 40 inches thick. The upper 23 inches of the subsoil is dark yellowish-brown silty clay loam; the next 10 inches is brown to dark-brown, mainly silty clay loam; and the lower 7 inches is dark reddish-brown gravelly sandy clay loam. The underlying material is yellowish-brown gravelly sand.

These soils are moderately permeable and have high available water capacity. Internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is more than 5 feet below the surface.

These soils are well suited to intensive use for row crops, and they have few limitations for other uses.

Representative profile of Bowes silt loam, 0 to 2 percent slopes, 51 feet north of railroad fence and 357 feet east of center of road in NW $\frac{1}{4}$ SE $\frac{1}{4}$  sec. 23, T. 42 N., R. 4 E.

- Ap—0 to 7 inches, very dark grayish-brown (10YR 3/2) silt loam; weak, very fine and fine, granular structure; friable; neutral; abrupt, smooth boundary.
- B1—7 to 11 inches, dark yellowish-brown (10YR 4/4) light silty clay loam; weak, very fine, subangular blocky structure parting to weak, very fine and fine, granular; friable; medium acid; clear, smooth boundary.
- B21t—11 to 18 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, very fine, subangular blocky structure; firm; thin, discontinuous, dark-brown (10YR 3/3) clay films; medium acid; clear, smooth boundary.
- B22t—18 to 24 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate to strong, very fine and fine, angular blocky structure; firm; thin, continuous, dark-brown (10YR 3/3) clay films; strongly acid; clear, smooth boundary.
- B23t—24 to 30 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, fine and medium, angular blocky structure; firm; thin, continuous,

- dark-brown (10YR 3/3) clay films; medium acid; clear, smooth boundary.
- B24t—30 to 37 inches, brown to dark-brown (10YR 4/3) silty clay loam; weak, medium, subangular blocky structure; friable; slightly acid; clear, smooth boundary.
- IIB31t—37 to 40 inches, brown to dark-brown (7.5YR 4/4) loam; weak, coarse, subangular blocky structure; firm; slightly acid; abrupt, smooth boundary.
- IIB32t—40 to 47 inches, dark reddish-brown (5YR 3/4) gravelly sandy clay loam; massive; firm; neutral; abrupt, smooth boundary.
- IIC—47 to 60 inches, yellowish-brown (10YR 5/4) gravelly sand; single grained; loose; mildly alkaline.

The A horizon ranges from 6 to 10 inches in thickness. The B horizon is 30 to 53 inches thick and is strongly acid to neutral. The IIB horizon ranges from 4 to 12 inches in thickness and from clay loam to loam or gravelly clay loam or sandy clay loam in texture. The C horizon ranges from sand to gravelly sand.

Bowes soils are associated with Rush and Dresden soils. They have a darker colored A horizon than Rush soils and a thicker silty layer over sand and gravel than Dresden soils.

**Bowes silt loam, 0 to 2 percent slopes (792A).**—This soil is on broad outwash flats along the Kishwaukee River. It has the profile described as representative of the series. Included with this soil in mapping were small areas of somewhat poorly drained soils and areas that have a thicker, darker colored surface layer.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Some areas are used for gravel pits. This soil has few limitations for most other uses. Management group I-2.

**Bowes silt loam, 2 to 4 percent slopes (792B).**—This soil is within or adjacent to areas of the nearly level Bowes soils. It has a profile similar to the one described as representative of the series, but the surface layer is thinner and lighter in color. Included with this soil in mapping were small areas of Dresden silt loam and areas that have a thicker, darker colored surface layer.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Some areas are used for gravel pits. Control of erosion is a necessary part of good management. This soil has few limitations for most other uses. Management group IIe-2.

## Camden Series

The Camden series consists of deep, nearly level to sloping, well drained and moderately well drained soils. These soils are on uplands near the major streams. They formed in silty material and in the underlying stratified loamy outwash. In some areas sand and gravel are below a depth of 5 feet.

In a representative profile the surface layer is very dark grayish-brown silt loam about 5 inches thick. The subsurface layer is dark grayish-brown silt loam about 3 inches thick. The subsoil is about 47 inches thick. The upper 21 inches of the subsoil is brown to dark-brown silty clay loam; the next 17 inches is olive-gray and strong-brown clay loam and silt loam that has a noticeable amount of sand; and the lower 9 inches is yellowish-brown and olive-gray clay loam.

These soils are moderately permeable and have high

available water capacity. Internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are well suited to row crops, and they have few limitations for most uses.

Representative profile of Camden silt loam, 0 to 2 percent slopes, 220 feet east and 260 feet south of the center of NE $\frac{1}{4}$  sec. 25, T. 41 N., R. 4 E.

- Ap—0 to 5 inches, very dark grayish-brown (10YR 3/2) silt loam; weak, very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A2—5 to 8 inches, dark grayish-brown (10YR 4/2) silt loam; weak, thick, platy structure; friable; neutral; abrupt, smooth boundary.
- B21t—8 to 13 inches, brown to dark-brown (10YR 4/3) silty clay loam; moderate, very fine and fine, angular blocky structure; firm; slightly acid; clear, smooth boundary.
- B22t—13 to 21 inches, brown to dark-brown (10YR 4/3) silty clay loam; moderate, fine, angular blocky structure; firm; medium acid; clear, smooth boundary.
- B23t—21 to 29 inches, brown to dark-brown (10YR 4/3) silty clay loam; common, fine, distinct, strong-brown (7.5YR 5/8) mottles; moderate, medium, subangular blocky structure; firm; grayish-brown (10YR 5/2) silt coatings; medium acid; clear, smooth boundary.
- IIB24t—29 to 42 inches, mixed strong-brown (7.5YR 5/8) and olive-gray (5Y 5/2) clay loam; weak, coarse, prismatic structure; friable; discontinuous dark-gray (10YR 4/1) clay films; neutral; abrupt, smooth boundary.
- IIB31—42 to 46 inches, mixed strong-brown (7.5YR 5/8) and olive-gray (5Y 5/2) silt loam that has a noticeable amount of sand; weak, coarse, prismatic structure; friable; discontinuous dark-gray (10YR 4/1) clay films; neutral; abrupt, smooth boundary.
- IIB32—46 to 55 inches, yellowish-brown (10YR 5/6) and olive-gray (5Y 5/2) clay loam; massive; friable; some dark-gray (10YR 4/1) clay films; neutral.

The A horizon ranges from 6 to 12 inches in thickness. The A2 horizon in most places is mixed with the A1 horizon because of plowing. The B horizon ranges from 20 to 50 inches in thickness and from medium acid to neutral. The IIB horizon is 10 to 30 inches thick. It is clay loam, silt loam that has a noticeable amount of sand, loam, or sandy loam in texture. In some places mottles are in the lower part of the B horizon.

Camden soils are associated with Rush, Dresden, and Millbrook soils. Camden soils formed in medium-textured outwash. Rush and Dresden soils have sand and gravel at a depth of less than 5 feet. Camden soils are better drained than Millbrook soils.

**Camden silt loam, 0 to 2 percent slopes (134A).**—This soil is in broad outwash areas or on terraces along the major streams. It has the profile described as representative of the series.

Included with this soil in mapping were small areas of somewhat poorly drained soils and areas of soils that are calcareous at a depth of less than 40 inches. Also included were small areas where sand or gravel is below a depth of 5 feet.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. It has few limitations for most other uses. Management group I-1.

**Camden silt loam, 2 to 6 percent slopes (134B).**—This soil is in outwash areas or on terraces along the major streams. It has a profile similar to the one described as

representative of the series, but the surface layer is thinner and lighter in color.

Included with this soil in mapping were small areas where the silty material is thicker than 40 inches and where the finer textured subsoil is mixed into the plow layer. Also included were small areas where sand or gravel is below a depth of 5 feet.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has few limitations for most other uses. Management group Iie-1.

### Catlin Series

The Catlin series consists of deep, nearly level to sloping soils that are well drained and moderately well drained. These soils are in higher positions on the landscape on uplands. They formed in loess and in the underlying loam glacial till.

In a representative profile the surface layer is very dark brown and dark-brown silt loam about 11 inches thick. The subsoil is about 54 inches thick. The upper 32 inches of the subsoil is brown to dark-brown, yellowish-brown, and dark yellowish-brown silty clay loam. The lower 22 inches is dark yellowish-brown, brown to dark-brown, and strong-brown clay loam. The underlying material is light yellowish-brown loam.

These soils are moderately permeable and have high available water capacity. Internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are well suited to row crops, and they have few limitations for many other uses.

Representative profile of Catlin silt loam, 2 to 4 percent slopes, 330 feet south and 70 feet west of the NE corner of SE $\frac{1}{4}$  sec. 14, T. 38 N., R. 4 E.

- Ap—0 to 8 inches, very dark brown (10YR 2/2) silt loam; weak, very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A3—8 to 11 inches, dark-brown (10YR 3/3) silt loam; weak, very fine, subangular blocky structure parting to weak, very fine, granular; friable; some patchy, very dark brown (10YR 2/2) coatings; neutral; clear, smooth boundary.
- B1—11 to 18 inches, brown to dark-brown (10YR 4/3) silty clay loam; moderate, very fine, subangular blocky structure; firm; some dark-brown (10YR 3/3) clay films; slightly acid; gradual, smooth boundary.
- B21t—18 to 29 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, fine, subangular blocky structure; firm; some dark-brown (10YR 3/3) clay films; slightly acid; gradual, smooth boundary.
- B22t—29 to 43 inches, yellowish-brown (10YR 5/4) silty clay loam; common, medium, distinct, strong-brown (7.5YR 5/6) mottles; moderate, medium, subangular blocky structure; firm; discontinuous dark yellowish-brown (10YR 4/4) clay films; medium acid; abrupt, smooth boundary.
- IIB23t—43 to 50 inches, dark yellowish-brown (10YR 4/4) clay loam; few, fine, distinct, strong brown (7.5YR 5/6) mottles; weak, medium, prismatic structure; firm; discontinuous brown to dark-brown (7.5YR 4/2) clay films; neutral; clear, smooth boundary.
- IIB24—50 to 65 inches, mixed brown to dark-brown (7.5YR 4/4) and strong-brown (7.5YR 5/6 and 5/8) clay loam; weak, medium, prismatic structure;

friable; some brown to dark-brown (7.5YR 4/2) clay films; mildly alkaline; abrupt, smooth boundary.

IIC—65 to 70 inches, light yellowish-brown (10YR 6/4) loam till; massive; friable; moderately alkaline.

The A horizon ranges from 9 to 14 inches in thickness. The B horizon ranges from 30 to 55 inches in thickness. The IIB horizon is 10 to 25 inches thick. The B and IIB horizons range from medium acid to mildly alkaline.

Catlin soils are associated with Flanagan and Saybrook soils. They are better drained than Flanagan soils and have a thicker mantle of loess than Saybrook soils.

**Catlin silt loam, 0 to 2 percent slopes (171A).**—This soil is on uplands. Included with it in mapping were areas where the surface layer is less than 10 inches thick and small areas of somewhat poorly drained soils. Also included were areas where a layer of stratified sand and silt is between the loess and the underlying glacial till and small areas where more than 5 feet of loess is over the till.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. It has few limitations for most other uses. Management group I-2.

**Catlin silt loam, 2 to 4 percent slopes (171B).**—This soil is on uplands. It has the profile described as representative of the series. Included with this soil in mapping were areas where the surface layer is less than 10 inches thick and small areas of somewhat poorly drained soils. Also included were areas where stratified sand and silt is between the loess and the underlying glacial till and small areas where more than 5 feet of loess is over the till.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has few limitations for most other uses. Management group Iie-2.

**Catlin silt loam, 4 to 7 percent slopes, eroded (171C2).**—This soil is in narrow areas downslope from gently sloping Catlin soils. It has a profile similar to the one described as representative of the series, but some of the brown to dark-brown, finer textured subsoil has been mixed into the plow layer.

Included with this soil in mapping were small areas in which all of the original surface layer has been removed through erosion, and small areas that have not been eroded. Also included were small areas of La Rose soils.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses because of slope. Management group Iie-4.

### Cut and Fill Land

Cut and fill land consists of areas from which soil material has been removed. The amount removed ranges from only the surface layer to as much as several feet of soil material. This land also consists of

low areas that have been covered with 1 to several feet of soil material. In many places the cuts and fills have been made during land leveling. Areas of this land are shown on the detailed soil map by an appropriate symbol.

The soil material is generally low in fertility and organic-matter content. Calcareous, compact glacial till is exposed in most places. This till, either in place or as fill material, supports little vegetation, and establishing a plant cover is difficult. Topdressing, special fertility practices, and other special management practices are necessary. Not placed in a management group.

### Dodge Series

The Dodge series consists of deep, nearly level to sloping, well drained and moderately well drained soils. These soils are on uplands near the major streams. They formed in loess and in the underlying loam glacial till.

In a representative profile the surface layer is brown to dark-brown silt loam about 6 inches thick. The sub-surface layer is dark grayish-brown silt loam about 3 inches thick. The subsoil is about 22 inches thick. The upper 16 inches of the subsoil is dark yellowish-brown, yellowish-brown, and pale-brown silty clay loam. The lower 6 inches is brown to dark-brown clay loam. The underlying material is brown to dark-brown loam.

These soils are moderately permeable and have high available water capacity. Internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are well suited to row crops, and they have moderate limitations for some other uses.

Representative profile of Dodge silt loam, 2 to 4 percent slopes, 40 feet south and 150 feet east of the NE corner of SE $\frac{1}{4}$ SW $\frac{1}{4}$  sec. 18, T. 41 N., R. 5 E.

- Ap—0 to 6 inches, brown to dark-brown (10YR 4/3) silt loam; weak, fine, granular structure; very friable; neutral; abrupt, smooth boundary.
- A2—6 to 9 inches, dark grayish-brown (10YR 4/2) silt loam; weak, thick, platy structure parting to very fine, subangular blocky; friable; neutral; abrupt, smooth boundary.
- B1—9 to 13 inches, dark yellowish-brown (10YR 4/4) light silty clay loam; moderate, very fine, subangular blocky structure; friable; strongly acid; abrupt, smooth boundary.
- B21t—13 to 19 inches, yellowish-brown (10YR 5/4) silty clay loam; weak, fine and medium, subangular blocky structure parting to weak, fine, angular blocky; firm; brown to dark-brown (10YR 4/3) clay films; strongly acid; abrupt, smooth boundary.
- B22t—19 to 25 inches, pale-brown (10YR 6/3) silty clay loam; weak, medium, prismatic structure; firm; very dark grayish-brown (10YR 3/2) patchy clay films; few concretions; slightly acid; abrupt, smooth boundary.
- IIB23t—25 to 31 inches, brown to dark-brown (7.5YR 4/4) clay loam; weak, medium, prismatic structure; firm; very dark grayish-brown (10YR 3/2) patchy clay films; few concretions; slightly acid; abrupt, smooth boundary.
- IIC—31 to 60 inches, brown to dark-brown (7.5YR 4/4) and brown (7.5YR 5/4) loam; few, fine, faint, yellowish-brown (10YR 5/8) mottles; massive; friable; black (10YR 2/1) channel fillings to a depth of 45 inches; moderately alkaline.

The A horizon ranges from 7 to 12 inches in thickness. In many places the A2 horizon has been mixed into the plow layer. The B horizon is 15 to 30 inches thick and is strongly acid to neutral. The IIB horizon ranges from 4 to 10 inches in thickness. In some places mottles are in the lower part of the B horizon.

Dodge soils are associated with Octagon and Miami soils. They have a lighter colored A horizon than Octagon soils and a thicker mantle of loess than Octagon and Miami soils.

**Dodge silt loam, 0 to 2 percent slopes (24A).**—This soil is on uplands above the gently sloping Dodge soils. Included with this soil in mapping were small areas of somewhat poorly drained soils and some areas where the silty material is more than 40 inches thick.

Surface runoff is slow, and the hazard of erosion is slight.

Most areas of this soil are intensively farmed and are well suited to row crops, but some areas remain in native woods. This soil has moderate limitations for some other uses. Management group I-1.

**Dodge silt loam, 2 to 4 percent slopes (24B).**—This soil is on uplands along the major streams and is generally associated with Miami soils. It has the profile described as representative of the series. Included with this soil in mapping were some areas where the silty material is more than 40 inches thick.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops, but some areas remain in native woods. This soil has moderate limitations for some other uses. Management group IIe-1.

**Dodge silt loam, 4 to 7 percent slopes, eroded (24C2).**—This soil is on uplands near the other Dodge soils. It has a profile similar to the one described as representative of the series, but the dark yellowish-brown, finer textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas of Miami soils and some areas where none of the subsoil has been mixed into the surface layer.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIe-4.

### Dresden Series

The Dresden series consists of gently sloping to sloping, well-drained soils that are moderately deep over sand and gravel. These soils are in outwash areas along the drainageways. They formed in loamy material and in the underlying sand and gravel.

In a representative profile the surface layer is very dark grayish-brown and dark-brown silt loam about 7 inches thick. The subsoil is about 31 inches thick. The upper 7 inches of the subsoil is dark yellowish-brown silty clay loam; the next 15 inches is yellowish-brown clay loam; and the lower 9 inches is brown to dark-brown gravelly clay loam. The underlying material is mixed brown to dark-brown and yellowish-brown and light yellowish-brown gravelly sand.

These soils are moderately permeable and have moderate available water capacity. Internal drainage

is rapid, and the soils remain stable with changes in moisture content. The water table is more than 5 feet below the surface.

These soils are suited to row crops. They have few limitations for most other uses. Because of the shallowness to sand and gravel, pollution is a risk for some uses.

Representative profile of Dresden silt loam, 2 to 4 percent slopes, 348 feet east and 60 feet north of the SW. corner of SW $\frac{1}{4}$  sec. 33, T. 40 N., R. 5 E.

- Ap—0 to 7 inches, very dark grayish-brown (10YR 3/2) and dark-brown (10YR 3/3) silt loam; moderate, fine and very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- B1t—7 to 14 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, medium, subangular blocky structure parting to weak, fine, granular; friable; dark-brown (10YR 3/3) clay films; neutral; gradual, smooth boundary.
- B2t—14 to 29 inches, yellowish-brown (10YR 5/4) clay loam; moderate, medium, subangular blocky structure; friable; dark yellowish-brown (10YR 3/4) clay films; slightly acid; gradual, smooth boundary.
- IIB3—29 to 38 inches, brown to dark-brown (7.5YR 4/4) gravelly clay loam; weak, coarse, subangular blocky structure, but tending toward single grained; friable; slightly acid; clear, smooth boundary.
- IIC—38 to 60 inches, mixed brown to dark-brown (7.5YR 4/4), yellowish-brown (10YR 5/4), and light yellowish-brown (10YR 6/4) gravelly sand; single grained; loose; moderately alkaline.

The A horizon ranges from 6 to 10 inches in thickness. The B horizon is 15 to 30 inches thick and is medium acid to neutral. The IIB horizon ranges from 4 to 10 inches in thickness and from clay loam to gravelly clay loam in texture.

Dresden soils are associated with Rush and Bowes soils. They have a darker colored A horizon than Rush soils and thinner silty material over sand and gravel than Rush and Bowes soils.

**Dresden silt loam, 2 to 4 percent slopes (325B).**—This soil is in outwash areas or on terraces along the major streams or drainageways. It has the profile described as representative of the series. Included with this soil in mapping were small areas that have a thicker, darker colored surface layer and small areas that have a thinner subsoil over sand and gravel.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is suited to row crops. Control of erosion is a necessary part of good management. Some areas are used for gravel pits. This soil has few limitations for most other uses. Management group IIE-3.

**Dresden silt loam, 4 to 7 percent slopes, eroded (325C2).**—This soil is in outwash areas or on terraces along the major streams. It has a profile similar to the one described as representative of the series, but the dark yellowish-brown, finer textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas that have a thinner subsoil over sand and gravel and small areas in which none of the original surface layer remains.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses because of slope. Management group IIIe-2.

## Drummer Series

The Drummer series consists of deep, nearly level, poorly drained soils. These soils are in drainageways and in broad, flat areas on uplands. They formed in silty material and in the underlying stratified loamy material.

In a representative profile the surface layer is black and very dark gray silty clay loam about 18 inches thick. The subsoil, about 32 inches thick, is olive-gray and gray to light-gray silty clay loam mottled with strong brown. The underlying material is gray and strong-brown sandy clay loam and silt loam.

These soils are moderately permeable and have high available water capacity. Surface drainage and internal drainage are very slow, and when wet, the soils have low stability. The soils are subject to frequent waterlogging and flooding or ponding during periods of runoff (fig. 7). Seasonally the water table is at a depth of less than 2 feet.

These soils are well suited to row crops if they are adequately tile drained. Frequent waterlogging and potential for flooding or ponding are severe limitations for most other uses.

Representative profile of Drummer silty clay loam, 480 feet south and 75 feet west of the NE. corner of SE $\frac{1}{4}$  sec. 24, T. 39 N., R. 3 E.

- A11—0 to 11 inches, black (10YR 2/1) silty clay loam; moderate, fine and very fine, granular structure; friable; neutral; gradual, smooth boundary.
- A12—11 to 18 inches, black (10YR 2/1) and very dark gray (10YR 3/1) silty clay loam; weak, fine, subangular blocky structure parting to moderate, fine, granular; friable; neutral; gradual, smooth boundary.
- B21g—18 to 29 inches, olive-gray (5Y 5/2) silty clay loam; few, fine, faint, strong-brown (7.5YR 5/6) mottles; moderate, fine, prismatic structure parting to moderate, fine, subangular blocky; firm; neutral; gradual, smooth boundary.
- B22g—29 to 50 inches, gray to light-gray (5Y 6/1) and strong-brown (7.5YR 5/6) silty clay loam; moderate, medium, prismatic structure; firm; neutral; abrupt, smooth boundary.
- IIC1g—50 to 60 inches, strong-brown (7.5YR 5/8) and gray to light-gray (5Y 6/1) sandy clay loam; massive; friable; mildly alkaline; abrupt, smooth boundary.
- IIC2g—60 to 78 inches, gray to light-gray (5Y 6/1) and some pinkish-gray (5YR 6/2), brown to dark-brown (7.5YR 4/4), and gray to light-gray (7.5YR 6/0) silt loam; massive; friable; moderately alkaline.

The A horizon ranges from 8 to 20 inches in thickness. The B horizon ranges from 15 to 40 inches in thickness and from silty clay loam to clay loam in texture in the lower part. It is neutral to mildly alkaline. The C horizon ranges from loam or silt loam to sandy clay loam.

Drummer soils are associated with Peotone and Flanagan soils. They have a thinner A horizon than Peotone soils and are more poorly drained than Flanagan soils.

**Drummer silty clay loam (152).**—This nearly level soil is in low areas of large upland flats and of drainageways. Included with this soil in mapping were small areas that have sand and gravel below a depth of 5 feet, small areas of poorly drained soils in depressions, and small knobs of better drained soils. Also included were small spots of Harpster, Peotone, and Houghton soils, most of which are shown on the maps by symbols.

Surface runoff is very slow to ponded, and the hazard of erosion is slight.

If this soil is artificially drained, it is intensively



Figure 7.—An area of Drummer soils after heavy rains.

farmed and is well suited to row crops. It has severe limitations for most other uses. Management group IIw-1.

### Elburn Series

The Elburn series consists of deep, nearly level, somewhat poorly drained soils in broad outwash areas or on unique, circular, raised, flat-topped mounds. These soils formed in silty material and in the underlying stratified loamy material.

In a representative profile the surface layer is very dark brown to very dark grayish-brown silt loam and silty clay loam about 14 inches thick. The subsoil is about 46 inches thick. The upper 34 inches is dark grayish-brown to yellowish-brown and light brownish-gray silty clay loam mottled with yellowish brown, and the lower 12 inches is yellowish-brown and strong-brown light clay loam.

These soils are moderately permeable and have high available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high runoff, and they are also subject to periodic waterlogging. Stability decreases when the soils are wet. In spring the water table is generally 1 foot to 3 feet below the surface.

These soils are well suited to intensive use for row crops. The periodic waterlogging is a limitation for most other uses.

Representative profile of Elburn silt loam, 300 feet south of center of road and 185 feet west of the NE. corner of NW $\frac{1}{4}$ SW $\frac{1}{4}$ . sec. 36, T. 37 N., R. 3 E .

- Ap—0 to 10 inches, very dark brown (10YR 2/2) silt loam; moderate, fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A3—10 to 14 inches, very dark grayish-brown (10YR 3/2) light silty clay loam; weak, very fine, subangular blocky structure; friable; slightly acid; clear, smooth boundary.
- B21t—14 to 21 inches, dark grayish-brown (10YR 4/2) silty clay loam; few, fine, faint, yellowish-brown (10YR 5/8) mottles; medium, very fine and fine, subangular blocky structure; firm; slightly acid; gradual, smooth boundary.
- B22t—21 to 30 inches, yellowish-brown (10YR 5/4) silty clay loam; few, fine, faint, yellowish-brown (10YR 5/8) mottles; moderate, fine and medium, subangular blocky structure; firm; discontinuous dark grayish-brown (10YR 4/2) clay films; slightly acid; gradual, smooth boundary.
- B23t—30 to 38 inches, yellowish-brown (10YR 5/8) and light brownish-gray (2.5Y 6/2) light silty clay loam; moderate, coarse, subangular blocky structure; friable; patchy dark grayish-brown (10YR 4/2) clay films; neutral; clear, smooth boundary.
- B24t—38 to 48 inches, mixed yellowish-brown (10YR 5/6 and 5/8) and light brownish-gray (2.5Y 6/2) light silty clay loam; weak, medium, prismatic structure; friable; patchy dark yellowish-brown (10YR 4/4) clay films; neutral; abrupt, smooth boundary.
- IIB3—48 to 60 inches, yellowish-brown (10YR 5/4) and strong-brown (7.5YR 5/6) light clay loam; weak, coarse, prismatic structure; friable; mildly alkaline.

The A horizon ranges from 12 to 20 inches in thickness. The B horizon ranges from 30 to 50 inches in thickness and from slightly acid to mildly alkaline. The IIB horizon ranges from 5 to 15 inches in thickness and is clay loam, loam, or sandy clay loam in texture.

Elburn soils are associated with Batavia, Plano, and Drummer soils. They have a thicker A horizon than Batavia soils. They are more poorly drained than Batavia and Plano soils, but they are better drained than Drummer soils.

**Elburn silt loam** (198).—This nearly level soil is in outwash areas or in the depressed centers of large, raised, circular mounds. Included with this soil in mapping were areas where glacial till is below a thin outwash layer. Also included were areas that have more than 5 feet of silt and small areas of poorly drained soils.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Artificial drainage is beneficial in some areas. This soil has limitations for most other uses. Management group I-3.

### Flanagan Series

The Flanagan series consists of deep, nearly level, somewhat poorly drained soils on upland flats and near drainageways. These soils formed in loess and in the underlying loam glacial till.

In a representative profile the surface is black silt loam in the upper 13 inches and very dark grayish brown light silty clay loam in the lower 3 inches. The subsoil is about 29 inches thick. The upper 19 inches of the subsoil is brown to dark-brown, yellowish-brown, and light olive-brown silty clay loam mottled with yellowish brown and grayish brown. The lower 10 inches is mostly grayish-brown, yellowish-brown, and light olive-brown heavy silt loam. The underlying material is brown, strong-brown, light brownish-gray, and yellowish-brown loam.

These soils are moderately permeable and have high available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high runoff, and they are also subject to periodic waterlogging. Stability decreases when the soils are wet. In spring the water table is generally 1 foot to 3 feet below the surface.

These soils are well suited to intensive use for row crops. The periodic waterlogging is a limitation for many other uses.

Representative profile of Flanagan silt loam, 804 feet south and 384 feet west of center of sec. 23, T. 39 N., R. 3 E.

- A1—0 to 13 inches, black (10YR 2/1) silt loam; weak, very fine, granular structure; friable; neutral; clear, smooth boundary.
- A3—13 to 16 inches, very dark grayish-brown (10YR 3/2) light silty clay loam; weak, very fine, subangular blocky structure parting to weak, very fine, granular; friable; black (10YR 2/1) coatings; slightly acid; clear, smooth boundary.
- B21t—16 to 21 inches, brown to dark-brown (10YR 4/3) silty clay loam; moderate, fine, subangular blocky structure; friable; continuous dark grayish-brown (10YR 4/2) clay films; medium acid; clear, smooth boundary.
- B22t—21 to 28 inches, yellowish-brown (10YR 5/4) silty clay loam; few, fine, distinct, yellowish-brown (10YR 5/6) mottles and few, fine, faint, grayish-brown (10YR 5/2) mottles; strong, medium, subangular blocky structure; friable; continuous dark grayish-brown (2.5Y 4/2) clay films; medium acid; gradual, smooth boundary.
- B23t—28 to 35 inches, light olive-brown (2.5Y 5/4) with some very dark gray (10YR 3/1) light silty clay loam; common, fine, distinct, yellowish-brown (10YR 5/6) mottles; weak, coarse, prismatic

structure parting to moderate, medium, subangular blocky; friable; continuous dark grayish-brown (10YR 4/2) clay films; slightly acid; clear, smooth boundary.

B31—35 to 42 inches, grayish-brown (10YR 5/2), yellowish-brown (10YR 5/6), and light olive-brown (2.5Y 5/4) heavy silt loam; weak, coarse, blocky structure; friable; very dark gray (10YR 3/1) root channel fillings; neutral; clear, smooth boundary.

IIB32—42 to 45 inches, mixed brown (7.5YR 5/4), strong-brown (7.5YR 5/6), light brownish-gray (10YR 6/2), and yellowish-brown (10YR 5/6) loam; weak, coarse, blocky structure; friable; neutral; abrupt, smooth boundary.

IIC—45 to 60 inches, mixed brown (7.5YR 5/4), strong-brown (7.5YR 5/6), light brownish-gray (10YR 6/2), and yellowish-brown (10YR 5/6) loam; massive; friable; moderately alkaline.

The A horizon ranges from 12 to 18 inches in thickness. The B horizon ranges from 24 to 40 inches in thickness and from medium acid to neutral. The IIB horizon ranges from 3 to 10 inches in thickness and from loam to clay loam in texture. Reaction of the B horizon ranges from medium acid to neutral.

Flanagan soils are associated with Catlin and Drummer soils. They are more poorly drained than Catlin soils and better drained than Drummer soils.

**Flanagan silt loam** (154).—This nearly level soil is on uplands at the upper ends of drainageways or down-slope from gently sloping, well drained and moderately well drained soils. Included with this soil in mapping were areas where a layer of stratified sand and silt is between the loess and the underlying glacial till and small areas where more than 5 feet of loess is over the till. Also included were areas that have a lighter colored surface layer that is less than 10 inches thick.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Artificial drainage is beneficial in some areas. This soil has moderate to severe limitations for many other uses. Management group I-3.

### Gravel Pits

Gravel pits consists of areas from which sand and gravel have been excavated or mined. Most of the pits are in the valley of the Kishwaukee River and are mined occasionally or regularly. Some are filled with water. Not placed in a management group.

### Harpster Series

The Harpster series consists of deep, nearly level, poorly drained soils. These soils are generally near depressions on uplands. They formed in silty material and in the underlying stratified loamy material.

In a representative profile the surface layer is black and very dark gray silty clay loam about 12 inches thick. The subsoil is about 22 inches thick. The upper 18 inches is dark-gray, gray, and olive-gray silty clay loam mottled with yellowish brown, and the lower 4 inches is olive-gray and yellowish-brown silt loam. The underlying material is olive-gray, light-brown, and light yellowish-brown loam mottled with yellowish brown.

These soils are moderately permeable and have high available water capacity. Surface drainage and internal drainage are very slow. The soils are subject to con-

centrations of water during periods of runoff, and they are also subject to frequent or continuous waterlogging. They have low stability when wet. In spring the water table is generally less than 2 feet below the surface.

These soils are well suited to row crops. Because the content of carbonates is high, practices that increase the fertility of the soil are needed. The frequent waterlogging and potential for flooding or ponding are severe limitations for most other uses.

Representative profile of Harpster silty clay loam, 600 feet west of center of old road and 192 feet south of center of baseline road, in the NE $\frac{1}{4}$ NW $\frac{1}{4}$ NE $\frac{1}{4}$  sec. 2, T. 41 N., R. 5 E.

- Apca—0 to 8 inches, black (10YR 2/1) and very dark gray (10YR 3/1) silty clay loam; moderate, very fine and fine, granular structure; friable; few fine shell fragments; moderately alkaline; abrupt, smooth boundary.
- Alca—8 to 12 inches, very dark gray (10YR 3/1) silty clay loam; moderate, very fine and fine, granular structure; friable; few fine shell fragments; moderately alkaline; clear, smooth boundary.
- B1g—12 to 19 inches, dark-gray (5Y 4/1) silty clay loam; few, fine, distinct, yellowish-brown (10YR 5/6) and olive (5Y 5/6) mottles; weak, very fine and fine, subangular blocky structure; firm; very few shell fragments; mildly alkaline; clear, smooth boundary.
- B21g—19 to 24 inches, gray (5Y 5/1) silty clay loam; few, fine, distinct, yellowish-brown (10YR 5/6) mottles; weak, very fine and fine, prismatic structure parting to moderate, fine, subangular blocky; firm; few iron and lime concretions; mildly alkaline; clear, smooth boundary.
- B22g—24 to 30 inches, olive-gray (5YR 5/2) light silty clay loam; few, fine, distinct, yellowish-brown (10YR 5/6) mottles and common, fine, distinct, light yellowish-brown (2.5Y 6/4) mottles; few lime concretions and common iron concretions; weak, coarse, prismatic structure; friable; mildly alkaline; clear, smooth boundary.
- B3g—30 to 34 inches, olive-gray (5Y 5/2) and yellowish-brown (10YR 5/6 and 5/8) heavy silt loam; weak, coarse, prismatic structure; friable; moderately alkaline; abrupt, smooth boundary.
- IIC1gc—34 to 41 inches, olive-gray (5Y 5/2) and yellowish-brown (10YR 5/6 and 5/8) loam; massive; friable; moderately alkaline; clear, smooth boundary.
- IIC2ca—41 to 48 inches, light-brown (7.5YR 6/4) loam; common, fine, prominent, greenish-gray (5GY 6/1) mottles and few, fine, distinct, yellowish-brown (10YR 5/8) mottles; massive; friable, moderately alkaline; clear, smooth boundary.
- IIC3—48 to 58 inches, light yellowish-brown (10YR 6/4) loam; common, fine, faint, yellowish-brown (10YR 5/6 and 5/8) mottles; massive; moderately alkaline.

The A horizon ranges from 12 to 20 inches in thickness. The B horizon ranges from 15 to 30 inches in thickness and from silty clay loam or clay loam to silt loam in texture. It ranges from mildly alkaline to moderately alkaline.

Harpster soils are associated with Peotone and Drummer soils. They are more calcareous in the A horizon than Drummer and Peotone soils, and they have free carbonates throughout the profile. They have a thinner A horizon than Peotone soils.

**Harpster silty clay loam (67).**—This nearly level soil is on large upland flats and in drainageways. It is commonly in slightly higher areas that are surrounded by other poorly drained soils. Included in mapping were small areas of soils that are not calcareous in the surface layer.

Surface runoff is slow to very slow, and the hazard of erosion is slight.

If artificially drained, this soil is intensively farmed and is well suited to row crops. Special practices are needed to improve fertility. This soil has severe limitations for most other uses. Management group IIw-1.

## Harvard Series

The Harvard series consists of deep, nearly level to sloping, well drained and moderately well drained soils. These soils are on uplands or on the rims of raised, circular mounds. They formed in loess and in the underlying stratified loamy material.

In a representative profile the surface layer is black silt loam about 8 inches thick. The subsurface layer is brown to dark-brown silt loam about 4 inches thick. The subsoil is about 32 inches thick. The upper 15 inches of the subsoil is brown to dark-brown, dark yellowish-brown, and yellowish-brown silty clay loam that is mottled with yellowish brown in the lower part. The lower 17 inches of the subsoil is mainly grayish-brown and yellowish-brown clay loam. The underlying material is light olive-brown loam about 12 inches thick, yellowish-brown sand about 7 inches thick, and dark-gray and gray clay loam to a depth of 68 inches.

These soils are moderately permeable and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are well suited to intensive use for row crops. They have few limitations for most other uses.

Representative profile of Harvard silt loam, 0 to 2 percent slopes, 520 feet south and 105 feet west of the NE. corner of NW $\frac{1}{4}$ SW $\frac{1}{4}$  sec. 12, T. 42 N., R. 5 E.

- Ap—0 to 8 inches, black (10YR 2/1) silt loam; weak, very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A2—8 to 12 inches, brown to dark-brown (10YR 4/3) silt loam; weak, fine, subangular blocky structure parting to weak, very fine, granular; friable; medium acid; clear, smooth boundary.
- B1t—12 to 16 inches, brown to dark-brown (10YR 4/3) light silty clay loam; moderate, very fine and fine, subangular blocky structure; friable; medium acid; clear, smooth boundary.
- B21t—16 to 21 inches, dark yellowish-brown (10YR 4/4) light silty clay loam; moderate, fine, subangular blocky structure; friable; strongly acid; clear, smooth boundary.
- B22t—21 to 27 inches, yellowish-brown (10YR 5/4) silty clay loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; moderate, medium, subangular blocky structure; friable; discontinuous dark yellowish-brown (10YR 4/4) clay films; few concretions; strongly acid; clear, smooth boundary.
- IIB23t—27 to 39 inches, grayish-brown (10YR 5/2) and yellowish-brown (10YR 5/6) clay loam; weak, medium, prismatic structure parting to weak, medium, subangular blocky; friable; patchy dark grayish-brown (10YR 4/2) clay films; concretions; medium acid; abrupt, smooth boundary.
- IIB3—39 to 44 inches, brown to dark-brown (10YR 4/3) clay loam; weak, coarse, prismatic structure; friable; neutral; abrupt, smooth boundary.
- IIC1—44 to 56 inches, light olive-brown (2.5Y 5/4) loam; common, fine, distinct, yellowish-brown (10YR 5/6) mottles; massive; friable; moderately alkaline; abrupt, smooth boundary.
- IIC2—56 to 63 inches, yellowish-brown (10YR 5/4) sand; single grained; loose; moderately alkaline; abrupt, smooth boundary.

IIC3—63 to 68 inches, dark-gray (10YR 4/1) and gray (10YR 5/1) clay loam; massive; friable; moderately alkaline.

The A horizon ranges from 7 to 12 inches in thickness. In some places an A2 horizon occurs, but in many places it has been mixed with the A1 horizon in plowing. The B horizon ranges from 15 to 40 inches in thickness and from strongly acid to neutral. The IIB horizon is 10 to 20 inches thick and is clay loam, loam, or sandy clay loam.

Harvard soils are associated with Batavia and Millbrook soils. They are better drained than Millbrook soils, and they have a thinner silty layer over the stratified sediment than Batavia soils.

**Harvard silt loam, 0 to 2 percent slopes (344A).**—This soil is in outwash areas. Included with it in mapping were small areas of somewhat poorly drained soils and areas of soils that are calcareous at a depth of less than 40 inches. Also included were small areas in which sand or gravel is below a depth of 5 feet.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. It has few limitations for most other uses. Management group I-2.

**Harvard silt loam, 2 to 4 percent slopes (344B).**—This soil is in outwash areas that are associated with the nearly level Harvard soils. This soil has the profile described as representative of the series.

Included with this soil in mapping were small areas of soils that are calcareous at a depth of less than 40 inches. Also included were small areas where sand or gravel is below a depth of 5 feet.

Surface runoff is medium, and the erosion hazard is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has few limitations for most other uses. Management group IIe-2.

**Harvard silt loam, 4 to 7 percent slopes, eroded (344C2).**—This soil is on the narrow rims and edges of raised, circular mounds. This soil has a profile similar to the one described as representative of the series, but the brown to dark-brown, finer textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas in which none of the original surface layer remains and small areas of soils that are calcareous at a depth of less than 40 inches.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some uses because of slope. Management group IIe-4.

### Herbert Series

This series consists of deep, nearly level, somewhat poorly drained soils on uplands near drainageways. These soils formed in loess and in the underlying loam glacial till.

In a representative profile the surface layer is very dark brown silt loam about 8 inches thick. The subsurface layer, about 4 inches thick, is dark grayish-brown silt loam mottled with brown. The subsoil is about 24 inches thick. The upper 8 inches of the subsoil is brown

to dark-brown, firm silty clay loam. The next 6 inches is multicolored, firm silty clay loam. The lower 10 inches is multicolored, firm clay loam. The underlying material is brown heavy loam.

These soils are moderately permeable and have high available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high runoff, and they are also subject to periodic waterlogging. When they are wet, their stability decreases. In spring, the water table is generally 1 foot to 3 feet below the surface.

These soils are well suited to intensive use for row crops. The periodic waterlogging is a limitation for most other uses.

Representative profile of Herbert silt loam, 147 feet north of center of blacktop road and 306 feet east of section line, in the NW $\frac{1}{4}$ NW $\frac{1}{4}$  sec. 14, T. 42 N., R. 4 E.

Ap—0 to 8 inches, very dark brown (10YR 2/2) silt loam; moderate, very fine, granular structure; friable; many roots; slightly acid; abrupt, smooth boundary.

A2—8 to 12 inches, dark grayish-brown (10YR 4/2) silt loam; few, fine, faint, brown (10YR 5/3) mottles; weak, medium and thick, platy structure parting to moderate, fine, granular; friable; many roots; slightly acid; clear, smooth boundary.

B21t—12 to 16 inches, brown to dark-brown (10YR 4/3) silty clay loam; few, fine, faint, brown (10YR 5/3) mottles; moderate, very fine, subangular blocky structure; firm; thin, discontinuous, dark grayish-brown (10YR 4/2) clay films; many roots; slightly acid; clear, smooth boundary.

B22t—16 to 20 inches, brown to dark-brown (10YR 4/3) silty clay loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; moderate, fine, subangular blocky structure; firm; thin, continuous, grayish-brown (10YR 5/2) clay films; many roots; few iron-manganese concretions; medium acid; clear, smooth boundary.

B23t—20 to 26 inches, mixed grayish-brown (10YR 5/2), brown (10YR 5/3), yellowish-brown (10YR 5/6), and strong-brown (7.5YR 5/6) silty clay loam; moderate, medium, subangular blocky structure; firm; thin, continuous, grayish-brown (2.5Y 5/2) clay films; common roots; very dark brown (10YR 2/2) channel fillings; medium acid; clear, smooth boundary.

IIB24t—26 to 33 inches, mixed brown (7.5YR 5/4), strong-brown (7.5YR 5/6), and yellowish-brown (10YR 5/6) clay loam and some gravel; moderate, medium, angular and subangular blocky structure; firm; thin, discontinuous, brown to dark-brown (7.5YR 4/2) clay films; very dark brown (10YR 2/2) channel fillings; common roots; medium acid; clear, smooth boundary.

IIB3t—33 to 36 inches, brown (10YR 5/3) clay loam; few, fine, distinct, light brownish-gray (10YR 6/2) and strong-brown (7.5YR 5/6) mottles; weak, coarse, angular blocky structure; firm; few, thin, discontinuous, dark grayish-brown (10YR 4/2) clay films; some very dark brown (10YR 2/2) channel fillings; common roots; neutral; clear, smooth boundary.

IIC—36 to 58 inches, brown (10YR 5/3) heavy loam and some gravel; few, fine, faint, very pale brown (10YR 7/3) mottles and few, fine, prominent, gray to light-gray (5Y 6/1) mottles; some streaks of light gray (10YR 7/1); massive; friable; moderately alkaline.

The A horizon ranges from 9 to 14 inches in thickness. The B horizon ranges from 8 to 33 inches in thickness and from medium acid to mildly alkaline. The IIB horizon ranges from 4 to 12 inches in thickness.

Herbert soils are associated with Saybrook and Dodge soils. They have a thicker, darker colored A horizon than Dodge soils, and they are more poorly drained than Saybrook and Dodge soils.

**Herbert silt loam** (62).—This nearly level soil is in the uplands at the upper end of drainageways near poorly drained soils. Included with this soil in mapping were areas of soils that have a thinner, lighter colored surface layer and areas of soils that are more than 40 inches deep over glacial till.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Artificial drainage is beneficial in some areas. This soil has moderate to severe limitations for many other uses. Management group I-3.

### Houghton Series

The Houghton series consists of deep, level to depressional, very poorly drained soils in low areas. These soils formed in fibrous plant remains deposited in swampy areas.

In a representative profile the surface layer is black muck about 15 inches thick. The underlying material is layered dark reddish-brown, dark olive-gray, and dark-gray muck about 40 inches thick.

These soils have moderately rapid permeability and high available water capacity. The surface is ponded, and the soils are very unstable. If drained, the soils are subject to subsidence, and if under a load, they are highly compressible. The water table is at or near the surface.

These soils are suited to row crops or specialty farming if they are drained. They have severe limitations for most other uses, because they are subject to ponding and are very unstable. Fire is a hazard in some places if the soil has been drained.

Representative profile of Houghton muck, 200 feet east and 70 feet north of the SW. corner of SE $\frac{1}{4}$ SE $\frac{1}{4}$  sec. 11, T. 40 N., R. 3 E.

- Oa1—0 to 15 inches, black (N 2/0, broken face and rubbed) sapric material; about 10 percent fiber, less than 5 percent rubbed; massive; neutral; abrupt, smooth boundary.
- Oa2—15 to 25 inches, dark reddish-brown (5YR 3/4, broken face and rubbed) sapric material; about 15 percent fiber, less than 5 percent rubbed; massive; neutral; abrupt, smooth boundary.
- Oa3—25 to 33 inches, dark olive-gray (5Y 3/2, broken face) or very dark gray (5Y 3/1 rubbed) sapric material; about 15 percent fiber, less than 5 percent rubbed; massive; neutral; abrupt, smooth boundary.
- Oa4—33 to 55 inches, dark-gray (5Y 4/1, broken face and rubbed) sapric material; less than 10 percent fiber, less than 5 percent rubbed; massive; neutral.

The surface layer varies considerably because of the degree of decomposition of the organic material. Some areas also have mineral soils washed in on top of and mixed with the organic material. Thickness and color in different layers vary considerably. The depth to underlying mineral material also varies considerably.

Houghton soils are associated with Drummer soils. They formed in organic material, but Drummer soils are mineral.

**Houghton muck** (103).—This nearly level soil is in depressions. In most places it is surrounded by poorly drained soils. Included with this soil in mapping were small areas of mineral soils and small areas where a thin organic layer is over the mineral material.

Surface runoff is ponded, and the hazard of erosion is slight. Where this soil is dry, it is subject to severe soil blowing.

If artificially drained, this soil is commonly farmed along with associated soils. Drainage systems are generally a maintenance concern. Special practices that increase fertility are needed. This soil has very severe limitations for most other uses. Management group IIIw-2.

### Kendall Series

The Kendall series consists of deep, nearly level, somewhat poorly drained soils. These soils are on uplands near the major streams. They formed in silty material and in the underlying strata of fine sand and silt.

In a representative profile the surface layer is dark grayish-brown silt loam about 8 inches thick. The sub-surface layer is grayish-brown silt loam about 6 inches thick. The subsoil is about 43 inches thick. In sequence from the top, it is 17 inches of grayish-brown and light olive-brown silty clay loam mottled with yellowish brown, 21 inches of mixed grayish-brown and yellowish-brown silty clay loam, and 5 inches of light brownish-gray and yellowish-brown loam. The underlying material is strata of yellowish-brown and light brownish-gray loam and silt loam.

These soils are moderately permeable and have high available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high runoff, and they are also subject to periodic waterlogging. Where the soils are wet, their stability decreases. In spring, the water table is generally 1 to 3 feet below the surface.

These soils are well suited to intensive use for row crops. The periodic waterlogging is a limitation for most other uses.

Representative profile of Kendall silt loam, 225 feet south and 36 feet west of the NE. corner of the NW $\frac{1}{4}$ -SW $\frac{1}{4}$  sec. 11, T. 37 N., R. 5 E.

- Ap—0 to 8 inches, dark grayish-brown (10YR 4/2) silt loam; weak, very fine, granular structure; very friable; neutral; abrupt, smooth boundary.
- A2—8 to 14 inches, grayish-brown (10YR 5/2) silt loam; weak, thick, platy structure parting to weak, very fine, granular; friable; slightly acid; clear, smooth boundary.
- B21t—14 to 20 inches, grayish-brown (2.5Y 5/2) silty clay loam; few, fine, distinct, yellowish-brown (10YR 5/6) mottles; moderate, very fine, angular blocky structure; firm; thin olive-brown (2.5Y 4/4) clay films; few iron-manganese concretions; slightly acid; clear, smooth boundary.
- B22t—20 to 31 inches, light olive-brown (2.5Y 5/4) silty clay loam; common, fine, distinct, yellowish-brown (10YR 5/6) mottles; weak, coarse, prismatic structure parting to moderate, fine, angular blocky; firm; thin grayish-brown (2.5Y 5/2) clay films; few iron-manganese concretions; medium acid; clear, smooth boundary.
- B23t—31 to 52 inches, mixed grayish-brown (2.5Y 5/2) and yellowish-brown (10YR 5/6 and 5/8) silty clay loam; weak, medium, prismatic structure; friable; discontinuous dark-gray (10YR 4/1) clay films; few iron-manganese concretions; slightly acid; gradual, smooth boundary.
- IIB3—52 to 57 inches, mixed light brownish-gray (2.5Y 6/2) and yellowish-brown (10YR 5/6) loam; weak, coarse, prismatic structure; friable; discontinuous dark-gray (10YR 4/1) clay films; neutral; abrupt, smooth boundary.

IIC—57 to 65 inches, strata of mixed stratified yellowish-brown (10YR 5/4 and 5/6) and light brownish-gray (2.5Y 6/2) loam and silt loam; massive; friable; moderately alkaline.

The A horizon ranges from 7 to 14 inches in thickness. The B horizon ranges from 20 to 45 inches in thickness and from medium to neutral. The IIB horizon is 5 to 15 inches thick and is loam, clay loam, or sandy loam.

Kendall soils are associated with St. Charles and Drummer soils. They are more poorly drained than St. Charles soils, and they are better drained than Drummer soils.

**Kendall silt loam** (242).—This nearly level soil is on uplands near the major streams. Included with this soil in mapping were areas in which glacial till is below a thin layer of outwash and areas in which stratified sand and silt are at a depth of less than 40 inches. Also included were small areas of poorly drained soils.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Artificial drainage is beneficial in some areas. The soil has moderate to severe limitations for most other uses. Management group I-3.

### Knight Series

The Knight series consists of deep, nearly level, poorly drained soils in depressions in the uplands. These soils formed in silty material that is less than 5 feet deep over stratified loamy outwash material.

In a representative profile the surface layer is very dark brown silt loam about 12 inches thick. The sub-surface layer is very dark grayish-brown and dark grayish-brown silt loam about 14 inches thick. The subsoil is about 34 inches thick. The upper 9 inches of the subsoil is mixed brown to dark brown, dark yellowish-brown, and dark grayish-brown silt loam; the next 19 inches is mixed light-gray to gray, light brownish-gray, light olive-gray, and yellowish-brown silty clay loam; and the lower 6 inches is mixed olive-gray, strong-brown, yellowish-brown, and dark yellowish-brown clay loam.

These soils are moderately slowly permeable and have high available water capacity. The internal drainage is very slow, and the soils have low stability when wet. They are subject to frequent waterlogging and flooding or ponding during periods of runoff. Seasonally the water table is less than 2 feet below the surface.

These soils are well suited to row crops if they are adequately drained. The frequent waterlogging and potential for flooding or ponding are limitations for other uses.

Representative profile of Knight silt loam, 325 feet east and 50 feet north of the SW. corner of NW $\frac{1}{4}$  sec. 35, T. 37 N., R. 5 E.

- A1—0 to 12 inches, very dark brown (10YR 2/2) silt loam; weak, fine and very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A2—12 to 26 inches, very dark grayish-brown (10YR 3/2) and some dark grayish-brown (10YR 4/2) silt loam; weak, thick, platy structure; friable; slightly acid; clear, smooth boundary.
- B1—26 to 35 inches, mixed brown to dark-brown (10YR 4/3), dark yellowish-brown (10YR 4/4), and dark grayish-brown (10YR 4/2) silt loam few, fine, faint, yellowish-brown (10YR 5/8) mottles; weak, medium and fine, subangular blocky structure; friable; medium acid; abrupt, smooth boundary.

B21t—35 to 44 inches, light-gray to gray (5Y 6/1) silty clay loam; common, fine, distinct, yellowish-brown (10YR 5/8) mottles; moderate, coarse, subangular blocky structure; firm; brown to dark-brown (10YR 4/3) clay films in root channels; medium acid; clear, smooth boundary.

B22t—44 to 54 inches, mixed light brownish-gray (2.5Y 6/2), light olive-gray (5Y 6/2), and yellowish-brown (10YR 5/8) silty clay loam; weak, coarse, prismatic structure; firm; few iron-manganese concretions; medium acid; abrupt, smooth boundary.

IIB3—54 to 60 inches, mixed olive-gray (5Y 5/2), strong-brown (7.5YR 5/6), yellowish-brown (10YR 5/8), and dark yellowish-brown (10YR 3/4) clay loam; weak, coarse, prismatic structure; friable; medium acid.

The A1 horizon ranges from 10 to 15 inches in thickness, and the A2 horizon from 12 to 20 inches. The B horizon is 20 to 40 inches thick. It is medium acid to neutral.

Knight soils are associated with Elburn and Plano soils. They are more poorly drained than Elburn and Plano soils, and they have a thick A2 horizon, which Elburn and Plano soils do not have.

**Knight silt loam** (191).—This nearly level soil is in depressions in the uplands or outwash areas. Included with this soil in mapping were small areas of soils that have a clay subsoil and small areas that have recent deposition on the surface.

Surface runoff is ponded, and the hazard of erosion is slight.

If artificially drained, this soil is intensively farmed and is well suited to row crops. It has severe limitations for most other uses. Management group IIw-1.

### La Rose Series

The La Rose series consists of deep, gently sloping to strongly sloping, well drained and moderately well drained soils. These soils are on rolling parts of the major glacial moraines. They formed in less than 2 feet of loess and in the underlying loam glacial till.

In a representative profile the surface layer is very dark grayish-brown silt loam about 7 inches thick. The subsoil is about 13 inches thick. The upper 7 inches of the subsoil is dark-brown, brown and yellowish-brown clay loam, and the lower 6 inches is yellowish-brown and pale-brown heavy loam. The underlying material is yellowish-brown and strong-brown loam.

These soils are moderately permeable and have moderate available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are suited to row crops. They have moderate limitations for some other uses.

Representative profile of La Rose silt loam, 4 to 7 percent slopes, eroded, 540 feet west and 250 feet north of the SE. corner of the NE $\frac{1}{4}$ NW $\frac{1}{4}$  sec. 33, T. 38 N., R. 3 E.

- Ap—0 to 7 inches, very dark grayish-brown (10YR 3/2) silt loam; weak, fine, granular structure; friable; neutral; abrupt, smooth boundary.
- B21t—7 to 10 inches, dark-brown (10YR 3/3) and brown to dark-brown (10YR 4/3) clay loam; moderate, very fine, subangular blocky structure parting to moderate, fine, granular; friable; neutral; clear, smooth boundary.
- B22t—10 to 14 inches, yellowish-brown (10YR 5/4) and brown to dark-brown (10YR 4/3) clay loam; moderate, fine and medium, subangular blocky

structure; firm; mildly alkaline; abrupt, smooth boundary.

B3—14 to 20 inches, yellowish-brown (10YR 5/4) and pale-brown (10YR 6/3) heavy loam; moderate, fine, subangular blocky structure; friable; mildly alkaline; clear, smooth boundary.

C—20 to 60 inches, yellowish-brown (10YR 5/4) and some strong-brown (7.5YR 5/6) loam; massive; friable; moderately alkaline.

The A horizon ranges from 7 to 10 inches in thickness. The B horizon ranges from 8 to 17 inches in thickness and from slightly acid to mildly alkaline. The solum is less than 20 inches thick in many places.

La Rose soils are associated with Saybrook and Octagon soils. They have a thinner solum than Saybrook and Octagon soils.

**La Rose silt loam, 2 to 4 percent slopes (60B).**—This soil is on glacial moraines. It has a profile similar to the one described as representative of the series, but the surface layer has not been mixed into the subsoil. Included with this soil in mapping were small areas where calcareous till is at a depth of more than 24 inches.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIe-2.

**La Rose silt loam, 4 to 7 percent slopes, eroded (60C2).**—This soil is on glacial moraines. It has the profile described as representative of the series. Included with this soil in mapping were small areas where the subsoil has not been mixed into the surface layer and small areas where all of the original surface layer has been removed through erosion.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is farmed and is suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIe-4.

**La Rose silt loam, 7 to 12 percent slopes, eroded (60D2).**—This soil is generally on glacial end moraines. Included with it in mapping were small areas where none of the original surface layer remains and small areas where calcareous till is exposed at the surface. Also included were small areas of less sloping La Rose soils.

Surface runoff is rapid, and the hazard of erosion is severe.

This soil is commonly farmed, but it has limitations for intensive cropping. Control of erosion by either mechanical means or vegetation is necessary. This soil has moderate limitations for some other uses. Management group IIIe-1.

## Limestone Quarries

**Limestone quarries** consists of two areas where limestone bedrock is mined. One quarry is south of Fairdale in Franklin Township, and the other is northeast of Cortland in Cortland Township. The quarry in Cortland Township is operated in conjunction with a gravel pit. Not placed in a management group.

## Lisbon Series

The Lisbon series consists of deep, nearly level, somewhat poorly drained soils. These soils are on uplands near drainageways. They formed in loess and in the underlying loam glacial till.

In a representative profile the surface layer is about 17 inches thick. The upper 12 inches of the surface layer is black silt loam, and the lower 5 inches is very dark grayish-brown light silty clay loam. The subsoil is about 13 inches thick. In sequence from the top, it is 9 inches of brown to dark-brown, firm silty clay loam mottled with dark grayish brown and 4 inches of brown to dark-brown, firm clay loam mottled with brown. The underlying material is brown to dark-brown loam mottled with strong brown.

These soils are moderately permeable and have high available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high runoff and to periodic waterlogging. Where the soils are wet, their stability decreases. In spring the water table is generally 1 foot to 3 feet below the surface.

These soils are well suited to intensive use for row crops, but the periodic waterlogging is a limitation for most other uses.

Representative profile of Lisbon silt loam, 75 feet north and 235 feet west of the SE. corner of NE $\frac{1}{4}$  sec. 11, T. 41 N., R. 4 E.

Ap—0 to 8 inches, black (10YR 2/1) silt loam; moderate, fine and medium, granular structure; friable; many roots; neutral; abrupt, smooth boundary.

A12—8 to 12 inches, black (10YR 2/1) silt loam; weak, medium, granular structure; friable; many roots; neutral; clear, smooth boundary.

A3—12 to 17 inches, very dark grayish-brown (2.5Y 3/2) light silty clay loam; moderate, fine, subangular blocky structure; friable; black (10YR 2/1) and very dark gray (10YR 3/1) coatings; many roots; neutral; clear, smooth boundary.

B21t—17 to 23 inches, brown to dark-brown (10YR 4/3) silty clay loam; few, fine, faint, dark grayish-brown (10YR 4/2) mottles; moderate, fine, subangular blocky structure; firm; thin, continuous, dark grayish-brown (2.5Y 4/2) clay films; black (10YR 2/1) and very dark gray (10YR 3/1) fillings in root channels; common roots; neutral; clear, smooth boundary.

B22t—23 to 26 inches, brown (10YR 4/3) silty clay loam; few, fine, faint, dark grayish-brown (10YR 4/2) mottles; moderate, fine and medium, subangular blocky structure; firm; thin, discontinuous, dark grayish-brown (2.5Y 4/2) clay films; black (10YR 2/1) and very dark gray (10YR 3/1) fillings in root channels; neutral; clear, smooth boundary.

IIB23t—26 to 30 inches, brown to dark-brown (7.5YR 4/4) clay loam that has some fine gravel; few, fine, faint, brown to dark-brown (7.5YR 4/2) mottles; moderate, medium, prismatic structure parting to moderate, medium, angular blocky; firm; thin, discontinuous, dark grayish-brown (10YR 4/2) clay films; common roots; mildly alkaline; clear, smooth boundary.

IIC—30 to 60 inches, brown to dark-brown (7.5YR 4/4) loam that has some pebbles; few, medium, faint, strong-brown (7.5YR 5/6) mottles; massive; firm; few roots; moderately alkaline.

The A horizon ranges from 12 to 18 inches in thickness. The B horizon ranges from 9 to 20 inches in thickness and from slightly acid to mildly alkaline. The IIB horizon ranges from 4 to 10 inches in thickness.

Lisbon soils are associated with Saybrook and Drummer soils. They have a thicker, darker colored A horizon and are more poorly drained than Saybrook soils, and they are better drained than Drummer soils.

**Lisbon silt loam (59).**—This nearly level soil is on slight rises surrounded by poorly drained soils or at the upper end of upland drainageways. Included with this soil in mapping were small areas of poorly drained soils.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Artificial drainage is beneficial in some areas. The soil has moderate to severe limitations for many other uses. Management group I-3.

### Lorenzo Series

The Lorenzo series consists of sloping to moderately steep, excessively drained soils that are shallow over sand and gravel. These soils are on narrow, gravelly ridges on uplands and on moraines. They formed in loamy material over sand and gravel.

In a representative profile the surface layer is very dark grayish-brown loam about 5 inches thick. The subsoil is brown to dark-brown gravelly clay loam about 9 inches thick. The underlying material is mainly yellowish-brown gravelly sand.

These soils have moderate permeability and low available water capacity. The internal drainage is very rapid, and the soils remain stable with changes in moisture content.

These soils have low productivity and are not well suited to row crops. Slope is the main limitation for other uses. Because of the shallowness to sand and gravel, pollution is a risk for some uses.

Representative profile of Lorenzo loam, 6 to 15 percent slopes, eroded, 800 feet east and 45 feet north of the SW. corner of NW $\frac{1}{4}$  sec. 24, T. 40 N., R. 5 E.

Ap—0 to 5 inches, very dark grayish-brown (10YR 3/2) loam; weak, fine, granular structure; friable; neutral; clear, smooth boundary.

B2t—5 to 14 inches, brown to dark-brown (7.5YR 4/4) gravelly clay loam; weak, fine, subangular blocky structure; friable; patchy brown to dark-brown (7.5YR 4/2) clay films; slightly acid; clear, smooth boundary.

IIC1—14 to 18 inches, brown to dark-brown (7.5YR 4/4) gravelly sand; single grained; loose; mildly alkaline; abrupt, smooth boundary.

IIC2—18 to 60 inches, yellowish-brown (10YR 5/4 and 5/6) fine and medium gravelly sand; single grained; loose; moderately alkaline.

The A horizon ranges from 5 to 10 inches in thickness. The B horizon ranges from 5 to 15 inches in thickness and from gravelly clay loam to gravelly sandy clay loam in texture. It is medium acid to neutral. The solum ranges from 12 to 24 inches thick.

Lorenzo soils are associated with Dresden soils. They have a thinner layer of silty material over gravel and sand than Dresden soils.

**Lorenzo loam, 6 to 15 percent slopes, eroded (318D2).**—This soil is on narrow, gravelly ridges on uplands and on narrow breaks adjacent to other soils that are underlain by sand and gravel.

Included with this soil in mapping were small areas of eroded soils that have a subsoil less than 5 inches thick. Also included were small spots where the gravel-

ly substratum is exposed at the surface.

Surface runoff is rapid, and the hazard of erosion is severe.

This soil is commonly farmed along with associated soils, but it has severe limitations for intensive cropping. Control of erosion by either mechanical means or vegetation is a necessary part of good management. This soil is droughty and is low in fertility. Permanent vegetation is difficult to establish. The soil has moderate limitations for some other uses because of slope. Management group IVE-1.

### Miami Series

The Miami series consists of deep, gently sloping to strongly sloping soils that are well drained and moderately well drained. These soils are on the rolling uplands of the major glacial moraines and along the major streams. They formed in less than 2 feet of loess and in the underlying loam glacial till.

In a representative profile the surface layer is dark grayish-brown silt loam about 6 inches thick. The subsoil is about 25 inches thick. The upper 7 inches of the subsoil is dark yellowish-brown silty clay loam, and the lower 18 inches is mainly dark yellowish-brown and brown to dark-brown clay loam. The underlying material is yellowish-brown loam.

These soils are moderately permeable in the subsoil and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are suited to row crops. They have moderate limitations for many other uses.

Representative profile of Miami silt loam, 4 to 7 percent slopes, eroded, in the center of SW $\frac{1}{4}$ SW $\frac{1}{4}$ -SE $\frac{1}{4}$  sec. 7, T. 41 N., R. 5 E.

Ap—0 to 6 inches, dark grayish-brown (10YR 4/2) silt loam; weak, fine, granular structure; friable; neutral; abrupt, smooth boundary.

B21t—6 to 13 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, fine, angular blocky structure; firm; medium acid; clear, smooth boundary.

IIB22t—13 to 18 inches, dark yellowish-brown (10YR 4/4) clay loam; moderate, medium, angular blocky structure; firm; medium acid; clear, smooth boundary.

IIB23t—18 to 24 inches, brown to dark-brown (7.5YR 4/4) clay loam; moderate, coarse, angular blocky structure; firm; thin brown to dark-brown (7.5YR 4/2) clay films; neutral; gradual, smooth boundary.

IIB24t—24 to 31 inches, brown (7.5YR 5/4) clay loam; moderate, coarse, angular blocky structure; firm; thin brown to dark-brown (7.5YR 4/2) clay films; neutral; clear, smooth boundary.

IIC—31 to 60 inches, yellowish-brown (10YR 5/4) loam till; massive; friable; moderately alkaline.

The A horizon ranges from 6 to 12 inches in thickness. The B horizon ranges from 15 to 30 inches in thickness. In some places the lower part of the B horizon is loam, and in some places it has mottles. The B horizon ranges from medium acid to mildly alkaline.

Miami soils are associated with Dodge and Herbert soils. They have a thinner mantle of loess than Dodge and Herbert soils and are better drained than Herbert soils.

**Miami silt loam, 2 to 4 percent slopes (27B).**—This soil is in the uplands on the glacial moraines, in areas above steeper soils. It has the profile described as rep-

representative of the series. Included with this soil in mapping were small areas of Dodge silt loam.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is suited to row crops. Most areas are intensively farmed, but some areas remain in native hardwoods. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIe-1.

**Miami silt loam, 4 to 7 percent slopes, eroded (27C2).**—This soil is in the uplands on the glacial moraines. It has a profile similar to the one described as representative of the series, but some of the dark yellowish-brown, finer textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas where none of the subsoil has been mixed into the surface layer.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is suited to row crops. Most of it is farmed, but some areas remain in native hardwoods. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIe-4.

**Miami silt loam, 7 to 12 percent slopes, eroded (27D2).**—This soil is on the glacial moraines. It has a profile similar to the one described as representative of the series, but some of the dark yellowish-brown, finer textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas where none of the subsoil has been mixed into the surface layer and small areas of Strawn soils.

Surface runoff is rapid, and the hazard of erosion is severe.

This soil is farmed, but it has limitations for intensive cropping. Control of erosion by either mechanical means or vegetation is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIIe-1.

### Millbrook Series

The Millbrook series consists of deep, nearly level, somewhat poorly drained soils on uplands near drainage ways. These soils formed in silty material and in the underlying stratified loamy outwash. In some areas sand and gravel are below a depth of 5 feet.

In a representative profile the surface layer is black silt loam about 8 inches thick. The subsurface layer is dark grayish-brown and brown to dark-brown silt loam about 4 inches thick. The subsoil is about 29 inches thick. The upper 14 inches of the subsoil is brown to dark-brown and dark grayish-brown silty clay loam mottled with yellowish brown, and the lower 15 inches is dark grayish-brown and yellowish-brown loam and sandy loam mottled with yellowish brown. The underlying material is gray, light brownish-gray and yellowish-brown, stratified loam, sandy loam, and silt loam.

These soils have a moderately permeable subsoil and have high available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high run-

off, and they are subject to periodic waterlogging. Where the soils are wet, their stability decreases. In spring the water table is generally 1 foot to 3 feet below the surface.

These soils are well suited to intensive use for row crops. The periodic waterlogging is a limitation for most other uses.

Representative profile of Millbrook silt loam, 40 feet east and 150 feet south of the NW. corner of the NE $\frac{1}{4}$ -NW $\frac{1}{4}$  sec. 12, T. 42 N., R. 5 E.

Ap—0 to 8 inches, black (10YR 2/1) silt loam; moderate, fine, granular structure; friable; strongly acid; abrupt, smooth boundary.

A2—8 to 12 inches, dark grayish-brown (10YR 4/2) and some brown to dark-brown (10YR 4/3) silt loam; weak, thin, platy structure parting to moderate, fine, crumb; friable; strongly acid; clear, smooth boundary.

B21t—12 to 18 inches, brown to dark-brown (10YR 4/3) silty clay loam; few, fine, distinct, yellowish-brown (10YR 5/6) mottles; moderate, fine, subangular blocky structure; friable; few, patchy, dark grayish-brown (10YR 4/2) clay films; medium acid; clear, smooth boundary.

B22t—18 to 26 inches, mixed dark grayish-brown (10YR 4/2) and brown to dark-brown (10YR 4/3) silty clay loam; common, fine, prominent, yellowish-brown (10YR 5/6) mottles; weak, fine and medium, prismatic structure parting to moderate, fine and medium, subangular blocky; firm; few very dark brown (10YR 2/2) fillings in root channels and wormholes; few very dark grayish-brown (10YR 3/2) iron-manganese concretions; discontinuous grayish-brown (2.5Y 5/2) clay films; slightly acid; clear, smooth boundary.

IIB23t—26 to 34 inches, mixed dark grayish-brown (10YR 4/2) and yellowish-brown (10YR 5/6 and 5/8) loam; weak, medium, prismatic structure parting to moderate, medium, subangular blocky; firm; few very dark brown iron-manganese concretions; discontinuous grayish-brown (2.5Y 5/2) clay films; neutral; clear, smooth boundary.

IIB3—34 to 41 inches, dark grayish-brown (10YR 4/2) sandy loam; few, fine, distinct, yellowish-brown (10YR 5/6) mottles; weak, coarse, subangular blocky structure; very friable; few, discontinuous, grayish-brown (10YR 5/2) clay films; neutral; clear, smooth boundary.

IIC—41 to 68 inches, stratified gray (5Y 6/1) silt loam and mixed light brownish-gray (2.5Y 6/2) and yellowish-brown (10YR 5/6 and 5/8) loam and sandy loam; massive; very friable; very dark brown (10YR 2/2) iron-manganese concretions; moderately alkaline.

The A horizon ranges from 9 to 14 inches in thickness. In places the A2 horizon has been mixed with the A1 horizon by plowing. The B horizon ranges from medium acid to neutral. The IIB horizon ranges from 5 to 15 inches in thickness and from clay loam to sandy loam or loam in texture.

Millbrook soils are associated with Harvard and Drummer soils. They are more poorly drained than Harvard soils and better drained than Drummer soils.

**Millbrook silt loam (219).**—This nearly level soil is in broad outwash areas or on terraces along the major streams. Included with it in mapping were areas where the silty material is more than 40 inches thick. Also included were areas where the surface layer is thinner than in this soil and small areas of poorly drained soils. In some places the depth to carbonates is less than 40 inches.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to

row crops. Artificial drainage is beneficial in some places. It has moderate to severe limitations for most other uses. Management group I-3.

### Muscatine Series

The Muscatine series consists of deep, nearly level, somewhat poorly drained soils. These soils are in the uplands or near drainageways. They formed in more than 5 feet of loess.

In a representative profile the surface layer is black silt loam about 14 inches thick. The subsoil, about 28 inches thick, is multicolored silty clay loam mottled with yellowish brown. The underlying material is multicolored silt loam.

These soils have a moderately permeable subsoil and have high available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high runoff, and they are also subject to periodic waterlogging. Where they are wet, their stability decreases. In spring the water table is generally 1 foot to 3 feet below the surface.

These soils are well suited to intensive use for row crops. The periodic waterlogging is a limitation for most other uses.

Representative profile of Muscatine silt loam, 200 feet north and 15 feet west of the SE. corner of the SW $\frac{1}{4}$ SE $\frac{1}{4}$  sec. 30, T. 37 N., R. 3 E.

- Ap—0 to 9 inches, black (10YR 2/1) silt loam; weak, very fine, granular structure; friable; slightly acid; abrupt, smooth boundary.
- A3—9 to 14 inches, black (10YR 2/1) silt loam that has some very dark grayish brown (10YR 3/2) in the lower part; weak, very fine, subangular blocky structure parting to weak, very fine, granular; friable; slightly acid; clear, smooth boundary.
- B1—14 to 17 inches, dark grayish-brown (10YR 4/2) silty clay loam; medium, very fine and fine, subangular blocky structure; firm; medium acid; clear, smooth boundary.
- B21t—17 to 22 inches, brown to dark-brown (10YR 4/3) silty clay loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; medium, fine, subangular blocky structure; firm; dark grayish-brown (10YR 4/2) clay films; slightly acid; clear, smooth boundary.
- B22t—22 to 29 inches, mixed dark grayish-brown (10YR 4/2 and 2.5Y 4/2) and grayish-brown (2.5Y 5/2) silty clay loam; common, fine, faint, yellowish-brown (10YR 5/6) mottles; moderate, medium, subangular blocky structure; firm; slightly acid; clear, smooth boundary.
- B23t—29 to 36 inches, mixed dark grayish-brown (2.5Y 4/2) and olive-gray (5Y 5/2) silty clay loam; common, medium, distinct, yellowish-brown (10YR 5/6) mottles; weak, coarse, subangular blocky structure; friable; neutral; clear, smooth boundary.
- B3—36 to 42 inches, mixed grayish-brown (2.5Y 5/2), yellowish-brown (10YR 5/6), and light olive-gray (5Y 6/2) silty clay loam; weak, coarse, prismatic structure; friable; neutral; abrupt, smooth boundary.
- C—42 to 60 inches, mixed yellowish-brown (10YR 5/6), light olive-gray (5Y 6/2), and grayish-brown (2.5Y 5/2) silt loam; massive; friable; moderately alkaline.

The A horizon ranges from 12 to 18 inches in thickness. The B horizon ranges from 20 to 40 inches in thickness and from medium acid to neutral.

Muscatine soils are associated with Sable and Catlin soils. They are better drained than Sable soils, and they are more

poorly drained and have a thicker mantle of loess than Catlin soils.

**Muscatine silt loam (41).**—This nearly level soil is in areas downslope from the well drained and moderately well drained Catlin soils. Included with this soil in mapping were small areas where the underlying glacial till is less than 5 feet deep. Also included were small areas of poorly drained Sable soils.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is very well suited to row crops. Artificial drainage is beneficial in some places. The soil has moderate to severe limitations for many other uses. Management group I-3.

### Octagon Series

This series consists of deep, gently sloping to sloping, well drained and moderately well drained soils. These soils are on the rolling uplands of the major glacial moraines. They formed in less than 2 feet of loess and in the underlying loam glacial till.

In a representative profile the surface layer is mainly very dark grayish-brown silt loam about 7 inches thick. The subsoil is brown, dark-brown, dark yellowish-brown, and yellowish-brown clay loam about 21 inches thick. The underlying material is multicolored loam.

These soils are moderately permeable and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are suited to row crops and have moderate limitations for some other uses.

Representative profile of Octagon silt loam, 4 to 7 percent slopes, eroded, 474 feet west and 151 feet north of the SE. corner of sec. 7, T. 39 N., R. 5 E.

- Ap—0 to 7 inches, very dark grayish-brown (10YR 3/2) and some brown to dark-brown (10YR 4/3) silt loam; weak, very fine, granular structure; very friable; strongly acid; abrupt, smooth boundary.
- IIB21t—7 to 14 inches, brown to dark-brown (10YR 4/3) clay loam; weak, prismatic structure parting to moderate, fine, subangular blocky; firm; dark-brown (10YR 3/3) patchy clay films; medium acid; clear, smooth boundary.
- IIB22t—14 to 20 inches, dark yellowish-brown (10YR 4/4) clay loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; weak, prismatic structure parting to weak, fine, angular blocky; friable; medium acid; clear, smooth boundary.
- IIB3t—20 to 28 inches, yellowish-brown (10YR 5/4 and 5/6) and brownish-yellow (10YR 6/6) clay loam; weak, medium, subangular blocky structure; friable; mildly alkaline; abrupt, smooth boundary.
- IIC—28 to 60 inches, yellowish-brown (10YR 5/4) and some light brownish-gray (10YR 6/2) and brown to dark-brown (7.5YR 4/4) loam; massive; friable; moderately alkaline.

The A horizon ranges from 7 to 10 inches in thickness. Generally, the A1 and A2 horizons have been mixed by plowing. The B horizon ranges from 15 to 30 inches in thickness. In some places the lower part of the B horizon is loam. This horizon ranges from medium acid to mildly alkaline.

Octagon soils are associated with and have natural drainage similar to that of Saybrook and La Rose soils. They have a thinner mantle of loess than Saybrook soils and a thicker solum than La Rose soils.

**Octagon silt loam, 2 to 4 percent slopes (656B).**—This soil is on uplands on the glacial moraines. It has the profile described as representative of the series. Included with this soil in mapping were small areas of La Rose soils.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIe-2.

**Octagon silt loam, 4 to 7 percent slopes, eroded (656C2).**—This soil is on uplands on the glacial moraines downslope from less sloping Octagon soils. This soil has a profile similar to the one described as representative of the series, but the brown to dark-brown, finer-textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas where all of the original surface layer has been removed through erosion and small areas where the subsoil is thinner than is typical for this soil.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is farmed and is suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group IIe-4.

### Peotone Series

The Peotone series consists of deep, nearly level, very poorly drained soils in upland depressions. These soils formed in silty material that is more than 5 feet thick over stratified sand and silt.

In a representative profile the surface layer is black silty clay loam about 20 inches thick. The subsoil is about 18 inches thick. The upper 6 inches of the subsoil is black and dark-gray silty clay loam, and the lower 12 inches is gray, dark-gray, and olive-yellow silty clay loam mottled with yellowish brown. The underlying material is gray, olive-gray, and strong-brown silty clay loam that extends to a depth of about 60 inches.

These soils have moderately slow permeability and high available water capacity. Surface drainage is ponded, and internal drainage is very slow. The soils have low stability if wet. They are subject to frequent water-logging and flooding or ponding during periods of runoff. Seasonally the water table is at or near the surface.

These soils are suited to row crops if they are adequately drained. The frequent waterlogging and potential for flooding are severe limitations for other uses.

Representative profile of Peotone silty clay loam, 75 feet east and 120 feet south of the junction of Lee Road and the Chicago and Northwestern Railroad tracks, in the NW $\frac{1}{4}$ NE $\frac{1}{4}$  sec. 7, T. 38 N., R. 4 E.

Ap—0 to 8 inches, black (10YR 2/1) silty clay loam; moderate, very fine and fine, granular structure; friable; neutral; abrupt, smooth boundary.

A12—8 to 20 inches, black (10YR 2/1) silty clay loam; weak, coarse, subangular blocky structure parting to moderate, very fine and fine, subangular blocky; friable; neutral; gradual, smooth boundary.

B21g—20 to 26 inches, black (10YR 2/1) and some dark-gray (10YR 4/1) heavy silty clay loam; weak, fine, prismatic structure parting to moderate, fine and medium, subangular blocky; friable; neutral; clear, smooth boundary.

B22g—26 to 38 inches, gray (5Y 5/1), dark-gray (5Y 4/1), and olive-yellow (5Y 6/6) silty clay loam; few, fine, faint, yellowish-brown (10YR 5/8) mottles; weak, medium, prismatic structure parting to moderate, fine, prismatic; friable; neutral; gradual, smooth boundary.

C—38 to 60 inches, gray (5Y 5/1), olive-gray (5Y 5/2), and strong-brown (7.5YR 5/6) silty clay loam; massive; friable; neutral.

The A horizon ranges from 20 to 35 inches in thickness. The B horizon is 10 to 20 inches thick. It ranges from neutral to mildly alkaline.

Peotone soils are associated with Drummer soils. They have a thicker A horizon than Drummer soils.

**Peotone silty clay loam (330).**—This nearly level soil is in depressions that are generally surrounded by other poorly drained soils. Included with this soil in mapping were small areas that have recently deposited material on the surface and small areas where the surface layer is thinner than typical.

Surface runoff is ponded, and the hazard of erosion is slight.

If artificially drained, this soil is intensively farmed and is suited to row crops. It has severe limitations for most other uses. Management group IIIw-1.

### Plano Series

The Plano series consists of deep, nearly level to gently sloping, well drained and moderately well drained soils. These soils are in broad areas in the uplands. They formed in silty material and in the underlying stratified loamy material.

In a representative profile the surface layer is very dark grayish-brown and dark-brown silt loam about 13 inches thick. The subsoil is about 49 inches thick. In sequence from the top, it is 10 inches of brown to dark-brown and dark yellowish-brown silty clay loam, 13 inches of brown to dark-brown and light olive-gray silty clay loam mottled with strong brown, 15 inches of dark yellowish-brown and yellowish-brown and some pale-olive silty clay loam mottled with strong brown, and 11 inches of dark yellowish-brown and yellowish-brown clay loam. The underlying material is yellowish-brown loam.

These soils are moderately permeable and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are well suited to intensive use for row crops. They have few limitations for a variety of other uses.

Representative profile of Plano silt loam, 0 to 2 percent slopes, 40 feet west and 150 feet north of the center of SE $\frac{1}{4}$ SE $\frac{1}{4}$  sec. 35, T. 37 N., R. 5 E.

Ap—0 to 9 inches, very dark grayish-brown (10YR 3/2) silt loam; weak, fine and very fine, granular structure; friable; neutral; abrupt, smooth boundary.

A3—9 to 13 inches, very dark grayish-brown (10YR 3/2) and dark-brown (10YR 3/3) silty clay loam; weak, fine, subangular blocky structure parting to weak,

- fine and very fine, granular; friable; slightly acid; clear, smooth boundary.
- B21t—13 to 23 inches, brown to dark-brown (10YR 4/3) and dark yellowish-brown (10YR 4/4) silty clay loam; moderate, fine, subangular blocky structure; firm; slightly acid; clear, smooth boundary.
- B22t—23 to 36 inches, brown to dark-brown (10YR 4/3) and some light olive-gray (5Y 6/2) silty clay loam; few, fine, distinct, strong-brown (7.5YR 5/6) mottles; moderate, fine and medium, subangular blocky structure; firm; patchy dark-brown (10YR 3/3) clay films; slightly acid; clear, smooth boundary.
- B3t—36 to 51 inches, dark yellowish-brown (10YR 4/4) and yellowish-brown (10YR 5/4) and some pale-olive (5Y 6/3) silty clay loam; common, fine, distinct, strong-brown (7.5YR 5/6) mottles; weak, medium, prismatic structure; firm; patchy brown to dark-brown (10YR 4/3) clay films; iron-manganese concretions; neutral; abrupt, smooth boundary.
- IIB4—51 to 62 inches, dark yellowish-brown (10YR 4/4) and yellowish-brown (10YR 5/4) clay loam; weak, coarse, prismatic structure; firm; neutral; abrupt, smooth boundary.
- IIC—62 to 74 inches, yellowish-brown (10YR 5/4) loam; massive; friable; moderately alkaline.

The A horizon ranges from 10 to 16 inches in thickness. The B horizon ranges from 30 to 50 inches in thickness and from medium acid to neutral. The IIB horizon ranges from 5 to 15 inches in thickness and is clay loam, loam, or sandy clay loam in texture.

Plano soils are associated with the drained Elburn and Drummer soils.

**Plano silt loam, 0 to 2 percent slopes (199A).**—This soil is in outwash areas or on the tops of raised circular mounds. It has the profile described as representative of the series.

Included with this soil in mapping were areas where glacial till is below a thin layer of outwash material and areas where the silt is more than 5 feet thick. Also included were small areas of somewhat poorly drained soils.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. It has few limitations for most other uses. Management group I-2.

**Plano silt loam, 2 to 4 percent slopes (199B).**—This soil is in outwash areas or on the higher parts of raised circular mounds. It has a profile similar to the one described as representative of the series, but the surface layer is thinner and lighter colored. Included with this soil in mapping were areas where glacial till is below a thin layer of outwash material and some areas where the silt is more than 5 feet thick.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has few limitations for most other uses. Management group IIe-2.

## Proctor Series

The Proctor series consists of deep, nearly level to gently sloping, well drained and moderately well drained soils on uplands. These soils formed in silty material and in the underlying stratified loamy out-

wash. In some areas sand and gravel are below a depth of 5 feet.

In a representative profile the surface layer is very dark brown silt loam about 12 inches thick. The subsoil is about 48 inches thick. The upper 13 inches of the subsoil is brown to dark-brown silty clay loam, the next 7 inches is yellowish-brown clay loam, the next 12 inches is mostly yellowish-brown sandy loam mottled with yellowish brown, and the lower 16 inches is brown to dark-brown and yellowish-brown light loam to sandy loam mottled with yellowish brown.

These soils are moderately permeable and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is more than 5 feet below the surface.

These soils are well suited to row crops, and they have few limitations for many other uses.

Representative profile of Proctor silt loam, 2 to 4 percent slopes, in the NW $\frac{1}{4}$ , SW $\frac{1}{4}$ , SW $\frac{1}{4}$  sec. 12, T. 42 N., R. 5 E.

- A1—0 to 12 inches, very dark brown (10YR 2/2) silt loam; weak, very fine, granular structure; very friable; neutral; abrupt, smooth boundary.
- B21t—12 to 19 inches, brown to dark-brown (10YR 4/3) light silty clay loam; weak, very fine and fine, subangular blocky structure; very friable; some dark-brown (10YR 3/3) patches; neutral; clear, smooth boundary.
- B22t—19 to 25 inches, brown to dark-brown (10YR 4/3) silty clay loam; weak, fine, prismatic structure parting to weak, very fine and fine, subangular blocky; friable; dark-brown (10YR 3/3) clay films; slightly acid; clear, smooth boundary.
- IIB23t—25 to 32 inches, yellowish-brown (10YR 5/4) clay loam; weak, coarse, prismatic structure parting to weak, fine and medium, subangular blocky; friable; brown to dark-brown (10YR 4/3) clay films; neutral; clear, smooth boundary.
- IIB24t—32 to 40 inches, yellowish-brown (10YR 5/4) sandy loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; weak, coarse, prismatic structure; friable; brown to dark-brown (10YR 4/3) clay films; neutral; abrupt, smooth boundary.
- IIB25—40 to 44 inches, gray (10YR 5/1) and yellowish-brown (10YR 5/6) loam; few, fine, distinct, yellowish-brown (10YR 5/8) mottles; weak, medium and coarse, subangular blocky structure; friable; slightly acid; abrupt, smooth boundary.
- IIB3—44 to 60 inches, brown to dark-brown (7.5YR 4/4) and yellowish-brown (10YR 5/4) light loam to sandy loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; weak, coarse, blocky structure; very friable; mildly alkaline.

The A horizon ranges from 10 to 16 inches in thickness. The B horizon and the IIB horizon are 18 to 50 inches thick. The IIB horizon ranges from clay loam to sandy loam in texture and from 20 to 40 inches in thickness. The B horizon ranges from slightly acid to mildly alkaline.

Proctor soils are associated with Elburn and Drummer soils. They are better drained than Elburn and Drummer soils, and they have a thinner silty layer over the stratified underlying material than Elburn soils.

**Proctor silt loam, 0 to 2 percent slopes (148A).**—This soil is in outwash areas along the major drainageways. It has the profile described as representative of the series.

Included with this soil in mapping were small areas where the surface layer and subsoil contain more sand than is typical for this soil. Also included were small areas of somewhat poorly drained soils and small areas

where sand and gravel are below a depth of 5 feet.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. It has few limitations for most other uses. Management group I-2.

**Proctor silt loam, 2 to 4 percent slopes (148B).**—This soil is in outwash areas along the major drainageways downslope from the other Proctor soils. It has a profile similar to the one described as representative of the series, but the surface layer is thinner and lighter colored.

Included with this soil in mapping were small areas where the surface layer and subsoil are more sandy than is typical for this soil. Also included were small areas where sand and gravel are below a depth of 5 feet.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has few limitations for most other uses. Management group IIe-2.

### Rush Series

The Rush series consists of deep, nearly level to gently sloping, well-drained soils. These soils are on uplands near the Kishwaukee River. They formed in silty material and in the underlying sand and gravel.

In a representative profile the surface layer is dark grayish-brown silt loam about 6 inches thick. The subsurface layer is dark grayish-brown and brown to dark-brown silt loam about 7 inches thick. The subsoil is about 43 inches thick. The upper 33 inches of the subsoil is dark yellowish-brown silty clay loam. The lower 10 inches is brown to dark-brown and strong-brown gravelly clay loam. The underlying material is yellowish-brown gravelly sand.

These soils have a moderately permeable subsoil and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is more than 5 feet below the surface.

These soils are well suited to intensive use for row crops. They have few limitations for a variety of other uses. Because of the underlying sand and gravel, pollution is a risk if the soil is used for some purposes.

Representative profile of Rush silt loam, 0 to 2 percent slopes, in the center of the SE $\frac{1}{4}$ NE $\frac{1}{4}$  sec. 12, T. 41 N., R. 4 E.

- Ap—0 to 6 inches, dark grayish-brown (10YR 4/2) silt loam; weak, very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A2—6 to 13 inches, dark grayish-brown (10YR 4/2) and brown to dark-brown (10YR 4/3) silt loam; moderate, thin and medium, platy structure parting to fine, granular; friable; neutral; clear, smooth boundary.
- B1—13 to 17 inches, dark yellowish-brown (10YR 4/4) light silty clay loam; weak, fine, subangular blocky structure; friable; neutral; clear, smooth boundary.
- B2t—17 to 26 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, very fine and fine, subangular blocky structure; firm; thin, continuous, brown to dark-brown (10YR 4/3) clay films;

- medium acid; clear, smooth boundary.
- B22t—26 to 37 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, fine, subangular blocky structure; firm; thin, continuous, brown to dark-brown (10YR 4/3) clay films; medium acid; clear, smooth boundary.
- B23t—37 to 46 inches, dark yellowish-brown (10YR 4/4) silty clay loam; weak, coarse, prismatic structure parting to weak, coarse, subangular blocky; firm; thin, continuous, brown to dark-brown (7.5YR 4/4) clay films; medium acid; abrupt, smooth boundary.
- IIB3t—46 to 56 inches, brown to dark-brown (7.5YR 4/4) and strong-brown (7.5YR 5/8) gravelly clay loam; weak, coarse, prismatic structure; firm; moderately thick, continuous, dark-brown (7.5YR 3/2) clay films; medium acid; abrupt, wavy boundary.
- IIC—56 to 60 inches, yellowish-brown (10YR 5/4) gravelly sand; single grained; loose; moderately alkaline.

The A horizon ranges from 7 to 15 inches in thickness. The B horizon ranges from 28 to 52 inches in thickness and from medium acid to neutral. The IIB horizon ranges from 4 to 12 inches in thickness and from clay loam to gravelly clay loam in texture.

Rush soils are associated with Bowes and Dresden soils. They have a thinner, lighter colored Ap horizon than Bowes and Dresden soils, and they have a thicker silty layer over sand and gravel than Dresden soils.

**Rush silt loam, 0 to 2 percent slopes (791A).**—This soil is on broad outwash flats along the Kishwaukee River. It has the profile described as representative of the series. Included with this soil in mapping were small areas of somewhat poorly drained soils and areas where the silt is more than 5 feet deep.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Some areas are used for gravel pits. The soil has few limitations for most other uses. Management group I-1.

**Rush silt loam, 2 to 4 percent slopes (791B).**—This soil is on outwash flats along the Kishwaukee River. It has a profile similar to the one described as representative of the series, but the surface layer is thinner and lighter colored. Included with this soil in mapping were small areas where gravel is at a depth of less than 40 inches and areas where there is more than 5 feet of silt over the gravel.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. Some areas are used for gravel pits. This soil has few limitations for most other uses. Management group IIe-1.

### Sable Series

The Sable series consists of deep, nearly level, poorly drained soils in drainageways in the uplands. These soils formed in more than 5 feet of loess.

In a representative profile the surface layer is about 17 inches thick. It is black silty clay loam in the upper 14 inches and very dark gray silty clay loam in the lower 3 inches. The subsoil, about 27 inches thick, is mainly olive-gray silty clay loam mottled with yellowish brown. The underlying material is gray, olive-gray, and yellowish-brown silt loam.

These soils are moderately permeable and have high

available water capacity. Surface drainage is very slow to ponded, and the internal drainage is very slow. The soils have low stability when wet. They are subject to frequent waterlogging and flooding or ponding during periods of runoff. Seasonally the water table is less than 2 feet from the surface.

These soils are well suited to row crops if they are adequately drained. The frequent waterlogging and potential for flooding are severe limitations for many other uses.

Representative profile of Sable silty clay loam, 84 feet north of the SE. corner of SW $\frac{1}{4}$  sec. 30, T. 37 N., R. 3 E.

- A1—0 to 14 inches, black (N 2/0) silty clay loam; weak, very fine, granular structure; friable; slightly acid; clear, smooth boundary.
- A3—14 to 17 inches, very dark gray (5Y 3/1) silty clay loam; weak, very fine, subangular blocky structure; friable; neutral; clear, smooth boundary.
- B21g—17 to 22 inches, olive-gray (5Y 5/2) silty clay loam; few, fine, distinct, yellowish-brown (10YR 5/6) mottles; weak, medium, subangular blocky structure; friable; neutral; clear, smooth boundary.
- B22g—22 to 30 inches, olive-gray (5Y 5/2) silty clay loam; common, medium, distinct, yellowish-brown (10YR 5/6) mottles; weak, medium, prismatic structure; friable; thin, discontinuous, very dark gray (10YR 3/1) coatings on faces of peds; neutral; gradual, smooth boundary.
- B23g—30 to 44 inches, mixed gray (5Y 5/1) and olive-gray (5Y 5/2) silty clay loam; common, medium, distinct, yellowish-brown (10YR 5/6) mottles; weak, coarse, prismatic structure; friable; neutral; clear, smooth boundary.
- C—44 to 60 inches, mixed gray (5Y 5/1), olive-gray (5Y 5/2), and yellowish-brown (10YR 5/6) silt loam; massive; friable; neutral.

The A horizon ranges from 12 to 18 inches in thickness. The B horizon ranges from 20 to 40 inches in thickness and from slightly acid to neutral.

Sable soils are associated with Muscatine soils. They are more poorly drained than Muscatine soils.

**Sable silty clay loam (68).**—This nearly level soil is in upland drainageways. Included with this soil in mapping were areas where stratified sandy and silty material is less than 5 feet deep.

Surface runoff is very slow to ponded, and the hazard of erosion is slight.

If artificially drained, this soil is intensively farmed and is well suited to row crops. It has severe limitations for most other uses. Management group IIw-1.

### St. Charles Series

The St. Charles series consists of deep, nearly level to gently sloping, well drained and moderately well drained soils. These soils are in broad upland areas. They formed in silty material and in the underlying stratified loamy material.

In a representative profile the surface layer is dark grayish-brown silty loam about 7 inches thick. The subsurface layer is dark grayish-brown and grayish-brown silt loam about 3 inches thick. The subsoil is about 39 inches thick. The upper 20 inches of the subsoil is dark yellowish-brown silty clay loam, the next 11 inches is dark yellowish-brown and yellowish-brown silty clay loam, and the lower 8 inches is yellowish-brown clay loam. The underlying material is yellowish-

brown and some light olive-gray, stratified loam and silt loam.

These soils are moderately permeable and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are well suited to intensive use for row crops. They have few limitations for a variety of other uses.

Representative profile of St. Charles silt loam, 0 to 2 percent slopes, 80 feet east of road center and 145 feet north of woodlot fence, in the NE $\frac{1}{4}$ NW $\frac{1}{4}$ SW $\frac{1}{4}$  sec. 11, T. 37 N., R. 5 W.

- Ap—0 to 7 inches, dark grayish-brown (10YR 4/2) silt loam; weak, very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A2—7 to 10 inches, dark grayish-brown (10YR 4/2) and grayish-brown (10YR 5/2) silt loam; weak, medium, platy structure; friable; neutral; abrupt, smooth boundary.
- B1—10 to 16 inches, dark yellowish-brown (10YR 4/4) silty clay loam; weak, very fine, angular blocky structure; friable; neutral; clear, smooth boundary.
- B21t—16 to 21 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, fine, angular blocky structure; firm; slightly acid; clear, smooth boundary.
- B22t—21 to 30 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, medium, angular blocky structure; firm; thin brown to dark-brown (7.5YR 4/4) clay films; medium acid; gradual, smooth boundary.
- B23t—30 to 41 inches, mixed dark yellowish-brown (10YR 4/4) and yellowish-brown (10YR 5/6) silty clay loam; weak, medium, subangular blocky structure; friable; discontinuous brown to dark-brown (7.5YR 4/4) clay films; few iron-manganese concretions; medium acid; clear, smooth boundary.
- IIB3—41 to 49 inches, yellowish-brown (10YR 5/6 and 5/8) clay loam; weak, fine, prismatic structure; friable; discontinuous brown to dark-brown (7.5YR 4/4) clay films; slightly acid; abrupt, smooth boundary.
- IIC—49 to 60 inches, yellowish-brown (10YR 5/6) and some light olive-gray (5Y 6/2) silt loam that has strata of loam; massive; friable; neutral.

The A horizon ranges from 8 to 12 inches in thickness. The B horizon ranges from 20 to 40 inches in thickness and from medium acid to neutral. The IIB horizon ranges from 5 to 15 inches in thickness and is clay loam, loam, or sandy clay loam.

St. Charles soils are associated with Batavia and Kendall soils. St. Charles soils have a thinner, lighter colored Ap horizon than Batavia soils. They are better drained than Kendall soils.

**St. Charles silt loam, 0 to 2 percent slopes (243A).**—This soil is on uplands near the major streams. It has the profile described as representative of the series. Included with this soil in mapping were areas where glacial till is below a thin layer of outwash material and small areas of somewhat poorly drained soils.

Surface runoff is slow, and the hazard of erosion is slight.

Most of this soil is intensively farmed and is well suited to row crops. Some areas remain in native hardwoods. This soil has few limitations for most other uses. Management group I-1.

**St. Charles silt loam, 2 to 4 percent slopes (243B).**—This soil is on uplands near the major streams. It has a profile similar to the one described as representative of the series, but the surface layer is thinner and lighter colored.

Included with this soil in mapping were areas where glacial till is below a thin layer of outwash material and small areas of somewhat poorly drained soils. Also included were small areas of Dodge soils.

Surface runoff is medium, and the hazard of erosion is moderate.

Most of this soil is intensively farmed and is well suited to row crops. Some areas remain in native hardwoods. Control of erosion is a necessary part of good management. The soil has few limitations for most other uses. Management group IIe-1.

### Sawmill Series

The Sawmill series consists of deep, nearly level, poorly drained soils. These soils are on bottom lands along the major streams. They formed in moderately fine textured alluvium.

In a representative profile the surface layer is black and very dark gray silty clay loam about 31 inches thick. At a depth of 15 inches is a 9-inch layer of black and very dark gray silty clay loam that is mottled with yellowish brown. The subsoil is very dark gray and dark-gray silty clay loam about 13 inches thick. The underlying material, about 11 inches thick, is very dark gray clay loam that has lenses of dark-gray sand and gravel.

These soils are moderately permeable and have high available water capacity. Surface drainage is very slow to ponded, and internal drainage is very slow. The soils have low stability when wet. They are subject to frequent waterlogging and flooding. Seasonally, the water table is less than 2 feet below the surface.

These soils are suited to use for row crops if they are adequately drained. They have severe limitations for most other uses, because they are frequently saturated with water and are subject to periodic flooding.

Representative profile of Sawmill silty clay loam in an area of Sawmill soils, 27 feet north and 250 feet east of the NE. corner of bridge, in the SW $\frac{1}{4}$ SW $\frac{1}{4}$ NW $\frac{1}{4}$  sec. 30, T. 41 N., R. 5 E.

- Ap—0 to 9 inches, black (10YR 2/1) silty clay loam; weak, very fine, granular structure; firm; neutral; abrupt, smooth boundary.
- A11—9 to 15 inches, black (10YR 2/1) and some very dark gray (10YR 3/1) silty clay loam; weak, very fine, granular structure; firm; neutral; abrupt, smooth boundary.
- A12—15 to 24 inches, black (10YR 2/1) and very dark gray (5Y 3/1) silty clay loam; many, fine, distinct, yellowish-brown (10YR 5/6) mottles; weak, coarse, prismatic structure; firm; neutral; abrupt, smooth boundary.
- A13—24 to 31 inches, black (10YR 2/1) silty clay loam; weak, coarse, prismatic structure; firm; neutral; clear, smooth boundary.
- Bg—31 to 44 inches, very dark gray (10YR 3/1) and dark gray (10YR 4/1) silty clay loam; weak, coarse, prismatic structure; firm; some thin black (N 2/0) lenses; neutral; clear, smooth boundary.
- C—44 to 55 inches, very dark gray (10YR 3/1) clay loam; massive; firm; some dark-gray (10YR 4/1) lenses of sand and gravel; neutral.

The A horizon ranges from 24 to 36 inches in thickness. The B horizon ranges from 10 to 20 inches in thickness and from slightly acid to neutral. In places it is clay loam in the lower part, and in some places it contains strata of sand,

loam, or silt loam. The C horizon is clay loam or silty clay loam that generally contains strata of sand, loam, sandy loam, or silt loam.

Sawmill soils are associated with Drummer soils. They have a thicker A horizon than Drummer soils.

**Sawmill soils (V107).**—This nearly level soil is on the flood plains of the major streams. It is in the low-lying area that is subject to stream overflow on either side of the watercourse. The surface layer ranges from silt loam to silty clay loam.

Included with this soil in mapping were areas that are calcareous at the surface, areas where less clay occurs throughout the profile, areas where the surface layer is thinner, and some areas of soils that are less poorly drained. Also included were small areas of organic soils and areas covered by recent deposition.

Surface runoff is very slow to ponded, and the hazard of erosion is slight.

If artificially drained, this soil is suited to row crops. Some areas are used for pasture, and some remain in trees. The soil has severe limitations for most other uses. Management group IIw-1.

### Saybrook Series

The Saybrook series consists of deep, gently sloping to sloping, well drained and moderately well drained soils. These soils are on uplands away from streams and moraines. They formed in loess and in the underlying loam glacial till.

In a representative profile the surface layer is very dark brown and dark-brown silt loam about 12 inches thick. The subsoil is about 32 inches thick. The upper 13 inches of the subsoil is dark yellowish-brown silty clay loam, the next 13 inches is dark yellowish-brown and brown to dark-brown clay loam, and the lower 6 inches is brown to dark-brown loam. The underlying material is brown loam.

These soils are moderately permeable and have high available water capacity. The internal drainage is medium, and the soils remain stable with changes in moisture content. The water table is generally more than 5 feet below the surface.

These soils are well suited to row crops. They have moderate limitations for some other uses.

Representative profile of Saybrook silt loam, 2 to 4 percent slopes, 125 feet south and 150 feet west of the NE. corner of the NE $\frac{1}{4}$  sec. 17, T. 40 N., R. 3 E.

- A1—0 to 9 inches, very dark brown (10YR 2/2) silt loam; moderate, very fine and fine, granular structure; friable; slightly acid; clear, smooth boundary.
- A3—9 to 12 inches, dark-brown (10YR 3/3) silt loam; moderate, fine, granular structure; friable; slightly acid; gradual, smooth boundary.
- B1t—12 to 17 inches, dark yellowish-brown (10YR 3/4) light silty clay loam; moderate, fine and medium, granular structure; firm; discontinuous dark-brown (10YR 3/3) clay films; slightly acid; gradual, smooth boundary.
- B21t—17 to 25 inches, dark yellowish-brown (10YR 4/4) silty clay loam; moderate, very fine, subangular blocky structure; firm; discontinuous dark-brown (10YR 3/3) clay films; slightly acid; clear, smooth boundary.
- IIB22t—25 to 32 inches, dark yellowish-brown (10YR 4/4) clay loam; few, fine, faint, yellowish-brown (10YR 5/6) mottles; weak, fine and medium, prismatic structure parting to moderate, very fine and

- fine, subangular blocky; firm; discontinuous dark-brown (10YR 3/3) clay films; neutral; clear, smooth boundary.
- IIB31t—32 to 38 inches, brown to dark-brown (7.5YR 4/4) clay loam; weak, medium, prismatic structure parting to moderate, fine and medium, subangular blocky; firm; patchy very dark grayish-brown (10YR 3/2) clay films; many fine pores; mildly alkaline; clear, smooth boundary.
- IIB32—38 to 44 inches, brown to dark-brown (7.5YR 4/4) heavy loam; weak, medium and coarse, prismatic structure; friable; mildly alkaline; clear, smooth boundary.
- IIC—44 to 60 inches, brown (7.5YR 5/4) loam till; common, fine, faint, strong-brown (7.5YR 5/6 and 5/8) mottles; massive; friable; moderately alkaline.

The A horizon ranges from 9 to 14 inches in thickness. The B horizon ranges from 13 to 35 inches in thickness and from slightly acid to mildly alkaline. In some places the lower part is mottled. The IIB horizon is 10 to 20 inches thick.

Saybrook soils are associated with Flanagan and Catlin soils. They have a thinner mantle of loess over glacial till than Catlin and Flanagan soils; and they are better drained than Flanagan soils.

#### Saybrook silt loam, 2 to 4 percent slopes (145B).—

This soil is on ridgetops or side slopes on uplands. It has the profile described as representative of the series.

Included with this soil in mapping were areas where the surface layer is less than 10 inches thick. Also included were small areas of less sloping soils and small areas of Catlin soils.

Surface runoff is medium, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group Iie-2.

**Saybrook silt loam, 4 to 7 percent slopes, eroded (145C2).—**This soil is on uplands. It has a profile similar to the one described as representative of the series, but the dark yellowish-brown, finer textured subsoil has been mixed into the plow layer. Included with this soil in mapping were small areas where none of the original surface layer remains and small areas of La Rose soils.

Surface runoff is rapid, and the hazard of erosion is moderate.

This soil is intensively farmed and is well suited to row crops. Control of erosion is a necessary part of good management. This soil has moderate limitations for some other uses. Management group Iie-4.

### Strawn Series

The Strawn series consists of deep, strongly sloping to steep, well drained and moderately well drained soils. These soils are on the rolling uplands of the major glacial moraines and along the major drainageways. They formed in less than 2 feet of loess and in the underlying loam glacial till.

In a representative profile the surface layer is dark grayish-brown silt loam about 4 inches thick. The subsoil is brown to dark-brown heavy loam and clay loam about 15 inches thick. The underlying material is yellowish-brown loam.

These soils are moderately permeable and have moderate available water capacity. Internal drainage is medium, and the soils remain stable with changes in

moisture content. The water table is generally more than 5 feet below the surface.

These soils have low productivity and are poorly suited to row crops. Slope is the main limitation for other uses.

Representative profile of Strawn silt loam in an area of Strawn soils, 10 to 20 percent slopes, eroded, 120 feet north and 120 feet west of the NE. corner of the SE $\frac{1}{4}$ , SE $\frac{1}{4}$  sec. 34, T. 37 N., R. 5 E.

- Ap—0 to 4 inches, dark grayish-brown (10YR 4/2) silt loam; weak, very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- B1—4 to 8 inches, brown to dark-brown (10YR 4/3) heavy loam; weak, very fine, subangular blocky structure; friable; neutral; clear, smooth boundary.
- B21t—8 to 14 inches, brown to dark-brown (10YR 4/3) clay loam; moderate, very fine and fine, angular blocky structure; firm; discontinuous brown to dark-brown (7.5YR 4/2) clay films; neutral; clear, smooth boundary.
- B22t—14 to 19 inches, brown to dark-brown (10YR 4/3) clay loam; weak, medium, angular blocky structure; firm; discontinuous brown to dark-brown (7.5YR 4/2) clay films; mildly alkaline; abrupt, smooth boundary.
- C—19 to 60 inches, yellowish-brown (10YR 5/4) loam till; massive; friable; moderately alkaline.

The A horizon ranges from 4 to 8 inches in thickness. The B horizon ranges from 10 to 20 inches in thickness and from slightly acid to mildly alkaline.

Strawn soils are associated with Octagon and Miami soils. They have a thinner solum than Octagon and Miami soils.

#### Strawn soils, 10 to 20 percent slopes, eroded (224E2).—

This soil is on narrow breaks along streams or drainageways and around large depressions in the southeastern corner of the county. Included with this soil in mapping were areas of Miami and Dodge soils and small eroded spots where all of the original surface layer and subsoil have been removed through erosion.

Surface runoff is very rapid, and the hazard of erosion is severe.

Some areas of this soil are farmed, but it has severe limitations for intensive cropping. Control of erosion by either mechanical or vegetative means is a necessary part of good management. In places this soil is used for trees or grasses, and it is suited to these uses. It has limitations for most other uses. Management group VIe-1.

### Virgil Series

The Virgil series consists of deep, nearly level, somewhat poorly drained soils. These soils are in broad outwash areas and on circular, raised, flat-topped mounds on uplands. They formed in loess and in the underlying stratified fine sand and silt.

In a representative profile the surface layer is very dark gray silt loam about 8 inches thick. The sub-surface layer is dark grayish-brown silt loam about 3 inches thick. The subsoil is about 36 inches thick. The upper 28 inches of the subsoil is multicolored silty clay loam, and the lower 8 inches is light olive-gray silt loam. The subsoil is mottled with yellowish brown. The underlying material is multicolored gravelly loam and silt loam.

These soils are moderately permeable and have high

available water capacity. Surface drainage and internal drainage are slow. The soils are subject to concentrations of water during periods of high runoff, and they are subject to periodic waterlogging. Where these soils are wet, their stability decreases. In spring the water table is generally 1 foot to 3 feet below the surface.

These soils are well suited to intensive use for row crops. The periodic waterlogging is a limitation for other uses.

Representative profile of Virgil silt loam, 70 feet north and 50 feet east of the SW. corner of the NW $\frac{1}{4}$  sec. 26, T. 40 N., R. 3 E.

- Ap—0 to 8 inches, very dark gray (10YR 3/1) silt loam; moderate, very fine, granular structure; friable; neutral; abrupt, smooth boundary.
- A2—8 to 11 inches, dark grayish-brown (10YR 4/2) silt loam; weak, medium, platy structure; friable; slightly acid; abrupt, smooth boundary.
- B1—11 to 16 inches, dark grayish-brown (10YR 4/2) silty clay loam; weak, very fine, subangular blocky structure; friable; slightly acid; clear, smooth boundary.
- B21t—16 to 24 inches, dark grayish-brown (10YR 4/2) silty clay loam; few, faint, yellowish-brown (10YR 5/8) mottles; moderate, fine, subangular blocky structure; firm; slightly acid; clear, smooth boundary.
- B22t—24 to 32 inches, light brownish-gray (2.5Y 6/2) silty clay loam; common, fine, distinct, yellowish-brown (10YR 5/8) mottles; moderate, medium, subangular blocky structure; firm; thin grayish-brown (2.5Y 5/2) clay films; neutral; clear, smooth boundary.
- B23t—32 to 39 inches, light olive-gray (5Y 6/2) light silty clay loam; common, medium, distinct, yellowish-brown (10YR 5/8) mottles; weak, coarse, prismatic structure; friable; discontinuous grayish-brown (2.5Y 5/2) clay films; neutral; abrupt, smooth boundary.
- B31—39 to 43 inches, light olive-gray (5Y 6/2) silt loam; common, medium, distinct, yellowish-brown mottles; weak, coarse, prismatic structure; friable; mildly alkaline; gradual, smooth boundary.
- IIB32—43 to 47 inches, light olive-gray (5Y 6/2) silt loam that has a noticeable amount of sand; common, medium, distinct, yellowish-brown (10YR 5/8) mottles; weak, coarse, prismatic structure; friable; mildly alkaline; abrupt, smooth boundary.
- IIC1—47 to 52 inches, mixed light olive-gray (5Y 6/2), yellowish-brown (10YR 5/8), and brown to dark-brown (7.5YR 4/4) gravelly loam; massive; friable; moderately alkaline; abrupt, smooth boundary.
- IIC2—52 to 60 inches, mixed pale-olive (5Y 6/3) and strong-brown (7.5YR 5/8) silt loam; massive; friable; moderately alkaline.

The A horizon ranges from 6 to 11 inches in thickness. The B horizon ranges from 24 to 32 inches in thickness and from medium acid to mildly alkaline. The IIB horizon ranges from 4 to 10 inches in thickness.

Virgil soils are associated with Batavia and Drummer soils. They are more poorly drained than Batavia soils, and they are better drained than Drummer soils.

**Virgil silt loam** (104).—This nearly level soil is on the flat to slightly depressed centers of raised circular mounds or at the upper ends of waterways in outwash areas.

Included with this soil in mapping were small areas of poorly drained soils and small areas of soils that are calcareous at a depth of less than 40 inches. Also included were small areas where the silt is more than 5 feet deep.

Surface runoff is slow, and the hazard of erosion is slight.

This soil is intensively farmed and is well suited to row crops. Artificial drainage is beneficial in some areas. The soil has moderate to severe limitations for many other uses. Management group I-3.

### Will Series

The Will series consists of nearly level, poorly drained soils that are moderately deep over sand and gravel. These soils are in broad flat outwash areas in the northeastern corner of the county. They formed in loamy outwash material over calcareous sand and gravel.

In a representative profile the surface layer is very dark brown clay loam about 9 inches thick. The subsoil is mostly very dark gray clay loam about 19 inches thick. The underlying material is light brownish-gray to dark-gray gravelly sandy loam over gravelly sand.

These soils have a moderately permeable subsoil and have moderate available water capacity. Surface drainage is very slow to ponded, and internal drainage is very slow. The soils have moderate stability when wet. They are subject to frequent waterlogging and flooding or ponding during periods of runoff. Seasonally, the water table is less than 2 feet below the surface.

These soils are well suited to intensive use for row crops if they are adequately drained. The frequent waterlogging and potential for flooding are limitations for other uses.

Representative profile of Will clay loam, 420 feet south and 90 feet west of the NW. corner of the SE $\frac{1}{4}$  sec. 1, T. 42 N., R. 5 E.

- Ap—0 to 9 inches, very dark brown (10YR 2/2) light clay loam; weak, very fine, granular structure; friable; moderately alkaline; abrupt, smooth boundary.
- B1—9 to 15 inches, very dark gray (10YR 3/1) clay loam; weak, coarse, prismatic structure; firm; some strong-brown (7.5YR 5/6) iron stainings; moderately alkaline; gradual, smooth boundary.
- B21—15 to 20 inches, very dark gray (2.5YR 3/0) and strong-brown (7.5YR 5/6) clay loam; weak, coarse, prismatic structure; firm; moderately alkaline; gradual, smooth boundary.
- B22—20 to 28 inches, very dark gray (5Y 3/1) clay loam; common, fine, prominent, strong-brown (7.5YR 5/6) mottles; weak, coarse, prismatic structure; firm; moderately alkaline; clear, smooth boundary.
- IIC1—28 to 34 inches, light brownish-gray (2.5Y 6/2) gravelly sandy loam; massive; very friable; moderately alkaline; clear, smooth boundary.
- IIC2—34 to 40 inches, dark-gray (5Y 4/1) gravelly sandy loam; massive; very friable; moderately alkaline; clear, smooth boundary.
- IIC3—40 to 60 inches, dark-gray (5Y 4/1) gravelly sand; single grained; loose; moderately alkaline.

The A horizon ranges from 8 to 15 inches in thickness. The B horizon ranges from 15 to 25 inches in thickness. The upper part of the B horizon ranges from clay loam to silty clay loam, and the lower part ranges from clay loam to gravelly clay loam.

Will soils are associated with the better drained Proctor soils.

**Will clay loam** (329).—This nearly level soil is in a broad, flat area in the northeastern corner of the county. Included with it in mapping were small areas of somewhat poorly drained soils and some areas of soils that have a thinner subsoil than is typical for this soil and in most places are calcareous at the surface.

Surface runoff is very slow to ponded, and the hazard of erosion is slight.

If artificially drained, this soil is intensively farmed and is well suited to row crops. It has severe limitations for most other uses. Because of the shallowness to sand and gravel, pollution is a risk for some uses. Management group IIw-1.

## Use and Management of the Soils

This section discusses the general management of crops, explains the capability grouping used by the Soil Conservation Service, and discusses the management of the soils in DeKalb County by management (capability) groups. Predicted yields of the principal crops grown in the county under a high level of management are given. Also discussed are the use of soils for ornamental trees and shrubs and for wildlife. The properties and features that affect recreation and engineering practices are listed, mainly in tables.

### General Management of Crops

About 90 percent of DeKalb County is cultivated. Corn and soybeans are the principal crops. Oats and grass-legume hay are the other important crops. Several thousand acres are used each year for the production of hybrid seed corn and many kinds of vegetable crops for the canning industry.

The main consideration in managing cultivated soils in the county are controlling erosion, overcoming the hazard of wetness, protecting the soils from flooding, conserving moisture, and maintaining tilth and fertility.

Among the measures that help control erosion are terracing, contour farming, minimum tillage, cover crops, grassed waterways, and the use of crop residue. Generally, a combination of several measures is used.

Measures that help overcome wetness include the use of tile drains, inlets to tile drains, ditches, and diversions.

Conserving moisture generally means reducing evaporation, limiting runoff, increasing infiltration, and controlling weeds. Practices that help conserve moisture include minimum tillage, use of crop residue, contour farming, stripcropping, and use of field windbreaks.

Among the measures that help maintain tilth and fertility are the application of chemical fertilizer, green manure, and barnyard manure and the inclusion in the cropping system of cover crops, grasses, and legumes. Crops respond well to fertilizer on all soils used as cropland. Application of lime is needed periodically, especially on most of the soils on uplands. Control of erosion also helps conserve fertility and maintain good tilth.

### Capability Grouping

Capability grouping shows, in a general way, the suitability of soils for most kinds of field crops. The grouping is based on permanent limitations of the soils when used for field crops, the risk of damage

when they are used, and the way they respond to treatment. The grouping does not take into account major and generally expensive landforming that would change slope, depth, or other characteristics of the soils; does not take into consideration possible but unlikely major reclamation projects; and does not apply to rice, cranberries, horticultural crops, or other crops that require special management.

Those familiar with the capability classification can infer from it much about the behavior of soils when used for other purposes, but this classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for engineering.

In the capability system, all kinds of soils are grouped at three levels, the capability class, the subclass, and the unit (management group in this soil survey). These categories are discussed in the following paragraphs.

CAPABILITY CLASSES, the broadest groups, are designated by Roman numerals I to VIII. The numerals indicate progressively greater limitations and narrower choices for practical use, defined as follows:

Class I soils have few limitations that restrict their use.

Class II soils have moderate limitations that reduce the choice of plants or require moderate conservation practices.

Class III soils have severe limitations that reduce the choice of plants, require special conservation practices, or both.

Class IV soils have very severe limitations that reduce the choice of plants, require very careful management, or both.

Class V soils are not likely to erode but have other limitations, impractical to remove, that limit use largely to pasture, range, woodland, or wildlife habitat. (None in DeKalb County.)

Class VI soils have severe limitations that make them generally unsuited to cultivation and limit their use largely to pasture or range, woodland, or wildlife habitat.

Class VII soils have very severe limitations that make them unsuited to cultivation and restrict their use largely to pasture or range, woodland, or wildlife habitat. (None in DeKalb County.)

Class VIII soils and landforms have limitations that preclude their use for commercial plants and restrict their use to recreation, wildlife habitat, water supply, or esthetic purposes. (None in DeKalb County.)

CAPABILITY SUBCLASSES are soil groups within one class; they are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral; for example, IIe. The letter *e* shows that the main limitation is risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and *c*, used in only some parts of the United States but not in DeKalb County, shows

that the main limitation is climate that is too cold or too dry.

Class I contains no subclasses, because the soils of this class have few limitations. Class V can contain, at the most, only the subclasses indicated by *w*, *s*, and *c*, because the soils in class V are subject to little or no erosion, although they have other limitations that restrict their use largely to pasture, range, woodland, wildlife, habitat, or recreation.

MANAGEMENT GROUPS are soil groups within the subclass. The soils in one group are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity and other responses to management. Management groups are generally designated by adding an Arabic numeral to the subclass symbol; for example, IIw-1 or IIe-1.

The capability classification of the soils in DeKalb County is shown in the Guide to Mapping Units at the back of this survey.

#### *Management groups*

In the following pages the management groups in DeKalb County are described, and suggestions for use and management for all the soils of each group are given. On soils used for cultivated crops, applications of lime and fertilizer are generally needed. The amounts to apply on any given soil should be determined by soil tests. The names of soil series represented are mentioned in the description of each management group, but this does not mean that all soils of a given series appear in the group. To find the names of all the soils in any given management group, refer to the "Guide to Mapping Units" at the back of this survey.

#### MANAGEMENT GROUP I-1

This group consists of deep, well-drained soils of the Camden, Dodge, Rush, and St. Charles series. These soils are nearly level and uneroded. They have a surface layer of silt loam and a subsoil of silty clay loam.

Permeability is moderate, and the available water capacity is high. The organic-matter content is low.

Maintaining the organic-matter content and maintaining good tilth and fertility are the main management needs. Returning crop residue to the soil and plowing under a green-manure catch crop about every fourth year maintain the organic-matter content and good tilth. Minimum tillage reduces compaction and increases the infiltration of water. Winter cover crops or crop residue left on the surface help to control soil blowing.

The soils in this group are used mainly for cash-grain crops. They are well suited to the commonly grown crops, which are mainly corn, soybeans, oats, and alfalfa. They are also well suited to grasses.

#### MANAGEMENT GROUP I-2

This group consists of deep, well-drained soils of the Batavia, Bowes, Catlin, Harvard, Plano, and Proctor series. These uneroded soils have a surface layer of silt loam and a subsoil of silty clay loam.

Permeability is moderate, and the available water

capacity is high. The organic-matter content is medium to high.

Maintaining the organic-matter content and maintaining good tilth and fertility are slight management concerns. Returning crop residue to the soil and plowing under a green-manure catch crop about every fourth year are ways to maintain the organic-matter content and good tilth. Minimum tillage reduces compaction and increases infiltration. Winter cover crops or crop residue left on the surface help to control soil blowing.

The soils in this group are used mainly for cash-grain crops. They are well suited to the commonly grown crops, which are mainly corn, soybeans, oats, and alfalfa. They are also well suited to grasses.

#### MANAGEMENT GROUP I-3

This group consists of deep, somewhat poorly drained soils of the Elburn, Flanagan, Herbert, Lisbon, Kendall, Millbrook, Muscatine, and Virgil series. These soils are nearly level and uneroded. They have a surface layer of silt loam and a subsoil of silty clay loam.

Permeability is moderate, and the available water capacity is high. The organic-matter content is medium to high.

Maintaining the organic-matter content and maintaining good tilth and fertility are a necessary part of good management. In some areas artificial drainage is needed. Minimum tillage reduces compaction and increases water infiltration. Winter cover crops or crop residue left on the surface help to control soil blowing.

The soils in this group are used mainly for cash-grain crops. They are well suited to the commonly grown crops. Corn and soybeans are the main crops, but oats and alfalfa are also suited.

#### MANAGEMENT GROUP IIe-1

This group consists of deep, moderately well drained and well drained soils of the Camden, Dodge, Miami, Rush, and St. Charles series. These soils are gently sloping and uneroded. They have a surface layer of silt loam and a subsoil of silty clay loam.

Permeability is moderate, and the available water capacity is high. The organic-matter content is low.

Control of erosion, the major concern of management, can be done easily. It can be done by using a suitable cropping system or such mechanical practices as contouring and terracing. Grassed waterways remove excess surface water safely. Minimum tillage helps maintain the organic-matter content and good tilth and also helps control erosion. Tilth is harder to maintain in eroded areas. Winter cover crops or crop residue left on the surface help to control soil blowing.

The soils in this group are generally used for cash crops. Corn and soybeans are the main crops, but oats and alfalfa or grasses are also well suited.

#### MANAGEMENT GROUP IIe-2

This group consists of deep, moderately well drained and well drained soils of the Batavia, Bowes, Catlin, Harvard, La Rose, Octagon, Plano, Proctor, and Saybrook series. These soils are gently sloping and uneroded. They have a surface layer of silt loam and,

except for the La Rose soil, a subsoil of silty clay loam. The La Rose soil has a subsoil of clay loam.

Permeability is moderate, surface runoff is medium, and the available water capacity is high. The organic-matter content is medium to high.

Control of erosion, the major concern of management, can be done easily. It can be done by using a suitable cropping system or by such mechanical practices as contouring and terracing. Minimum tillage reduces compaction, increases water infiltration, and helps to control runoff and erosion. Winter cover crops or crop residue left on the surface help to control soil blowing.

The soils in this group are suited to crops commonly grown in the county. Corn and soybeans are the main crops, but oats and alfalfa or grasses are also suited.

#### MANAGEMENT GROUP IIe-3

Dresden silt loam, 2 to 4 percent slopes, the only soil in this group, is a well-drained, gently sloping, un-eroded soil that is moderately deep over sand and gravel. It has a surface layer of silt loam and a subsoil of clay loam.

Permeability is moderate in the subsoil and rapid to very rapid in the underlying sand and gravel. The available water capacity is moderate. The organic-matter content is medium. This soil is medium acid to slightly acid in the subsoil.

Controlling erosion is the main concern of management. Maintaining the organic-matter content and good tilth is a slight concern. If such conservation practices as terracing and contouring are not used, erosion should be controlled by the cropping system. The return of all crop residue to the soil maintains the organic-matter content and good tilth. Winter cover crops, manure, and crop residue left on the surface help to maintain the organic-matter content and good tilth. These practices along with minimum tillage, grassed waterways, and contouring help to control erosion and runoff.

Most areas of this soil are used for cultivated crops. This soil is suited to the commonly grown crops, such as corn and soybeans, but it is also suited to oats and grasses or legumes.

#### MANAGEMENT GROUP IIe-4

This group consists of deep, moderately well drained and well drained soils of the Batavia, Catlin, Dodge, Harvard, La Rose, Miami, Octagon, and Saybrook series. These soils are moderately sloping and eroded. They have a thin surface layer of silt loam and a subsoil of silty clay loam or clay loam.

Permeability is moderate, and surface runoff is rapid. The available water capacity is mainly high but is less than that of the soils in management groups IIe-1 and IIe-2 because of increased runoff and removal of the original surface layer by erosion. The organic-matter content is low to medium.

Controlling erosion is the major concern of management. The hazard of erosion is greater because of past erosion. Erosion can be controlled by using a suitable cropping system or such mechanical practices as contouring and terracing. Tilth is poor and is difficult to

maintain, especially where tillage extends into the subsoil. Minimum tillage reduces compaction, increases water infiltration, and helps to control runoff and erosion. Winter cover crops or crop residue left on the surface help to control soil blowing.

The soils in this group are suited to crops commonly grown in the county. Corn and soybeans are the main crops, but oats and alfalfa or grasses are also suited.

#### MANAGEMENT GROUP IIw-1

This group consists of mostly deep, poorly drained soils of the Drummer, Harpster, Knight, Sable, Sawmill, and Will series. These soils are nearly level. Except for the Will soil, they have a surface layer and subsoil of silty clay loam. The moderately deep Will soil has a surface layer and subsoil of clay loam.

Permeability is moderately slow in the Knight soil and moderate in the other soils. The available water capacity is moderate in the Will soil and high in the others. The organic-matter content is high. All of these soils are medium acid to moderately alkaline in the subsoil. The Harpster and Will soils are calcareous.

Providing drainage is the main management needed. Tile drains are effective where adequate outlets are available. Flooding is a hazard annually in most areas of the Sawmill soils, and ponding and slow surface runoff are limitations in some areas of the other soils. Shallow surface drains are beneficial in some of these areas. Winter cover crops or crop residue left on the surface help to control soil blowing.

Where drained, the soils in this group are well suited to crops grown in the county. Corn and soybeans are the main crops, but oats and grasses or legumes are also suited.

#### MANAGEMENT GROUP IIIe-1

This group consists of deep, well drained and moderately well drained soils of the La Rose and Miami series. These soils are strongly sloping and eroded. They have a surface layer of silt loam and a subsoil of clay loam. In some areas the present plow layer consists of material from the clay loam subsoil.

Permeability is moderate in both soils. The available water capacity is high in the Miami soil and moderate in the La Rose soil. The organic-matter content is low in the Miami soil and medium in the La Rose soil. These soils are medium acid to mildly alkaline in the subsoil.

Water erosion, the main hazard on these soils in cultivated areas, can be controlled by suitable cropping practices or such mechanical practices as contouring or terracing. Tilth is poor and is difficult to maintain, especially where tillage extends into the clay loam subsoil. Minimum tillage maintains the organic-matter content and tilth and helps to control erosion.

The soils in this group are not well suited to intensive row cropping, but they are suited to crops grown in the county. Corn and soybeans should be grown in rotation with grasses and legumes. These soils are suited to use for woodland.

#### MANAGEMENT GROUP IIIe-2

Dresden silt loam, 4 to 7 percent slopes, eroded, the only soil in this group, is a well-drained soil that is

moderately deep to sand and gravel. This soil is moderately sloping and eroded. It has a surface layer of silt loam and a subsoil of clay loam. In some areas the present plow layer consists of material from the clay loam subsoil.

Permeability and the available water capacity are moderate. The organic-matter content is medium. This soil is medium acid to neutral.

Water erosion, the main hazard on this soil in cultivated areas, can be controlled by suitable cropping practices or such mechanical practices as contouring or terracing. Tilth is poor and is difficult to maintain, especially where tillage extends into the clay loam subsoil. This soil has a hazard of droughtiness because the depth to sand and gravel is only moderate and because of the increased runoff caused by slope. Minimum tillage maintains the organic-matter content and the tilth, reduces runoff, and helps to control erosion.

This soil is not well suited to intensive row cropping. It is suited to the crops grown in the county. Corn and soybeans should be grown in rotation with grasses and legumes. This soil is suited to use for woodland.

#### MANAGEMENT GROUP IIIw-1

Peotone silty clay loam, the only soil in this group, is a very poorly drained, nearly level to depressional soil. It has a surface layer and subsoil of silty clay loam.

Permeability is moderately slow, and the available water capacity is high. The organic-matter content is high. This soil is neutral to mildly alkaline in the subsoil.

Wetness is a major hazard on this soil. Ponding and slow surface runoff are concerns of management in many areas. Shallow surface drains or surface inlet tile drains may help in these areas. Tile drains are beneficial where adequate outlets are available. Winter cover crops or crop residue left on the surface help to control soil blowing.

Where drained, this soil is well suited to crops grown in the county. Corn and soybeans are the main crops, but oats, grasses, and legumes are suited.

#### MANAGEMENT GROUP IIIw-2

Houghton muck, the only soil in this group, is a deep, very poorly drained, organic soil. This soil is nearly level to depressional. It formed in organic material.

Permeability is moderately rapid, and the available water capacity is high. The organic-matter content is high, but natural fertility is somewhat low because of deficiencies of mineral elements. Reaction is neutral throughout.

Providing drainage is the main concern of management. Tile lines are difficult to install and maintain. Ponding and slow surface runoff are other concerns of management on much of this soil. Shallow surface drains are beneficial in some areas. Suitable outlets for tile drains may be difficult to obtain. Overdrainage can result in a hazard of soil blowing or fire.

If adequately drained, this soil is suited to corn and soybeans. Special fertility practices are needed. This

soil is suited to vegetables and other specialty crops. It is not well suited to oats or hay.

#### MANAGEMENT GROUP IVe-1

Lorenzo loam, 6 to 15 percent slopes, eroded, the only soil in this group, is an excessively drained, strongly sloping, eroded soil that is shallow over sand and gravel. This soil has a surface layer of loam and a subsoil of clay loam or gravelly clay loam. In some areas the present plow layer consists mainly of subsoil material.

Permeability is moderately rapid, and the available water capacity is low. The organic-matter content is low. This soil is medium acid to neutral in the subsoil.

Water erosion is a hazard and the low available water capacity is a limitation in cultivated areas. Tilth is difficult to maintain, especially where tillage extends into the subsoil. Moisture frequently is insufficient for good crop growth.

This soil is poorly suited to cultivated crops because of droughtiness and the hazard of erosion. Where erosion is controlled and fertility and tilth are maintained, most crops common to the county can be grown. This soil is suited to trees, and some areas remain in native woods.

#### MANAGEMENT GROUP VIe-1

Strawn soils, 10 to 20 percent slopes, eroded, the only soil in this group, are deep, well drained and moderately well drained soils. They are moderately steep and eroded. They have a surface layer of silt loam and a subsoil of clay loam. In some areas the present plow layer consists mainly of subsoil material.

Permeability and the available water capacity are moderate. The organic-matter content is low. This soil is slightly acid to mildly alkaline in the subsoil.

Water erosion is a severe hazard, and slope is also a limiting factor. Water runoff is rapid, especially in areas that have poor cover of vegetation.

The soil in this group is not suited to cultivated crops. It is suited to use for woodland, and some areas remain in native woods. Some areas are used for pasture, but moderately steep slopes make maintenance of pasture difficult and the operation of machinery dangerous.

### Predicted Yields

Table 4 shows predicted yields of the principal crops grown on arable soils in DeKalb County under a high level of management. These predictions are based on yields for the period 1954-63, on soil tests, and on the experience and records of farmers, agronomists, conservationists, and farm advisers (9). The predictions are adjusted to reflect the trend toward higher yields during the period 1963-68. Average yields are expected to increase. A few farmers obtain yields as high as 200 bushels of corn per acre in some years, but yields this high are still uncommon.

Management was determined on the basis of farming techniques, crop varieties, and fertilizers commonly

TABLE 4.—*Predicted average acre yields of principal crops*

[Yields are those to be expected under a high level of management. Absence of a yield figure indicates that the crop is not suited to the soil or is not commonly grown on it]

Soil	Corn	Soybeans	Alfalfa	Pasture
	Bu	Bu	Tons	AUD <sup>1</sup>
Batavia silt loam, 0 to 2 percent slopes	120	42	5.2	260
Batavia silt loam, 2 to 4 percent slopes	120	42	5.2	260
Batavia silt loam, 4 to 7 percent slopes, eroded	110	38	4.8	240
Bowes silt loam, 0 to 2 percent slopes	120	40	5.0	250
Bowes silt loam, 2 to 4 percent slopes	120	40	5.0	250
Camden silt loam, 0 to 2 percent slopes	110	38	5.0	250
Camden silt loam, 2 to 6 percent slopes	110	38	5.0	250
Catlin silt loam, 0 to 2 percent slopes	130	45	5.5	275
Catlin silt loam, 2 to 4 percent slopes	130	45	5.5	275
Catlin silt loam, 4 to 7 percent slopes, eroded	115	40	5.0	250
Dodge silt loam, 0 to 2 percent slopes	105	38	4.8	240
Dodge silt loam, 2 to 4 percent slopes	105	38	4.8	240
Dodge silt loam, 4 to 7 percent slopes, eroded	95	35	4.2	210
Dresden silt loam, 2 to 4 percent slopes	90	32	3.8	190
Dresden silt loam, 4 to 7 percent slopes, eroded	80	28	3.5	175
Drummer silty clay loam	130	45	3.4	170
Elburn silt loam	140	48	5.8	290
Flanagan silt loam	140	48	5.8	290
Harpster silty clay loam	120	42	4.5	225
Harvard silt loam, 0 to 2 percent slopes	115	40	5.0	250
Harvard silt loam, 2 to 4 percent slopes	115	40	5.0	250
Harvard silt loam, 4 to 7 percent slopes, eroded	105	35	4.5	225
Herbert silt loam	120	42	5.0	250
Houghton muck	110	38		
Kendall silt loam	115	40	5.0	250
Knight silt loam	110	38	4.2	210
La Rose silt loam, 2 to 4 percent slopes	110	38	5.0	250
La Rose silt loam, 4 to 7 percent slopes, eroded	100	35	4.5	225
La Rose silt loam, 7 to 12 percent slopes, eroded	95	32	4.2	210
Lisbon silt loam	130	45	5.5	275
Lorenzo loam, 6 to 15 percent slopes, eroded	65	22	2.5	125
Miami silt loam, 2 to 4 percent slopes	100	35	4.5	225
Miami silt loam, 4 to 7 percent slopes, eroded	90	32	4.0	200
Miami silt loam, 7 to 12 percent slopes, eroded	85	30	3.8	190
Millbrook silt loam	125	45	5.2	260
Muscatine silt loam	140	48	5.8	290
Octagon silt loam, 2 to 4 percent slopes	115	38	5.0	250
Octagon silt loam, 4 to 7 percent slopes, eroded	105	34	4.6	230
Peotone silty clay loam	105	38	3.5	175
Plano silt loam, 0 to 2 percent slopes	135	48	5.5	275
Plano silt loam, 2 to 4 percent slopes	135	48	5.5	275
Proctor silt loam, 0 to 2 percent slopes	125	45	5.2	260
Proctor silt loam, 2 to 4 percent slopes	125	45	5.2	260
Rush silt loam, 0 to 2 percent slopes	115	38	5.0	250
Rush silt loam, 2 to 4 percent slopes	115	38	5.0	250
Sable silty clay loam	130	45	5.2	260
St. Charles silt loam, 0 to 2 percent slopes	115	40	4.8	240
St. Charles silt loam, 2 to 4 percent slopes	115	40	4.8	240
Sawmill soils	110	38	4.2	210
Saybrook silt loam, 2 to 4 percent slopes	120	42	5.2	260
Saybrook silt loam, 4 to 7 percent slopes, eroded	110	38	4.8	240
Strawn soils, 10 to 20 percent slopes, eroded			3.0	150
Virgil silt loam	125	45	5.2	260
Will clay loam	105	38	4.5	225

<sup>1</sup> AUD is animal-unit-days, a term used to express the carrying capacity of pasture. It is the number of days 1 acre can carry 1 animal unit during a single grazing season without injury to the sod. One animal unit is defined as 1 cow, 2 yearling calves, 1 horse, 7 sheep, or 4 brood sows. For example, 8 calves can graze about 25 days in a pasture that has a capacity of 100 animal-unit-days.

used in 1968. Differences in weather from year to year may result in annual yields that range 20 percent above or below those shown in the table. Hay and pasture yields are predicted for varieties of legumes adapted to the soil.

Under a high level of management, adequate drainage, control of flooding, and control of erosion are provided; the proper number of plants is grown; high-quality seed is used; tillage is kept to a minimum and is done when soil moisture is favorable; weeds, plant diseases, and harmful insects are controlled; favorable soil reaction and near-optimum levels of nitrogen, phosphorus, and potassium are maintained; efficient use is

made of available crop residue, barnyard manure, and green-manure crops; crops are harvested with the smallest possible loss; the combination of practices used is efficient; and all operations are timely.

### Use of the Soils for Ornamental Trees and Shrubs

Use of the soils for woodland is not important in DeKalb County. Most trees are planted for their ornamental, shade, or esthetic value. Not all trees grow equally well on all soils. Many trees cannot survive on certain kinds of soil, for example, wet soils. To get optimum value from plantings and to increase their survival rate, a planting guide for trees and shrubs has

TABLE 5.—Guide for tree and shrub plantings

Tree and shrub planting group, soil series, and map symbols	Tree species	Shrub species
<p>Group 1. Silt loams; well drained and moderately well drained; moderate permeability in the subsoil; reaction ranges from medium acid to neutral. Roots penetrate these soils easily.</p> <p>Batavia: 105A, 105B, 105C2            Bowes: 792A, 792B            Camden: 134A, 134B            Catlin: 171A, 171B, 171C2            Dodge: 24A, 24B, 24C2            Dresden: 325B, 325C2            Harvard: 344A, 344B, 344C2            Plano: 199A, 199B            Proctor: 148A, 148B            Rush: 791A, 791B            St. Charles: 243A, 243B            Saybrook: 145B, 145C2</p>	<p>Yellow-poplar, black walnut, white pine, white oak, red oak, Austrian pine, sugar maple, white ash, hackberry, Kentucky coffeetree, Norway spruce, Douglas-fir, red pine, Scotch pine.</p>	<p>Witchhazel, aromatic sumac, winged sumac, smooth sumac, staghorn sumac, hazel, Amur maple, autumn-olive, gray dogwood.</p>
<p>Group 2. Silt loams; well drained and moderately well drained; generally moderately well drained; generally moderate permeability in the subsoil, but the Lorenzo soil has moderately rapid permeability; reaction ranges from medium acid to neutral. Free carbonates occur at a depth of less than 40 inches, commonly at a depth of 15 to 30 inches; root development may be restricted to the zone above the carbonates.</p> <p>La Rose: 60B, 60C2, 60D2            Lorenzo: 318D2            Miami: 27B, 27C2, 27D2            Octagon: 656B, 656C2            Strawn: 224E2</p>	<p>Black walnut, white oak, red oak, black maple or sugar maple, white pine, Austrian pine, American basswood, white ash, Kentucky coffeetree, hackberry, bur oak, shingle oak, Scotch pine.</p>	<p>Witchhazel, aromatic sumac, winged sumac, smooth sumac, staghorn sumac, hazel, Amur maple, autumn-olive, gray dogwood.</p>
<p>Group 3. Silt loams; somewhat poorly drained; moderate permeability in the subsoil; reaction ranges from slightly acid to neutral. These soils are subject to periodic waterlogging; water table seasonally at a depth of 1 to 3 feet.</p> <p>Elburn: 198            Flanagan: 154            Herbert: 62            Kendall: 242            Lisbon: 59            Millbrook: 219            Muscatine: 41            Virgil: 104</p>	<p>Red maple, green ash, white pine, bur oak, shingle oak, swamp white oak, American hornbeam, American basswood, redcedar, Austrian pine, northern red oak, pin oak.</p>	<p>Silky dogwood, indigobush, autumn-olive, wild plum, gray dogwood, coralberry, black haw.</p>

TABLE 5.—Guide for tree and shrub plantings—Continued

Tree and shrub planting group, soil series, and map symbols	Tree species	Shrub species
<p>Group 4. Silty clay loams; poorly drained; moderate to moderately slow permeability in the subsoil; reaction is neutral. These soils are subject to frequent or continuous waterlogging unless tile drained, and to flooding or ponding; water table seasonally at a depth of less than 2 feet.</p> <p>Drummer: 152                      Knight: 191                      Peotone: 330                      Sable: 68                      Sawmill: V107                      Will: 329</p>	<p>Pin oak, swamp white oak, green ash, eastern cottonwood, shingle oak, black ash, silver maple, European alder, black willow, tamarack, river birch.</p>	<p>Winterberry, wax myrtle, red-osier dogwood, groundcherry, Tatarian honeysuckle, highbush cranberry, Siberian crabapple.</p>
<p>Group 4A: Same as group 4, except calcareous at the surface and free carbonates throughout the soil; reaction ranges from mildly alkaline to moderately alkaline.</p> <p>Harpster: 67</p>	<p>Black willow, eastern cottonwood, flowering dogwood, tamarack.</p>	<p>Medium purple willow, Lambert purple willow, dwarf purple willow.</p>
<p>Group 5. Mucks and peats; very poorly drained. These soils are frequently ponded and frequently or continuously saturated with water; water table seasonally at a depth of less than 1 foot.</p> <p>Houghton: 103</p>	<p>Leave in natural state -----</p>	<p>Leave in natural state. Medium purple willow, Lambert purple willow, dwarf purple willow.</p>

been developed for the soils of DeKalb County. Table 5 lists the species to be favored for the different kinds of soil. This list is not intended to include all known species, but rather to represent some of the common species that are grown or can be grown in the county. The tree and shrub planting group in which each soil has been placed is listed in the "Guide to Mapping Units" at the back of this survey.

**Use of the Soils for Wildlife**

The soils in DeKalb County are used mainly for the production of farm crops. However, the soil resources provide good potential for development of wildlife habitat.

Three major kinds of wildlife are recognized in DeKalb County. These are openland wildlife, woodland wildlife, and wetland wildlife. The potential is high for developing habitat for openland wildlife and woodland wildlife throughout most of the county. Only small, localized areas have a suitable potential for kinds of wildlife that prefer wetland habitat. The three major kinds of wildlife are defined as follows:

**Openland wildlife** are birds, mammals, and reptiles that normally frequent cropland, pasture, and hayland that is overgrown with grasses, herbs, and shrubs. Examples of openland wildlife are rabbit, red fox,

skunk, quail, and meadowlark. The ratings for elements of wildlife habitat are based on the ability of the soils to grow grain and seed crops, grasses and legumes, wild herbaceous plants, and hardwood plants.

**Woodland wildlife** are mammals and birds that frequent areas of hardwood and coniferous trees, shrubs, or a combination of the vegetation. Examples of woodland wildlife are squirrel, deer, raccoon, woodpecker, and nuthatch. The ratings for elements of wildlife habitat are based on the ability of the soils to grow grasses and legumes, wild herbaceous plants, hardwood plants, and coniferous plants.

**Wetland wildlife** are mammals, birds, and reptiles that frequent wet areas, such as ponds, marshes, and swamps. Examples of wetland wildlife are muskrat, wild duck and geese, kingfisher, and redwing blackbird. The ratings for elements of wildlife habitat are based on the ability of the soils to grow wetland food and cover plants and grain and seed crops and limitations for shallow-water developments and excavated ponds.

The soils of DeKalb County have been placed in three wildlife groups. In table 6, they are rated for their suitability for the improvement, maintenance, or creation of specific wildlife habitat elements and kinds of wildlife. The wildlife habitat elements rated are grain and seed crops, domestic grasses and legumes, wild herbaceous plants, hardwood plants, wetland plants, and shallow-water areas.

TABLE 6.—Suitability of the soils for elements

Wildlife groups, soil series, and map symbols	Elements of wildlife habitat		
	Grain and seed crops	Grasses and legumes	Wild herbaceous plants
<p>Group 1. Well drained and moderately well drained, nearly level to moderately steep soils on uplands and terraces.</p> <p>Batavia: 105A, 105B, 105C2.  Bowes: 792A, 792B.  Camden: 134A, 134B.  Catlin: 171A, 171B, 171C2.  Dodge: 24A, 24B, 24C2.  Dresden: 325B, 325C2.  Harvard: 344A, 344B, 344C2.  La Rose: 60B, 60C2, 60D2.  Lorenzo: 318D2.  Miami: 27B, 27C2, 27D2.  Octagon: 656B, 656C2.  Plano: 199A, 199B.  Proctor: 148A, 148B.  Rush: 791A, 791B.  St. Charles: 243A, 243B.  Saybrook: 145B, 145C2.  Strawn: 224E2.</p>	<p>Good where slopes are 0 to 7 percent.  Fair where slopes are more than 7 percent.</p>	<p>Good, except the Strawn soil is fair where slopes are more than 12 percent.</p>	<p>Good.....</p>
<p>Group 2. Somewhat poorly drained, nearly level soils on uplands and terraces.</p> <p>Elburn: 198.  Flanagan: 154.  Herbert: 62.  Kendall: 242.  Lisbon: 59.  Millbrook: 219.  Muscatine: 41.  Virgil: 104.</p>	<p>Good.....</p>	<p>Good.....</p>	<p>Good.....</p>
<p>Group 3. Poorly drained and very poorly drained, nearly level to depressional soils on uplands, terraces, and bottom lands.</p> <p>Drummer: 152.  Harpster: 67.  Houghton: 103.  Knight: 191.  Peotone: 330.  Sable: 68.  Sawmill: V107.  Will: 329.</p>	<p>Good where the soils are drained.  Poor where the soils are not drained: wetness hazard; flooding or ponding limits growth.  Very poor for the Houghton soil.</p>	<p>Good where the soils are drained.  Poor where the soils are not drained: wetness hazard; flooding or ponding limits growth.  Very poor for the Houghton soil.</p>	<p>Good where the soils are drained.  Fair where the soils are not drained: wetness hazard; flooding or ponding limits growth.  Poor for the Houghton soil.</p>

### Recreational Uses of the Soils

In table 7 the soils of DeKalb County are placed in groups, and the degree of limitation for and the soil features affecting recreational uses are given. The degree of limitation for each group is based on soil characteristics that affect use, such as natural drainage, seasonal high water table, flooding hazard, permeability, slope, texture of the surface layer, and stoniness or rockiness. The "Guide to Mapping Units" at the back of this survey lists the recreational group in which each soil has been placed.

The ratings used are *slight*, *moderate*, or *severe*. A rating of *slight* means that the soil has few or no limitations for the use specified in the table or that the limitations can be easily overcome. A rating of *moderate* indicates that the limitations can be overcome by careful planning and maintenance. A rating of *severe* indicates that the soil is poorly suited to the

use specified or that the limitations can be overcome only by intensive, costly engineering practices. The soil properties that determine moderate and severe limitations are shown in table 7. The recreational uses given in the table are discussed in the following paragraphs.

**Utility buildings.**—Among these are washrooms and bathrooms, picnic shelters, and service buildings that are used seasonally or all year. The degree of limitation is based mainly on soil features that contribute to the adequate support of these structures. Information on soil limitations for septic tank filter fields is in the section "Engineering Uses of the Soils."

**Campsites.**—These are areas suitable for tents and trailers and for living outdoors for a period of 1 week or longer. Little site preparation should be required. The soils are rated according to their limitations for unsurfaced parking areas for cars and camp trailers and for heavy traffic by people, horses, and such small vehicles as bicycles.

*of wildlife habitat and kinds of wildlife*

Elements of wildlife habitat—Continued			Kinds of wildlife		
Hardwood plants	Wetland plants	Shallow-water developments	Openland	Woodland	Wetland
Good-----	Very poor: no suitable plant species for food and cover.	Very poor: water table too deep.	Good, except the Strawn soil is fair.	Good, except the Lorenzo and Strawn soils are fair.	Very poor: no suitable wetland food and cover plants; water supply limited.
Good-----	Fair: limited number of suitable species for food and cover.	Fair: water table not high enough to maintain water level all year.	Good-----	Good-----	Fair: water supply and number of suitable wetland food and cover plants moderately limited.
Good where the soils are drained. Fair where the soils are not drained. Poor for the Houghton soil.	Poor where the soils are drained: number of suitable species limited. Good where the soils are not drained.	Fair where the soils are drained: water supply limited. Good where the soils are not drained: soils on bottom lands are subject to flooding.	Good where the soils are drained. Fair where the soils are not drained: grain and seed crops severely limited.	Good where the soils are drained. Fair where the soils are not drained: growth of food and cover plants is limited.	Poor where the soils are drained: number of suitable wetland food and cover plants severely limited. Good where the soils are not drained.

**Picnic areas.**—Soils used for picnic areas support intensive foot traffic. Features that affect the desirability of a site, such as trees or ponds, are not considered in the ratings.

**Playgrounds.**—These areas are developed for intensive play and for organized games, such as baseball, football, and tennis. They are subject to intensive foot traffic.

**Paths and trails.**—Soils used for paths and trails support intensive traffic by people on foot or on horseback. Little preparation is required. Paths and trails on sloping soils should be contoured to control erosion.

**Golf fairways.**—Only limitations for fairways have been considered. Greens, traps, and hazards are generally made from transported soil material. Soils used for fairways should support intensive traffic by people on foot or driving golf carts. Turf and various kinds of trees and shrubs grow well on these soils.

**Engineering Uses of the Soils <sup>3</sup>**

This section is useful to those who need information about soils used as structural material or as foundation upon which structures are built. Among those who can benefit from this section are planning commissions, town and city managers, land developers, engineers, contractors, and farmers.

Among the soil properties that are highly important in engineering are permeability, strength, compaction characteristics, soil drainage, shrink-swell potential, grain-size distribution, plasticity, and soil reaction. Also important are depth to the water table, depth to bedrock, and soil slope. These properties, in various degrees and combinations, affect construction and maintenance of roads, airports, pipelines, foundations

<sup>3</sup> JOHN T. WHELAN, engineer, Soil Conservation Service, helped prepare this section.

TABLE 7.—*Recreational*

Recreational group, soil series, and map symbols	Degree of limitation and soil features affecting use for—	
	Utility buildings	Campsites
<b>Group 1. Well drained and moderately well drained, nearly level to sloping soils on uplands.</b> Batavia: 105A, 105B, 105C2. Bowes: 792A, 792B. Camden: 134A, 134B. Catlin: 171A, 171B, 171C2. Dodge: 24A, 24B, 24C2. Dresden: 325B, 325C2. Harvard: 344A, 344B, 344C2. La Rose: 60B, 60C2. Miami: 27B, 27C2. Octagon: 656B, 656C2. Plano: 199A, 199B. Proctor: 148A, 148B. Rush: 791A, 791B. St. Charles: 243A, 243B. Saybrook: 145B, 145C2.	Slight -----	Slight -----
<b>Group 2. Well drained and moderately well drained, strongly sloping to steep soils on uplands.</b> La Rose: 60D2. Lorenzo: 318D2. Miami: 27D2. Strawn: 224E2.	Moderate for 60D2, 318D2, and 27D2: slope. Severe for 224E2: slope-----	Moderate where slopes are 7 to 12 percent. Severe where slopes are more than 12 percent: bare soil is slippery and sticky when wet.
<b>Group 3. Somewhat poorly drained, nearly level soils on uplands.</b> Elburn: 198. Flanagan: 154. Herbert: 62. Kendall: 242. Lisbon: 59. Millbrook: 219. Muscatine: 41. Virgil: 104.	Moderate: seasonal water table at a depth of 1 foot to 3 feet.	Moderate: seasonal water table at a depth of 1 foot to 3 feet.
<b>Group 4. Poorly drained, nearly level and depressional soils on uplands and bottom lands.</b> Drummer: 152. Harpster: 67. Knight: 191. Peotone: 330. Sable: 68. Sawmill: V107. Will: 329.	Severe: seasonal water table near surface; subject to ponding; soils dry slowly; V107 subject to flooding.	Severe: seasonal water table near surface; subject to ponding; soils dry slowly; bare soils are slippery and sticky when wet; V107 subject to flooding.
<b>Group 5. Very poorly drained, nearly level to depressional, organic soils.</b> Houghton: 103.	Very severe: seasonal water table at or near the surface; subject to ponding; unstable; high compressibility; subject to subsidence when drained.	Very severe: seasonal water table at or near the surface; subject to ponding; unstable; high compressibility; subject to subsidence when drained; bare soil subject to blowing.

for small buildings, irrigation systems, ponds and small dams, and systems for disposal of sewage and refuse.

Information in this section of the soil survey can be helpful to those who—

1. Select potential residential, industrial, commercial, and recreational areas.
2. Evaluate alternate routes for roads, highways, pipelines, and underground cables.
3. Seek sources of gravel, sand, or clay.
4. Plan farm drainage systems, irrigation systems, ponds, terraces, and other structures for controlling water and conserving soil.

5. Correlate performance of structures already built with properties of the kinds of soil on which they are built, for the purpose of predicting performance of structures on the same or similar kinds of soil in other locations.
6. Predict the trafficability of soils for cross-country movement of vehicles and construction equipment.
7. Develop preliminary estimates pertinent to construction in a particular area.

Most of the information in this section is presented in tables. Table 8 shows several estimated soil proper-

*uses of the soils*

Degree of limitation and soil features affecting use for—Continued			
Picnic areas	Playgrounds	Paths and trails	Golf fairways
Slight	Slight where slopes are 0 to 2 percent. Moderate where slopes are 2 to 7 percent.	Slight	Slight.
Moderate where slopes are 7 to 12 percent. Severe where slopes are more than 12 percent: bare soil is slippery and sticky when wet; 318D2 is droughty.	Severe: slopes of more than 7 percent; 318D2 is droughty.	Moderate: bare soil is slippery and sticky when wet.	Moderate: slopes of more than 7 percent; difficult to maintain good turf on 318D2 because of droughtiness.
Moderate: seasonal water table at a depth of 1 foot to 3 feet.	Moderate: seasonal water table at a depth of 1 foot to 3 feet.	Moderate: seasonal water table at a depth of 1 foot to 3 feet.	Moderate: seasonal water table at a depth of 1 foot to 3 feet.
Severe: seasonal water table near surface; subject to ponding; soils dry slowly; bare soils are slippery and sticky when wet; V107 subject to flooding.	Severe: seasonal water table near surface; subject to ponding; soils dry slowly; bare soils are slippery and sticky when wet; V107 subject to flooding.	Severe: seasonal water table near surface; subject to ponding; soils dry slowly; bare soils are slippery and sticky when wet; V107 subject to flooding.	Severe: seasonal water table near surface; subject to ponding; soils dry slowly; turf damaged easily when wet; V107 subject to flooding.
Very severe: seasonal water table at or near the surface; subject to ponding; unstable; high compressibility; subject to subsidence when drained; bare soil subject to blowing.	Very severe: seasonal water table at or near the surface; subject to ponding; unstable; high compressibility; subject to subsidence when drained; bare soil subject to blowing.	Very severe: seasonal water table at or near the surface; subject to ponding; unstable; high compressibility; subject to subsidence when drained.	Very severe: seasonal water table at or near the surface; subject to ponding; unstable; high compressibility; subject to subsidence when drained; bare soil subject to blowing.

ties significant in engineering; table 9 gives interpretations for various engineering uses; and table 10 gives the results of engineering laboratory tests on soil samples.

This information, along with the soil map and other parts of this publication, can be used to make interpretations in addition to those given in tables 8 and 9, and it also can be used to supplement information obtained from other published maps, records, and aerial photographs for the purpose of making maps and reports that can be used readily by engineers.

This information, however, does not eliminate the need for further investigations at sites selected for

engineering works, especially works that involve heavy loads or require excavations to depths greater than those shown in the tables, generally depths of more than 6 feet. Also, inspection of sites, especially the small ones, is needed because many delineated areas of a given soil mapping unit may contain small areas of other kinds of soil that have strongly contrasting properties and different suitabilities or limitations for soil engineering.

Some of the terms used in this soil survey have special meaning in soil science that is not used in engineering. The Glossary defines many of these terms as they are commonly used in soil science.

TABLE 8.—*Estimated soil properties*

[The symbol &gt; means more than;

Soil series and map symbols	Depth of seasonal high water table	Depth from surface	USDA texture	Classification	
				Unified	AASHTO
	<i>Feet</i>	<i>Inches</i>			
Batavia: 105A, 105B, 105C2-----	>5	0-10 10-42 42-60	Silt loam----- Silty clay loam----- Silt loam, loam-----	CL or ML CL ML or SM	A-6 or A-4 A-6 A-2 or A-4
Bowes: 792A, 792B-----	>5	0-7 7-37 37-47 47-60	Silt loam----- Silty clay loam----- Gravelly sandy clay loam, loam. Stratified sand and gravel.	ML or CL CL SM or CL SP, GP, SW, or GW	A-4 or A-6 A-6 or A-7 A-2 or A-6 A-1
Camden: 134A, 134B-----	>5	0-8 8-29 29-55	Silt loam----- Silty clay loam----- Stratified clay loam, silt loam.	ML or CL CL SM, SC, or CL	A-4 A-6 A-2, A-4, or A-6
Catlin: 171A, 171B, 171C2-----	>5	0-11 11-43 43-70	Silt loam----- Silty clay loam----- Clay loam, loam-----	CL, ML, or OL CL CL	A-7, A-4, or A-6 A-6 or A-7 A-4
Dodge: 24A, 24B, 24C2-----	>5	0-9 9-25 25-31 31-60	Silt loam----- Silty clay loam----- Clay loam----- Loam-----	ML or CL CL CL CL	A-4 or A-6 A-6 or A-7 A-6 or A-7 A-4 or A-6
Dresden: 325B, 325C2-----	>5	0-7 7-29 29-38 38-60	Silt loam----- Silty clay loam, clay loam. Gravelly clay loam----- Stratified sand and gravel.	ML or CL CL SM or CL GP or SP	A-4 or A-6 A-6 or A-7 A-2, A-4, or A-6 A-1
Drummer: 152-----	>2	0-18 18-50 50-78	Silty clay loam----- Silty clay loam----- Silt loam, sandy clay loam.	CL or OH CL ML, CL, SC	A-7 A-6 or A-7 A-2, A-4, or A-6
Elburn: 198-----	1-3	0-14 14-48 48-60	Silt loam, silty clay loam. Silty clay loam----- Clay loam-----	ML or CL CL CL	A-4 or A-6 A-6 or A-7 A-6 or A-4
Flanagan: 154-----	1-3	0-16 16-35 35-60	Silt loam----- Silty clay loam----- Loam, silt loam-----	CL, ML, MH, or OL CL or CH CL or ML	A-7, A-4, or A-6 A-6 or A-7 A-4 or A-6
Harpster: 67-----	>2	0-12 12-30 30-58	Silty clay loam----- Silty clay loam----- Silt loam, loam-----	CL or OH CL ML or CL	A-7 A-6 or A-7 A-4 or A-6
Harvard: 344A, 344B, 344C2-----	>5	0-12 12-27 27-68	Silt loam----- Silty clay loam----- Stratified clay loam, loam, and sand.	ML or CL CL CL, SM, SC	A-4 or A-6 A-6 or A-7 A-2, A-4, A-6, or A-7
Herbert: 62-----	1-3	0-12 12-26 26-36 36-58	Silt loam----- Silty clay loam----- Clay loam----- Loam-----	ML or CL CL CL CL	A-4 or A-6 A-6 or A-7 A-6 or A-7 A-4

significant in engineering  
the symbol < means less than]

Percentage less than 3 inches passing sieve <sup>1</sup> —			Liquid limit	Plasti- city index	Permea- bility	Available water capacity	Reaction	Shrink-swell potential
No. 4 (4.7 mm)	No. 10 (2.0 mm)	No. 200 (0.074 mm)						
95-100	90-100	80-100	30-40	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
95-100	90-100	85-100	40-55	20-35	0.6-2.0	0.18-0.20	5.6-7.3	Moderate.
90-100	80-90	25-60	15-25	5-15	0.6-6.3	0.11-0.19	7.4-8.4	Low.
100	100	80-95	30-40	10-15	0.6-2.0	0.22-0.24	6.6-7.3	Low.
100	95-100	70-95	30-50	15-30	0.6-2.0	0.18-0.20	5.1-6.5	Moderate.
80-100	65-90	25-65	20-30	10-20	2.0-6.3	0.10-0.16	6.1-7.3	Low.
40-80	30-70	0-10	-----	NP	6.3-20	0.04-0.06	7.4-8.4	Low.
100	95-100	80-95	30-40	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
95-100	95-100	60-90	40-55	20-35	0.6-2.0	0.16-0.20	5.6-6.5	Moderate.
90-100	80-95	30-80	30-55	NP-15	0.6-6.3	0.12-0.16	6.6-7.3	Low.
100	100	95-100	30-45	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
100	95-100	95-100	40-55	20-35	0.6-2.0	0.19-0.21	5.6-6.5	Moderate.
95-100	85-95	55-95	30-55	5-15	0.2-1.0	0.16-0.20	6.6-8.4	Low.
100	95-100	90-100	30-40	10-15	0.6-2.0	0.19-0.23	6.6-7.3	Low.
95-100	90-100	80-95	40-55	20-35	0.6-2.0	0.16-0.19	5.1-5.5	Moderate.
95-100	90-100	55-75	30-55	15-30	0.6-2.0	0.16-0.19	6.1-6.5	Moderate.
95-100	85-95	55-75	15-25	5-15	0.2-1.0	0.14-0.18	7.4-8.4	Low.
100	100	80-95	20-35	5-15	0.6-2.0	0.22-0.24	6.6-7.3	Low.
100	95-100	70-95	25-55	15-30	0.6-2.0	0.18-0.20	6.1-7.3	Moderate.
80-100	65-90	25-65	15-25	5-15	2.0-6.3	0.10-0.16	6.1-6.6	Low.
40-90	30-90	0-15	-----	NP	6.3-20	0.04-0.06	7.4-8.4	Low.
95-100	95-100	85-100	40-55	20-40	0.6-2.0	0.21-0.23	6.6-7.3	Moderate.
95-100	95-100	85-100	40-50	20-40	0.6-2.0	0.19-0.21	6.6-7.3	Moderate.
90-100	80-100	30-75	35	NP-15	0.6-2.0	0.19-0.21	7.4-8.4	Low.
100	95-100	90-100	30-50	10-15	0.6-2.0	0.22-0.24	6.1-7.3	Low.
100	95-100	80-95	40-55	20-35	0.6-2.0	0.18-0.20	6.1-7.3	Moderate.
90-100	60-100	50-80	30-55	5-15	0.6-6.3	0.14-0.18	7.4-7.8	Low.
100	100	95-100	30-55	10-15	0.6-2.0	0.20-0.25	6.1-7.3	Low.
100	95-100	95-100	40-55	20-35	0.6-2.0	0.19-0.21	5.6-6.5	Moderate.
95-100	85-95	55-75	15-25	5-15	0.2-1.0	0.18-0.20	6.6-8.4	Low.
100	95-100	90-100	20-60	15-25	0.6-2.0	0.19-0.23	7.4-8.4	Moderate.
95-100	90-100	80-100	35-60	20-35	0.6-2.0	0.16-0.19	7.4-8.4	Moderate.
90-100	80-100	60-90	25	NP-15	0.6-2.0	0.10-0.19	7.4-8.4	Low.
100	95-100	80-95	30-40	10-15	0.6-2.0	0.20-0.25	5.6-7.3	Low.
95-100	90-100	60-90	40-55	20-35	0.6-2.0	0.16-0.20	5.1-6.0	Moderate.
90-100	80-95	30-80	25	NP-15	0.6-6.3	0.12-0.16	6.6-8.4	Low.
100	95-100	90-100	30-50	15-25	0.6-2.0	0.22-0.24	6.1-6.5	Low.
95-100	90-100	80-95	40-55	20-35	0.6-2.0	0.18-0.20	5.6-6.5	Moderate.
95-100	90-100	55-75	30-55	15-30	0.6-2.0	0.17-0.19	5.6-7.3	Moderate.
95-100	90-100	55-75	15-35	5-25	0.6-2.0	0.17-0.19	7.4-8.4	Low.

TABLE 8.—Estimated soil properties

Soil series and map symbols	Depth of seasonal high water table	Depth from surface	USDA texture	Classification	
				Unified	AASHTO
	Feet	Inches			
Houghton: 103-----	0-1	0-55	Muck (sapric material).	Pt	-----
Kendall: 242-----	1-3	0-14	Silt loam-----	ML or CL	A-4 or A-6
		14-52	Silty clay loam-----	CL	A-6 or A-7
		52-65	Loam, silt loam-----	ML or CL	A-4 or A-6
Knight: 191-----	<2	0-35	Silt loam-----	CL or OL	A-6 or A-7
		35-54	Silty clay loam-----	CL	A-6 or A-7
		54-60	Clay loam-----	CL	A-6 or A-7
La Rose: 60B, 60C2, 60D2-----	>5	0-7	Silt loam-----	ML or CL	A-4 or A-6
		7-20	Clay loam, loam-----	CL	A-6 or A-7
		20-60	Loam-----	CL	A-4
Lisbon: 59-----	1-3	0-12	Silt loam-----	CL, ML, OL	A-4 or A-6
		12-26	Silty clay loam-----	CL	A-6 or A-7
		26-60	Loam, clay loam-----	CL	A-4
Lorenzo: 318D2-----	>5	0-5	Loam-----	CL	A-4
		5-14	Clay loam-----	CL	A-6
		14-60	Stratified sand and gravel.	SP or GP	A-1
Miami: 27B, 27C2, 27D2-----	>5	0-6	Silt loam-----	ML or CL	A-4 or A-6
		6-31	Clay loam, silty clay loam.	CL	A-6 or A-7
		31-60	Loam-----	CL	A-4
Millbrook: 219-----	1-3	0-12	Silt loam-----	CL or ML	A-6 or A-7
		12-34	Silty clay loam, loam-----	CL	A-6 or A-7
		34-68	Stratified loam, silt loam, and sandy loam.	SM, SC, ML, or CL	A-2, A-4, or A-6
Muscatine: 41-----	1-3	0-14	Silt loam-----	ML or CL	A-4 or A-6
		14-42	Silty clay loam-----	CL or CH	A-7
		42-60	Silt loam-----	CL	A-6 or A-4
Octagon: 656B, 656C2-----	>5	0-7	Silt loam-----	ML or CL	A-4 or A-6
		7-28	Clay loam-----	CL	A-6 or A-7
		28-60	Loam-----	CL	A-4
Peotone: 330-----	0-1	0-20	Silty clay loam-----	CL, OH, or CH	A-7
		20-38	Silty clay loam-----	CL or CH	A-7
		38-60	Silty clay loam-----	CL	A-7
Plano: 199A, 199B-----	>5	0-9	Silt loam-----	CL or ML	A-6 or A-4
		9-51	Silty clay loam-----	CL	A-6
		51-74	Stratified clay loam, loam.	ML or CL	A-4 or A-6
Proctor: 148A, 148B-----	>5	0-12	Silt loam-----	CL or ML	A-6 or A-4
		12-25	Silty clay loam-----	CL	A-6 or A-7
		25-60	Stratified clay loam, loam, and sandy loam.	SM, SC, or CL	A-2, A-4, A-6, or A-7
Rush: 791A, 791B-----	>5	0-13	Silt loam-----	ML or CL	A-4 or A-6
		13-46	Silty clay loam-----	CL	A-6 or A-7
		46-56	Gravelly clay loam-----	CL	A-4 or A-6
		56-60	Stratified sand and gravel.	SP or GP	A-1
Sable: 68-----	<2	0-17	Silty clay loam-----	CH, CL, or OH	A-7
		17-44	Silty clay loam-----	CL or CH	A-6 or A-7
		44-60	Silt loam-----	CL	A-6
St. Charles: 243A, 243B-----	>5	0-10	Silt loam-----	CL or ML	A-6 or A-4
		10-41	Silty clay loam-----	CL	A-6
		41-60	Clay loam, silt loam-----	ML or CL	A-4 or A-6

significant in engineering—Continued

Percentage less than 3 inches passing sieve <sup>1</sup> —			Liquid limit	Plasti- city index	Permea- bility	Available water capacity	Reaction	Shrink-swell potential
No. 4 (4.7 mm)	No. 10 (2.0 mm)	No. 200 (0.074 mm)						
			Percent		Inches per hour	Inches per inch of soil	pH	
				NP	2.0-6.3	>0.25	6.6-7.3	Not applicable
100	95-100	90-100	30-40	10-15	0.6-2.0	0.22-0.24	6.1-7.3	Low.
100	95-100	80-95	40-55	20-35	0.6-2.0	0.18-0.20	5.1-6.0	Moderate.
90-100	60-100	50-80	15-25	5-15	0.6-6.3	0.14-0.18	6.6-8.4	Low.
95-100	90-100	80-90	30-40	10-15	0.6-2.0	0.22-0.24	6.1-7.3	Low.
95-100	90-100	75-90	40-55	20-35	0.02-0.2	0.18-0.20	5.6-6.0	Moderate.
90-100	85-95	60-90	30-55	15-30	0.6-2.0	0.14-0.16	5.6-6.0	Moderate.
100	95-100	85-100	35-55	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
95-100	90-100	65-85	30-50	15-30	0.6-2.0	0.16-0.18	6.1-7.8	Moderate.
95-100	85-95	60-80	15-25	5-15	0.2-1.0	0.14-0.18	7.4-8.4	Low.
100	95-100	90-100	35-55	15-25	0.6-2.0	0.20-0.25	6.6-7.3	Low.
95-100	90-100	70-95	30-50	15-30	0.6-2.0	0.16-0.19	6.6-7.3	Moderate.
95-100	85-95	60-80	30-55	5-15	0.2-1.0	0.14-0.18	7.4-8.4	Low.
100	95-100	80-95	20-65	5-30	0.6-2.0	0.16-0.20	6.6-7.3	Low.
90-100	80-100	60-80	30-55	15-25	0.6-2.0	0.16-0.18	6.1-6.5	Moderate.
50-70	30-45	0-10		NP	6.3-20	0.02-0.04	7.4-8.4	Low.
100	95-100	85-95	30-40	10-15	0.6-2.0	0.16-0.20	6.6-7.3	Low.
95-100	90-100	65-85	30-55	15-30	0.6-2.0	0.16-0.19	5.6-7.3	Moderate.
95-100	85-95	60-80	15-25	5-15	0.2-1.0	0.14-0.18	7.4-8.4	Low.
100	95-100	90-100	30-40	10-15	0.6-2.0	0.20-0.25	5.1-5.5	Low.
100	90-100	60-90	40-55	20-35	0.6-2.0	0.18-0.20	5.6-7.3	Moderate.
95-100	90-100	30-80	25	NP-15	0.6-6.3	0.12-0.18	6.6-8.4	Low.
100	100	95-100	30-40	10-15	0.6-2.0	0.20-0.25	6.1-6.5	Low.
100	100	95-100	40-55	20-35	0.6-2.0	0.19-0.21	5.6-7.3	Moderate.
100	100	95-100	15-25	5-15	0.6-2.0	0.18-0.23	7.4-8.4	Low.
100	95-100	75-95	25-50	10-15	0.6-2.0	0.20-0.25	5.1-5.5	Low.
95-100	90-100	65-85	30-55	15-30	0.6-2.0	0.16-0.19	5.6-7.8	Moderate.
95-100	85-95	60-80	20-36	5-15	0.2-1.0	0.14-0.18	7.4-7.8	Low.
95-100	90-100	85-100	40-80	30-55	0.6-2.0	0.19-0.23	6.6-7.3	Moderate.
95-100	90-100	85-100	40-80	30-55	0.2-0.6	0.19-0.21	6.6-7.3	Moderate.
95-100	90-100	85-100	30-55	20-40	0.2-0.6	0.19-0.21	6.6-7.3	Moderate.
100	100	95-100	30-40	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
100	100	95-100	40-55	20-35	0.6-2.0	0.19-0.21	6.1-7.3	Moderate.
90-100	80-90	50-75	30-55	5-15	0.6-6.3	0.10-0.14	6.6-8.4	Low.
100	95-100	80-95	30-40	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
95-100	90-100	60-90	40-55	20-35	0.6-2.0	0.18-0.20	6.1-7.3	Moderate.
80-100	50-95	25-80	35	NP-15	0.6-6.3	0.10-0.14	6.1-7.8	Low.
100	100	80-95	30-40	10-15	0.6-2.0	0.22-0.24	6.6-7.3	Low.
100	95-100	85-95	30-50	15-30	0.6-2.0	0.18-0.20	5.6-7.3	Moderate.
80-100	65-90	50-65	30-40	15-25	2.0-6.3	0.14-0.16	5.6-6.0	Moderate.
40-80	30-70	0-10		NP	6.3-20	0.04-0.06	7.4-8.4	Low.
100	100	95-100	40-55	20-40	0.6-2.0	0.15-0.20	6.1-7.3	Moderate.
100	100	95-100	40-50	20-40	0.6-2.0	0.13-0.19	6.1-7.3	Moderate.
100	100	95-100	15-25	5-15	0.6-2.0	0.18-0.23	6.6-7.3	Low.
100	100	95-100	30-40	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
100	100	95-100	40-55	20-35	0.6-2.0	0.18-0.20	5.6-7.3	Moderate.
90-100	80-90	50-75	30-55	5-15	0.6-6.3	0.10-0.14	6.1-7.3	Low.

TABLE 8.—Estimated soil properties

Soil series and map symbols	Depth of seasonal high water table	Depth from surface	USDA texture	Classification	
				Unified	AASHTO
	Feet	Inches			
Sawmill: V107 -----	<2	0-31 31-44 44-55	Silty clay loam----- Silty clay loam----- Clay loam -----	CL, CH, or OH CL CL	A-7 A-6 or A-7 A-6 or A-7
Saybrook: 145B, 145C2-----	>5	0-12 12-25 25-38 38-60	Silt loam ----- Silty clay loam----- Clay loam ----- Loam -----	CL or OL CL CL CL	A-6 or A-7 A-7 or A-6 A-7 or A-6 A-4
Strawn: 224E2 -----	>5	0-4 4-19 19-60	Silt loam ----- Heavy loam, clay loam----- Loam -----	ML or CL CL CL	A-4 or A-6 A-6 or A-7 A-4
Virgil: 104 -----	1-3	0-11 11-39 39-60	Silt loam ----- Silty clay loam----- Silt loam, loam-----	ML or CL CL CL or SM	A-4 or A-6 A-6 or A-7 A-2, A-4, or A-6
Will: 329-----	<2	0-9 9-28 28-60	Clay loam ----- Clay loam ----- Gravelly sandy loam, stratified sand and gravel.	CL or OH CL GM, SP, SM, or SC	A-7 A-7 A-1, A-2, A-3, or A-4

<sup>1</sup> Most of the soils in DeKalb County have little or no coarse fraction greater than 3 inches in diameter in the upper layers. Some of the soils, such as Bowes, Dodge, Dresden, Herbert, La Rose, Lisbon, Lorenzo, Miami, Octagon, Rush, Saybrook, Strawn, and Will

TABLE 9.—Interpretations of engineering

Soil series and map symbols	Degree and kind of limitation for—				Suitability as a source of—
	Septic tank absorption fields	Shallow excavations	Dwellings with basements	Roads and streets	Road fill
Batavia: 105A, 105B, 105C2.	Slight-----	Slight-----	Slight-----	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Bowes: 792A, 792B --	Slight <sup>1</sup> -----	Slight-----	Slight-----	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Good for underlying material: stratified sand and gravel.
Camden: 134A, 134B--	Slight-----	Slight-----	Slight-----	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.

*significant in engineering—Continued*

Percentage less than 3 inches passing sieve <sup>1</sup> —			Liquid limit	Plasti- city index	Permea- bility	Available water capacity	Reaction	Shrink-swell potential
No. 4 (4.7 mm)	No. 10 (2.0 mm)	No. 200 (0.074 mm)						
			Percent		Inches per hour	Inches per inch of soil	pH	
100	95-100	90-100	40-55	20-40	0.6-2.0	0.19-0.23	6.6-7.3	Moderate.
95-100	90-100	80-100	40-50	20-40	0.6-2.0	0.19-0.21	6.1-7.3	Moderate.
80-100	70-90	60-90	30-55	NP-15	0.6-2.0	0.10-0.19	6.6-7.3	Low.
100	95-100	90-100	30-45	10-20	0.6-2.0	0.20-0.25	6.1-6.5	Low.
95-100	90-100	80-95	40-55	20-35	0.6-2.0	0.16-0.19	6.1-6.5	Moderate.
95-100	90-100	55-75	30-55	15-30	0.6-2.0	0.16-0.19	6.6-7.8	Moderate.
95-100	85-95	55-75	15-25	5-15	0.2-1.0	0.14-0.18	7.4-8.4	Low.
100	90-100	80-100	30-50	10-15	0.6-2.0	0.20-0.25	6.6-7.3	Low.
95-100	90-100	65-85	30-55	15-30	0.6-2.0	0.16-0.18	6.6-7.3	Moderate.
95-100	85-95	60-80	15-25	5-15	0.2-1.0	0.14-0.18	7.4-8.4	Low.
100	95-100	90-100	30-45	5-15	0.6-2.0	0.22-0.24	6.1-7.3	Low.
100	95-100	80-100	35-50	15-30	0.6-2.0	0.18-0.20	6.1-7.3	Moderate.
90-100	60-100	40-95	10-30	5-15	0.6-6.3	0.14-0.18	7.4-8.4	Low.
100	85-90	80-95	50-70	20-30	0.6-2.0	0.21-0.23	7.4-8.4	Moderate.
95-100	70-90	60-90	35-60	20-35	0.6-2.0	0.18-0.20	7.4-8.4	Moderate.
40-100	30-80	0-40	15	NP-6	0.6-2.0	0.02-0.10	7.4-8.4	Low.

soils, have 5 to 20 percent coarse fraction greater than 3 inches in the lower layers.

<sup>1</sup> NP means nonplastic.

*properties of the soils*

Suitability as a source of—Continued		Soil features affecting—			
Sand and gravel	Topsoil	Drainage for crops and pasture	Irrigation	Terraces and diversions	Grassed waterways
Poor to fair source of sand below a depth of 4 feet: generally contains considerable fines.	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; high available water capacity; sloping areas subject to runoff and erosion.	Soil properties are generally favorable; irregular topography in places.	Soil properties are favorable.
Good source of sand and gravel below a depth of 4 to 5 feet.	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Deep cuts should be avoided because of sand and gravel at a depth of 4 to 5 feet.	Soil properties are favorable.
Generally not suitable in upland areas; on stream terraces there may be thick deposits below a depth of 5 feet.	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Soil properties are generally favorable; irregular topography in places.	Soil properties are favorable.

TABLE 9.—*Interpretations of engineering*

Soil series and map symbols	Degree and kind of limitation for—				Suitability as a source of—
	Septic tank absorption fields	Shallow excavations	Dwellings with basements	Roads and streets	Road fill
Dodge: 24A, 24B, 24C2.	Moderate: permeability of the subsoil is dominantly 0.63 to 1.00 inch per hour.	Slight	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material; plasticity index generally less than 15.
Dresden: 325B, 325C2.	Slight <sup>1</sup>	Moderate: sand and gravel below a depth of 2 to 3 feet may slump.	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Good for underlying material: stratified sand and gravel.
Drummer: 152	Severe <sup>1</sup> : seasonally saturated with water near the surface.	Severe: seasonal high water table.	Severe: seasonal high water table at a depth of less than 2 feet.	Severe: generally has a high water table; high susceptibility to frost action.	Poor for subsoil: high water table.
Elburn: 198	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally more than 15. Fair for underlying material: plasticity index generally less than 15; seasonally saturated with water.
Flanagan: 154	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally more than 15. Fair for underlying material: plasticity index generally less than 15; seasonally saturated with water.
Harpster: 67	Severe: seasonally saturated with water near the surface.	Severe: seasonal high water table.	Severe: seasonally saturated with water close to the surface.	Severe: poorly drained; seasonal high water table; high susceptibility to frost action.	Poor for subsoil: plasticity index generally more than 15. Variable in underlying material: high water table.
Harvard: 344A, 344B, 344C2.	Slight	Slight	Slight	Severe for subsoil: plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.

*properties of the soils*—Continued

Suitability as a source of—Continued		Soil features affecting—			
Sand and gravel	Topsoil	Drainage for crops and pasture	Irrigation	Terraces and diversions	Grassed waterways
Not suitable-----	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Soil properties are generally favorable; irregular topography a limitation in places.	Soil properties are favorable.
Good source of gravel and sand below a depth of 2 to 3 feet.	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Gravel at a depth of 2 to 3 feet; droughty; difficult to vegetate.	Gravelly substratum; may be difficult to vegetate.
Generally not suited in upland areas; may have sand or gravel in some areas along streams.	Poor: high clay content; seasonal high water table.	Additional drainage generally needed; tile drains function well if outlets are available.	Not generally irrigated.	Not needed because of topography.	Not needed because of topography.
Poor to fair source of sand below a depth of 4 feet; contains many fines.	Good-----	Seasonal high water table at a depth of 1 foot to 3 feet; tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed because of topography.	Soil properties are generally favorable; needs subsurface drainage.
Not suitable-----	Good-----	Seasonal high water table at a depth of 1 foot to 3 feet; tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed because of topography.	Soil properties are generally favorable; needs subsurface drainage.
Not suitable in upland areas; some terrace areas have occasional pockets or thin layers of sand below a depth of 5 feet.	Poor: high clay content; seasonal high water table.	Drainage generally needed; tile drains function well if outlets are available.	Not generally irrigated.	Not needed because of topography.	Not needed because of topography.
Generally not suitable in upland areas; thick deposits on terraces below a depth of 5 feet in some places.	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Soil properties are generally favorable; irregular topography a limitation in some places.	Soil properties are favorable.

TABLE 9.—*Interpretations of engineering*

Soil series and map symbols	Degree and kind of limitation for—				Suitability as a source of—
	Septic tank absorption fields	Shallow excavations	Dwellings with basements	Roads and streets	Road fill
Catlin: 171A, 171B, 171C2.	Slight	Slight	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Herbert: 62	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally 15 or more. Fair for underlying material: plasticity index generally less than 15; seasonally saturated with water.
Houghton: 103	Very severe: continuously saturated with water; organic material has poor supporting strength for septic tank and tile.	Severe: high water table; organic material.	Very severe: continuously saturated with water; organic material has poor supporting strength.	Very severe: organic material.	Not suitable: organic material.
Kendall: 242	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally greater than 15. Fair for underlying material: plasticity index generally less than 15; seasonally saturated with water.
Knight: 191	Severe: seasonally saturated with water near the surface; ponding may occur in some places.	Severe: seasonal high water table.	Severe: seasonally saturated with water near the surface.	Severe: seasonal high water table; subject to ponding.	Poor for subsoil: plasticity index generally greater than 15. Fair to good for underlying material: plasticity index generally less than 15.
La Rose: 60B, 60C2, 60D2.	Moderate: permeability dominantly 0.60 to 1.00 inch per hour.	Slight where slopes are less than 7 percent. Moderate where slopes are more than 7 percent.	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Lisbon: 59	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally more than 15. Fair for underlying material: plasticity index generally less than 15; seasonally saturated with water.

*properties of the soils*—Continued

Suitability as a source of—Continued		Soil features affecting—			
Sand and gravel	Topsoil	Drainage for crops and pasture	Irrigation	Terraces and diversions	Grassed waterways
Not suitable -----	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Soil properties are generally favorable; irregular topography in places.	Soil properties are favorable.
Not suitable -----	Fair: 8 to 16 inches of suitable material.	Seasonal high water table 1 foot to 3 feet below the surface; tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed because of topography.	Soil properties are generally favorable; needs sub-surface drainage.
Not suitable: organic material.	Poor: high water table.	Water table at or near surface; tile drains are difficult to maintain; material subsides when water table is lowered.	Not generally irrigated, unless used for specialized crops; not suitable for furrow irrigation.	Not needed because of topography.	Not needed.
Poor to fair source below a depth of 4 feet; generally contains considerable fines.	Fair: 8 to 16 inches of suitable material.	Seasonal high water table at a depth of 1 foot to 3 feet; tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed because of topography.	Not needed because of topography.
Generally not suitable.	Fair: high water table.	Artificial drainage generally needed; tile drains function well if outlets are available.	Not generally irrigated.	Not needed-----	Not needed.
Not suitable-----	Fair: 8 to 16 inches of suitable material. Not suited if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Calcareous glacial material at a depth of 2 to 3 feet; difficult to vegetate.	Calcareous substratum may be difficult to vegetate.
Not suitable-----	Fair: 8 to 16 inches of suitable material.	Seasonal high water table at a depth of 1 foot to 3 feet; tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed -----	Not needed because of topography.

TABLE 9.—*Interpretations of engineering*

Soil series and map symbols	Degree and kind of limitation for—				Suitability as a source of—
	Septic tank absorption fields	Shallow excavations	Dwellings with basements	Roads and streets	Road fill
Lorenzo: 318D2-----	Moderate <sup>1</sup> : slopes are 6 to 15 percent.	Moderate: sand and gravel below a depth of 1 foot to 2 feet; subject to slumping.	Moderate: slopes are 6 to 15 percent.	Moderate: slopes are 6 to 15 percent.	Poor for subsoil: plasticity index generally more than 15. Good for underlying material: stratified sand and gravel.
Miami: 27B, 27C2, 27D2.	Moderate: permeability of subsoil is 0.63 to 1.00 inch per hour.	Slight in most places. Moderate where slopes are 7 to 12 percent.	Slight in most places. Moderate where slopes are 7 to 12 percent.	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Millbrook: 219-----	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally greater than 15. Fair to good for underlying material: plasticity index generally less than 15.
Muscatine: 41-----	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally more than 15. Fair to poor for underlying material: plasticity index may be more than 15; seasonally saturated with water.
Octagon: 656B, 656C2.	Moderate: permeability dominantly 0.60 to 1.00 inch per hour.	Slight-----	Slight-----	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Peotone: 330-----	Severe: generally saturated with water near the surface; subject to ponding.	Severe: seasonal high water table.	Severe: generally saturated with water near the surface; subject to ponding.	Severe: generally high water table; susceptible to frost action; plastic material.	Poor for subsoil: plasticity index generally more than 15. Poor for underlying material: plasticity index generally more than 15.
Plano: 199A, 199B-----	Slight-----	Slight-----	Slight-----	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.

*properties of the soils*—Continued

Suitability as a source of—Continued		Soil features affecting—			
Sand and gravel	Topsoil	Drainage for crops and pasture	Irrigation	Terraces and diversions	Grassed waterways
Good source of stratified sand and gravel below a depth of 1 foot to 2 feet.	Not suited: thin surface layer; gravelly.	Natural drainage is excessive.	Not suited; steep --	Gravel at a depth 1 foot to 2 feet; difficult to vegetate; low productivity.	Gravelly substratum difficult to vegetate.
Not suitable-----	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; subject to runoff and erosion.	Soil properties are generally favorable; irregular topography is a limitation in some areas.	Exposed subsoil difficult to vegetate.
Generally not suitable in upland areas; stream terraces have a thick deposit below a depth of 5 feet in places.	Fair: 8 to 16 inches of suitable material.	Seasonal high water table at a depth of 1 foot to 3 feet; tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed -----	Not needed because of topography.
Not suitable-----	Good-----	Seasonal high water table at a depth of 1 foot to 3 feet; tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed -----	Not needed.
Not suitable-----	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; subject to runoff and erosion.	Soil properties are generally favorable; irregular topography is a limitation in some areas.	Soil properties are favorable.
Generally not suitable.	Poor: high clay content; high water table.	Generally high water table; difficult to obtain outlets; depressional; surface drains may be needed to supplement tile.	Not generally irrigated.	Not needed -----	Not needed.
Poor to fair source of sand below a depth of 4 feet; generally contains considerable fines.	Fair: 8 to 16 inches of suitable material.	Natural drainage is adequate.	Medium intake rate; moderate permeability.	Soil properties are favorable.	Soil properties are favorable.

TABLE 9.—*Interpretations of engineering*

Soil series and map symbols	Degree and kind of limitation for—				Suitability as a source of—
	Septic tank absorption fields	Shallow excavations	Dwellings with basements	Roads and streets	Road fill
Proctor: 148A, 148B	Slight	Slight	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Rush: 791A, 791B	Slight	Slight	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Good for underlying material: sand and gravel below a depth of 4 to 5 feet.
Sable: 68	Severe: seasonally saturated with water near the surface.	Severe: seasonal high water table.	Severe: seasonally saturated with water near the surface.	Severe: generally has a high water table; high frost action potential.	Poor for subsoil: plasticity index generally more than 15. Variable in underlying material: high water table.
St. Charles: 243A, 243B.	Slight	Slight	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Sawmill: V107	Severe: seasonally saturated with water; subject to flooding.	Severe: seasonal high water table; subject to flooding.	Severe: seasonal high water table; subject to flooding.	Severe: seasonal high water table; subject to flooding.	Poor: plasticity index generally greater than 15; subject to flooding.
Saybrook: 145B, 145C2.	Moderate: permeability dominantly 0.60 to 1.00 inch per hour.	Slight	Slight	Severe: subsoil has plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.
Strawn: 224E2	Severe: too steep	Severe: too steep	Severe: too steep	Severe: plasticity index greater than 15.	Poor for subsoil: plasticity index generally more than 15. Fair to good for underlying material: plasticity index generally less than 15.

*properties of the soils*—Continued

Suitability as a source of—Continued		Soil features affecting—			
Sand and gravel	Topsoil	Drainage for crops and pasture	Irrigation	Terraces and diversions	Grassed waterways
Poor to fair source of sand below a depth of 4 feet; generally contains considerable fines.	Fair: 8 to 16 inches of suitable material.	Natural drainage is adequate.	Medium intake rate; moderate permeability.	Soil properties are favorable.	Soil properties are favorable.
Good source of sand and gravel below a depth of 4 to 5 feet.	Fair: 8 to 16 inches of suitable material. Poor if eroded.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Deep cuts should be avoided because of sand and gravel at a depth of 4 to 5 feet.	Soil properties are favorable.
Not suitable -----	Poor: high clay content; high water table.	Additional drainage generally needed; tile drains function well if outlets are available.	Not generally irrigated.	Not needed -----	Not needed.
Poor to fair source below a depth of 4 feet; generally contains considerable fines.	Fair: 8 to 16 inches of suitable material.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Soil properties are favorable.	Soil properties are favorable.
Generally not suitable.	Poor: high clay content; high water table; subject to flooding.	Additional drainage generally needed; adequate outlets may be difficult to obtain.	Not generally irrigated.	Not needed -----	Not needed.
Not suitable -----	Fair: 8 to 16 inches of suitable material.	Natural drainage is adequate.	Medium intake rate; moderate permeability; sloping areas subject to runoff and erosion.	Soil properties are generally favorable; irregular topography is a limitation in some places.	Soil properties are favorable.
Not suitable -----	Not suitable: steep; eroded; thin over calcareous material.	Natural drainage is adequate.	Not suited: steep	Not suitable on slopes of more than 12 percent; thin over calcareous material.	Calcareous substratum may be difficult to vegetate.

TABLE 9.—*Interpretations of engineering*

Soil series and map symbols	Degree and kind of limitation for—				Suitability as a source of—
	Septic tank absorption fields	Shallow excavations	Dwellings with basements	Roads and streets	Road fill
Virgil: 104 -----	Severe: periodically saturated with water at a depth of 1 foot to 3 feet.	Severe: seasonal high water table.	Severe: seasonally saturated with water at a depth of 1 foot to 3 feet.	Severe: high susceptibility to frost action.	Poor for subsoil: plasticity index generally more than 15. Fair for underlying material: plasticity index generally less than 15; seasonally saturated with water.
Will: 329 -----	Severe <sup>1</sup> : seasonally saturated with water near the surface.	Severe: seasonal high water table.	Severe: seasonally saturated with water near the surface.	Severe: poorly drained; seasonal high water table; susceptible to frost heave.	Poor for subsoil: plasticity index generally more than 15. Poor for underlying material: high water table.

<sup>1</sup> Pollution is a hazard because of permeability in the substratum or a high water table.

TABLE 10.—*Engineering*

[Tests performed by the Illinois Department of

Soil name and location	Parent material	Report No. 71- IL-019	Depth	Moisture-density <sup>1</sup>		
				Maximum dry density	Optimum moisture	
			Inches	Lb per cu ft	Percent	
Batavia silt loam: 260 feet south and 20 feet east of the center of the NW¼NE¼ sec. 26, T. 40 N., R. 3 E. (Modal)	Water-deposited silts and sands.		7-1	0-7	102	20
			7-2	14-23	101	20
			7-3	48-68	131	9
			7-4	68-96	119	14
Catlin silt loam: 188 feet north and 206 feet east of the SW. corner of the NW¼NW¼ sec. 28, T. 40 N., R. 3 E. (Modal)	Loess over loam till.		12-1	0-6	97	22
			12-2	13-38	100	20
			12-3	50-70	128	10
			11-1	0-7	99	21
510 feet west and 540 feet south of the NE. corner of sec. 34, T. 38 N., R. 5 E. (Thinner solum than modal profile)	Loess over loam till.		11-2	11-40	100	21
			11-3	40-50	132	8
			11-4	50-65	118	13
Dodge silt loam: 40 feet south and 150 feet east of the NE. corner of the SE¼SW¼ sec. 18, T. 41 N., R. 5 E. (Modal)	Loess over loam till.		1-1	0-6	105	18
			1-2	13-25	102	20
			1-3	25-31	113	15
			1-4	31-60	123	11
Dresden silt loam: 348 feet east and 60 feet north of the SW. corner of sec. 33, T. 40 N., R. 5 E. (Modal)	Outwash of sand and gravel.		9-1	0-7	107	17
			9-2	14-29	106	17
			9-3	38-60	126	11
Elburn silt loam: 155 feet south and 20 feet east of the NW¼NE¼ sec. 26, T. 40 N., R. 3 E. (Modal)	Water-deposited silts and sands.		8-1	0-6	102	21
			8-2	16-38	101	21
			8-3	56-76	126	10
			8-4	76-102	121	12
Flanagan silt loam: 804 feet south and 384 feet west of the center of sec. 23, T. 39 N., R. 3 E. (Modal)	Loess over loam till.		10-1	0-13	92	25
			10-2	16-35	103	20
			10-3	45-60	126	11

properties of the soils—Continued

Suitability as a source of—Continued		Soil features affecting—			
Sand and gravel	Topsoil	Drainage for crops and pasture	Irrigation	Terraces and diversions	Grassed waterways
Poor to fair source below a depth of 4 feet; generally contains considerable fines.	Fair: 8 to 16 inches of suitable material.	Tile drains are suited and are needed during wet seasons.	Not generally irrigated.	Not needed because of topography.	Not needed because of topography.
Good source of stratified sand and gravel below a depth of 2 1/2 feet.	Poor: calcareous; high water table.	Artificial drainage generally needed; tile function well if outlets are available; keep tile above sand and gravel if possible; use sealed joints in sand and gravel	Not generally irrigated.	Not needed because of topography.	Not needed.

test data

Transportation, Bureau of Materials, Springfield]

Mechanical analysis <sup>2</sup>											Liquid limit	Plasticity index	Classification	
Percentage passing sieve—						Percentage smaller than—				AASHTO <sup>3</sup>			Unified	
1 in.	3/4 in.	3/8 in.	No. 4 (4.7 mm)	No. 10 (2.0 mm)	No. 40 (0.42 mm)	No. 200 (0.074 mm)	0.05 mm	0.02 mm	0.005 mm	0.002 mm	Percent			
100	99	96	92	100	99	97	85	58	34	27	35	13	A-6(14)	CL
				100	100	99	94	67	38	31	43	23	A-7-6(25)	CL
				85	70	43	32	21	3	2	15	2	A-4(0)	SM
				100	99	96	88	60	30	18	23	6	A-4(4)	CL-ML
				100	99	96	64	51	34	26	42	15	A-7-6(16)	ML
100	99	98	96	100	99	98	90	69	40	33	44	23	A-7-6(25)	CL
				94	89	62	57	46	28	22	20	7	A-4(1)	CL-ML
				100	99	97	87	55	25	17	39	14	A-6(15)	ML
				100	99	98	90	67	34	43	20	20	A-7-6(22)	CL
100	100	97	93	86	71	46	45	33	17	12	16	2	A-4(0)	SC
	99	98	97	95	92	82	75	52	19	13	28	11	A-6(7)	CL
				100	99	94	89	50	12	7	33	7	A-4(7)	ML
100	99	99	98	100	99	97	84	57	30	21	46	24	A-7-6(25)	CL
'99	98	97	96	97	93	67	59	40	24	16	34	22	A-6(11)	CL
				94	88	64	55	39	27	23	26	13	A-6(15)	CL
	100	99	97	94	86	76	70	55	30	24	34	12	A-6(6)	CL
	99	99	97	92	85	76	74	63	43	36	41	21	A-7-6(15)	CL
	100	98	96	90	54	14	12	10	6	3			A-2-4(0)	SP
				100	99	98	90	62	32	24	48	28	A-7-6(30)	CL
				100	98	96	81	60	36	28	38	13	A-6(14)	CL
	100	99	97	93	85	61	39	26	7	3	18	3	A-4(0)	CL
				100	98	90	82	55	30	23	21	4	A-4(1)	CL-ML
				100	98	86	80	58	29	14	51	19	A-7-6(20)	MH
100	99	98	97	100	99	95	89	33	33	23	54	30	A-7-6(33)	CH
				95	89	66	60	46	26	16	20	6	A-4(1)	CL-ML

TABLE 10.—Engineering

Soil name and location	Parent material	Report No. 71- IL-019	Depth	Moisture-density <sup>1</sup>		
				Maximum dry density	Optimum moisture	
			Inches	Lb per cu ft	Percent	
Herbert silt loam: 147 feet north of road and 306 feet east of section line in the NW¼NW¼ sec. 14, T. 42 N., R. 4 E. (Modal)	Loess over loam till.		3-1	0-8	101	20
			3-2	12-16	101	20
			3-3	36-58	113	14
Octagon silt loam: 1,133 feet west and 79 feet north of the intersection of Base Line and Lloyd Road, sec. 32, T. 42 N., R. 5 E. (Modal)	Loess over loam till.		2-1	0-7	110	16
			2-2	7-18	110	15
			2-3	24-60	124	11
Peotone silty clay loam: 75 feet east and 120 feet south of the junction of Lee Road and CNWRR tracks, NW¼NE¼ sec. 7, T. 38 N., R. 4 E. (Modal)	Silty clay loam sediments.		4-1	0-8	82	32
			4-2	8-34	100	21
			4-3	34-45	109	17
Saybrook silt loam: 95 feet west and 238 feet south of the NE corner of SE¼NW¼ sec. 27, T. 41 N., R. 3 E. (Modal)	Loess over loam till.		5-1	0-8	101	21
			5-2	15-26	99	21
			5-3	26-31	110	15
			5-4	44-58	128	10
Virgil silt loam: 1,000 feet north and 330 feet west of the intersection of Loves Road and Pleasant Street, sec. 18, T. 40 N., R. 5 E. (Modal)	Water-deposited silts and sands.		6-1	0-9	109	16
			6-2	11-46	104	19
			6-3	64-92	118	14

<sup>1</sup> Based on AASHTO Designation T 99, Method A (1).

<sup>2</sup> Mechanical analysis according to AASHTO Designation T 88-57 (1). Results by this procedure may differ somewhat from the results obtained by the soil survey procedure of the Soil Conservation Service (SCS). In the AASHTO procedure, the fine material is analyzed by the hydrometer method, and the various grain-size fractions are calculated on the basis of all material up to and including that 3 inches in diameter. In the SCS soil survey procedure, the fine material is analyzed by the pipette method, and the

### Engineering soil classification systems

The two systems most commonly used in classifying samples of soils for engineering are the Unified system (2) used by the Soil Conservation Service, Department of Defense, and other agencies, and the AASHTO system (1) adopted by the American Association of State Highway and Transportation Officials.

In the Unified system soils are classified according to particle-size distribution, plasticity, liquid limit, and organic matter. Soils are grouped in 15 classes. There are eight classes of coarse-grained soils, identified as GW, GP, GM GC, SW SP, SM, and SC; six classes of fine-grained soils, identified as ML, CL, OL, MH, CH, and OH; and one class of highly organic soils, identified as Pt. Soils on the borderline between two classes are designated by symbols for both classes; for example, CL-ML.

The AASHTO system is used to classify soils according to those properties that affect use in highway construction and maintenance. In this system a soil is placed in one of seven basic groups that range from

A-1 through A-7 on the basis of grain-size distribution, liquid limit, and plasticity index. In group A-1 are gravelly soils of high bearing strength, or the best soils for subgrade (foundation). At the other extreme, in group A-7, are clay soils that have low strength when wet and are the poorest soils for subgrade. Where laboratory data are available to justify a further breakdown, the A-1, A-2, and A-7 groups are divided as follows: A-1-a, A-1-b; A-2-4, A-2-5, A-2-6, A-2-7; and A-7-5 and A-7-6. As additional refinement, the engineering value of a soil material can be indicated by a group index number. Group indexes range from 0 for the best material to 20 or more for the poorest. The AASHTO classification for tested soils, with group index numbers in parentheses, is shown in table 10; the estimated classification, without group index numbers, is shown in table 8 for all soils mapped in the county.

### Soil properties significant in engineering

Several estimated soil properties significant in en-

test data—Continued

Mechanical analysis <sup>2</sup>											Liquid limit	Plasticity index	Classification	
Percentage passing sieve—						Percentage smaller than—				AASHTO <sup>3</sup>			Unified	
1 in.	¾ in.	⅜ in.	No. 4 (4.7 mm)	No. 10 (2.0 mm)	No. 40 (0.42 mm)	No. 200 (0.074 mm)	0.05 mm	0.02 mm	0.005 mm		0.002 mm	Percent		
100	99	99	100	98	98	93	89	66	30	21	36	14	A-6 (24)	CL
		100	99	99	98	93	89	76	48	41	49	26	A-7-6 (26)	CL
			99	99	95	63	59	49	32	28	35	23	A-6 (11)	CL
	100	99	99	98	95	75	66	48	22	16	29	11	A-6 (6)	CL
	100	99	98	96	93	67	61	49	29	21	36	19	A-6 (10)	CL
	100	99	97	95	90	69	63	49	15	10	22	7	A-4 (2)	CL
				100	96	89	87	79	48	37	72	37	A-7-6 (40)	OH
				100	99	99	96	84	48	38	53	30	A-7-6 (39)	CH
				100	99	99	94	70	36	30	51	33	A-7-6 (36)	CH
				100	99	95	86	64	34	22	42	20	A-7-6 (21)	CL
			100	99	97	89	84	64	38	32	48	28	A-7-6 (26)	CL
	100	99	99	97	88	59	50	33	22	16	41	24	A-7-6 (11)	CL
100	99	97	95	92	86	62	59	48	30	24	22	9	A-4 (2)	CL
			100	99	95	73	58	36	17	11	30	9	A-4 (5)	CL
				100	100	99	93	57	33	24	46	26	A-7-6 (28)	CL
			100	99	99	92	85	48	27	20	29	13	A-6 (11)	CL

material coarser than 2 millimeters in diameter is excluded from the calculation of grain-size fractions. The mechanical analysis data used in this table are not suitable for use in naming textural classes of soil.

<sup>2</sup> Based on AASHTO Designation M 145-66-I (I).

<sup>3</sup> 100 percent passed the 1½ inch sieve.

<sup>4</sup> NP means nonplastic.

gineering are shown in table 8. These estimates are made for typical soil profiles, by layers sufficiently different to have different significance for soil engineering. The estimates are based on field observations made in the course of mapping, on test data for these and similar soils, and on experience with the same kinds of soil in other counties. Following are explanations of some of the columns in table 8.

Depth to bedrock is not shown in table 8, because most soils in the county are deep enough that bedrock generally does not affect their use.

Depth to seasonal high water table is the distance from the surface of the soil to the highest level that ground water reaches in the soil in most years.

Soil texture is described in table 8 in the standard terms used by the Department of Agriculture (?). These terms take into account the relative percentages of sand, silt, and clay in soil material that is less than 2 millimeters in diameter. "Loam," for example, is soil material that contains 7 to 27 percent clay, 28 to

50 percent silt, and less than 52 percent sand. If the soil contains gravel or other particles coarser than sand, an appropriate modifier is added, as for example, "gravelly loamy sand." "Sand," "silt," "clay," and some of the other terms used in USDA textural classification are defined in the Glossary.

Liquid limit and plasticity index indicate the effect of water on the strength and consistence of soil material. As the moisture content of a clayey soil is increased from a dry state, the material changes from a semisolid to a plastic. If the moisture content is further increased, the material changes from a plastic to a liquid. The plastic limit is the moisture content at which the soil material changes from a semisolid to a plastic; and the liquid limit, from a plastic to a liquid. The plasticity index is the numerical difference between the liquid limit and the plastic limit. It indicates the range of moisture content within which a soil material is plastic. Liquid limit and plasticity index are estimated in table 8, but in table 10 the data on

liquid limit and plasticity index are based on tests of soil samples.

Permeability is that quality of a soil that enables it to transmit water or air. It is estimated on the basis of those soil characteristics observed in the field, particularly structure and texture. The estimates in table 8 do not take into account lateral seepage or such transient soil features as plowpans and surface crusts.

Available water capacity is the ability of soils to hold water for use by most plants. It is commonly defined as the difference between the amount of water in the soil at field capacity and the amount at the wilting point of most crop plants.

Reaction is the degree of acidity or alkalinity of a soil, expressed as a pH value. The pH value and terms used to describe soil reaction are explained in the Glossary.

Shrink-swell potential is the relative change in volume to be expected of soil material with changes in moisture content, that is, the extent to which the soil shrinks as it dries out or swells when it gets wet. Extent of shrinking and swelling is influenced by the amount and kind of clay in the soil. Shrinking and swelling of soils cause much damage to building foundations, roads, and other structures. A *high* shrink-swell potential indicates a hazard to maintenance of structures built in, on, or with material that has this rating.

#### **Engineering interpretations of soils**

The estimated interpretations in table 9 are based on the engineering properties of soils shown in table 8, on test data for soils in this survey area and in others nearby or adjoining, and on the experience of engineers and soil scientists with the soils of DeKalb County. In table 9, ratings are used to summarize limitation or suitability of the soils for all listed purposes other than drainage for crops and pasture, irrigation, terraces and diversions, and grassed waterways. For these particular uses, table 9 lists those soil features not to be overlooked in planning, installation, and maintenance.

Soil limitations are indicated by the ratings slight, moderate, and severe. *Slight* means soil properties are generally favorable for the rated use, or in other words, limitations are minor and easily overcome. *Moderate* means some soil properties are unfavorable but can be overcome or modified by special planning and design. *Severe* means soil properties are so unfavorable and so difficult to correct or overcome that they require major soil reclamation and special design. For some uses, the rating of severe is divided to obtain ratings of severe and very severe. *Very severe* means one or more soil properties are so unfavorable for a particular use that overcoming the limitations is most difficult and costly and commonly not practical for the rated use.

Soil suitability is rated by the terms *good*, *fair*, and *poor*, which have meanings approximately parallel to the terms slight, moderate, and severe, respectively.

Following are explanations of the columns in table 9.

Septic tank absorption fields are subsurface systems of tile or perforated pipe that distribute effluent from a septic tank into natural soil. The soil material between depths of 18 inches and 6 feet is evaluated. The

soil properties considered are those that affect both absorption of effluent and construction and operation of the system. Properties that affect absorption are permeability, depth to water table or rock, and susceptibility to flooding. Slope is a soil property that affects difficulty of layout and construction and also the risk of soil erosion, lateral seepage, and downslope flow of effluent. Large rocks or boulders increase construction costs.

Shallow excavations are those that require digging or trenching to a depth of less than 6 feet, for example, excavations for pipelines, sewerlines, phone and power transmission lines, basements, open ditches, and cemeteries. Desirable soil properties are good workability, moderate resistance to sloughing, gentle slopes, absence of rock outcrops or large stones, and freedom from flooding or a high water table.

Dwellings with basements, as rated in table 9, are no more than three stories high and are supported by foundation footings placed in undisturbed soil. The features that affect the ratings of a soil for dwellings are those that relate to capacity to support load and resist settlement under load and those that relate to ease of excavation. Soil properties that affect capacity to support load are wetness, susceptibility to flooding (fig. 8), density, plasticity, texture, and shrink-swell potential. Those that affect excavation are wetness, slope, depth to bedrock, and content of stones and rocks.

Roads and streets have an all-weather surface that is expected to carry automobile traffic all year, but not fast-moving, heavy trucks. They have a subgrade of underlying soil material; a base consisting of gravel, crushed rock, or soil material stabilized with lime or cement; and a flexible or rigid surface, commonly asphalt or concrete. These roads are graded to shed water and have ordinary provisions for drainage. They are built mainly from soil at hand, and most cuts and fills are less than 6 feet deep.

Soil properties that most affect design and construction of roads and streets are load-supporting capacity and stability of the subgrade and the workability and quantity of cut and fill material available. The AASHTO and Unified classifications of the soil material, and also the shrink-swell potential, indicate traffic-supporting capacity. Wetness and flooding affect stability of the material. Slope, depth to hard rock, content of stones and rocks, and wetness affect the ease of excavation and amount of cut and fill needed to reach an even grade.

Road fill is soil material used in embankments for roads. The suitability ratings reflect (1) the predicted performance of soil after it has been placed in an embankment that has been properly compacted and provided with adequate drainage and (2) the relative ease of excavating the material at borrow areas.

Sand and gravel are used in great quantities in many kinds of construction. The ratings in table 9 provide guidance about where to look for probable sources. A soil rated as a *good* or *fair* source of sand or gravel generally has a layer at least 3 feet thick, the top of which is within a depth of 6 feet. The ratings do not take into account thickness of overburden, location of the water table, or other factors that affect mining of



Figure 8.—Flooding in a developed area on the flood plain of the Kishwaukee River.

the materials, and neither do they indicate quality of the deposit.

Topsoil is used for topdressing an area where vegetation is to be established and maintained. Suitability is affected mainly by ease of working and spreading the soil material, as for preparing a seedbed; natural fertility of the material, or the response of plants when fertilizer is applied; and absence of substances toxic to plants. Texture of the soil material and its content of stone fragments are characteristics that affect suitability, but also considered in the ratings is damage that will result at the area from which topsoil is taken.

Drainage for crops and pasture is affected by such soil properties as permeability, texture, and structure; depth to claypan, rock, or other layers that influence rate of water movement; depth to the water table; slope; stability in ditchbanks; susceptibility to stream overflow; salinity or alkalinity; and availability of outlets for drainage.

Irrigation of a soil is affected by such features as slope; susceptibility to stream overflow, water erosion, or soil blowing; soil texture; content of stones; accumulations of salts and alkali; depth of root zone; rate of water intake at the surface; permeability of soil layers below the surface layer and in fragipans or other layers that restrict movement of water; amount of water held available to plants; and need for drainage, or depth to water table or bedrock.

The factors considered for terraces, diversions, and grassed waterways are those features and qualities of soils that affect layout and construction and the establishment, growth, and maintenance of plants.

Features considered include soil slope, texture and reaction of the subsoil, depth to gravel or other limiting layers, and natural soil drainage.

#### Soil test data

Table 10 contains engineering test data for some of the major soil series in DeKalb County. These tests were made to help evaluate the soils for engineering purposes. The engineering classifications shown are based on data obtained by mechanical analyses and by tests to determine liquid limits and plastic limits. The mechanical analyses were made by combined sieve and hydrometer methods.

Moisture-density data are important in earthwork. If a soil material is compacted at successively higher moisture content, assuming that the compactive effort remains constant, the density of the compacted material increases until the *optimum moisture content* is reached. After that, density decreases with increase in moisture content. The highest dry density obtained in the compactive test is termed the *maximum dry density*. As a rule, maximum strength of earthwork is obtained if the soil is compacted to the maximum dry density.

Mechanical analysis refers to the measurement of the amounts of various size classes of soil grains (sand, silt, or clay) in a sample. Proportions of the size classes determine the textural class of the material. Names used by engineers for various size classes of particles differ from those used by soil scientists. For example, fine sand in engineering terminology consists of particles 0.42 to 0.074 millimeter in diameter; but fine

sand, as determined by the soil scientist, consists of particles 0.25 to 0.10 millimeter in diameter.

## *Formation and Classification of the Soils*

This section describes how the factors of soil formation have affected the development of soils in DeKalb County. It also explains the system of soil classification currently used and places each soil series in the classes of that system.

### **Factors of Soil Formation**

The principal factors of soil formation are parent material, climate, plant and animal life, relief and drainage, and time. The kinds of soils in an area and how they formed are the result of the interaction of these five factors. The relative importance of each factor differs from place to place, and each modifies the effect of the other four. A change in any one of the five major factors of soil formation produces different soils, and each soil has its own characteristics and thus can be distinguished from other soils. The differences in all soils are in some measure related to the variation in and the interaction of the five soil-forming factors. Many times the exact role of each factor is difficult to determine, but the combined effect always produces a soil profile that has its own distinguishing characteristics. In some cases one factor may dominate the formation of a soil.

Man, in such activities as clearing forests and cultivating and fertilizing fields, also changes the course of soil formation, but so far he has had little effect on the overall development of the soils in DeKalb County.

### *Parent material*

In DeKalb County the parent materials of the soils are glacial till, glacial outwash, loess, and alluvium. The three ice advances each deposited its own till, which can be distinguished by physical and chemical properties, namely texture or grain size and clay mineralogy. Not only can the soils that formed in loess be distinguished from those that formed in till, but also soils that formed in different tills can be distinguished from each other. This comes about because different tills have distinguishing physical and chemical properties that the soils may inherit. Thus, soils can be established for each geologic unit (4). The detailed soil map shows associations of soils not only because of their proximity to one another on the landscape, but also because of the relationship of some of their characteristics.

After the county was covered by three major ice advances of the late Wisconsinan age (10), it was covered by a mantle of windblown silt, called loess, which regionally thins from west to east. A small proportion of the soils formed entirely in loess, but a large proportion formed in 2 to 5 feet of loess and in the

underlying glacial till. For example, Muscatine and Sable soils formed entirely in loess, and Catlin, Saybrook, and Flanagan soils formed in loess and in the underlying glacial till.

About 8 percent of the soils in the county formed mostly in till, and only the surface layer formed in loess. Examples are Miami and Octagon soils. Soils that formed in loess have distinctly different characteristics from those that formed in till.

The differences between soils that formed in different geologic units may not be great enough to justify separations for farming uses, but many of these differences are significant to other soil uses (fig. 9).

About 20 percent of the soils formed in outwash. The outwash areas are along the major streams or in front of the glacial moraines. Elburn, Camden, and Millbrook soils are examples of soils that formed in outwash.

In addition to the typical outwash areas, detailed soil mapping and aerial photographic interpretation have revealed a field of more than 500 highly symmetrical, circular to elliptical, raised, flat-topped mounds in the area around DeKalb and Malta. The mound field, which covers approximately 60 square miles, is confined to the low-relief, dominantly ground-moraine area between the Bloomington moraine on the west and north and the Shabbona, Arlington, and Mendota moraines on the east and south. The mounds, because of their relatively low relief, are most evident on aerial photographs.

The higher mounds rise 5 to 15 feet above the adjacent areas of gently rolling soils. The surfaces of the mounds tend to be flat or slightly depressed toward their center. In many areas where surface drainage is undeveloped, the mounds are surrounded by a moat or an annular depression.


Many of the mounds are closely spaced and are separated by depressions only a few feet wide. In many places one or more satellite mounds overlap the flanks of the larger mound. The mounds generally contain more than 20 feet of silt interbedded with sand underlain by till. The mounds apparently represent the deposit of lakes that existed in the region during part of the Woodfordian time. Examples of soils that formed on the mounds are those of the Batavia and Virgil series.

In the southeastern corner of the county, the outwash, which is water-worked drift material, has been deposited locally on the till. The outwash generally does not extend to a depth of more than 7 or 8 feet.

The soils of one series formed in water-deposited material or alluvium. These soils are on the flood plains of the major streams. They are the Sawmill soils. The soils of another series formed in partly decomposed organic material. They are the Houghton soils.

### *Climate*

Climate affects the formation of soils through its influence on the rate of weathering of parent material. The humid, temperate climate of DeKalb County is conducive to the relatively rapid breakdown or



**Figure 9.**—Distribution of parent materials in DeKalb County. Numbered areas are as follows: 1, mainly 2 to 4 feet of loess over more than 10 feet of outwash, lacustrine deposits, or alluvium; 2, raised circular mounds that consist of more than 20 feet of stratified, water-laid, silty sediment and are entirely covered by 0 to 4 feet of loess; 3, glacial till plain made up of 2 to 5 feet of stratified, water-worked glacial material covered with 2 to 4 feet of loess; 4, more than 5 feet of loess; 5, Capron till covered with 5 to 4 feet of loess; 6, Tiskilwa till covered with 0 to 4 feet of loess; 7, Malden till covered with 0 to 4 feet of loess.

weathering of soil minerals, to the formation of clay, and to the movement of these materials downward in the soil profile. In most of the soils on uplands of the county, there is more clay in the subsoil than in the surface layer.

#### **Plants and animals**

Plants have had a greater effect than animals on the formation of soils in DeKalb County, but the animals

and organisms that live on and in the soils also have been important. The changes they bring about depend mainly on the kind of life processes peculiar to each. The kinds of plants and animals that live on and in the soil are affected, in turn, by climate, parent material, relief, and the length of time that the soil has been forming.

Some soils in the county formed under trees and some under prairie grasses. Many of the sloping soils formed under trees, mainly oak and hickory. Dodge and Miami soils formed under trees. The soils that formed under prairie grasses are in the more nearly level areas of the county. Saybrook and Flanagan soils are examples of soils that formed under prairie grasses. They have a thicker and darker colored surface layer and are higher in organic-matter content than the soils that formed under forest. Organic matter has accumulated in all the soils.

#### **Relief and drainage**

Relief influences the amount of runoff, the degree of erosion, and the amount of water that infiltrates and percolates through the soil profile. Where the soils formed in uniform, permeable parent material, such as loess, natural drainage is closely associated with slope. The moderately well drained and well drained soils are in the more rolling areas, and the somewhat poorly drained to very poorly drained soils are mainly on flats or in depressions. Slopes in the county range from less than 2 percent in drainageways and on the nearly level uplands to more than 15 percent on the steeper parts of the uplands bordering streams.

In the poorly drained soils, iron compounds have been reduced and moved downward in the profile, causing the subsoil to have a gray color. Some of this iron has accumulated as concretions, or small, rounded pellets. In the well-drained soils, the iron compounds are oxidized and are generally more diffuse. They give a brown or yellowish-brown color to the subsoil.

#### **Time**

Time is necessary for the formation of soil from parent material, but the length of time required is largely dependent on the combined action of the other soil-forming factors.

Time is the effective age of a soil, as expressed by the degree of horizon differentiation. Recent deposits that have little or no horizon differentiation are considered young soils.

Soils that formed in parent material low in content of lime develop more rapidly and become more acid than soils that formed in material high in content of lime. Permeable soils are leached of lime and other soluble minerals much more rapidly than slowly permeable soils. Soils develop more rapidly under forest than under prairie vegetation. Steep soils develop more slowly than less strongly sloping soils because of greater runoff and less movement of water through

the soil. Soils generally develop more rapidly in a humid climate than in a dry one. They are generally more strongly developed or have greater horizon differentiation where they have been exposed to weathering processes over a long period of time.

Most of the soils on uplands in DeKalb County are moderately well developed. The alluvial soils have weak horizon differentiation or none at all, because of continuing deposition.

### Classification of the Soils

Soils are classified so that we can more easily remember their significant characteristics. Classification enables us to assemble knowledge about the soils, to see their relationship to one another and to the whole environment, and to develop principles that help us to understand their behavior and their response to manipulation. First through classification, and then through use of soil maps, we can apply our knowledge of soils to specific fields and other tracts of land.

The narrow categories of classification, such as those used in detailed soil surveys, allow us to organize and

apply knowledge about soils in managing farms, fields, and woodlands; in developing rural areas; in engineering work; and in many other ways. Soils are placed in broad classes to facilitate study and comparison in large areas, such as countries and continents.

The system of soil classification currently used was adopted by the National Cooperative Soil Survey in 1965 (8). Because this system is under continual study, readers interested in developments of the current system should search the latest literature available.

The current system of classification has six categories. Beginning with the broadest, these categories are the order, the suborder, the great group, the subgroup, the family, and the series. In this system the criteria used as a basis for classification are soil properties that are observable and measurable. The properties are chosen, however, so that the soils of similar genesis, or mode of origin, are grouped. The same property or subdivisions of this property may be used in several different categories. In table 11 the soil series of DeKalb County are placed in four categories of the current system. Classes of the current system are briefly defined in the following paragraphs.

TABLE 11.—*Classification of soils*

Series	Family	Subgroup	Order
Batavia	Fine-silty, mixed, mesic	Mollic Hapludalfs	Alfisols.
Bowes	Fine-silty, mixed, mesic	Mollic Hapludalfs	Alfisols.
Camden	Fine-silty, mixed, mesic	Typic Hapludalfs	Alfisols.
Catlin	Fine-silty, mixed, mesic	Typic Argiudolls	Mollisols.
Dodge	Fine-silty, mixed, mesic	Typic Hapludalfs	Alfisols.
Desden	Fine-loamy over sandy or sandy-skeletal, mixed, mesic	Mollic Hapludalfs	Alfisols.
Drummer	Fine-silty, mixed, mesic	Typic Haplaquolls	Mollisols.
Elburn	Fine-silty, mixed, mesic	Aquic Argiudolls	Mollisols.
Flanagan	Fine, montmorillonitic, mesic	Aquic Argiudolls	Mollisols.
Harpster	Fine-silty, mesic	Typic Calciquolls	Mollisols.
Harvard	Fine-silty, mixed, mesic	Mollic Hapludalfs	Alfisols.
Herbert	Fine-silty, mixed, mesic	Udolic Ochraqualfs	Alfisols.
Houghton	Euic, mesic	Typic Medisaprists	Histosols.
Kendall	Fine-silty, mixed, mesic	Aeric Ochraqualfs	Alfisols.
Knight	Fine-silty, mixed, mesic	Argiaquic Argialbolls	Mollisols.
La Rose	Fine-loamy, mixed, mesic	Typic Argiudolls	Mollisols.
Lisbon	Fine-silty, mixed, mesic	Aquic Argiudolls	Mollisols.
Lorenzo	Fine-loamy over sandy or sandy-skeletal, mixed, mesic	Typic Argiudolls	Mollisols.
Miami	Fine-loamy, mixed, mesic	Typic Hapludalfs	Alfisols.
Millbrook	Fine-silty, mixed, mesic	Udolic Ochraqualfs	Alfisols.
Muscatine	Fine-silty, mixed, mesic	Aquic Argiudolls	Mollisols.
Octagon	Fine-loamy, mixed, mesic	Mollic Hapludalfs	Alfisols.
Peotone	Fine, montmorillonitic, mesic	Cumulic Haplaquolls	Mollisols.
Plano	Fine-silty, mixed, mesic	Typic Argiudolls	Mollisols.
Proctor	Fine-silty, mixed, mesic	Typic Argiudolls	Mollisols.
Rush	Fine-silty, mixed, mesic	Typic Hapludalfs	Alfisols.
Sable	Fine-silty, mixed, mesic	Typic Haplaquolls	Mollisols.
St. Charles	Fine-silty, mixed, mesic	Typic Hapludalfs	Alfisols.
Sawmill	Fine-silty, mixed, mesic	Cumulic Haplaquolls	Mollisols.
Saybrook	Fine-silty, mixed, mesic	Typic Argiudolls	Mollisols.
Strawn	Fine-loamy, mixed, mesic	Typic Hapludalfs	Alfisols.
Virgil	Fine-silty, mixed, mesic	Udolic Ochraqualfs	Alfisols.
Will <sup>1</sup>	Fine-loamy over sandy or sandy-skeletal, mixed, mesic	Typic Haplaquolls	Mollisols.

<sup>1</sup> The Will soils in this county are taxadjuncts to the Will series, because they are calcareous rather than neutral in the upper part of the profile.

**Order.**—Ten soil orders are recognized. The properties used to differentiate among soil orders are those that tend to give broad climatic groupings of soils. Three exceptions to this are the Entisols, Histosols, and Vertisols, which occur in many different climates. Each order is named with a word of three or four syllables, ending in *sol* (Moll-i-sol).

**Suborder.**—Each order is divided into suborders, based on those soil characteristics that seem to produce classes that have the greatest genetic similarity. The suborders are more narrowly defined than the orders. The soil properties used to separate suborders are mainly those that reflect either the presence or absence of a high water table; soil climate; the accumulation of clay, iron, or organic carbon in the upper part of the solum; cracking of soils caused by a decrease in soil moisture; and fine stratification. The names of suborders have two syllables. The last syllable indicates the order. An example is *Aquoll* (*Aqu*, meaning water or wet; and *oll*, from Mollisol).

**Great group.**—Soil suborders are separated into great groups on the basis of uniformity in the kinds and sequence of soil horizons and features. The horizons used to make separations are those in which clay, carbonates, and other constituents have accumulated or been removed and those that have pans that interfere with growth of roots, movement of water, or both. Some features used are acidity, climate, composition, and color of the soil. The names of great groups have three or four syllables and are made by adding a prefix to the name of the suborder. An example is *Haplaquoll* (*Hapl*, meaning simple horizons; *aqu*, for wetness or water; and *oll*, from Mollisols).

**Subgroup.**—Great groups are divided into subgroups, one that represents the central (typic) segment of the group and others, called intergrades, that have properties of the group and also one or more properties of another great group, suborder, or order. Other subgroups may have soil properties unlike those of any other great group, suborder, or order. The names of subgroups are derived by placing one or more adjectives in front of the name of the great group. An example is *Typic Haplaquolls* (a typical Haplaquoll).

**Family.**—Soil families are established within a subgroup mainly on the basis of properties important to the growth of plants or to the behavior of soils when used for engineering. Among the properties considered are texture, mineralogy, reaction, soil temperature, permeability, soil depth, and consistence. A family name consists of a series of adjectives preceding the subgroup name. The adjectives are the class names for texture, mineralogy, and so on, that are used to differentiate families. An example is the fine-silty, mixed, mesic family of *Typic Haplaquolls*.

**Series.**—The series is a group of soils that have major horizons that, except for texture of the surface layer, are similar in important characteristics and in arrangement in the profile.

## Literature Cited

(1) American Association of State Highway [and Transportation] Officials. 1970. Standard specifications for highway

- materials and methods of sampling and testing. Ed. 10, 2 vol., illus.
- (2) American Society for Testing and Materials. 1974. Method for classification of soils for engineering purposes. ASTM Stand. D 2487-69. In 1974 Annual Book of ASTM Standards, Part 19, 464 pp., illus.
- (3) Barger, G. L., R. H. Shaw, and R. F. Dale. 1959. Chances of receiving selected amounts of precipitation in the north-central region of the United States. Iowa State Univ. Agric. Exp. Stn., 277 pp.
- (4) Flemel, R. C., K. C. Hinkley, and J. L. Hessler. 1973. The DeKalb mounds—a possible Wisconsin pingo field in north central Illinois, United States. In The Wisconsin Stage. Geol. Soc. Am., Inc. 327 pp., illus.
- (5) Huff, F. A. and S. A. Changnon. 1959. Hail climatology of Illinois. Ill. State Water Surv. Rep. Invest. 38, 46 pp.
- (6) Joos, L. A. 1960. Freeze probabilities in Illinois. Ill. Agric. Exp. Stn. in coop. Natl. Weather Serv., U.S. Dep. Commer. Bull. 650, 16 pp., illus.
- (7) United States Department of Agriculture. 1951. Soil survey manual. U.S. Dep. Agric. Handb. 18, 503 pp., illus. [Supplement issued May 1962]
- (8) ———. 1960. Soil classification, a comprehensive system, 7th approximation. Soil Conserv. Serv., 265 pp., illus. [Supplements issued March 1967, September 1968, April 1969]
- (9) University of Illinois Agricultural Experiment Station. 1965. Productivity of soils in the north-central region of the United States. North-Central Reg. Res. Publ. 166, Bull. 710, 27 pp., map.
- (10) William, H. B., and J. C. Frye. 1970. Pleistocene stratigraphy of Illinois. Ill. State Geol. Surv. Bull. 94, 204 pp., illus.

## Glossary

- Alluvium.** Soil material, such as sand, silt, or clay, that has been deposited on land by streams.
- Available water capacity** (also termed available moisture capacity). The capacity of soils to hold water available for use by most plants. It is commonly defined as the difference between the amount of soil water at field capacity and the amount at wilting point. It is commonly expressed as inches of water per inch of soil.
- Bottom land.** Nearly level land on the bottom of a valley that has a stream flowing through it. It is subject to flooding and is often referred to as a flood plain.
- Calcareous soil.** A soil that contains enough calcium carbonate (often with magnesium carbonate) to effervesce (fizz) visibly when treated with cold, dilute hydrochloric acid.
- Clay.** As a soil separate, the mineral soil particles less than 0.002 millimeter in diameter. As a soil textural class, soil material that is 40 percent or more clay, less than 45 percent sand, and less than 40 percent silt.
- Consistence, soil.** The feel of the soil and the ease with which a lump can be crushed by the fingers. Terms commonly used to describe consistence are—
- Loose.*—Noncoherent when dry or moist; does not hold together in a mass.
- Friable.*—When moist, crushes easily under gentle pressure between thumb and forefinger; can be pressed together into a lump.
- Firm.*—When moist, crushed under moderate pressure between thumb and forefinger, but resistance is distinctly noticeable.
- Plastic.*—When wet, readily deformed by moderate pressure but can be pressed into a lump; will form a “wire” when rolled between thumb and forefinger.
- Sticky.*—When wet, adheres to other material, and tends to stretch somewhat and pull apart, rather than to pull free from other material.
- Hard.*—When dry, moderately resistant to pressure; can be broken with difficulty between thumb and forefinger.
- Soft.*—When dry, breaks into powder or individual grains under very slight pressure.
- Cemented.*—Hard and brittle; little affected by moistening.
- Contour farming.** Plowing, planting, cultivating, and harvesting in rows that are at right angles to the natural direction of the slope or are parallel to terrace grade.
- Cover crop.** A close-growing crop grown primarily to improve the soil and protect it between periods of regular crop

- production; or grown between trees in orchards or between vines in vineyards.
- Crop residue.** The part of a plant, or crop, left in the field after harvest.
- Depth, soil.** Thickness of soil over a specified layer, generally one that does not permit the growth of roots.
- Classes used in this survey are—
- |                                  |                                   |
|----------------------------------|-----------------------------------|
| Deep—36 inches or more.          | Shallow—10 to 20 inches.          |
| Moderately deep—20 to 36 inches. | Very shallow—less than 10 inches. |
- Diversion, or diversion terrace.** A ridge of earth, generally a terrace, that is built to divert runoff from its natural course and, thus, to protect areas downslope from the effects of such runoff.
- Erosion.** The wearing away of the land surface by wind (sandblast), running water, and other geological agents.
- Fertility, soil.** The quality of a soil that enables it to provide compounds, in adequate amounts and in proper balance, for the growth of specified plants when other growth factors, such as light, moisture, temperature, and the physical condition of the soil, are favorable.
- Glacial till (geology).** Unsorted, nonstratified glacial drift that consists of clay, silt, sand, gravel, and boulders transported and deposited by glacial ice.
- Green manure crop (agronomy).** A crop of grasses or legumes grown for the purpose of being turned under in an early stage of maturity or soon after maturity to improve the soil.
- Horizon, soil.** A layer of soil, approximately parallel to the surface, that has distinct characteristics produced by soil-forming processes. These are the major horizons:
- O horizon.*—The layer of organic matter on the surface of a mineral soil. This layer consists of decaying plant residues.
- A horizon.*—The mineral horizon at the surface or just below an O horizon. This horizon is the one in which living organisms are most active and therefore is marked by the accumulation of humus. The horizon may have lost one or more of soluble salts, clay, and sesquioxides (iron and aluminum oxides).
- B horizon.*—The mineral horizon below an A horizon. The B horizon is in part of a layer of change from the overlying A to the underlying C horizon. The B horizon also has distinctive characteristics caused (1) by accumulation of clay, sesquioxides, humus, or some combination of these; (2) by prismatic or blocky structure; (3) by redder or stronger colors than the A horizon; or (4) by some combination of these. Combined A and B horizons are usually called the solum, or true soil. If a soil lacks a B horizon, the A horizon alone is the solum.
- C horizon.*—The weathered rock material immediately beneath the solum. In most soils this material is presumed to be like that from which the overlying horizons were formed. If the material is known to be different from that in the solum, a Roman numeral precedes the letter C.
- Leached soil.** A soil from which most of the soluble constituents have been removed from the entire profile or have been removed from one part of the profile and have accumulated in another part.
- Loess.** A fine-grained eolian deposit consisting dominantly of silt-sized particles.
- Mottled.** Irregularly marked with spots of different colors that vary in number and size. Mottling in soils usually indicates poor aeration and lack of drainage. Descriptive terms are as follows: Abundance—*few, common, and many*; size—*fine, medium, and coarse*; and contrast—*faint, distinct, and prominent*. The size measurements are these: *fine*, less than 5 millimeters (about 0.2 inch) in diameter along the greatest dimension; *medium*, ranging from 5 millimeters to 15 millimeters (about 0.2 to 0.6 inch) in diameter along the greatest dimension; and *coarse*, more than 15 millimeters (about 0.6 inch) in diameter along the greatest dimension.
- Munsell notation.** A system for designating color by degrees of the three simple variables—hue, value, and chroma. For example, a notation of 10YR 6/4 is a color with a hue of 10YR, a value of 6, and a chroma of 4.
- Natural soil drainage.** Refers to the conditions of frequency and duration of periods of saturation or partial saturation that existed during the development of the soil, as opposed to altered drainage, which is commonly the result of artificial drainage or irrigation but may be caused by the sudden deepening of channels or the blocking of drainage outlets. Seven different classes of natural soil drainage are recognized.
- Excessively drained* soils are commonly very porous and rapidly permeable and have a low water-holding capacity.
- Somewhat excessively drained* soils are also very permeable and are free from mottling throughout their profile.
- Well-drained* soils are nearly free from mottling and are commonly of intermediate texture.
- Moderately well drained* soils have uniform color in the A and upper B horizons and have mottling in the lower B and the C horizons.
- Somewhat poorly drained* soils are wet for significant periods but not all the time. They commonly have mottlings in the B and C horizons and the lower part of the A horizon in some soils.
- Poorly drained* soils are wet for long periods and are light gray and generally mottled from the surface downward, although mottling may be absent or nearly so in some soils.
- Very poorly drained* soils are wet nearly all the time. They have a dark gray or black surface layer and are gray or light gray, with or without mottling, in the deeper parts of the profile.
- Organic-matter content.** The content of plant and animal material, in or on the soil, in all stages of decomposition. Ratings used in this survey have the following limits: low—below 2 percent of volume; moderate—2 to 4 percent; and high—more than 4 percent.
- Permeability, soil.** The quality of a soil horizon that enables water or air to move through it. The permeability of a soil may be limited by the presence of one nearly impermeable horizon even though the others are permeable. Terms used to describe permeability are as follows: *Very slow, slow, moderately slow, moderate, moderately rapid, rapid, and very rapid*.
- Reaction, soil.** The degree of acidity or alkalinity of a soil expressed in pH values. A soil that tests to pH 7.0 is precisely neutral in reaction because it is neither acid nor alkaline. An acid, or "sour," soil is one that gives an acid reaction; an alkaline soil is one that is alkaline in reaction. In words, the degrees of acidity or alkalinity are expressed thus:
- |                    |            |                        |                |
|--------------------|------------|------------------------|----------------|
|                    | pH         |                        | pH             |
| Extremely acid     | Below 4.5  | Mildly alkaline        | 7.4 to 7.8     |
| Very strongly acid | 4.5 to 5.0 | Moderately alkaline    | 7.9 to 8.4     |
| Strongly acid      | 5.1 to 5.5 | Strongly alkaline      | 8.5 to 9.0     |
| Medium acid        | 5.6 to 6.0 | Very strongly alkaline | 9.1 and higher |
| Slightly acid      | 6.1 to 6.5 |                        |                |
| Neutral            | 6.6 to 7.3 |                        |                |
- Sand.** Individual rock or mineral fragments in soils having diameters ranging from 0.05 to 2.0 millimeters. Most sand grains consist of quartz, but they may be any mineral composition. The textural class name of any soil that contains 85 percent or more sand and not more than 10 percent clay.
- Silt.** Individual mineral particles in a soil that range in diameter for the upper limit of clay (0.002 millimeter) to the lower limit of very fine sand (0.05 millimeter). Soil of the silt textural class is 80 percent or more silt and less than 12 percent clay.
- Soil.** A natural, three-dimensional body on the earth's surface that supports plants and that has properties resulting from the integrated effect of climate and living matter acting on earthy parent material, as conditioned by relief over periods of time.
- Solum.** The upper part of a soil profile, above the parent material, in which the processes of soil formation are active. The solum in mature soil formation are active. The solum in mature soil includes the A and B horizons. Generally, the characteristics of the material in these horizons are unlike those of the underlying material. The living roots and other plant and animal life characteristic of the soil are largely confined to the solum.
- Stratified.** Composed of, or arranged in, strata, or layers, such as stratified alluvium. The term is confined to geological materials. Layers in soils that result from the processes of soil formation are called horizons; those inherited from the parent material are called strata.
- Structure, soil.** The arrangement of primary soil particles into compound particles or clusters that are separated from

adjoining aggregates and have properties unlike those of an equal mass of unaggregated primary soil particles. The principal forms of soil structure are—*platy* (laminated), *prismatic* (vertical axis of aggregates longer than horizontal), *columnar* (prisms with rounded tops), *blocky* (angular or subangular), and *granular*. *Structureless* soils are (1) *single grained* (each grain by itself, as in dune sand) or (2) *massive* (the particles adhering together without any regular cleavage, as in many claypans and hardpans).

**Subsoil.** Technically, the B horizon; roughly, the part of the solum below plow depth.

**Substratum.** Technically the part of the soil below the solum.

**Subsurface layer.** The horizon between the surface layer and the subsoil, generally, the A2 horizon.

**Surface layer.** A term used in nontechnical soil descriptions for one or more layers above the subsoil. Includes the A horizon and, in places, part of the B horizon; has no depth limit.

**Terrace (geological).** An old alluvial plain, ordinarily flat or undulating, bordering a river, lake, or the sea. Stream terraces are frequently called second bottoms, as contrasted

to flood plains, and are seldom subject to overflow. Marine terraces were deposited by the seas and are generally wide.

**Texture, soil.** The relative proportions of sand, silt, and clay particles in a mass of soil. The basic textural classes, in order of increasing proportion of fine particles, are *sand*, *loamy sand*, *sandy loam*, *loam*, *silt loam*, *silt*, *sandy clay loam*, *clay loam*, *silty clay loam*, *sandy clay*, *silty clay* and *clay*. The sand, loamy sand, and sandy loam classes may be further divided by specifying "coarse," "fine," or "very fine."

**Tilth, soil.** The condition of the soil in relation to the growth of plants, especially soil structure. Good tilth refers to the friable state and is associated with high noncapillary porosity and stable, granular structure. A soil in poor tilth is nonfriable, hard, nonaggregated, and difficult to till.

**Upland (geology).** Land consisting of material unworked by water in recent geologic time and lying, in general, at a higher elevation than the alluvial plain or stream terrace. Land above the lowlands along rivers.

**Water table.** The highest part of the soil or underlying rock material that is wholly saturated with water. In some places an upper, or perched, water table may be separated from a lower one by a dry zone.

GUIDE TO MAPPING UNITS

For a complete description of a mapping unit, read both the description of the mapping unit and the description of the soil series to which the mapping unit belongs. Other information is given in tables as follows:

Acreage and extent, table 3,  
page 10.  
Predicted yields, table 4, page 37.  
Guide for tree and shrub plantings,  
table 5, page 38.

Wildlife, table 6, page 40.  
Recreational uses of the soils, table 7,  
page 42.  
Engineering uses of the soils, tables 8, 9,  
and 10, pages 44 through 61.

Map symbol	Mapping unit	De- scribed on page	Management group		Tree and shrub planting group	Recreation group
			Symbol	Page	Number	Number
24A	Dodge silt loam, 0 to 2 percent slopes-----	14	I-1	34	1	1
24B	Dodge silt loam, 2 to 4 percent slopes-----	14	IIe-1	34	1	1
24C2	Dodge silt loam, 4 to 7 percent slopes, eroded-----	14	IIe-4	35	1	1
27B	Miami silt loam, 2 to 4 percent slopes-----	23	IIe-1	34	2	1
27C2	Miami silt loam, 4 to 7 percent slopes, eroded-----	24	IIe-4	35	2	1
27D2	Miami silt loam, 7 to 12 percent slopes, eroded-----	24	IIIe-1	35	2	2
41	Muscatine silt loam-----	25	I-3	34	3	3
59	Lisbon silt loam-----	23	I-3	34	3	3
60B	La Rose silt loam, 2 to 4 percent slopes-----	22	IIe-2	34	2	1
60C2	La Rose silt loam, 4 to 7 percent slopes, eroded-----	22	IIe-4	35	2	1
60D2	La Rose silt loam, 7 to 12 percent slopes, eroded-----	22	IIIe-1	35	2	2
62	Herbert silt loam-----	20	I-3	34	3	3
67	Harpster silty clay loam-----	18	IIw-1	35	4A	4
68	Sable silty clay loam-----	29	IIw-1	35	4	4
103	Houghton muck-----	20	IIIw-2	36	5	5
104	Virgil silt loam-----	32	I-3	34	3	3
105A	Batavia silt loam, 0 to 2 percent slopes-----	11	I-2	34	1	1
105B	Batavia silt loam, 2 to 4 percent slopes-----	11	IIe-2	34	1	1
105C2	Batavia silt loam, 4 to 7 percent slopes, eroded-----	11	IIe-4	35	1	1
V107	Sawmill soils-----	30	IIw-1	35	4	4
134A	Camden silt loam, 0 to 2 percent slopes-----	12	I-1	34	1	1
134B	Camden silt loam, 2 to 6 percent slopes-----	12	IIe-1	34	1	1
145B	Saybrook silt loam, 2 to 4 percent slopes-----	31	IIe-2	34	1	1
145C2	Saybrook silt loam, 4 to 7 percent slopes, eroded-----	31	IIe-4	35	1	1
148A	Proctor silt loam, 0 to 2 percent slopes-----	27	I-2	34	1	1
148B	Proctor silt loam, 2 to 4 percent slopes-----	28	IIe-2	34	1	1
152	Drummer silty clay loam-----	15	IIw-1	35	4	4
154	Flanagan silt loam-----	17	I-3	34	3	3
171A	Catlin silt loam, 0 to 2 percent slopes-----	13	I-2	34	1	1
171B	Catlin silt loam, 2 to 4 percent slopes-----	13	IIe-2	34	1	1
171C2	Catlin silt loam, 4 to 7 percent slopes, eroded-----	13	IIe-4	35	1	1
191	Knight silt loam-----	21	IIw-1	35	4	4
198	Elburn silt loam-----	17	I-3	34	3	3
199A	Plano silt loam, 0 to 2 percent slopes-----	27	I-2	34	1	1
199B	Plano silt loam, 2 to 4 percent slopes-----	27	IIe-2	34	1	1
219	Millbrook silt loam-----	24	I-3	34	3	3
224E2	Strawn soils, 10 to 20 percent slopes, eroded-----	31	VIe-1	36	2	2
242	Kendall silt loam-----	21	I-3	34	3	3
243A	St. Charles silt loam, 0 to 2 percent slopes-----	29	I-1	34	1	1
243B	St. Charles silt loam, 2 to 4 percent slopes-----	29	IIe-1	34	1	1
318D2	Lorenzo loam, 6 to 15 percent slopes, eroded-----	23	IVe-1	36	2	2
325B	Dresden silt loam, 2 to 4 percent slopes-----	15	IIe-3	35	1	1
325C2	Dresden silt loam, 4 to 7 percent slopes, eroded-----	15	IIIe-2	35	1	1
329	Will clay loam-----	32	IIw-1	35	4	4
330	Peotone silty clay loam-----	26	IIIw-1	36	4	4
344A	Harvard silt loam, 0 to 2 percent slopes-----	19	I-2	34	1	1
344B	Harvard silt loam, 2 to 4 percent slopes-----	19	IIe-2	34	1	1
344C2	Harvard silt loam, 4 to 7 percent slopes, eroded-----	19	IIe-4	35	1	1

GUIDE TO MAPPING UNITS--Continued

Map symbol	Mapping unit	De- scribed on page	Management group		Tree and shrub planting group	Recreation group
			Symbol	Page	Number	Number
656B	Octagon silt loam, 2 to 4 percent slopes-----	26	Iie-2	34	2	1
656C2	Octagon silt loam, 4 to 7 percent slopes, eroded-----	26	Iie-4	35	2	1
791A	Rush silt loam, 0 to 2 percent slopes-----	28	I-1	34	1	1
791B	Rush silt loam, 2 to 4 percent slopes-----	28	Iie-1	34	1	1
792A	Bowes silt loam, 0 to 2 percent slopes-----	12	I-2	34	1	1
792B	Bowes silt loam, 2 to 4 percent slopes-----	12	Iie-2	34	1	1



# NRCS Accessibility Statement

---

This document is not accessible by screen-reader software. The Natural Resources Conservation Service (NRCS) is committed to making its information accessible to all of its customers and employees. If you are experiencing accessibility issues and need assistance, please contact our Helpdesk by phone at 1-800-457-3642 or by e-mail at [ServiceDesk-FTC@ftc.usda.gov](mailto:ServiceDesk-FTC@ftc.usda.gov). For assistance with publications that include maps, graphs, or similar forms of information, you may also wish to contact our State or local office. You can locate the correct office and phone number at <http://offices.sc.egov.usda.gov/locator/app>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.