

QUAILBUSH

Atriplex lentiformis (Torr.)

S. Wats

plant symbol = ATLE

Contributed By: USDA, NRCS, National Plant Data Center



© Barbara J. Collins
California Lutheran University

Alternative Names

big saltbush

Uses

Ethnobotanic: The Native American Pima groups eat quailbush seeds. They ground the seeds into a meal and used them as a thickener in soups or added them to flour for making bread. Most of this shrub is edible, young shoots are suitable for greens. Several tribes used this shrub for its salty taste. The crushed leaves and roots were used as soap for washing clothes (Moerman 1998).

Native Americans tribes ground the roots and flowers and applied it to ant bites. The leaves were chewed to treat head colds. The crushed flowers and stems can be steamed and inhaled to treat nasal congestion (Moerman 1998).

Wildlife: Rabbits, lizards, rattlesnakes, coyotes, quails, and other birds use the seeds and foliage for food and habitat. The foliage and twigs provide shelter for many small mammals and livestock.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

General: Quailbush is a large, fast growing shrub. It occurs in river floodplains, on roadsides, and in the borders of drainage. The branches are widespread, slender, and flexible. The leaves are alternate, triangular or ovate to oblong, and are 1 ½ to 2 inches long. The plants are male or female and have the ability to alter their sex due to environmental conditions.

Distribution: It ranges from the Upper San Joaquin and Salinas Valley southward to lower California, in Lower and Upper Sonoran Life Zones (McMinn 1939). It extends eastward into Nevada, Utah, Colorado, and New Mexico. For current distribution, please consult the Plant profile page for this species on the PLANTS Web site

Adaptation

Quailbush grows best with full sunlight in any well-drained but not too fertile soil. It tolerates very alkaline soils and can succeed in hot and dry climates. This species is not often found in colder areas of the country but it can tolerate temperatures between -5 and -10° C.

Establishment

Propagation from Seed: The seed is best sown in April or May and placed in containers or seed trays containing a compost of peat and sand to which a slow-release fertilizer has been added. Firm the medium gently, sow the seed thinly and evenly on top, and cover with its own depth of medium (Heuser 1997). Place the pots in a cold frame at 13° C and the seed should germinate between one to three weeks. The seedlings should be placed into individual pots and grown in a greenhouse for the first winter.

Management

Atriplex lentiformis will defoliate under extreme drought conditions. They need to be under some form of water stress, salt stress, or drought stress. The salt they accumulate in their leaves allows them to extract water from the soil. They tolerate and remove the excess salts by bladders in their leaves that act as salt sinks, keeping the salt from the plant cells.

Cultivars, Improved and Selected Materials (and area of origin)

'Casa' - Released 1979 by the Lockeford Plant Materials Center, CA, this cultivar has exhibited excellent performance as a conservation plant on various critical areas, for upland game cover and for environmental enhancement on deep, medium, or fine-textured soils that are well to poorly drained. It can be grown on slightly acidic to strongly alkaline soils (pH 6-8.5) and survive on an annual precipitation of 20-25 cm (8-10 inches) when irrigated for initial establishment. This species occurs from the upper San Joaquin and Salinas valleys south to lower California. It extends eastward into Nevada, Utah, and New Mexico.

References

- Collins, B.J. 2001. *Wildflowers of Southern California*. California Lutheran University, Thousand Oaks, California. Accessed: 11Jan02. <<http://www1.clunet.edu/wf/index.htm>>
- Heuser, C.W. 1997 *The complete book of plant propagation*. The Taunton Press, Newtown, Connecticut.
- Junah, S., T. Ayers, R. Scott, D. Wilken, & D. Young 1995. *A flora of San Cruz Island*. Santa Barbara Botanic Garden, Santa Barbara, California.
- Kearney, T.H., R.H. Peebles, J.H. Howell, & E. McClintock 1960. *Arizona flora*. 2nd ed. University of California Press, Berkeley, California.
- McMinn, H.E. 1939. *An illustrated manual of California shrubs*. University of California Press, Berkeley, Los Angeles, & London.
- Moerman, D. 1998. *Native American ethnobotany*. Timber Press, Oregon.
- Mozingo, H.N. 1987. *Shrubs of the Great Basin: a natural history*. University of Nevada Press, Reno, Las Vegas & London.

Thomas, J.H. 1961. *Flora of the San Cruz mountains of California: a manual of the vascular plants*. Stanford University Press, Stanford, California.

Van Dersal, W.R. 1939. *Native woody plants of the United States, their erosion control & wildlife values*. United States Department of Agriculture, Washington, D.C.

Vines, R.A. 1960. *Trees, shrubs, and woody vines of the southwest*. University of Texas Press, Austin, Texas.

Prepared By

Jammie Favorite
formerly, USDA, NRCS, National Plant Data Center
Baton Rouge, Louisiana

Species Coordinator

Lincoln M. Moore
USDA, NRCS, National Plant Data Center
Baton Rouge, Louisiana

Edited: 09Jan02 jsp; 25Feb03 ahv; 14Mar05 rln

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.