

# TECHNICAL NOTE

---

USDA-Natural Resources Conservation Service  
Boise, Idaho – Salt Lake City, Utah

---

TN PLANT MATERIALS NO. 26

OCTOBER 2010


## Legume Inoculation

Ron Miller, State Agronomist, NRCS, Columbia, MO  
Jerry Kaiser, Plant Materials Specialist, NRCS, Elsberry, MO  
Daniel G. Ogle, Plant Materials Specialist, NRCS, Boise, ID

The legume family, *Fabaceae*, have the ability to react in a symbiotic (mutually beneficial) relationship with unique strains of *Rhizobia* bacteria to convert atmospheric nitrogen ( $N_2$ ) from the soil atmosphere to ammonia nitrogen ( $NH_3$ ). Plant available nitrogen created by this relationship between a growing plant and soil bacteria is virtually free to the producer. This additional nitrogen from *Rhizobia* bacteria improves soil fertility, produces higher protein forage, and increases forage yields in the host plant and in many cases for associated species and future crops.

The correct strain of bacteria will infect the roots of the legume to form nodules (see Figure 1). The bacteria work within the nodules to transform atmospheric nitrogen to plant available nitrogen. The interaction involves the exchange of water and nutrients for nitrogen with both the plant and bacteria benefitting. The host plant is the only plant that will benefit from the nodules growing on live roots. The roots must die and roots and nodules must decay to make nitrogen available to other plants and soil organisms.

The ability to acquire nitrogen from the atmosphere is a trait that allows legumes to be grown without the addition of nitrogen fertilizer. Growing legumes in a crop rotation or forage management system typically adds nitrogen to the soil for subsequent crops or production periods. In order for this symbiotic relationship to establish and develop; the appropriate *Rhizobia* bacteria must be present in the soil as the legume's roots grow and develop. Soil contains



*Figure 1 – Roots with nodules*

millions of bacteria, but *Rhizobia* bacteria proliferate best in soils that are planted with the correct host legume crop. When a legume is planted on a particular soil for the first time, there is a very good chance that the correct *Rhizobia* strain will not be present. Successful formation of nodules requires inoculation of the legume seed with the appropriate bacteria at planting and placing the seed in the soil in contact with the inoculant.

The producer should follow the manufacturer’s recommendations regarding the strain of inoculum and the amount to be added with the legume seed. The inoculant package will list the appropriate legume species. Due to the different formulations and carriers used in developing the inoculant, each manufacturer will have a different requirement for the amount needed.

**Benefits of Inoculation**

- Increased production – forage yields may increase from 10 to 100% depending on specific soil conditions; soils with average fertility may have yield increases from 15 to 25%.
- Improved protein content – inoculated alfalfa has tested 3.5% higher for protein.
- Increases soil nitrogen for future crops – nitrogen fixation will vary depending on plant species, suitable bacterium population, soil nitrogen content, soil fertility level, soil pH, moisture, and temperature. Average nitrogen values are shown below:

<u>Legume Species</u>	<u>Pounds of N Fixed/Ac/Yr</u>
Alfalfa	140 to 200
Ladino clover	130 to 185
Alsike clover	70 to 125
Red clover	62 to 119
Birdsfoot trefoil	44 to 100
Soybeans	80 to 128

**Types of Inoculation Procedures**

Inoculants may be available in dry, granular, or liquid formulations. Inoculants can be mixed and applied with the seed by one of the following methods:

1. Pre-inoculated seed – The inoculant is pre-applied to the seed prior to sale to the producer. Seed inoculated in this manner must be stored under cool conditions until immediately prior to planting to insure viability of the bacteria.
2. Slurry Method – The dry, granular, or liquid inoculant is mixed with the seed in a wash tub or barrel along with a liquid to form a slurry. Water is the most common liquid used. The slurry is transferred to the planter box for planting. Various commercial “stickers” are also available and can be used with the slurry method to aid in adhering bacteria to the seed. This method is generally considered to be the most effective of the in-field inoculation methods. Make sure that all seed and liquid is removed from the planter boxes at the end of the day.
3. Sprinkle Method – Seed is pre-moistened with water or a “sticker” product in the planter hopper, a wash tub, or a barrel prior to mixing in the dry or granular inoculant formulations; liquid inoculant can be used without additional water or sticker products. Allow the seed to dry in a cool- shady area before attempting to plant. This method is not considered as good as the slurry method but can be adequate under good planting conditions. Again, remove all seed from the planter boxes at the end of the day.

Refer to Table 1 for a list of manufacturers and the inoculants they produce. Refer to Table 2 for a list of the specific *Rhizobia* bacteria that are required for a legume species or group of species.

**IMPORTANT NOTE** - Inoculants are perishable. They should be transported and stored in a cool, dry place and never used past the date recommended by the manufacturer.

### **Points to Remember**

- Inoculation of legumes with the appropriate *Rhizobia* bacteria is needed to maximize biological nitrogen fixation.
- Based on the relatively inexpensive cost of inoculants and the high cost of nitrogen fertilizers, the addition of inoculants to a legume species is a wise investment in fertilizer management and crop production.
- Select the *Rhizobia* inoculant or mixture of inoculants based on the legume species that will be planted. Do not use an inoculant if the legume species is not listed on the package or company literature as appropriate.
- Do not allow the inoculant to be exposed to direct sunlight or heat when transporting.
- Store the inoculant according to the manufacturer's recommendations (refrigeration may be required but do not freeze).
- Mix the seed and inoculant in a shady location out of direct sunlight.
- Plant seed within 24 hours after mixing with inoculant or re-inoculate.
- Never mix the inoculant with any kind of fertilizer or pesticide unless specifically labeled for that use. Avoid contact with agricultural lime.
- Effective sticking agents for use with inoculants include 10 percent solutions in water of powdered milk, corn syrup, molasses, or sugar along with the "commercial sticker" products that are available. Using carbonated liquids or compounds that are acidic or basic will result in poor inoculant survival.
- Soil moisture is very important for successful inoculation and nodulation. Do not plant inoculated seed into dry soil.
- Soil temperature and soil pH can also affect the survival and ultimate effectiveness of inoculants.
- Not all root nodules fix nitrogen because there are other soil bacteria that form root nodules. Slice through a few nodules and look at the color of the cut surface. Red or pink color indicates the ability to fix nitrogen. White color indicates ineffective bacteria. Green color indicates nodules that are no longer producing.
- No simple test exists to determine whether suitable *Rhizobia* are already present in the soil prior to planting.
- A relatively small investment of time and money in inoculant and sticker products will ensure that the correct bacteria species is present for a legume to produce nitrogen.
- Because inoculants are cultures of living organisms, contact suppliers well in advance of the planned seeding dates to ensure viable inoculants will be available.

**TABLE 1 – Available Sources of Inoculants**

Seed dealers can supply compatible Rhizobia bacteria species for each legume listed in Table 1. The information in this table is presented merely to show various manufacturers of inoculants but may not list all manufacturers. Trade names are used solely to provide information and should not be considered a recommendation or endorsement by the Natural Resources Conservation Service.

Legume Species	Inoculant Manufacturers					
	Novozymes	Becker Underwood, Inc.	CelPril - Bayer Crop Science	EMD Crop Bioscience	INTX Microbials, LLC	Plant Probotics
Alfalfa	X	X	X	X	X	X
Birdsfoot trefoil		X			X	X
Cicer milkvetch						
Alsike clover		X		X	X	X
Ladino clover		X		X	X	X
Red clover		X		X	X	X
Kura clover						X
Lespedeza		X			X	X
Field peas						
Partridge pea		X			X	X
Sainfoin						
Soybeans						

**Addresses and contact information for the manufacturers in Table 1**

Philom Bios/Novozymes  
 3935 Thatcher Avenue  
 Saskatoon, SK Canada S7R1A3  
 Phone: 888.744.5662  
 Fax: 306.975.1215  
 Phone: 952.361.3495  
 Cell: 952.913.3941

Nitragin/EMD Crop Bioscience  
 13100 West Lisbon Avenue  
 Suite 600  
 Brookfield, WI 53005  
 Phone: 262.957.2000  
 Fax: 262.957.2121

Becker Underwood, Inc.  
 801 Dayton Avenue  
 Ames, IA 50010  
 Phone: 800.232.5907  
 FAX: 515.232.5961

INTX Microbials, LLC  
 200 W. Seymour  
 P.O. Box 62  
 Kentland, IN 47951  
 Phone: 219.474.5510  
 FAX: 219.474.3700

CelPril – Bayer Crop Sciences  
 251 Oak Street  
 Manteca, CA 95337  
 Phone: 209.823.1738  
 FAX: 209.823.8855

Plant Probotics  
 6835 Lindel Court  
 Indianapolis, IN 46268  
 Phone: 317.329.7731

**TABLE 2 - Legume Cross - Inoculation Groups**

<u>Legume</u>	<u>Inoculation Species</u>
Alfalfa Sweetclover (yellow and white)	<i>Sinorhizobium meliloti</i> or <i>Rhizobium mongolense</i>
Birdsfoot Trefoil	<i>Mesorhizobium loti</i> or <i>Rhizobium loti</i>
Cicer Milkvetch	<i>Astragalus spp.</i> – specific <i>Rhizobium</i>
Clovers ( <i>Trifolium spp.</i> ) Alsike clover Ladino and White clover Red clover Strawberry clover	<i>Rhizobium leguminosarum trifolii</i>
Crownvetch	<i>Mesorhizobium spp.</i>
Kura Clover	<i>Rhizobium spp.</i> for Kura Clover
Prairie Clover	<i>Rhizobium spp.</i> for Prairie Clover
Field Peas (Austrian, dry, green) Field Beans Lentils Hairy Vetch and other Vetch species	<i>Rhizobium leguminosarum viciae</i>
Medic species	<i>Sinorhizobium medicae</i>
Lespedeza Partridge Pea	<i>Bradyrhizobium spp.</i> for Lespedeza
Sainfoin	<i>Rhizobium spp.</i> for Sainfoin
Soybeans	<i>Bradyrhizobium japonicum</i> or <i>Bradyrhizobium elkanii</i> or <i>Bradyrhizobium liaoningense</i> or <i>Sinorhizobium fredii</i>

NOTE: An attempt has been made to include all *Rhizobia* manufacturers and their current contact information in this publication. Please notify Ron Miller at [ron.miller@mo.usda.gov](mailto:ron.miller@mo.usda.gov) or Dan Ogle at [dan.ogle@id.usda.gov](mailto:dan.ogle@id.usda.gov) with the necessary information on any manufacturer that was not included. Errors in this listing are not intentional.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.