

UNITED STATES DEPARTMENT OF AGRICULTURE
 NATURAL RESOURCES CONSERVATION SERVICE
 PLANT MATERIALS CENTER
 CORVALLIS, OREGON

and

INSTITUTE FOR APPLIED ECOLOGY
 CORVALLIS, OREGON

NOTICE OF RELEASE OF WILLAMETTE VALLEY GERMPLASM ROEMER'S
 FESCUE

SELECTED CLASS OF MANIPULATED TRACK GERMPLASM

The U.S. Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS) and the Institute for Applied Ecology announce the release of a selected class pre-variety of Roemer's fescue (*Festuca roemeri* var. *roemeri* [Pavlick] E.B. Alexeev, synonym: *Festuca idahoensis* Elmer ssp. *roemeri* [Pavlick] S. Aiken).

This population will be referred to as Willamette Valley Germplasm Roemer's fescue. It has been assigned USDA NRCS accession number 9079510 and PI 665693. The Oregon Seed Certification Service designated it as PVGOR102. Willamette Valley Germplasm is released as a selected class of certified seed and plants, manipulated track. There is a conservation need for this material at this level of development and additional selection is not warranted.

Origin: Parental G0 seed of Willamette Valley Germplasm was collected from nine naturally occurring stands (populations) of Roemer's fescue growing within the Willamette Valley Level III ecoregion (USEPA, 2011). At least eight plants were sampled from each location. Collections were first made in 2001-2002 under the coordination of the Institute for Applied Ecology. The nine collection sites ranged in elevation from 110 to 335 m (360–1100 ft) above sea level in an area where the average annual precipitation is between 102 and 127 cm (40–50 in).

Table 1. Origin of nine parent populations comprising Willamette Valley Germplasm

Abiqua Road	Marion Co., OR	Elevation 450 ft	Lat. 45.0076°N	Long. -122.6752°W
Bald Hill	Benton Co., OR	Elev. 460-500 ft	Lat. 44.5637- 44.5662°N	Long. -123.3283- 123.3396°W
Baskett Butte	Polk Co., OR	Elevation 360 ft	Lat. 44.9651°N	Long. -123.2579°W
Fire Knob	Marion Co., OR	Elevation 700 ft	Lat. 44.8925°N	Long. -122.8153°W
Kloster Mountain	Lane Co., OR	Elevation 875 ft	Lat. 43.867°N	Long. -122.8428°W
Rock Hill	Lane Co., OR	Elevation 800 ft	Lat. 44.1843°N	Long. -123.0531°W
Row Point	Lane Co., OR	Elevation 840 ft	Lat. 43.7882°N	Long. -122.9437°W
Spencer Butte	Lane Co., OR	Elevation 400 ft	Lat. 44.0581°N	Long. -123.0945°W
Weiss Road ridgeline	Lane Co., OR	Elevation 1100 ft	Lat. 43.8988°N	Long. 123.1329°W

Method of selection: From 2003 to 2006 a common-garden study of Roemer's fescue was conducted at the USDA NRCS, Corvallis Plant Materials Center (Benton County, Oregon; elevation 69 m [225 ft]; moderately well-drained Woodburn silt loam; mean annual precipitation 107 cm [42 in]). The study was a cooperative effort among the Institute for Applied Ecology, USDA NRCS, USDI Bureau of Land Management, and USDA Forest Service. Results documented substantial genetic variation within and among 47 populations from throughout the range of the species in the Pacific Northwest, USA, for growth, fitness, phenological, and morphological traits (Tables 2 and 3). Using climatic and physiographic variables, genetic patterns over the landscape were examined through principal component and regression analysis. Elevation and latitude of the seed source, and to a lesser extent temperature and precipitation, explained a significant proportion of the genetic variation, suggesting that observed variation was associated with adaptation to local environments. Roemer's fescue variation clustered into seed transfer zones corresponding to EPA Level III or Level IV ecoregions (Wilson et al., 2008).

In order to incorporate high genetic diversity, nine populations from a total of 17 found within the Willamette Valley seed transfer zone were selected to represent the Willamette Valley Germplasm (Table 1). Populations were eliminated for being too high in elevation (above 914 m or 3000 ft), or having atypical adaptation or morphology, low seed yield, or poor survival (Tables 2 and 3). A portion of individuals within each selected population overlapped in flowering periods despite a 12 day range in mean flowering dates among populations (Table 2. Data for individual families within each population are not shown). This suggests some plants within each population have the opportunity to randomly cross with certain individuals from all other populations. Random mating among the nine populations was promoted by using a Latin Square crossing block of G1 plants in order to form a genetically diverse polycross and potentially reduce or eliminate inbreeding depression suspected within smaller Willamette Valley populations (Darris et al., 2008; Wilson et al., 2008). Seed production, testing, and conservation use involves G2 seed and later generations.

Description: Willamette Valley Germplasm Roemer's fescue is a native cool season, perennial bunchgrass with mostly basal foliage. The foliage is fine textured and densely tufted (cespitose). Morphological diversity of plants within this germplasm is high. Leaves are sometimes glaucous (covered with a whitish waxy coating) and their color varies throughout a wide spectrum of green to pale blue. Plants rarely flower until the second full growing season. Stem color ranges from yellowish green to green, dark purple, or red, turning mostly straw colored at seed maturity. The stiff, erect, smooth culms grow from 70 to 115 cm (28–45 in) tall and the basal width is 10 to 18 cm (4–7 in) (Table 2).

The botanical traits for the polycross of nine populations that form Willamette Valley Germplasm Roemer's fescue fall within the taxonomic limits of *Festuca roemeri* var. *roemeri* (Wilson, 2007). The panicle (seed head or inflorescence) is loosely open to moderately contracted and 5–20 cm (2–8 in) long; branches are erect to slightly spreading. The spikelets are 9–13.5 mm (0.35–0.53 in) long with 4–6 florets. Glumes are exceeded by the upper florets. Lemma awns are 2–5 mm (0.08–0.20 in) long. Leaf sheaths are closed for less than half their length and collars are glabrous; ligules are 0.1–

0.5 mm (0.004–0.02 in) long; blades are 0.5–1.2 mm (0.02–0.05 in.) in diameter (Darbyshire and Pavlick, 2007). Roemer's fescue is an allotetraploid ($2n=28$) and is highly cross-pollinated by wind.

The mean anthesis and seed maturity periods for populations comprising Willamette Valley Germplasm range in Julian dates from 134 to 146 (May 14–May 26) and 169 to 177 (June 18–June 26), respectively. Julian date is the number of days counted from January 1st as day 1. Moderately low to high incidences of stem rust (14–51%) (*Puccinea* spp.) observed on G1 plants of the parent populations persisted among G2 plants (Table 2).

Suggested area of use: Based on results of the common garden study, the origin of parent populations and what is known about the species, Willamette Valley Germplasm Roemer's fescue is primarily recommended for use within the Willamette Valley Level III ecoregion of western Oregon and southwest Washington below an elevation of 762 m (2500 ft). Movement of the germplasm within this provisional seed transfer zone should pose minimal risk of maladaptation. This zone lies within USDA Plant Hardiness Zones 8a and 8b (U.S. National Arboretum, 2012). Pending further investigations on adaptation, area of use may extend into adjacent ecoregions for select purposes outside of restoration. Use should not extend beyond the species natural range.

As with the species in general, Willamette Valley Germplasm is likely to grow best in full sun and part shade near trees on moderately well to well-drained, medium to fine-textured soils that are moderately acidic to slightly basic. Although Roemer's fescue is considered drought tolerant and has extensive roots, it favors more mesic rather than xeric habitats within a site. While the species is known for tolerance to serpentine soils, moderate to low fertility conditions, and wildfire, the specific adaptation of Willamette Valley Germplasm to such extremes remains to be tested.

Anticipated use: Recommended uses for Willamette Valley Germplasm include restoration of upland prairies, oak savanna, and similar native plant communities, as well as wildlife habitat improvement. This includes food or cover for various song and game birds, small mammals, and beneficial insects. Roemer's fescue can be sown alone or in a mixture with forbs or other grasses. After further evaluation, other uses may include general revegetation and erosion control of disturbed sites and roadsides where a fine textured perennial grass is desired, low input turf, and cover crop for vineyards or young orchards. Roemer's fescue is a potential native alternative to introduced sheep, hard, and chewings red fescues in several applications. Potential as a forage grass may be similar to Idaho fescue but remains to be tested.

Ecological considerations and evaluation: Willamette Valley Germplasm is the result of minor purposeful selection and open crossing of nine populations for one generation for the limited purpose of improving fitness and genetic diversity over single wild populations. Many wild populations in the Willamette Valley have become small, have apparent lower diversity, and show reduced fitness possibly the result of inbreeding depression stemming from habitat loss and isolation of populations (Wilson et. al., 2008). While this germplasm may be genetically enhanced over such populations of Roemer's fescue, it is not anticipated to have substantial genetic gain in terms of ease of

establishment, reproductive capability, and plant growth over other naturally occurring populations found within the same ecoregion that remain fit and diverse.

This germplasm is recommended for use primarily within the ecoregion and elevation range (Willamette Valley seed transfer zone) from which the nine parent populations originated. Within this area and adjacent seed zones, the species is not considered weedy and is widely accepted as beneficial rather than detrimental to natural plant communities. According to the NRCS “Worksheet for Environmental Evaluation of Plant Releases” as applied to the intended area and type of use, this germplasm was deemed to have a low chance of adversely affecting the environment, be easy to control, have a moderate level of importance for conservation use, and a moderate chance to propagate and maintain itself under natural conditions. Willamette Valley Germplasm is not necessarily intended to replace on-site sources of Roemer’s fescue for ecological restoration plantings. Individuals with such concerns for a particular environment or ecosystem should make their decisions on a case by case basis. Uses beyond habitat restoration should be tested using standard precautions.

Willamette Valley Germplasm Roemer’s fescue is a known host of stem rust disease of grasses (*Puccinea graminis*). It may be also be host to other rusts (*Puccinea* spp.) and trace amounts of ergot (*Claviceps purpurea*) have been observed in similar populations of Roemer’s fescue. The rust and ergot potentially carried by this species occur commonly on other introduced and endemic grasses in the suggested area of use. Therefore, their occurrence on Willamette Valley Germplasm, if detected, is not considered to pose a unique problem or special risk to other new or already susceptible species. In the unlikely event of ergot becoming abundant, special precautions should be considered to avoid toxicity to grazing animals (such as mowing or temporary livestock exclusion).

Release production: Seed of Willamette Valley Germplasm Roemer’s fescue is readily produced using cultural practices similar to those for other fine fescues in the Pacific Northwest, USA. Post-harvest residue (straw, older foliage) should be removed and remaining stubble should be 4–7 cm (1.6–2.8 in) tall. From 2010 to 2012, seed yields ranged from 64 to 246 kg per ha (56–216 lb per acre). Low yields were due in part to atypical wide spacing among plants within and between rows.

In Oregon, seed certification requires a minimum isolation distance of 274 m (900 ft) between fields of different Roemer’s fescue populations, regardless of pre-variety class or generation. In addition, Roemer’s fescue can hybridize with fescues in the sheep fescue complex (sheep, hard, and Idaho fescues), so similar isolation distances from these species may be applicable. Roemer’s fescue does not cross with species in the red fescue complex.

Red fescue often contaminates Roemer’s fescue seed fields and can be difficult to distinguish for removal using anatomical differences alone. An ammonia fluorescence test conducted by a seed lab in conjunction with a seed lot germination test will detect red fescue contamination in a field. The roots of red fescue (and species in the red fescue complex) fluoresce yellow while the roots of Roemer’s fescue (and species in the sheep fescue complex) fluoresce green when sprayed with a 0.5% solution of ammonium hydroxide. For seed certification of pre-variety Roemer’s fescue germplasm in Oregon,

the maximum allowable red fescue seed content determined by this test is 1 percent for all classes and generations.

Availability of plant materials: The USDA NRCS Plant Materials Center, Corvallis, Oregon, will maintain G2 and G3 generation seed. Beginning in 2012, a limited quantity of selected class certified (green tag) seed will be available to qualified commercial growers upon request. G2, G3, and G4 seed are recommended for certification. There is no limitation on the number of years a field can be in production. Seed is expected to be commercially available by late 2013 or 2014.

References:

Darbyshire, S.J. and L.E. Pavlick. 2007. *Festuca*, modified by Barkworth from Barkworth et al. (eds.), Flora of North America vol. 24. <http://herbarium.usu.edu/webmanual> (accessed 03 July 2012).

Darris, D.C., B.L. Wilson, R. Fiegenger, R. Johnson, and M.E. Horning. 2008. Polycross populations of the native grass *Festuca roemerii* as pre-varietal germplasm, their derivation, release, increase, and use. *Native Plants Journal* 9:304–312.

USEPA. 2011. Level III ecoregions of the continental United States. US EPA, National Health and Environmental Effects Research Laboratory, Western Ecology Division, Corvallis, OR.

U.S. National Arboretum. 2012. USDA plant hardiness zone map. USDA Agricultural Research Service. Washington, DC. <http://planthardiness.ars.usda.gov/> (accessed 11 July 2012).

Wilson, B.L. 2007. A new variety of *Festuca roemerii* (Poaceae) from the California floristic province of North America. *Journal of the Botanical Research Institute of Texas* 1:59–67.

Wilson, B.L., D.C. Darris, R. Fiegenger, R. Johnson, M. Horning, and K. Kuykendall. 2008. Seed transfer zones for a native grass *Festuca roemerii*: genecological evidence. *Native Plants Journal* 9:287–301.

Prepared by: Dale Darris and Annie Young-Mathews, USDA NRCS Plant Materials Center, Corvallis, Oregon. (June, 2012).

Table 2. Morphology, phenology, and disease occurrence of Roemer's fescue. Selected populations for each germplasm (seed zone) are highlighted.																
Seed Zone or Group	Population	rust03	col03	ht04	wid04	til04	fol04	phen	anth	an-e	an-l	mat	surv05	fol05	ferl05	surv06
Coastal (Oregon)	Cummins Creek Trail	3%	1.0	110	17	6.1	4.7	2.4	130	127	135	170	98%	5.4	3.5	13%
Coastal (Oregon)	Cape Perpetua	2%	2.1	87	16	5.8	5.5	1.9	136	131	139	175	97%	6.1	4.0	28%
Coastal (California)	Mt. Tamalpais	5%	1.9	82	16	6.6	5.1	2.4	130	124	136	172	81%	5.3	3.2	8%
Coastal (California)	San Bruno Mt.	2%	2.1	68	13	5.2	4.8	1.8	133	130	137	179	84%	5.0	3.1	10%
Columbia Gorge	Memaloose Park	0%	1.3	65	15	5.8	3.4	2.9	136	126	145	172	67%	3.2	4.0	0%
Columbia Gorge	Drano Lake	19%	1.1	97	14	4.1	3.7	2.2	141	127	148	180	75%	3.3	0.6	6%
High elevation	Fairview Peak	31%	1.2	90	17	5.4	5.2	2.0	132	127	138	169	88%	4.4	3.2	8%
High elevation	Blue Mt. near Deer Park	6%	1.1	74	14	3.2	4.3	2.4	121	114	131	161	83%	3.8	0.3	13%
High elevation	Mary's Peak	11%	1.0	76	16	1.8	5.7	1.4	144	136	155	176	77%	5.0	0.6	13%
High elevation	Hurricane Ridge	5%	1.9	66	9	2.2	2.3	1.8	125	121	132	163	29%	2.0	0.5	2%
High elevation	Silver Fork Elliot Cr.	13%	1.0	81	14	5.1	4.5	3.2	125	122	131	160	81%	3.6	3.7	5%
High elevation	Horse Rock Ridge	20%	1.0	89	18	5.3	4.7	2.2	136	127	140	172	83%	4.3	1.1	0%
High elevation	Table Rock Wilderness	5%	1.1	81	15	3.1	4.4	1.8	134	130	142	169	75%	3.4	0.6	2%
Puget Lowland	Glacial Heritage Preserve	11%	1.1	103	15	3.6	5.0	2.1	134	128	138	167	95%	5.2	2.1	13%
Puget Lowland	Mortar Pt. 3	13%	1.1	106	14	4.0	4.5	2.2	135	133	140	167	89%	5.1	2.6	14%
Puget Lowland	Thirteenth Division Prairie	13%	1.0	102	14	3.6	4.5	2.1	134	131	142	168	92%	4.8	2.0	13%
Puget Lowland	Scatter Creek	9%	1.1	102	13	3.7	4.5	1.9	134	129	138	167	89%	4.8	2.3	17%
Puget Lowland	Mima Mounds	8%	1.0	101	11	3.2	3.6	1.9	134	131	142	167	80%	5.1	1.8	5%
Puget Lowland	Johnson Prairie	16%	1.0	99	13	3.5	4.1	2.0	136	129	144	167	86%	4.2	1.8	9%
Puget Lowland	Yelm	10%	1.0	94	11	3.4	4.0	2.0	136	133	141	169	59%	4.2	2.2	5%
Puget Lowland	Upper Weir Prairie	11%	1.0	102	11	3.2	3.5	2.1	135	129	141	168	89%	3.0	1.3	13%
Puget Lowland	Rocky Prairie N.A.P.	5%	1.0	103	11	3.0	3.9	1.8	141	134	145	169	81%	3.9	1.3	11%
San Juan/Olympic	Fidalgo Island; Ravine Trail	2%	2.6	88	14	4.3	3.9	2.0	136	131	141	172	66%	3.9	1.3	6%
San Juan/Olympic	Yellow Island	4%	1.4	88	14	4.1	6.0	2.0	136	130	143	172	96%	6.2	1.2	24%
San Juan/Olympic	Smith Prairie, Whidbey Is.	0%	2.7	86	12	3.0	4.9	1.6	136	130	145	170	77%	5.2	1.3	6%
Klamath Mountains	Pyramid Rock	11%	1.5	73	14	4.4	3.9	2.0	139	129	148	177	79%	4.0	2.9	2%
Klamath Mountains	Lodgepole Meadow	8%	2.8	79	12	4.0	3.3	1.6	142	127	149	182	81%	3.4	0.9	5%
Klamath Mountains	Hukill Hollow	2%	2.2	78	10	4.1	2.9	2.8	136	126	142	166	56%	2.4	1.8	3%
Klamath Mountains	Edgewood-Weed Road	13%	1.0	82	12	2.7	4.7	2.0	136	130	140	169	69%	3.9	2.3	0%
Klamath Mountains	Cobleigh Road	21%	1.1	74	10	2.7	2.9	1.7	144	137	152	178	34%	2.9	1.8	5%
Klamath Mountains	Ace Williams Mt.	19%	1.2	79	10	2.8	2.8	1.6	140	136	147	178	52%	2.5	0.4	2%
Klamath Co.	Duncan Springs	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*

Table 2 continued on next page.

Table 2. (cont.) Morphology, phenology, and disease occurrence of Roemer's fescue. Selected populations for each germplasm (zone) are highlighted.

Seed Zone or Group	Population	rust03	col03	ht04	wid04	til04	fol04	phen	anth	an-e	an-l	mat	surv05	fol05	fer05	surv06
Willamette Valley	Kloster Mountain	36%	1.0	105	17	5.3	4.6	2.0	139	136	143	177	83%	4.5	2.7	5%
Willamette Valley	Rock Hill	19%	1.0	94	17	4.4	5.5	2.1	134	130	138	169	83%	5.3	2.3	11%
Willamette Valley	Weiss Road ridgeline	51%	2.8	91	14	4.3	4.3	1.8	144	134	153	175	63%	4.0	1.2	0%
Willamette Valley	Morton Farm	42%	1.3	82	13	4.6	4.5	2.1	139	134	142	172	85%	4.8	1.3	0%
Willamette Valley	Bald Hill west ¹	36%	2.5	81	13	4.1	4.6	1.6	146	139	159	176	70%	5.2	1.4	7%
Willamette Valley	Kingston Prairie	13%	2.2	93	14	3.6	4.9	1.9	142	131	162	171	75%	5.6	1.2	2%
Willamette Valley	Spencer's Butte	30%	2.3	89	16	4.5	4.8	2.0	138	134	143	172	80%	5.0	1.9	2%
Willamette Valley	Row Point, Dorena Reservoir	17%	1.9	90	12	4.1	3.7	2.1	140	135	148	172	52%	2.8	0.4	0%
Willamette Valley	Bald Hill southeast ¹	38%	2.8	80	12	4.0	4.1	1.8	143	138	150	177	67%	4.6	1.4	3%
Willamette Valley	Novick Property	39%	2.5	83	13	3.4	3.9	1.5	142	134	152	175	56%	3.7	0.6	2%
Willamette Valley	Baskett Butte	23%	2.8	84	14	4.0	5.4	1.8	141	134	150	173	73%	5.7	1.2	2%
Willamette Valley	Abiqua Road	14%	1.1	108	14	3.2	3.6	1.8	144	134	157	175	65%	3.4	0.1	2%
Willamette Valley	Fire Knob	17%	1.5	84	13	2.7	4.3	1.5	141	138	145	169	64%	3.8	0.5	0%
Willamette Valley	Open Spaces Park	51%	2.2	69	10	2.6	3.2	1.4	145	133	153	178	58%	3.8	1.5	0%
Willamette Valley	Beazell	31%	1.4	77	10	2.5	3.6	1.4	148	136	155	178	45%	4.4	0.7	0%

¹Bald Hill west and Bald Hill southeast were treated as one population for the Willamette Valley germplasm

rust03 = % of leaf rust (*Puccinia* spp.) infection on plants, 2003
col03 = color of foliage score from 1=blue to 3=green, 2003
ht04 = culm height cm, 2004
wid04 = basal width cm, 2004
til04 = fertile tiller score, 1=fewest, 10=most, 2004
fol04 = foliage abundance score, 1=lowest, 10=highest, 2004
phen = phenology score on March 23, 2004 (1=vegetative, 2=boot, 3=jointing, etc.)
anth = Mean Julian date of anthesis for entire population
an-e = earliest Julian date of anthesis for any family in a population
an-l = latest Julian date of anthesis for any family in a population
mat = Mean Julian date of seed maturity
surv05 = % survival of plants in a population, 2005
fol05 = foliage abundance score, 1=lowest, 10=highest, 2005
fert05 = fertile tiller score, 1=fewest, 10=most, 2005
surv06 = % survival of plants in a population, 2006

Table 3. Seed Yield of Roemer's fescue. Selected populations for each germplasm (zone) are highlighted.

Seed Zone or Group	population	yield/plant (g)	Std.Dev.	Minimum	Maximum	Variance
Coastal (Oregon)	Cummins Creek Trail	34.69	5.55	26.38	43.50	30.76
Coastal (Oregon)	Cape Perpetua	34.25	10.24	21.13	51.14	104.87
Coastal (California)	Mt. Tamalpais	33.16	12.85	19.50	53.50	165.03
Coastal (California)	San Bruno Mt.	26.22	8.16	14.88	41.57	66.56
Columbia Gorge	Memaloose Park	19.56	15.90	2.00	33.00	252.93
Columbia Gorge	Drano Lake	17.86	7.44	4.67	31.25	55.40
High elevation	Fairview Peak	27.74	14.88	10.25	48.88	221.40
High elevation	Blue Mt. near Deer Park	9.69	4.77	4.38	19.57	22.73
High elevation	Mary's Peak	5.17	5.36	0.50	14.80	28.69
High elevation	Hurricane Ridge	4.02	1.88	1.40	6.40	3.54
High elevation	Silver Fork Elliot Cr.	24.81	11.28	12.67	43.43	127.31
High elevation	Horse Rock Ridge	24.26	9.64	13.00	41.71	92.91
High elevation	Table Rock Wilderness	10.65	8.12	1.25	26.67	65.91
Puget Lowland	Glacial Heritage Preserve	14.67	5.91	3.17	21.75	34.89
Puget Lowland	Mortar Pt. 3	12.29	4.14	6.50	17.50	17.14
Puget Lowland	Thirteenth Division Prairie	12.05	4.21	7.25	19.29	17.71
Puget Lowland	Scatter Creek	11.11	4.11	4.57	17.14	16.87
Puget Lowland	Mima Mounds	10.93	5.84	2.29	19.17	34.13
Puget Lowland	Johnson Prairie	10.73	4.61	4.80	16.50	21.27
Puget Lowland	Yelm	9.94	5.25	2.33	19.29	27.60
Puget Lowland	Upper Weir Prairie	9.14	4.42	2.20	15.67	19.52
Puget Lowland	Rocky Prairie N.A.P.	7.82	3.85	1.00	13.25	14.85
San Juan/Olympic	Fidalgo Island; Ravine Trail	14.09	4.02	7.00	19.13	16.19
San Juan/Olympic	Yellow Island	10.29	4.25	3.63	15.50	18.07
San Juan/Olympic	Smith Prairie, Whidbey Is.	8.64	2.69	4.60	12.00	7.23
Klamath Mountains	Pyramid Rock	15.80	8.74	1.25	25.50	76.31
Klamath Mountains	Lodgepole Meadow	12.30	4.75	4.50	20.60	22.54
Klamath Mountains	Hukill Hollow	10.51	5.61	4.00	19.88	31.47
Klamath Mountains	Edgewood-Weed Road	7.65	4.17	3.40	14.13	17.43
Klamath Mountains	Cobleigh Road	5.74	3.50	1.40	10.60	12.24
Klamath Mountains	Ace Williams Mt.	5.42	2.39	2.29	10.43	5.73
Klamath Co.	Duncan Spring	4.27	5.32	0.67	16.00	28.33
Willamette Valley	Kloster Mountain	24.24	9.50	11.43	42.00	90.26
Willamette Valley	Rock Hill	15.56	6.32	7.00	22.60	39.99
Willamette Valley	Weiss Road ridgeline	13.70	7.52	0.33	24.40	56.59
Willamette Valley	Morton Farm	13.05	10.65	4.67	28.50	113.47
Willamette Valley	Bald Hill west	12.53	5.32	4.00	20.38	28.33
Willamette Valley	Kingston Prairie	11.45	5.13	4.25	20.00	26.35
Willamette Valley	Spencer's Butte	10.86	6.86	2.86	21.38	47.05
Willamette Valley	Row Point, Dorena Reservoir	9.19	4.43	5.17	19.00	19.59
Willamette Valley	Bald Hill southeast	7.58	1.41	4.86	9.50	2.00
Willamette Valley	Novick Property	7.49	5.82	1.33	17.29	33.87
Willamette Valley	Baskett Butte	6.97	3.58	3.00	12.83	12.85
Willamette Valley	Abiqua Road	6.77	4.46	1.00	12.71	19.90
Willamette Valley	Fire Knob	6.15	6.21	1.20	16.17	38.51
Willamette Valley	Open Spaces Park	5.59	3.86	0.67	9.67	14.92
Willamette Valley	Beazell	5.07	4.60	0.50	14.50	21.16

**SIGNATURES for RELEASE of WILLAMETTE VALLEY GERMPLASM
ROEMER'S FESCUE (*Festuca roemerii*)**

Ron Alvarado
State Conservationist, Oregon
Natural Resources Conservation Service

Date: _____

Roylene Rides at the Door
State Conservationist, Washington
Natural Resources Conservation Service

Date: _____

Tom N. Kaye
Executive Director
Institute for Applied Ecology

Date: _____

Terrell Erickson
Director, Ecological Sciences Division
Natural Resources Conservation Service

Date: _____