

ZENECA Ag Products

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



This is a specimen label and may be inaccurate or out of date. It is intended as a guide in providing general information regarding use of this product. Always read and follow the EPA approved label on the product container.

ZENECA Ag Products

AMBUSH[®] INSECTICIDE

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

ACTIVE INGREDIENT

Permethrin

(3-Phenoxyphenyl)methyl(±)-*cis, trans*-3-(2,2-dichloroethenyl)-

2,2-dimethylcyclopropanecarboxylate*25.6%

INERT INGREDIENTS74.4%

TOTAL100.0%

**Cis/trans* ratio: Min. 35% (±) *cis* and max. 65% (±) *trans*. Contains 2 pounds active ingredient per gallon and is an emulsifiable concentrate.

EPA Reg. No. 10182-18

KEEP OUT OF REACH OF CHILDREN WARNING—AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages via aspiration of the solvent.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water.

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL
1-800-F-A-S-T-M-E-D (327-8633).**

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call
CHEMTREC, 1-800-424-9300.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber ≥ 14 mils or nitrile rubber ≥ 14 mils or Viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

For mushroom house use:

- For exposures in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber ≥ 14 mils or nitrile rubber ≥ 14 mils or Viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

GENERAL INSTRUCTIONS

Apply AMBUSH® insecticide as shown in "DIRECTIONS FOR APPLICATION." Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation. Do not apply more than 2.0 pounds active ingredient per acre per season. Crops in this label may be planted immediately after last treatment. Do not plant other crops within 60 days after last application. Any other crop may be planted if it is not harvested or grazed.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or State agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

CHEMIGATION

SPRINKLER IRRIGATION APPLICATION

Apply AMBUSH insecticide at rates and timing described elsewhere in this label. Apply with center pivot or continuous-move equipment distributing ½ acre-inch or less during treatment. If stationary (solid sets, handlines, or wheel lines other than continuous-move) are used, AMBUSH insecticide should be injected into no more than the last 20 to 30 minutes of the set. Do not apply when winds are greater than 10 to 15 mph to avoid drifts or wind skips. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

USE PRECAUTIONS—SPRINKLER IRRIGATION APPLICATION

- Apply this product only through sprinkler (including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Do not apply through chemigation systems connected to public water systems.

COMMERCIAL IMPREGNATION AND APPLICATION OF AMBUSH INSECTICIDE ON DRY BULK FERTILIZER

AMBUSH insecticide may be impregnated on dry bulk fertilizers. When applied as directed, AMBUSH insecticide/dry bulk fertilizer mixtures provide insect control equal to the same rate of AMBUSH insecticide applied in water.

Bulk fertilizer impregnated with AMBUSH insecticide should be applied immediately, not stored. The AMBUSH insecticide/fertilizer mixtures may be surface applied or shallow incorporated. If shallow incorporated, the higher rate of AMBUSH insecticide should be used.

Apply using a minimum of 200 pounds and a maximum of 450 pounds of dry bulk fertilizer per acre with the recommended amount of AMBUSH insecticide per acre.

For impregnation of AMBUSH insecticide on dry fertilizers, use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle (or nozzles) should be positioned inside of the mixer to provide uniform spray coverage of the tumbling fertilizer. AMBUSH insecticide should be sprayed uniformly onto the fertilizer using a fine spray pattern.

The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with AMBUSH insecticide provides a satisfactory dry mixture.

If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry, free-flowing mixture. Micro-Cel™ E (Manville Sales Corp.) is the recommended absorbent powder. It should be added separately and uniformly to the prepared AMBUSH insecticide fertilizer mixture in a quantity that is sufficient to provide a suitably free-flowing mixture. Generally less than 2% by weight of Micro-Cel I E is required.

DO NOT impregnate AMBUSH insecticide onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the insecticide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with AMBUSH insecticide.

The amount of AMBUSH insecticide actually required in the manufacture of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of AMBUSH insecticide actually contained in the mixture applied to the soil represents the correct rate of use.

All individual state regulations relating to bulk dry fertilizer blending, registration, labeling, and application of the mixtures are the responsibility of the individual and/or company selling the fertilizer and AMBUSH insecticide mixture.

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|---|--|------------------------------------|---|
| COLE CROPS—Brassica Leafy Vegetables | | | |
| BROCCOLI | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Armyworm spp. Plant Bug Thrips | 0.05 to 0.1 (3.2 to 6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply within 1 day of harvest. |
| BRUSSELS SPROUTS | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Armyworm spp. Plant Bug Thrips | 0.05 to 0.1 (3.2 to 6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply within 1 day of harvest. |
| CABBAGE | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Southern White Butterfly | 0.05 to 0.2 (3.2 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 to 10 days or as needed. Do not apply more than 1.0 pound active ingredient per acre per season. Do not apply within 1 day of harvest. |
| | Armyworm spp. Cutworm Flea Beetle | 0.1 to 0.2 (6.4 to 12.8) | |
| CAULIFLOWER | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Armyworm spp. Plant Bug Thrips | 0.05 to 0.1 (3.2 to 6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply within 1 day of harvest. |
| CHINESE BROCCOLI^b (gai lon, white flowering) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Armyworm spp. Plant Bug Thrips | 0.05 to 0.1 (3.2 to 6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply within 1 day of harvest. |
| CHINESE CABBAGE (Tight-heading varieties only) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Southern White Butterfly | 0.05 to 0.2 (3.2 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 to 10 days or as needed. Do not apply more than 1.0 pound active ingredient per acre per season. Do not apply within 1 day of harvest. |
| | Armyworm spp. Cutworm Flea Beetle | 0.1 to 0.2 (6.4 to 12.8) | |
| COLLARDS (AR, AZ, GA, IL, NC, OK, SC, TX only) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Beet Armyworm Corn Earworm Southern White Butterfly Leafminer Leafhopper European Corn Borer Fall Armyworm Cutworm Green Cloverworm Southern Armyworm Tobacco Budworm Vegetable Leafminer | 0.05 to 0.1 (3.2 to 6.4) | Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Spray using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 8 applications. Do not apply within 1 day of harvest. |

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|--|--|------------------------------------|--|
| CUCURBIT VEGETABLES | | | |
| CUCURBIT VEGETABLES† | Leafminer and Squash Bug Aphid | 0.2 (12.8) | Apply by air or ground equipment. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active ingredient per acre per season. AMBUSH insecticide may be applied up to harvest. |
| | Cabbage Looper Cucumber Beetle (adult) Cutworm Melon Worm Pickleworm Plant Bug (including Lygus and Stink Bug) Leafhopper Rindworm Squash Vine Borer | 0.1 to 0.2 (6.4 to 12.8) | |
| FRUITING VEGETABLES | | | |
| EGGPLANT | Colorado Potato Beetle | 0.2 (12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 2.0 pounds active ingredient per acre per season. Do not apply within 3 days of harvest. |
| | Vegetable Leafminer Cabbage Looper Flea Beetle | 0.1 to 0.2 (6.4 to 12.8) | |
| PEPPERS (Bell) | Vegetable Leafminer Cabbage Looper Flea Beetle Pepper Weevil Corn Earworm Cutworm | 0.1 to 0.2 (6.4 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.6 pounds active ingredient per acre per season. Do not apply within 3 days of harvest. |
| | European Corn Borer | 0.2 (12.8) | |
| TOMATOES | Tomato Fruitworm Cabbage Looper Tomato Pinworm Vegetable Leafminer Beet Armyworm Southern Armyworm Colorado Potato Beetle Granulate Cutworm Hornworm | 0.05 to 0.2 (3.2 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Do not apply more than 1.2 pounds active ingredient per acre per crop season. Can be applied up to the day of harvest. |
| LEAFY VEGETABLES | | | |
| LEAFY VEGETABLES* (Except Brassica) | Vegetable Leafminer Cabbage Looper Cutworm Complex Fall Armyworm Beet Armyworm Corn Earworm Green Cloverworm Leafhopper European Corn Borer Southern Armyworm Tobacco Budworm Aphid Alfalfa Looper | 0.1 to 0.2 (6.4 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 3 to 5 days or as needed. Do not apply more than 2.0 pounds active ingredient per acre per season. Can be applied up to 1 day of harvest. Spinach only—Do not apply more than 1.0 pound active ingredient per acre per season. |

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks | |
|--|--|---|---|------|
| ROOT, TUBER, AND BULB VEGETABLES | | | | |
| HORSERADISH | Imported Crucifer Weevil <i>(Baris lepidii)</i> control | 0.1% ai solution ^c in a spring pre- plant dip and 0.2 lb. ai per acre (12.8) as a foliar spray. | Soak sets for 30 minutes and air-dry before planting; make up to 3 foliar applications as needed to control weevil adults during oviposition. Apply by ground equipment in a minimum spray volume of 20 gallons per acre. Do not apply within 30 days of harvest. | |
| ONIONS (Dry Bulb and Garlic) | Onion Thrips Armyworm | 0.15 to 0.3 (9.6 to 19.2) | Apply by air or ground equipment. Apply in a minimum of 20 gallons of water per acre with ground equipment or in a minimum of 5 gallons per acre by air as foliar spray. Begin applications when pests appear and repeat as necessary to maintain control. Use the higher label rates as onion thrips population increases and avoid rescue situations. Do not apply more than 2.0 pounds active ingredient per acre per season. Do not apply within 1 day of harvest. | |
| | Cutworm Leafminer Onion Maggot (adult) Stink Bug | 0.1 to 0.3 (6.4 to 19.2) | | |
| POTATOES | Colorado Potato Beetle Potato Leafhopper Aster Leafhopper Potato Flea Beetle Potato Aphid Potato Tuberworm Potato Psyllid Cabbage Looper Beet Armyworm Cutworm European Corn Borer Tarnished Plant Bug | 0.05 to 0.2 (3.2 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 1.6 pounds active ingredient per acre per season. Can be applied up to 14 days prior to harvest. | |
| TURNIPS (FL, GA, IL, IN, OK, SC, TX, WA only) | Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a Beet Armyworm Corn Earworm Southern White Butterfly Leafminer Leafhopper European Corn Borer Fall Armyworm Cutworm Green Cloverworm Southern Armyworm Tobacco Budworm Vegetable Leafminer | 0.05 to 0.1 (3.2 to 6.4) | Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply within 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock. Do not exceed 8 applications. (TX, OK, IL, IN only) Do not exceed 4 applications. (SC, GA, FL, WA only) | |
| CEREAL GRAINS | | | | |
| FIELD CORN/POPCORN/ FIELD CORN GROWN FOR SEED (At plant use) | Armyworm Cutworm | 0.5 ounce per 1,000 linear feet of row | Apply as an in-furrow, band, or T-band treatment using a minimum 4" band. Use table below to determine AMBUSH insecticide needs for each acre. | |
| Row spacing (inches) | | 40 | 30 | 20 |
| AMBUSH insecticide (pounds active ingredient per acre) | | 0.10 | 0.15 | 0.20 |
| FIELD CORN/POPCORN/ FIELD CORN GROWN FOR SEED (Preemergent use)** | Armyworm Cutworm Stalk Borer | 0.1 to 0.2 (6.4 to 12.8) or 0.5 to 1.0 ounce per 1,000 linear feet of row (based on a 4" band and 40" row spacing) | AMBUSH insecticide may be applied as a preplant or pre- emergence application. For best results, apply at planting time. Apply as a broadcast spray by ground (minimum of 10 gallons of water per acre) or air (minimum of 2 gallons of water per acre) or 4" to 15" band using sufficient spray volume to achieve adequate coverage. Linear row calculations should be used proportional to the standard band width/row width formula to adjust rates for different band widths or row spacings. Use higher rates of AMBUSH insecticide when incorporating into the soil with- out exceeding the recommended dosage. | |

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|--|--|------------------------------------|--|
| FIELD CORN/POPCORN/ FIELD CORN GROWN FOR SEED (Foliar use)** | Armyworm (including Fall Armyworm) Cutworm European Corn Borer Southwestern Corn Borer Flea Beetle Corn Earworm Corn Rootworm Beetle Tobacco Budworm ^b Stalk Borer Hop Vine Borer | 0.1 to 0.2 (6.4 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Do not apply more than 0.6 pound active ingredient per acre per season. Allow a minimum of 6 days between treatments. Do not apply within 30 days of harvest of grain or fodder (stover). Forage may be harvested on the day of application. When treating for stalk borer, AMBUSH insecticide must be applied when or shortly before the stalk borer larvae are moving into the corn from surrounding weeds and grasses. Mowing or burndown herbicide is suggested to initiate movement. |
| | Western Bean Cutworm | 0.05 to 0.1 (3.2 to 6.4) | For control of corn earworm, apply just before silking and continue as necessary to maintain control. |
| SWEET CORN | Corn Earworm European Corn Borer Fall Armyworm Corn Rootworm (adult) Cutworm Flea Beetle ^b Hop Vine Borer Leafhopper Southern Armyworm Stalk Borer | 0.1 to 0.2 (6.4 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 1.2 pounds active ingredient per acre per season. Do not apply within 1 day of harvest of ears or forage or livestock grazing. AMBUSH insecticide may be mixed with Lannate [®] (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded. |
| SWEET CORN (Florida only) | Corn Earworm European Corn Borer Fall Armyworm Southern Armyworm Corn Rootworm (adult) Cutworm Aster Leafhopper | 0.1 to 0.25 (6.4 to 16.0) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 2.0 pounds active ingredient per acre per crop season. AMBUSH insecticide may be mixed with Lannate (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded. |
| TREE CROPS | | | |
| ALMONDS | Navel Orangeworm Peach Twig Borer | 0.2 to 0.4 (12.8 to 25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Can be applied up to 7 days before harvest. Do not apply more than 2.0 pounds active ingredient per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. For control of peach twig borer in dormant season almonds: Apply 0.1 to 0.2 pound active ingredient per acre (6.4 to 12.8 fluid ounces per acre). AMBUSH insecticide may be applied with or without dormant oil by ground equipment (dilute or concentrate sprays) or by aircraft. When combined with dormant oil and applied by air, observe the application rate and spray volume specified on the dormant oil label. When not using dormant oil, almonds may be treated by air with a minimum of 15 gallons of finished spray per acre. Apply by air only when soil conditions do not permit regular ground application. For maximum effectiveness, combine AMBUSH insecticide with labeled rates of dormant oil and apply with air blast ground sprayers in volumes of 100 to 400 gallons of finished spray per acre. |
| | Ants | 0.4 (25.6) | Apply by ground equipment in a minimum of 15 gallons of finished spray per acre. Applications should follow mowing of weed growth to ensure maximum coverage of the soil surface. Overhead moisture following application will enhance activity. |

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|---------------------------------|---|------------------------------------|---|
| APPLES | Apple Aphid Redbanded Leafroller Obliquebanded Leafroller Plum Curculio White Apple Leafhopper Spotted Tentiform Leafminer Tarnished Plant Bug Green Fruitworm | 0.1 to 0.4 (6.4 to 25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Do not apply after petal fall. Do not apply more than 0.6 pound active ingredient per acre per season. |
| AVOCADOS | Thrips Leafhopper Mirid Bug Lepidopterous Larvae Avocado Caterpillar Avocado Lace Bug Avocado Leafroller Orange Tortrix Avocado Looper Avocado Tree Girdler Avocado Whitefly Brown Soft Scale Spanworm Twig Borer Scale Crawler | 0.2 (12.8) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7- to 10-day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| CHERRIES | Lesser Peachtree Borer Rose Chafer Green Fruitworm Redbanded Leafroller Plum Curculio Tarnished Plant Bug | 0.1 to 0.2 (6.4 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Apply when insects appear. In the east, do not exceed 6 applications per season, with no more than 4 applications after petal fall. In the west, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not graze livestock in treated orchards. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| FILBERTS | Filbertworm Obliquebanded Leafroller | 0.2 to 0.4 (12.8 to 25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 pounds active ingredient per acre per season. Do not apply within 14 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| PAPAYA (Florida only) | Fruit Fly Papaya Webworm Papaya Whitefly Brown Soft Scale Scale Crawler Aphid Mealybug | 0.1 to 0.2 (6.4 to 12.8) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7- to 10-day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| PEACHES | Peach Twig Borer Oriental Fruit Moth Green Fruitworm Lesser Peachtree Borer Plum Curculio Tarnished Plant Bug Rose Chafer | 0.1 to 0.3 (6.4 to 19.2) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Can be applied up to 14 days before harvest. Do not apply more than 5 applications or 1.5 pounds active ingredient per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |

Continued on next page

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|---|--|------------------------------------|---|
| PEACHES (Cont'd.) | | | For control of peach twig borer in dormant season peaches: Apply 0.1 to 0.2 pound active ingredient per acre (6.4 to 12.8 fluid ounces per acre). AMBUSH insecticide may be applied with or without dormant oil by ground equipment (dilute or concentrate sprays) or by aircraft. When combined with dormant oil and applied by air, observe the application rate and spray volume specified on the dormant oil label. When not using dormant oil, peaches may be treated by air with a minimum of 15 gallons of finished spray per acre. Apply by air only when soil conditions do not permit regular ground application. For maximum effectiveness, combine AMBUSH insecticide with labeled rates of dormant oil and apply with air blast ground sprayers in volumes of 100 to 400 gallons of finished spray per acre. |
| PEARS (Post-harvest, dormant through pre-bloom) | Pear Psylla Codling Moth Green Fruitworm | 0.2 to 0.4 (12.8 to 25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply at a rate of 0.8 to 1.6 pints (0.2 to 0.4 pound active ingredient) per acre. May be combined with 2 to 8 gallons of spray oil per acre. Apply during the post-harvest through dormant through pre-bloom growth periods only. Do not apply more than 0.8 pound active ingredient per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. Pre-bloom: Apply two pre-bloom sprays. Pre-bloom sprays can be applied from dormant through bud burst stages. Post-harvest: For control of codling moth: Apply 0.2 to 0.4 pound active ingredient per acre (12.8 to 25.6 fluid ounces per acre). AMBUSH insecticide may be applied with or without dormant oil by ground equipment (dilute or concentrate sprays) or by aircraft. When combined with dormant oil and applied by air, observe the application rate and spray volume specified on the dormant oil label. When not using dormant oil, pears may be treated by air with a minimum of 15 gallons of finished spray per acre. Apply by air only when soil conditions do not permit regular ground application. For maximum effectiveness, combine AMBUSH insecticide with labeled rates of dormant oil and apply with air blast ground sprayers in volumes of 100 to 400 gallons of finished spray per acre. |
| PISTACHIOS | Peach Twig Borer Navel Orangeworm Chinch Bug Plant Bug Stink Bug Leaffooted Bug | 0.2 to 0.4 (12.8 to 25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply at a rate of 0.8 to 1.6 pints (0.2 to 0.4 pound active ingredient) per acre. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply after 10 percent hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |
| | Ants | 0.4 (25.6) | Apply by ground equipment in a minimum of 15 gallons of finished spray per acre. Applications should follow mowing of weed growth to ensure maximum coverage of the soil surface. Overhead moisture following application will enhance activity. |
| WALNUTS | Navel Orangeworm Codling Moth Walnut Husk Fly | 0.2 to 0.4 (12.8 to 25.6) | Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 pounds active ingredient per acre per season. Do not apply within 1 day of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. |

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|--|---|--|---|
| OTHER CROPS | | | |
| ALFALFA, Alfalfa grown for seed | Alfalfa Caterpillar Armyworm Cutworm Looper Webworm Potato Leafhopper Blue Alfalfa Aphid ^e Green Peach Aphid Pea Aphid ^e Green Cloverworm Spotted Alfalfa Aphid Velvetbean Caterpillar | 0.05 to 0.2 (3.2 to 12.8) | Apply by air or ground equipment. Use higher recommended dosage for increased pest pressure or for increased residual pest control. Apply in a minimum of 2 gallons of finished spray per acre by aircraft or 10 gallons of finished spray per acre with ground equipment. Use higher gallonage, 5 to 10 gallons by air, or 20 by ground, finished spray per acre when foliage is dense and/or when pest populations are high. Do not apply more than 0.2 pound active ingredient per cutting. When rates of 0.1 pound active ingredient per acre or less are used, application may be made on day of harvest. When rates greater than 0.1 pound active ingredient per acre are used, do not apply within 14 days of harvest. Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. |
| | Alfalfa Weevil ^e Egyptian Alfalfa Weevil ^e Cucumber Beetle Meadow Spittlebug Plant Bug (including <i>Lygus</i> spp.) ^e Stink Bug | 0.1 to 0.2 (6.4 to 12.8) | |
| ARTICHOKES | Artichoke Plume Moth Leafminer | 0.1 to 0.3 (6.4 to 19.2) | Apply by air or ground equipment. By air, apply in a minimum of 2 to 10 gallons of finished spray per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.5 pounds active ingredient per acre per season. Buds may be harvested on the day of application. |
| ASPARAGUS | Cutworm Complex Asparagus Beetle | 0.05 to 0.1 (3.2 to 6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of finished spray per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 0.4 pound active ingredient per acre per season. Do not apply within 1 day of harvest. |
| | Asparagus Beetle Tarnished Plant Bug Lygus Bug Japanese Beetle ^b (adult stage) | 0.1 (6.4) | For post-harvest application, apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the asparagus beetle, lygus bug, and the adult stage of the Japanese beetle are present. |
| MUSHROOMS (Mushroom houses and adjacent premise areas) | Mushroom Flies (Sciarid and Phorid adults) | Apply as a fogging or aerosol treatment at the rate of 3.2 to 4.0 ounces (0.05 to 0.0625 lb. ai) per 30 ounces of water or suitable diluent. Use 1 quart of solution per standard double house (35,000 cu. ft.: 8,000 sq. ft.) | Preparation of the building prior to fogging: (1) Close all doors, windows, and ventilators, (2) Lock or barricade all entrances, turn off pilot lights, post warning signs, and take precautions to prevent persons and animals from entering the area. Use prior to filling the house, during cooldown, during spawning, up to pinning, and between breaks. Do not use when mushrooms are present. Treat once daily or as needed when flies appear. Do not make more than 20 applications prior to pinning of first break; apply no more than 2 applications between each break. Do not apply more than 30 applications total per crop of 5 breaks. Do not apply within 3 days of harvest. Length of exposure time should be limited to 1 hour; then ventilate the house. Use fans to ventilate in houses that do not have forced air circulation. Wear a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). |
| RANGE GRASS (NM only) | Range Caterpillar | 0.01 (0.64) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient volume to obtain uniform coverage. Apply as needed. Do not apply more than once per year. Cattle may be present during application. Do not harvest or feed hay to livestock. |

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|--|--|---|--|
| SOYBEANS | Green Cloverworm Mexican Bean Beetle Bean Leaf Beetle Cabbage Looper Potato Leafhopper Velvetbean Caterpillar Saltmarsh Caterpillar Cutworm Flea Beetle Thistle Caterpillar | 0.05 to 0.1 (3.2 to 6.4) | Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Do not apply more than 0.4 pound active ingredient per acre per season. Can be applied up to 60 days before harvest. Do not graze treated areas or harvest for forage or hay. |
| | Corn Earworm Beet Armyworm Japanese Beetle Soybean Looper ^d Webworm | 0.1 to 0.2 (6.4 to 12.8) | |
| ORNAMENTALS | | | |
| CHRISTMAS TREES ^b (Container and field grown) | Nantucket Pine Tip Moth | 0.1 to 0.2 (6.4 to 12.8) | Apply first application when adults appear and repeat at 5- to 7-day intervals or as needed throughout the season. Use sufficient water to obtain full coverage of foliage. |
| CHRYSANTHEMUMS | Vegetable Leafminer Liromyza Leafminer Fly | 0.1 to 0.2 (6.4 to 12.8) | Apply by ground. Avoid spraying the blooms. AMBUSH insecticide may be applied on a weekly schedule. Caution: AMBUSH insecticide has demonstrated excellent plant safety; however, not all cultivars have been tested. Before treating large numbers of plants of a particular cultivar, treat a few plants and observe prior to full-scale applications. |
| CONIFERS (Container and field grown) | Nantucket Pine Tip Moth | 0.1 to 0.2 (6.4 to 12.8) | Apply first application when adults appear and repeat at 5- to 7-day intervals or as needed throughout the season. AMBUSH insecticide may be diluted in a nonvolatile vegetable oil or in water in a minimum of 1 gallon of finished spray per acre using equipment calibrated to give adequate coverage. |
| ORNAMENTALS Nursery Stock ^b (Field grown) | Beet Armyworm Cabbage Looper Citrus Thrips <i>Heliothis</i> spp. Leafhopper Leafminer Whitefly Cutworm | 0.1 to 0.2 (lb. ai/100 gal.) | Marginal leaf burn may occur on <i>Salvia</i> , <i>Dieffenbachia</i> , and <i>Pteris</i> Fern. Application to blooming plants may cause browning of petals. Not all species and varieties have been tested. Before treating large numbers of plants of a particular variety, treat a few plants and observe prior to full-scale application. Apply to nonbearing plants of fruiting species. |
| PINE SEED ORCHARD | Coneworm Seed Bug | 13 fl. oz. per 100 gals. water (0.025% dilution by wt.) for high-volume sprayers; 68 fl. oz. per 100 gallons water (0.125% dilution by wt.) for low-volume sprayers; 48 fl. oz. per acre application in not less than 10 gallons of water for aerial applications | Apply first application within 1 week of female flower closure or peak pollen flight for webbing coneworm control. For other coneworms and seed bugs, apply first application within 30 days following female flower closure. Repeat applications at intervals of 4 weeks, but do not apply more than 6 applications. Apply approximately 5 to 10 gallons of the 0.025% dilution per tree with high-volume sprayers. With low-volume sprayers, apply 100 gallons of the 0.125% dilution per acre. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. Avoid contact with open water. |
| HYBRID POPLAR TREES (Grown for pulp) (WA, OR, ID only) | Aphid Beetle Cutworm | 0.1 to 0.2 (6.4 to 12.8) | Apply by air in a minimum of 5 gallons of water. Apply when field counts indicate damaging insect populations are developing or present. Repeat as necessary to maintain control. |
| | Spotted Tentiform Leafminer Tarnished Plant Bug | 0.2 (12.8) | |

DIRECTIONS FOR APPLICATION

| Crop | Target Pests | Rate Lbs. ai/Acre (Oz./Acre) | Remarks |
|---|--|------------------------------------|--|
| ROSES^b (Field grown) | <i>Heliothis</i> spp. | 0.1 to 0.2 (6.4 to 12.8) | Apply by air or ground equipment. Apply in a minimum of 2 gallons per acre by air and 10 gallons of finished spray per acre with ground equipment. |
| ROSES^b (Greenhouse) | Beet Armyworm Cabbage Looper Omnivorous Leafroller | 8 fl. oz. per 100 gals. water | AMBUSH insecticide is not phytotoxic to the following varieties of greenhouse roses: Ballena, Bettina, Cara Mia, Coquette, Excitement, Forever Yours, G. Wave, Jack Frost, Jr. Bridesmaid, Matador, Paul's Pink, Samantha, Seventeen, Town Crier, Tropicana, and Visa. Other varieties may vary in their sensitivity to AMBUSH insecticide, and a small number of plants should be treated under local conditions to determine plant safety prior to commercial use. |

* LEAFY VEGETABLES (Except Brassica): Amaranth (leafy amaranth, Chinese spinach, tampala); arugula (Rocket); Cardoon; celery; Chinese celery; celtuce; chervil; corn salad; chrysanthemum, edible-leaved; chrysanthemum, garland; cress, garden; cress, upland (yellow rocket, winter cress); dandelion; dock (sorrel); endive (escarole); fennel, Florence; lettuce (head and leaf); orach; parsley; purslane, garden; purslane, winter; radicchio (red chicory); rhubarb; spinach; spinach, vine (Malabar spinach, Indian spinach); spinach, New Zealand; Swiss chard.

† CUCURBIT VEGETABLES: Chayote (fruit); Chinese Waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourds, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelons, including hybrids such as true cantaloupe, cantaloupe, casaba, crenshaw, golden pershaw melon, honeydew melons, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon; pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), and winter (butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon, including hybrids.

** AMBUSH insecticide may be applied in tank mixtures with the following herbicides in accordance with label limitations and precautions. Label dosage rates should not be exceeded. GRAMOXONE® SUPER; GRAMOXONE EXTRA, Bicep®, Lasso®/Atrazine Premix; Lasso 4E & ME; Dual® 8E; Atrazine 4L & 80W; Bladex® 80W & 4L; Princep® Caliber 90®; 2,4-D; Roundup®; Bronco®; Banvel®; Extrazine®.

^a Aids in suppression.

^b Currently not registered for use in California.

^c 0.1% active ingredient solution = 3½ pints formulation per 100 gallons of water.

^d Maximum use rate in California is 0.1 pound active ingredient per acre.

^e Under heavy insect pressure, use 0.2 pound active ingredient per acre. A second application of alternate chemistry may be necessary to reduce populations below economic thresholds.

| Rate Conversion Chart | | | |
|---------------------------------|----------------------|------------|--------------------------|
| Pound Active Ingredient/Acre | Fluid Ounces/Acre | Pints/Acre | Acres Treated/ Gallon |
| 0.05 | 3.2 | 0.2 | 40.0 |
| 0.10 | 6.4 | 0.4 | 20.0 |
| 0.125 | 8.0 | 0.5 | 16.0 |
| 0.15 | 9.6 | 0.6 | 13.3 |
| 0.175 | 11.2 | 0.7 | 11.4 |
| 0.20 | 12.8 | 0.8 | 10.0 |

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Plastic Containers: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

AMBUSH® and GRAMOXONE® are trademarks of a ZENECA Group Company. Banvel® is a trademark of BASF Corporation.

Bronco®, Lasso®, and Roundup® are trademarks of Monsanto Company.

Bicep®, Dual®, and Princep® Caliber 90® are trademarks of Novartis.

Bladex®, Extrazine®, and Lannate® are trademarks of E. I. du Pont de Nemours and Company.

Micro-Cel™ is a trademark of Manville Sales Corporation.

AMBUSH and GRAMOXONE EXTRA are restricted use pesticides.

© 1998. ZENECA Inc.

This specimen label is current as of the time of printing.

For current information, contact ZENECA Ag Products at 1-800-759-2500.

069608 (Rev. 11/97)

Made in U.S.A.

ZENECA Ag Products

1800 Concord Pike
Wilmington, DE 19850-5458