



CONSERVATION ENHANCEMENT ACTIVITY

E328D

CONSERVATION STEWARDSHIP PROGRAM

Leave standing grain crops unharvested to benefit wildlife

Conservation Practice 328: Conservation Crop Rotation

APPLICABLE LAND USE: Crop (Annual & Mixed)

RESOURCE CONCERN: ANIMALS

ENHANCEMENT LIFE SPAN: 1 Year

Enhancement Description

Implement a crop rotation which allows a portion of grain crops to be left in fields unharvested to provide food and cover for wildlife during winter months.

Criteria

- Crops must be grown in a planned sequence as outlined in the plan. The crop rotation shall include a minimum of three different crops. For this purpose, a cover crop is considered a different crop.
- Where applicable, plan suitable crop substitutions when the planned crop cannot be planted due to weather, soil conditions, or other local situations.
- Select the crops and crop management activities that provide food, cover, and shelter for the targeted wildlife species using an approved habitat evaluation procedure.
- Leave a minimum ½ acre of unharvested, standing grain crops for each 40 acres of cropland. Unharvested plots shall be located in a single location on the 40 acre unit and additional plots shall be located on different 40 acres. *This enhancement is to be planned, contracted, and implemented on an entire field, not just the unharvested acres.*

E328D-Leave standing grain crops unharvested to benefit wildlife	July 2019	Page 1
--	-----------	----------



CONSERVATION STEWARDSHIP PROGRAM

- Locate the unharvested plots adjacent to permanent cover such as brushy fence rows, field borders, forest land, or wetlands (this does not include newly established vegetation).
- Leave unharvested crops standing over winter until it is time to prepare the soil for planting the next crop.





CONSERVATION STEWARDSHIP PROGRAM

Documentation and Implementation Requirements

Participant will:

- Prior to implementation, provide NRCS with the planned crop rotation.

Field	Acres	Planned Crops (in sequence)	Length of Crop Rotation (years)

- Prior to implementation, develop a map showing planned location(s), crop type(s), and acreage of crops to be left unharvested.
- During implementation, notify NRCS of any planned changes in crops, crop rotation, or unharvested areas to verify the planned system meets the enhancement criteria.
- During implementation, take photos of all unharvested plots. Photos must indicate field location and date.
- After implementation, if changes to the rotation were made, complete the tables above to document the applied Conservation Crop Rotation for the contract period and provide to NRCS.
- After implementation, make a map showing implemented location(s), crop type(s), and acreage of crops that were left unharvested each year available for review by NRCS to verify implementation of the enhancement.
- After implementation, make photos of the unharvested plots available for review by NRCS to verify implementation of the enhancement.

NRCS will:

- As needed, provide technical assistance in selecting crop rotations or substitute crops that would meet the criteria of the enhancement.
- As needed, provide technical assistance in selecting crops for food, cover, and shelter according to the approved habitat evaluation procedure.
- As needed, provide additional assistance to the participant as requested.



CONSERVATION STEWARDSHIP PROGRAM

- During implementation, evaluate planned crop changes, crop rotation, or unharvested areas to verify the planned system meets the enhancement criteria.
- After implementation, review the map(s) showing implemented location(s), crop type(s), and acreage of crops that were left unharvested each year, to verify implementation of the enhancement.
- After implementation, review photos of unharvested plots to verify implementation of the enhancement.

NRCS Documentation Review:

I have reviewed all required participant documentation and have determined the participant has implemented the enhancement and met all criteria and requirements.

Participant Name _____ Contract Number _____

Total Amount Applied _____ Fiscal Year Completed _____

NRCS Technical Adequacy Signature

Date



SOUTH DAKOTA (SD) SUPPLEMENT TO CONSERVATION ENHANCEMENT ACTIVITY

CONSERVATION STEWARDSHIP PROGRAM

E328D

Additional Criteria for SD:

In addition to the criteria specified in the national job sheet E328D, the following additional criteria apply in SD:

- Any annually planted commodity crop left standing over winter, is adequate for wildlife.
- In SD the approved habitat evaluation guide is the SD-CPA-19 (WHEG).
- If using a cover crop, see table 1 below for acceptable species:



Table 1: Cover Crop - Common Species and Properties

Cover Crop	Full seeding rate lbs/acre/4	Seeding depth, inches	Reduce erosion	Increase soil organic matter	Savrange nutrients	Biological N fixation	Suppress weeds	Provide supplemental hay	Provide supplemental grazing	Rooting depth / Plant water use ^{1/}	Minimize / Reduce surface soil compaction	Minimize/ Reduce subsoil compaction	Seed size (Large or Fine)	Crop type and seeding dates /2 /3	Winter survival	Salinity Tolerance	CN Ratio	Mycorrhizal fungi association	Seeds/lb	Shade Tolerance
Alfalfa	6.5	.25 - .75	G	G	G	Y	G	G	F	DH	G	G	F	CB	Y	P	L	M	210,000	F
Barley	50	.75 - 2.0	G	G	G	N	G	G	G	MM	G	F	L	CG	N	G	M	M	14,000	F
Brassica hybrids	7	.25 - .5	F	F	G	N	G	F	G	MM	G	G	F	CB	N	G	L	N	180,000	P
Buckwheat /5	50	.5 - 1.5	P	P	F	N	F	P	P	SL	F	P	L	WB	N	P	L	N	19,000	G
Cabbage, African	5	.25 - .75	F	F	G	N	F	F	F	MM	G	G	F	CB	N	G	L	N	180,000	F
Camelina, Winter	3	.25 - .5	F	F	F	N	P	P	P	ML	P	F	F	CB	S	P	L	N	400,000	P
Canola	5	.25 - .75	F	F	G	N	G	F	F	MM	G	G	F	CB	S	G	L	N	140,000	F
Clover, Balansa	5	.25 - .75	F	P	F	Y	P	P	F	SL	P	P	F	CB	N	P	L	M	500,000	F
Clover, Crimson	15	.25 - .75	F	F	F	Y	P	F	F	SM	P	P	F	CB	S	P	L	M	150,000	F
Clover, Red	5	.25 - .75	G	F	F	Y	F	F	F	SL	F	F	F	CB	Y	P	L	M	275,000	G
Clover, Sweet	4	.25 - 1.0	G	G	F	Y	G	F	F	MM	G	G	F	CB	Y	F	L	M	260,000	G
Collards or Kale	5	.25 - .5	F	F	G	N	G	F	G	MM	G	G	F	CB	N	G	L	N	175,000	F
Corn	12	1 - 1.5	G	G	G	N	G	F	G	DH	G	G	L	WG	N	P	H	H	2,500	F
Cowpeas or Dry Beans	30	1 - 1.5	P	F	F	Y	P	P	F	SL	F	F	L	WB	N	P	L	M	4,000	F
Fava beans	75	1 - 1.5	F	F	F	Y	F	G	G	DM	F	F	L	CB	N	F	L	P	2,500	P
Flax	30	.25 - .75	F	F	F	N	P	P	P	SM	F	P	F	CB	N	P	H	H	80,000	P
Lentils	30	1 - 1.5	P	P	P	Y	P	P	P	SL	P	P	F	CB	N	P	L	M	20,000	P
Millet, hay	15	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	N	P	M	H	180,000	P
Millet, proso	25	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	N	P	M	H	80,000	P
Mustard	6	.25 - .75	F	F	F	N	G	F	P	MH	G	F	F	CB	N	P	L	N	140,000	P
Oats	70	.5 - 1.5	G	G	G	N	G	G	G	MM	G	F	L	CG	N	F	M	H	16,000	F
Peas	70	1.5 - 3.0	F	P	P	Y	F	G	G	SL	F	F	L	CB	N	P	L	M	3,500	F
Phacelia	4	.25 - .5	F	F	F	N	P	P	P	DH	F	P	F	CB	N	P	L	M	225,000	F
Radishes	8	.25 - .75	F	F	G	N	G	P	G	DH	G	G	F	CB	N	P	L	N	25,000	P
Rapeseed	5	.25 - .75	F	F	G	N	G	F	G	MM	G	G	F	CB	Y	G	L	N	140,000	F
Rye, Cereal	60	.75 - 2.0	G	G	G	N	G	G	G	MH	G	G	L	CG	Y	G	H	M	18,000	G
Ryegrass, Annual	15	.5 - 1.5	G	G	G	N	F	G	G	MM	G	F	F	CG	S	F	M	M	190,000	G
Safflowers	30	.5 - 1.0	F	F	G	N	F	P	P	DM	F	G	L	WB	N	F	M	M	15,000	P
Sorghum, Forage and Sudan Hybrids	15	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	17,000	P
Sorghum, Grain	5	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	17,000	P
Soybeans	35	1 - 1.5	F	P	F	Y	F	F	F	SM	F	F	L	WB	N	P	L	M	3,000	F
Sudangrass	20	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	25,000	P
Sugar beets	4	.25 - .5	F	P	G	N	F	P	G	DH	G	G	F	CB	N	G	L	N	22,000	P
Sunflowers	7	.5 - 1.0	F	F	G	N	F	P	G	DM	F	G	L	WB	N	F	M	M	8,000	P
Sunn hemp	15	1.5 - 2.0	F	F	F	Y	F	P	F	DM	F	F	L	WB	N	P	L	M	15,000	P
Teff grass	5	.13 - .25	G	G	F	N	F	G	G	SM	G	F	F	WG	N	P	M	H	1M	N
Triticale	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	M	M	15,000	F
Turnips	4	.25 - .5	F	P	G	N	G	P	G	DH	G	G	F	CB	S	P	L	N	175,000	P
Vetch, Chickling	50	.5 - 1.5	F	F	F	Y	F	F	P	SL	F	F	L	CB	N	P	L	M	2,500	F
Vetch, Common	25	.5 - 1.5	F	F	F	Y	F	F	G	SM	F	F	L	CB	N	P	L	M	8,000	F
Vetch, Hairy	15	.5 - 1.5	G	F	F	Y	F	F	F	SM	G	F	L	CB	Y	P	L	M	14,000	G
Wheat, Spring	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	N	G	M	M	15,000	F
Wheat, Winter	60	.75 - 2.0	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	M	M	15,000	F



SOUTH DAKOTA (SD) SUPPLEMENT TO CONSERVATION ENHANCEMENT ACTIVITY

CONSERVATION STEWARDSHIP PROGRAM

E328D

Additional Criteria for SD:

In addition to the criteria specified in the national job sheet E328D, the following additional criteria apply in SD:

- Any annually planted commodity crop left standing over winter, is adequate for wildlife.
- In SD the approved habitat evaluation guide is the SD-CPA-19 (WHEG).
- If using a cover crop, see table 1 below for acceptable species:



Table 1: Cover Crop - Common Species and Properties

Cover Crop	Full seeding rate lbs/acre/4	Seeding depth, inches	Reduce erosion	Increase soil organic matter	Savrange nutrients	Biological N fixation	Suppress weeds	Provide supplemental hay	Provide supplemental grazing	Rooting depth / Plant water use ^{1/}	Minimize / Reduce surface soil compaction	Minimize/ Reduce subsoil compaction	Seed size (Large or Fine)	Crop type and seeding dates ^{2/3}	Winter survival	Salinity Tolerance	CN Ratio	Mycorrhizal fungi association	Seeds/lb	Shade Tolerance
Alfalfa	6.5	.25 - .75	G	G	G	Y	G	G	F	DH	G	G	F	CB	Y	P	L	M	210,000	F
Barley	50	.75 - 2.0	G	G	G	N	G	G	G	MM	G	F	L	CG	N	G	M	M	14,000	F
Brassica hybrids	7	.25 - .5	F	F	G	N	G	F	G	MM	G	G	F	CB	N	G	L	N	180,000	P
Buckwheat ⁵	50	.5 - 1.5	P	P	F	N	F	P	P	SL	F	P	L	WB	N	P	L	N	19,000	G
Cabbage, African	5	.25 - .75	F	F	G	N	F	F	F	MM	G	G	F	CB	N	G	L	N	180,000	F
Camelina, Winter	3	.25 - .5	F	F	F	N	P	P	P	ML	P	F	F	CB	S	P	L	N	400,000	P
Canola	5	.25 - .75	F	F	G	N	G	F	F	MM	G	G	F	CB	S	G	L	N	140,000	F
Clover, Balansa	5	.25 - .75	F	P	F	Y	P	P	F	SL	P	P	F	CB	N	P	L	M	500,000	F
Clover, Crimson	15	.25 - .75	F	F	F	Y	P	F	F	SM	P	P	F	CB	S	P	L	M	150,000	F
Clover, Red	5	.25 - .75	G	F	F	Y	F	F	F	SL	F	F	F	CB	Y	P	L	M	275,000	G
Clover, Sweet	4	.25 - 1.0	G	G	F	Y	G	F	F	MM	G	G	F	CB	Y	F	L	M	260,000	G
Collards or Kale	5	.25 - .5	F	F	G	N	G	F	G	MM	G	G	F	CB	N	G	L	N	175,000	F
Corn	12	1 - 1.5	G	G	G	N	G	F	G	DH	G	G	L	WG	N	P	H	H	2,500	F
Cowpeas or Dry Beans	30	1 - 1.5	P	F	F	Y	P	P	F	SL	F	F	L	WB	N	P	L	M	4,000	F
Fava beans	75	1 - 1.5	F	F	F	Y	F	G	G	DM	F	F	L	CB	N	F	L	P	2,500	P
Flax	30	.25 - .75	F	F	F	N	P	P	P	SM	F	P	F	CB	N	P	H	H	80,000	P
Lentils	30	1 - 1.5	P	P	P	Y	P	P	P	SL	P	P	F	CB	N	P	L	M	20,000	P
Millet, hay	15	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	N	P	M	H	180,000	P
Millet, proso	25	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	N	P	M	H	80,000	P
Mustard	6	.25 - .75	F	F	F	N	G	F	P	MH	G	F	F	CB	N	P	L	N	140,000	P
Oats	70	.5 - 1.5	G	G	G	N	G	G	G	MM	G	F	L	CG	N	F	M	H	16,000	F
Peas	70	1.5 - 3.0	F	P	P	Y	F	G	G	SL	F	F	L	CB	N	P	L	M	3,500	F
Phacelia	4	.25 - .5	F	F	F	N	P	P	P	DH	F	P	F	CB	N	P	L	M	225,000	F
Radishes	8	.25 - .75	F	F	G	N	G	P	G	DH	G	G	F	CB	N	P	L	N	25,000	P
Rapeseed	5	.25 - .75	F	F	G	N	G	F	G	MM	G	G	F	CB	Y	G	L	N	140,000	F
Rye, Cereal	60	.75 - 2.0	G	G	G	N	G	G	G	MH	G	G	L	CG	Y	G	H	M	18,000	G
Ryegrass, Annual	15	.5 - 1.5	G	G	G	N	F	G	G	MM	G	F	F	CG	S	F	M	M	190,000	G
Safflowers	30	.5 - 1.0	F	F	G	N	F	P	P	DM	F	G	L	WB	N	F	M	M	15,000	P
Sorghum, Forage and Sudan Hybrids	15	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	17,000	P
Sorghum, Grain	5	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	17,000	P
Soybeans	35	1 - 1.5	F	P	F	Y	F	F	F	SM	F	F	L	WB	N	P	L	M	3,000	F
Sudangrass	20	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	25,000	P
Sugar beets	4	.25 - .5	F	P	G	N	F	P	G	DH	G	G	F	CB	N	G	L	N	22,000	P
Sunflowers	7	.5 - 1.0	F	F	G	N	F	P	G	DM	F	G	L	WB	N	F	M	M	8,000	P
Sunn hemp	15	1.5 - 2.0	F	F	F	Y	F	P	F	DM	F	F	L	WB	N	P	L	M	15,000	P
Teff grass	5	.13 - .25	G	G	F	N	F	G	G	SM	G	F	F	WG	N	P	M	H	1M	N
Triticale	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	M	M	15,000	F
Turnips	4	.25 - .5	F	P	G	N	G	P	G	DH	G	G	F	CB	S	P	L	N	175,000	P
Vetch, Chickling	50	.5 - 1.5	F	F	F	Y	F	F	P	SL	F	F	L	CB	N	P	L	M	2,500	F
Vetch, Common	25	.5 - 1.5	F	F	F	Y	F	F	G	SM	F	F	L	CB	N	P	L	M	8,000	F
Vetch, Hairy	15	.5 - 1.5	G	F	F	Y	F	F	F	SM	G	F	L	CB	Y	P	L	M	14,000	G
Wheat, Spring	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	N	G	M	M	15,000	F
Wheat, Winter	60	.75 - 2.0	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	M	M	15,000	F