



## CONSERVATION ENHANCEMENT ACTIVITY

### E340I

# CONSERVATION STEWARDSHIP PROGRAM

## Using cover crops for biological strip till

### Conservation Practice 340: Cover Crop

APPLICABLE LAND USE: Crop (Annual & Mixed)

RESOURCE CONCERN: Soil

ENHANCEMENT LIFE SPAN: 1 Year

### Enhancement Description

Establish alternating strips of cover crops in which one strip acts as a biological strip-tiller and the adjacent strip promotes soil health with high residue cover crops. This will facilitate planting of the subsequent cash crop into the biologically strip-tilled row without the need for mechanical disturbance.

### Criteria

- Plant species, seedbed preparation, seeding rates, seeding dates, seeding depths, fertility requirements, and planting methods will be consistent with applicable local criteria and soil/site conditions **(REFER TO STATE SPECIFIC LISTS)**.
- Determine method and timing of cover crop termination to meet grower's objective and current NRCS Cover Crop Termination Guidelines. Terminate the cover crop as late as practical to maximize plant biomass production and nutrient uptake.
- Select species that are compatible with other components of the cropping system.
- Use a precision guidance system to ensure seeding is placed in the existing cover crop rows.
- Do not burn cover crop residue.
- Do not harvest or graze cover crop.

E340I – Using cover crops for biological strip till	July 2019	Page   1
---	-----------	----------



# CONSERVATION STEWARDSHIP PROGRAM

## Documentation and Implementation Requirements

Participant will:

- Prior to implementation, provide NRCS with the current and planned crop rotation and field operation(s) used for each crop.

### Planned Management Rotation Including Cover Crop

Field	Planned Crops/Cover Crop (in sequence)	Planting Date	Harvest/Termination Date

### Cover Crop Mix and Seeding Rate

Species	Variety	Seed Size	Typical Seeding Depth	Seeding Rate (PLS lbs/acre)	Percent of Mix (%)

### Establishment and Management Considerations:

Task	Provide information and details
Seedbed Preparation	
Seeding Date	
Seeding Depth	
Seeding Method	
Fertilizer, as needed	
Weed Management, as needed	
Termination Date (window)	
Termination Method	



# CONSERVATION STEWARDSHIP PROGRAM

- Prior to implementation, read and follow current [NRCS Cover Crop Termination Guidelines](#).
- During implementation, cover crops must not be grazed, burned, harvested or biomass removed.
- During implementation, notify NRCS of any planned changes in crops, crop rotation, or unharvested areas to verify the planned system meets the enhancement criteria.
- After implementation, if changes to the cover crop and crop rotation were made, complete the tables above to document the applied Cover Crop for the contract period and provide to NRCS.

**NRCS will:**

- As needed, provide technical assistance in selecting cover crop mixes for the crop rotations or substitute species that would meet the criteria of the enhancement.
- As needed, provide additional assistance to the participant as requested.
- Prior to implementation, provide and explain the current [NRCS Cover Crop Termination Guidelines](#).
- During implementation, evaluate planned adjustments in cover crop selected, timing in crop rotation, management, or field operations to verify the new system meets the enhancement criteria.
- After implementation, evaluate the applied crop rotation or management using information provided from the participant, if any variation to planned evaluation, document that the applied rotation met the enhancement criteria.

**NRCS Documentation Review:**

I have reviewed all required participant documentation and have determined the participant has implemented the enhancement and met all criteria and requirements.

Participant Name \_\_\_\_\_ Contract Number \_\_\_\_\_

Total Amount Applied \_\_\_\_\_ Fiscal Year Completed \_\_\_\_\_

\_\_\_\_\_  
NRCS Technical Adequacy Signature

\_\_\_\_\_  
Date

E340I – Using cover crops for biological strip till	July 2019	Page   3
---	-----------	----------



## SOUTH DAKOTA (SD) SUPPLEMENT TO CONSERVATION ENHANCEMENT ACTIVITY

## CONSERVATION STEWARDSHIP PROGRAM

### E340I

#### **Additional Criteria for SD:**

In addition to the criteria specified in the National job sheet E340I, the following additional criteria apply in SD:

- Row or strip used to promote soil health using high residue cover crops:
  - Ninety percent (90%) of mix will be rated Good (G) or Fair (F) for Increase Soil Organic Matter on the attached Cover Crop Table 1.
- Row or strip used to act as a biological strip-tiller:
  - Ninety percent (90%) of mix will be broadleaf species identified as having a Medium (M) to Deep (D) Rooting Depth on the attached Cover Crop Table 1.



**Table 1: Cover Crop - Common Species and Properties**

Cover Crop	Full seeding rate lbs/acre/4	Seeding depth, inches	Reduce erosion	Increase soil organic matter	Scavenge nutrients	Biological N fixation	Suppress weeds	Provide supplemental hay	Provide supplemental grazing	Rooting depth / Plant water use <sup>1/</sup>	Minimize / Reduce surface soil compaction	Minimize/ Reduce subsoil compaction	Seed size (Large or Fine)	Crop type and seeding dates /2 /3	Winter Survival	Salinity Tolerance	CN Ratio	Mycorrhizal fungi/ association	Seeds/lb	Shade Tolerance
Alfalfa	6.5	.25 - .75	G	G	G	Y	G	G	F	DH	G	G	F	CB	Y	P	L	M	210,000	F
Barley	50	.75 - 2.0	G	G	G	N	G	G	G	MM	G	F	L	CG	N	G	M	M	14,000	F
Brassica hybrids	7	.25 - .5	F	F	G	N	G	F	G	MM	G	G	F	CB	N	G	L	N	180,000	P
Buckwheat /5	50	.5 - 1.5	P	P	F	N	F	P	P	SL	F	P	L	WB	N	P	L	N	19,000	G
Cabbage, African	5	.25 - .75	F	F	G	N	F	F	F	MM	G	G	F	CB	N	G	L	N	180,000	F
Camelina, Winter	3	.25 - .5	F	F	F	N	P	P	P	ML	P	F	F	CB	S	P	L	N	400,000	P
Canola	5	.25 - .75	F	F	G	N	G	F	F	MM	G	G	F	CB	S	G	L	N	140,000	F
Clover, Balansa	5	.25 - .75	F	P	F	Y	P	P	F	SL	P	P	F	CB	N	P	L	M	500,000	F
Clover, Crimson	15	.25 - .75	F	F	F	Y	P	F	F	SM	P	P	F	CB	S	P	L	M	150,000	F
Clover, Red	5	.25 - .75	G	F	F	Y	F	F	F	SL	F	F	F	CB	Y	P	L	M	275,000	G
Clover, Sweet	4	.25 - 1.0	G	G	F	Y	G	F	F	MM	G	G	F	CB	Y	F	L	M	260,000	G
Collards or Kale	5	.25 - .5	F	F	G	N	G	F	G	MM	G	G	F	CB	N	G	L	N	175,000	F
Corn	12	1 - 1.5	G	G	G	N	G	F	G	DH	G	G	L	WG	N	P	H	H	2,500	F
Cowpeas or Dry Beans	30	1 - 1.5	P	F	F	Y	P	P	F	SL	F	F	L	WB	N	P	L	M	4,000	F
Fava beans	75	1 - 1.5	F	F	F	Y	F	G	G	DM	F	F	L	CB	N	F	L	P	2,500	P
Flax	30	.25 - .75	F	F	F	N	P	P	P	SM	F	P	F	CB	N	P	H	H	80,000	P
Lentils	30	1 - 1.5	P	P	P	Y	P	P	P	SL	P	P	F	CB	N	P	L	M	20,000	P
Millet, hay	15	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	N	P	M	H	180,000	P
Millet, proso	25	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	N	P	M	H	80,000	P
Mustard	6	.25 - .75	F	F	F	N	G	F	P	MH	G	F	F	CB	N	P	L	N	140,000	P
Oats	70	.5 - 1.5	G	G	G	N	G	G	G	MM	G	F	L	CG	N	F	M	H	16,000	F
Peas	70	1.5 - 3.0	F	P	P	Y	F	G	G	SL	F	F	L	CB	N	P	L	M	3,500	F
Phacelia	4	.25 - .5	F	F	F	N	P	P	P	DH	F	P	F	CB	N	P	L	M	225,000	F
Radishes	8	.25 - .75	F	F	G	N	G	P	G	DH	G	G	F	CB	N	P	L	N	25,000	P
Rapeseed	5	.25 - .75	F	F	G	N	G	F	G	MM	G	G	F	CB	Y	G	L	N	140,000	F
Rye, Cereal	60	.75 - 2.0	G	G	G	N	G	G	G	MH	G	G	L	CG	Y	G	H	M	18,000	G
Ryegrass, Annual	15	.5 - 1.5	G	G	G	N	F	G	G	MM	G	F	F	CG	S	F	M	M	190,000	G
Safflowers	30	.5 - 1.0	F	F	G	N	F	P	P	DM	F	G	L	WB	N	F	M	M	15,000	P
Sorghum, Forage and Sudan Hybrids	15	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	17,000	P
Sorghum, Grain	5	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	17,000	P
Soybeans	35	1 - 1.5	F	P	F	Y	F	F	F	SM	F	F	L	WB	N	P	L	M	3,000	F
Sudangrass	20	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	H	25,000	P
Sugar beets	4	.25 - .5	F	P	G	N	F	P	G	DH	G	G	F	CB	N	G	L	N	22,000	P
Sunflowers	7	.5 - 1.0	F	F	G	N	F	P	G	DM	F	G	L	WB	N	F	M	M	8,000	P
Sunn hemp	15	1.5 - 2.0	F	F	F	Y	F	P	F	DM	F	F	L	WB	N	P	L	M	15,000	P
Teff grass	5	.13 - .25	G	G	F	N	F	G	G	SM	G	F	F	WG	N	P	M	H	1M	N
Triticale	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	M	M	15,000	F
Turnips	4	.25 - .5	F	P	G	N	G	P	G	DH	G	G	F	CB	S	P	L	N	175,000	P
Vetch, Chickling	50	.5 - 1.5	F	F	F	Y	F	F	P	SL	F	F	L	CB	N	P	L	M	2,500	F
Vetch, Common	25	.5 - 1.5	F	F	F	Y	F	F	G	SM	F	F	L	CB	N	P	L	M	8,000	F
Vetch, Hairy	15	.5 - 1.5	G	F	F	Y	F	F	F	SM	G	F	L	CB	Y	P	L	M	14,000	G
Wheat, Spring	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	N	G	M	M	15,000	F
Wheat, Winter	60	.75 - 2.0	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	M	M	15,000	F



/1 Rooting Depth/Water Use		/2 Crop types		Ratings		
SL=	Shallow rooted/Low water use	Shallow=	6 - 18 inches	CG = cool season grass	L= Low	G= Good
SM=	Shallow rooted/Medium water use	Medium=	18 - 24 inches	CB = cool season broadleaf	M= Medium	F= Fair
SH=	Shallow rooted/High water use	Deep=	24 + inches	WG = warm season grass	H= High	P= Poor
ML=	Medium rooted/Low water use			WB = warm season broadleaf	Y= Yes	
MM=	Medium rooted/Medium water use				N = No	
MH=	Medium rooted/High water use				S = Sporadic	
DL=	Deep rooted/Low water use				N/A= Not Applicable	
DM=	Deep rooted/Medium water use					
DH=	Deep rooted/High water use					
/3 Seeding Dates		/4 Full Seeding rates		/5 Buckwheat contamination		
May 1 through August 5 – warm season winter kill species		Multiply by the percent desired if mixtures are used.		To reduce chances of buckwheat contamination in wheat		
Early spring through August 20 – cool season winter kill species				do not rotate to wheat for grain for 2 years		
August 1 through Winter – species that do not winter kill						
Seeding dates fluctuate annually. Seeding dates may be adjusted up to 15 days by the District Conservationist, based on local weather and site conditions.						