

## **Introduction**

The Maple River 8-Digit Hydrologic Unit Code (HUC) (09020205) sub-basin is approximately 998,700 acres covering parts of five counties (Barnes, Cass, Ransom, Richland, and Steele) in the Souris-Red-Rainy Region - Red River Sub-Region. Of the 998,750 acres, Cass County contains 56%, Barnes 26%, Ransom 10%, Richland 1%, and Steele 7%. There are approximately 880 farms in the sub-basin.



This sub-basin encompasses commodities ranging from sugar beets, corn and soybeans, and multiple small grain crops to beef cattle and swine.

Conservation assistance is provided by five NRCS service centers, one soil survey office, and two Resource Conservation & Development offices.

Produced by the Natural Resources Planning Staff Bismarck, ND The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.



## **Physical Description**

The following table and map show land cover / land use within the sub-basin.

Land Cover/ Land Use (National Resources Inventory [NRI]) <sup>1</sup>	Acres	Percent of HUC			
Forestland	0	0%			
Cropland	861,800	86%			
Conservation Reserve Program (CRP) Land <sup>28</sup>	9,900	1%			
Tame Grass/Hayland	12,400	1%			
Pastureland	18,500	2%			
Rangeland	27,400	3%			
Urban/Farmstead/ Transportation Land	63,700	7%			
Water/Wetlands	2,400	1%*			
Federal Lands	2,600	1%*			
Minor Lands **	0	0%			
North Dakota HUC Totals 998,700100%*					
* Less than one percent of total acres. See below for special considerations. ** Minor land includes farmsteads, windbreaks, marshland, etc. a: Estimate from Farm Service Agency records and include CRP/CREP. b: Totals may not add due to rounding and small unknown acreages.					
Irrigated Land (ND State Water Commision Estimates)	3,000	<1%			





The above map was developed from U.S. Geologic Survey's (USGS) ND Gap Analysis Program data.<sup>3</sup>



The following map is a plot of 1961-1990 annual average precipitation contours from National Oceanic and Atmospheric Administration (NOAA) Cooperative stations and (where appropriate) USDA-NRCS Snowpack Telemetry (SNOTEL) Stations. Christopher Daly used the PRISM (Parameter-elevation Regressions on Independent Slopes Model) model to generate the gridded estimates from which this map was derived: the modeled grid was approximately 4x4 km latitude/longitude, and was resampled to 2x2 km using a Gaussian filter. Mapping was performed by Jenny Weisberg and Nathaniel DeYoung. Funding was provided by USDA-NRCS National Water and Climate Center. (4/20/98)





The sub-basin is part of the Souris-Red-Rainy River Region - Red River Sub-Region. All drainage patterns flow to the east ending at the Red River, which flows north into Canada. The following map shows the relief for the sub-basin.<sup>4</sup>





The North Dakota Department of Health collects water quality data on major water bodies. The following table shows the total miles of streams and acres of lakes/reservoirs within the sub-basin and also the miles and acres that have a water quality limitation. The second part of the table shows the livestock numbers, feeding operations, and permitted operations. Also included is the livestock numbers for all cattle, beef cows, dairy cows, hogs and pigs, and sheep and lambs. The livestock numbers were extrapolated from 2002 Agricultural Census county data to 8-digit HUC's.

		Units	North Dakota ₅	Maple River Sub-basin <sup>6</sup>	Maple River as percent of North Dakota	Impaired Water Quality (303d) <sup>7</sup>	Percent Impaired* Maple River
Water Quality	Total – Major Water bodies						
Data	Rivers/Streams	Miles	56,687	1,771	3.1%	41	2.3%
*Percent of Total Miles and acres in HUC	Lakes/Reservoirs	Acres	434,658	446	0.1%	None - ID	0%

Animal Feeding Facilities – North Dakota Department of Health <sup>8</sup>					
Animal Type	Dairy	Beef	Swine	Other	Total
Number of Animal Feeding Operations	2	7	23	0	32
Number of Animals	250	5,675	14,951	0	20,876
Number of Permitted Farms					22

Livestock Numbers (rounded to nearest 100) <sup>9</sup>					
	Cattle and Calves	Beef Cows	Dairy Cows	Hogs and Pigs	Sheep and Lambs
North Dakota	1,873,200	982,300	34,500	138,800	114,000
Maple River	21,700	10,200	300	10,500	2,200
Maple River as a percent of North Dakota	1.2%	1.0%	0.9%	7.6%	1.9%



Common Resource Areas (CRA's) are geographical areas where resource concerns, problems, or treatments are similar. Landscape conditions, soil, climate, human considerations, and other natural resource information were used to determine the geographic boundaries. CRA's are subsets of Major Land Resource Areas. The following map<sup>10</sup> shows the CRA's for Maple River sub-basin with the descriptions below.



**<u>55B.1 – Central Black Glaciated Drift Plain</u>**: The Central Black Glaciated Drift Plains are a gently rolling to undulating landscape with a thick layer of glacial till. Temporary and seasonal wetlands are numerous throughout the area. These soils are very fertile, but agricultural success is subject to annual climatic fluctuations. Most of the soils are deep, well drained and moderately well drained, sandy to clayey and have a frigid temperature regime.

**56.1 – Red River Valley:** The Red River Valley (Glaciated Lake Agassiz) is an extremely flat landscape composed of thick lacustrine sediments. Soils range from silty to clayey in texture. Most soils have a high water table and are very productive. Saline soils exist in places. Most areas are farmed with main crops being small grain, sugar beets, and soybeans. The native vegetation was tall grass prairie. Primary resource concerns are soil erosion and deposition by wind.

**56.3 – Sheyenne Delta of the Red River Valley**: The Sheyenne Delta landscape ranges from strongly rolling sand dunes in the northeastern edge, to nearly level high water table sandy soils of the south. A risk of wind erosion exists throughout this area. The area is used for range and cropland. A portion of the Sheyenne delta is in the Sheyenne National Grassland.



# Soil Productivity 11

The Maple River sub-basin is divided in productivity by the marginal to moderately productive soils of Glacial Lake Agassiz beaches. The soils to the west of the beaches are glacial drift and marginal to high in productivity, while the soils to the east of the beaches are the highly productive soils of Glacial Lake Agassiz.





## Common Land Unit

The entire sub-basin has the common land unit digitized by Farm Services Agency (FSA).

### **Resource Concerns**

One of the goals of the Natural Resource Conservation Service is to look at an area to help quantify the types and amounts of resources that may be of concern. This helps in identifying priority areas for the types and amounts of assistance to be given to a particular watershed.

- The acres of land above sustainable levels due to wind erosion have increased by 7,600 acres from 1987 to 1997. These levels of unsustainable wind erosion rates may be in part to intense row cropping systems in the sub-basin
- NRI estimates indicate water erosion rates on cultivated cropland were 1.3 tons/acre/year in 1997.
- Controlling erosion not only sustains the longterm productivity of the land, but also affects the amount of soil, pesticides, fertilizer, and other organic material that move into the basin's waters.
- Through NRCS programs, many farmers and ranchers have applied conservation practices to reduce the effects of erosion by water.



 NRI estimates indicate 25,500 acres of Highly Erodible Land (HEL) in 1997 compared to 23,000 acres in 1987. This is nearly an 11% increase in HEL being farmed.

- The two 303(d) listed stream segments were listed primarily for sediment and siltation. total fecal coliform was also listed on one of the streams segments. Stream reaches listed for sediment are affected by erosion on croplands and from stream banks.
- Conservation practices that can be used to address these water quality issues include erosion control, nutrient and pest management, grazing management, and riparian buffers.



## **Resource Concerns - Continued**

The following table shows the different projects, plans, studies, and assessments and their status that have been conducted within the sub-basin.

Watershed Projects, Plans, Studies and Assessments					
NRCS Watershed Projects		NRCS Watershed Plans, Studies & Assessments			
Name	Status	Name	Status		
None	NA	None	NA		
NDDH TMDLs		Soil Conservation District Assessments and Studies			
Number Listed		Name	Status		
Lakes/Reservoirs – 0	Streams – 2	Maple River Assessment – Cass Co.	Ongoing		
EPA 319 Watershed Projects					
Name		Status			
None		NA			

#### Soil

- Sandy soils and steep soils still require conservation practices to control excessive soil erosion.
- Windbreak plantings, reduced tillage systems, and improved cropping systems are still needed.
- Soil health, especially compaction on heavier or fine textures soils and organic matter on sandy soils are two resource concerns.

#### Water

- Aquifers<sup>12</sup> There are five aquifers (Galesburg/Page, Sheyenne Delta, McVille, Tower City, and Bantel) under the sub-basin.
- Sediment and nutrients are primary water quality pollutants impairing the watershed streams and lakes.
- The Maple River has a large number of livestock operations on or near the river, which are having some water quality impacts from nutrient loading and total fecal coliform.



• Flooding in the valley does occasionally occur and impact crop production.



## **Resource Concerns - Continued**

#### Air

• Blowing snow is a concern during the winter months.

#### Plants

- Major concerns are with controlling invasive weeds and maintaining good pasture condition.
- Direct seeding and annual cropping has limited success.
- Soil erosion and low organic matter remain resource concerns.
- Conventional tillage systems are still utilized.
- Water conservation and water quality (potential for pesticide contamination) are issues on irrigated cropland.
- Noxious weeds and poor range condition reduce productivity for livestock and wildlife.
- There is no real forest land present in the sub-basin. There are many field and farmstead windbreaks.

#### Animals

• Animals on threatened and endangered species list are noted in the following table.

Federally Listed Threatened And Endangered Species				
Species Category	Threatened	Endangered	Candidate	
Mammals	None	None	None	
Birds	Bald Eagle	Whooping Crane	None	
Fish	None	None	None	
Invertebrates	None	None	Dakota Skipper	
Plants	Western Prairie	None	None	
	Fringed Orchid			
Critical Habitat – None				



## Census and Social Data<sup>13</sup>

Number of Farms: 880

#### Number of Operators:

- Average Age: 53
- Full-Time Operators: 71%
- Part-Time Operators: 29%





# Estimated Level of Willingness and Ability to Participate in Conservation: MODERATE

#### Limited Resource and Beginning Farmer

Four percent of the operators are minority producers. Limited Resource Farmers are also estimated at 2.5 percent. Although rather low percentages these facts point to the potential need for special technical assistance targeted to reach people who (1) may lack experience with government farm programs, (2) have good stewardship intentions but lack management skills, and (3) lack the time to visit an NRCS field office and seek assistance.

All data is provided "as is." There are no warranties, express or implied, including warranty of fitness for a particular purpose, accompanying this document. Use for general planning purposes only.



## **References**

- <sup>2</sup> USDA-Farm Services Agency, Common Land Unit GIS data layer, 2005.
- <sup>3</sup> USDI-US Geologic Services, ND GAP analysis data, 2005.
- <sup>4</sup> USDA-NRCS, Natural Resources Planning Staff, 30 meter Relief Data GIS data layer, 2002.
- <sup>5</sup> ND Department of Health, Environmental Health Section, Water Quality Division, National Hydrography GIS layers, June 2006.
- <sup>6</sup> ND Department of Health, Environmental Health Section, Water Quality Division, National Hydrography GIS layers, June 2006.
- <sup>7</sup> ND Department of Health, Environmental Health Section, Water Quality Division, List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota, 2006.
- <sup>8</sup> ND Department of Health, Environmental Health Section, Water Quality Division, Animal Feeding Operations Program data, 2006.
- <sup>9</sup> 2002 Census of Agriculture, North Dakota, State and County Data Volume 1, Geographic Area Series Part 34, U.S. Department of Agriculture, National Agricultural Statistics Service, June 2004. (county data was prorated to HUC by the percent of a HUC in a county)
- <sup>10</sup> USDA-NRCS, Natural Resources Planning Staff, Common Resource Area GIS data layer, 2004.
- <sup>11</sup> USDA-NRCS, Natural Resources Planning Staff, Soils Productivity GIS data layer, 2005.
- <sup>12</sup> ND Department of Health, Environmental Health Section, Water Quality Division, Ambient Ground Water Monitoring Program data, 1997.
- <sup>13</sup> 2002 Census of Agriculture, North Dakota, State and County Data Volume 1, Geographic Area Series Part 34, U.S. Department of Agriculture, National Agricultural Statistics Service, June 2004. (county data was prorated to HUC by the percent of a HUC in a county)

<sup>&</sup>lt;sup>1</sup> USDA-NRCS, NRI data.