

CONSERVATION ENHANCEMENT ACTIVITY

E3281



Forage harvest to reduce water quality impacts by utilization of excess soil nutrients

Conservation Practice 328: Conservation Crop Rotation

APPLICABLE LAND USE: Crop (Annual & Mixed); Crop (Perennial)

RESOURCE CONCERN: Water

ENHANCEMENT LIFE SPAN: 1 Year

Enhancement Description

Establish a forage crop (single species or mix) following a primary annual crop to take up excess soil nutrients. Select forage known to effectively utilize and scavenge nutrients. Forage shall be harvested for forage, but not be grazed or burned.

Criteria

- This enhancement is applicable on fields where excess soil nutrients cause or increase
 water quality degradation concerns. Presence of excess nutrients must be identified
 in recent soil tests or increased risk to water quality documented by risk assessment
 tool. (Refer to state specific guidance of options to maximize nutrient uptake in
 local climate and cropping systems)
- Forage species, seedbed preparation, seeding rates, seeding dates, seeding depths, fertility requirements, and planting methods will be consistent with applicable local criteria and soil/site conditions. (Refer to state specific lists of forage crops known to effectively utilize and scavenge nutrients)
- Select forage crop (single species or mix of two or more species) and planting dates
 which will not compete with the other crop(s) yield or harvest. If legumes are part
 of the forage mix, consider that this may add nutrients to the system.

E328I - Forage harvest to reduce water	August 2019	Page 1
quality impacts by utilization of excess soil		
nutrients		



United States Department of Agriculture

- Select forage crop that is compatible with other components of the crop rotation and for its ability to efficiently scavenge and utilize excess soil nutrients, specifically nitrogen or phosphorous, whichever is identified as a potential risk to water quality. Nutrient uptake only occurs when a crop is actively growing. Therefore, it is imperative that the crops in rotation be planted as soon as possible after forage crop harvest (hay/balage/haylage/etc.) to maximize nutrient cycling and minimize offsite transport of nutrients.
- Determine method and timing of forage crop harvest to meet client objectives. Harvest the forage crop as late as practical to maximize plant biomass production and nutrient uptake.
- Ensure any herbicides used in the crop rotation are compatible with forage crop selections.
- Do not burn forage or residue.
- Do not graze forage crop.
- Reduce or maintain soil erosion from water and wind to below soil tolerance (T) level (average annual soil loss).



United States Department of Agriculture

CONSERVATION ENHANCEMENT ACTIVITY E328I OREGON SUPPLEMENT



Contact State Agronomist for scavenger crops and guidance of options to maximize nutrient uptake.

Enhancement E328I March 2020 Page | 1



E328I - Forage harvest to reduce water

nutrients

quality impacts by utilization of excess soil

<u>Documenta</u>	ation and Impleme	ntation Requirements		
		rovide NRCS with the curr	ent and planned cr	op rotation and field
-		entified in soil tests. Soil t	easts should be take	un as clasa to production
	st as possible.	entinea in son tests. 3011 t	ests siloulu de tuke	ii us ciose to production
Field	Soil Test Date	Nutrient (Nitrogen o	r Phosphorus)	Soil Test Nutrient Result (ppm or lbs/ac)
Current Ma	anagement Rotatio	n		
Field Current Cr		Crops (in sequence)	Planting Date	Harvest Date
			ri e	
Current Fie	eld Operations for E	ach Crop		
Field	Crop	Field Ope	eration	Tim <mark>ing of Field</mark> Operation (month/year)
		·		

August 2019

Page | 3

Planned Management Rotation including Forage Crop

Field	Planned Crops/Forage Crop (in sequence)	Planting Date	Harvest Date

Planned Field Operations for Each Crop

Field	Crop	Field Operation	Timing of Field Operation (month/year)

Planned Forage Crop and Seeding Rate (forage crop may be single species or mix of two or more species)

Species	Variety	Seed Size	Typic <mark>al</mark> Seeding D <mark>epth</mark>	Seeding Rate (PLS lbs/acre)	Percent of Mix (%)

Forage Crop Establishment and Management Considerations

Establish forage crop mix as soon as practical prior to or after harvest of the production crop.
During implementation, forage crop must not be grazed or burned.
During implementation, notify NRCS of any planned changes in forage crop mix or crop
rotation, or management to verify the planned system meets the enhancement criteria.

E328I - Forage harvest to reduce water	August 2019	Page 4
quality impacts by utilization of excess soil		
nutrients		



United States Department of Agriculture

☐ After implementation, if changes were made, update the tables above to document the applied crop rotation for the contract period and provide to NRCS.

After implementation, complete the table below and provide to NRCS

Task	Provide information and details
Seedbed Preparation	
Seeding Date	
Seeding Depth	
Seeding Method	
Fertilizer, as needed	
Weed Management, as needed	
Harvest Date (window)	
Harvest Method	

NRCS will:

As needed, provide technical assistance in selecting forage crop for the crop rotation or substitute species that would meet the criteria of the enhancement. Forage crop may consist a single species or mix of two or more species.	0
As needed, provide additional assistance to the participant as requested.	
Prior to implementation, verify the enhancement is being planned on fields where excess soil nutrients cause or increase water quality degradation concerns. Presence of excess nutrients must be identified in recent soil tests or increased risk to water quality documented by risk assessment tool. <refer guidance="" specific="" state="" to=""></refer>	
Prior to implementation, use information provided from the participant to calculate the average annual soil erosion value (water and wind) for each field using NRCS erosion prediction technologies.	
Benchmark Management Soil Loss = tons/acre/year	
Planned Management Soil Loss = tons/acre/year During implementation, evaluate any planned changes in forage crop selected, timing in crop rotation, management, or field operations to verify the new system meets the enhancement criteria.	

E328I - Forage harvest to reduce water	August 2019	Page 5
quality impacts by utilization of excess soil		
nutrients		



	After implementation, if there were any changes to planned rotation or management evaluate the applied crop rotation using information provided from the participant to calculate average annual erosion value to document that the applied rotation meets the enhancement criteria.			
	Applied Management Soil Loss = to	ns/acre/year		
<u>NF</u>	RCS Documentation Review:			
I have reviewed all required participant documentation and have determined the participant has implemented the enhancement and met all criteria and requirements.				
Pa	rticipant Name	Contract Number		
То	tal Amount Applied	Fiscal Year Completed		
NR	RCS Technical Adequacy Signature	Date		

E328I - Forage harvest to reduce water	August 2019	Page 6
quality impacts by utilization of excess soil		
nutrients		