



CONSERVATION ENHANCEMENT ACTIVITY

E328E

CONSERVATION STEWARDSHIP PROGRAM

Soil health crop rotation

Conservation Practice 328: Conservation Crop Rotation

APPLICABLE LAND USE: Crop (Annual & Mixed)

RESOURCE CONCERN: Soil

PRACTICE LIFE SPAN: 1 Year

Enhancement Description

Implement a crop rotation which addresses all four principle components of soil health: increases diversity of the cropping system; maintains residue throughout the year; keeps a living root; and minimizes soil chemical, physical and biological disturbance. The rotation will include at least 4 different crop and/or cover crop types (crop types include cool season grass, warm season grass, cool season broadleaf, warm season broadleaf) grown in a sequence that will produce a positive trend in the Organic Matter (OM) subfactor value over the life of the rotation, as determined by the Soil Conditioning Index (SCI). The current NRCS wind and water erosion prediction technologies must be used to document the rotation and SCI calculations.

Criteria

- Crops must be grown in a planned sequence as outlined in the plan. The crop rotation must include a minimum of four different crop types. For the purpose of this criteria a cover crop is considered a different crop.
- Where applicable, plan suitable crop substitutions when the planned crop cannot be planted due to weather, soil conditions, or other local situations.
- Grow crops that will produce a positive trend in the Organic Matter (OM) subfactor value over the life of the rotation, as determined by the Soil Conditioning Index (SCI). (management SCI value)



CONSERVATION STEWARDSHIP PROGRAM

- The crop rotation includes at least 2 years of high residue crops and/or cover crops per 3 years of the rotation. **(See STATE list of high residue crops)**
- For crop diversity, the planned crop sequence should contain four different crop types; for example, a mix of the following: warm season grass; warm season broadleaf; cool season grass; cool season broadleaf.
- Leave crop residue on the soil surface throughout the year.
- Keep a living root system established as much as practical for the given soil, cropping system, and climate area. Maximize root growth periods by planting the next crop or cover crop as soon as practical after the harvest and/or utilize perennial crops in the rotation. Aim to have living roots at least 90% of available growing days. **(See STATE provided guidance of options to maximize living root systems in local climate and cropping systems; determine available growing days and period of no growth, such as frozen periods in the north.)** Show before and after management files from current NRCS wind and water erosion prediction technologies to document benchmark and planned crop rotation to show increase in living root periods.
- Minimize all types of soil disturbance. No more than one crop-year in the rotation will have a Soil Tillage Intensity Rating (STIR) value greater than 20 (crop STIR value) and the rotation will have a positive trending SCI (management SCI value).



CONSERVATION STEWARDSHIP PROGRAM

Documentation and Implementation Requirements

Participant will:

- Prior to implementation, provide NRCS with the current and planned crop rotation and planned field operation(s) used for each crop.

Current Management – Crop Rotation

Field	Acres	Planned Crops (in sequence)	Length of Crop Rotation (years)	Crop Type (Warm Grass-WG, Cool Grass-CG, Warm Broadleaf-WB, Cool Broadleaf-CB)

Current Management – Field Operations

Field	Crop	Field Operation	Timing of Field Operation (month/year)

Planned Management – Crop Rotation *(Planned crop rotation must include at least 2 years of high residue crops and/or cover crops per 3 years of the rotation and at least 4 different crop types. Use STATE list of high residue crops.)*

Field	Acres	Planned Crops (in sequence)	Length of Crop Rotation (years)	Crop Type (Warm Grass-WG, Cool Grass-CG, Warm Broadleaf-WB, Cool Broadleaf-CB)



CONSERVATION STEWARDSHIP PROGRAM

Planned Management – Field Operations

Field	Crop	Field Operation	Timing of Field Operation (month/year)

- During implementation, notify NRCS of any planned changes in crops, crop rotation, or field operations to verify the planned system meets the enhancement criteria.
- During implementation, take dated pictures with field indicated at least every 3 months to show residue or growing crops.
- During implementation, leave crop residue on the soil surface throughout the year.
- After implementation, if changes to the rotation were made, complete the tables above to document the applied Conservation Crop Rotation for the contract period and provide to NRCS.
- After implementation, provide for review pictures showing residue or growing crops throughout the year.

NRCS will:

- As needed, provide technical assistance in selecting crop rotations or substitute crops that would meet the criteria of the enhancement.
- Prior to implementation, verify the planned crop rotation includes at least four different crop types.
- Prior to implementation, verify the crop rotation includes at least 2 years of high residue crops and/or cover crops per 3 years of the rotation. (Use STATE list of high residue crops.)
- Prior to implementation, use information provided from the participant to calculate the management Soil Conditioning Index (SCI) value for each field using current NRCS wind



CONSERVATION STEWARDSHIP PROGRAM

and water erosion prediction technologies. Crop rotation must produce a positive trend in the Organic Matter (OM) subfactor value.

Management SCI Value = _____

OM subfactor value = _____

- Prior to implementation, use NRCS wind and water erosion prediction technologies to document benchmark and planned crop rotation to show increase in living root periods.
- During implementation, evaluate planned changes in crops, crop rotation, or field operations to verify the planned system meets the enhancement criteria.
- After implementation, if the applied crop rotation is different than the planned crop rotation, use information provided from the participant to calculate SCI value to document that the applied rotation met the enhancement criteria.
Management SCI Value = _____ OM subfactor value = _____
- After implementation, review pictures showing residue or growing green crops throughout the year to verify the applied system meets the enhancement criteria.

NRCS Documentation Review:

I have reviewed all required participant documentation and have determined the participant has implemented the enhancement and met all criteria and requirements.

Participant Name _____ Contract Number _____

Total Amount Applied _____ Fiscal Year Completed _____

NRCS Technical Adequacy Signature

Date

ALABAMA – E328E Soil Health Crop Rotation

-All crops must be planted no till/strip till

-No more than 2 years of a 5-year rotation may be planted to the same crop (except corn)

1. **A perennial grass grown at least 2 years but no more than 4 years with at least two other crops in the rotation:**

Examples:

- a. Corn grain, 1 yr., cotton 1 yr., ***Small grain (rye, wheat, oat, barley, triticale)/legume cover crop not harvested (interseeded) prior to each cash crop, Corn grain, 1 yr., Fescue or Bahiagrass 2 yr. or more****.
- b. Corn grain, 1 yr., Soybean 1yr. ***Small grain/legume/brassica cover crop***, 1 yr. Corn grain, 1 yr., ***Fescue 2 yr. or more.***
- c. Corn grain, 1 yr. ***Small grain/legume/brassica cover crop***, Peanut 1yr. ***Small grain/legume/brassica cover crop***, 1 yr. Corn grain ***Small grain/legume/brassica cover crop***, 1 yr., ***Bahiagrass 2 yr. or more.***

OR;

2. **A legume that is grown at least 2 years but no more than 4 years with at least one other crop in the rotation:**

Examples:

- a. Any of the above rotations except substituting alfalfa or sericea lespedeza for the perennial grasses

OR;

3. **A legume-grass mixture that is grown at least 2 years but no more than 4 years with at least one other crop in the rotation:**

Examples:

- a. Any of the above rotations except substituting fescue/white clover or bahiagrass/white clover for the perennial grasses

OR;

4. **A small grain grown in combination with a legume, forb, or grass-forb mixture that is used as a green manure, whether interseeded or planted after small grain harvest with at least two other crops in the rotation.**

Examples:

- a. Corn (grain) 1 yr. , ***Wheat*** harvested rdl, ***Pearl Millet*** summer cover crop not harvest; ***Small grain/legume/brassica cover crop***; Cotton 1 yr., ***Small grain/legume/brassica cover crop***, Corn grain, 1 yr. ***wheat*** harvested for grain, soybean double-crop, Corn grain, 1 yr. ***Small grain/legume/brassica cover crop***. Cover crop must be at least 24 inches tall or 5000 lbs. dry matter prior to termination except prior to corn planting.
- b. Corn (grain) 1 yr. , ***Small grain/legume/brassica cover crop***; Peanut 1 yr., ***Small grain/legume/brassica cover crop***, Corn grain, 1 yr. ***wheat*** harvested for grain, soybean double-crop, Corn grain, 1 yr. ***Small grain/legume/brassica cover crop***.. Cover crop must be at least 24 inches tall or 5000 lbs. dry matter prior to termination except prior to corn planting.

* The Resource Conserving Crop in all examples rotations are ***italic and bold*** font.