

Cover Crop Table 1



Table 1: Cover Crop - Common Species and Properties																				
Cover Crop	Full seeding rate lbs/acre	Seeding depth, inches	Reduce erosion	Increase soil organic matter	Scavenge nutrients	Promote biological nitrogen fixation	Suppress weeds	Provide supplemental hay	Provide supplemental grazing	Rooting depth/Plant water use	Minimize/Reduce surface soil compaction	Minimize/Reduce subsoil compaction	Seed size (Large or Fine)	Crop type	Winter Survival	Salinity Tolerance	C:N Ratio	Mycorrhizal fungi association	Seeds/lb	Shade Tolerance
Alfalfa	6.5	.2575	G	G	G	Y	G	G	F	DH	G	G	F	СВ	Υ	Р	L	м	210,000	F
Barley	50	.75 - 2.0	G	G	G	N	G	G	G	MM	G	F	L	CG	N	G	м	м	14,000	F
Brassica hybrids	7	.255	F	F	G	N	G	F	G	MM	G	G	F	СВ	N	G	L	N	180,000	Р
Buckwheat	50	.5 - 1.5	Р	Р	F	N	F	Р	Р	SL	F	Р	L	WB	N	Р	L	N	19,000	G
Cabbage, African	5	.2575	F	F	G	N	F	F	F	ММ	G	G	F	СВ	N	G	L	N	180,000	F
Camelina, Winter	3	.255	F	F	F	N	Р	Р	Р	ML	Р	F	F	СВ	S	Р	L	N	400,000	Р
Canola	5	.2575	F	F	G	N	G	F	F	мм	G	G	F	СВ	S	G	L	N	140,000	F
Clover, Balansa	5	.2575	F	Р	F	Y	Р	Р	F	SL	Р	Р	F	СВ	Ν	Р	L	м	500,000	F
Clover, Crimson	15	.2575	F	F	F	Y	Р	F	F	SM	Р	Р	F	СВ	s	Р	L	м	150,000	F
Clover, Red	5	.2575	G	F	F	Y	F	F	F	SL	F	F	F	СВ	Y	Р	L	м	275,000	G
Clover, Sweet	4	.25 - 1.0	G	G	F	Y	G	F	F	MM	G	G	F	CB	Y	F	L	M	260,000	G
Collards or Kale	5	.255	F	F	G	N	G	F	G	MM	G	G	F	СВ	N	G	L	N	175,000	F
Corn	12	1 - 1.5	G	G	G	N	G	F	G	DH	G	G	L	WG	N	Р	н	н	2,500	F
																			,	
Cowpeas or Dry Beans	30	1 - 1.5	Р	F	F	Y	Р	Р	F	SL	F	F	L	WB	N	Р	L	м	4,000	F
Fava beans	75	1 - 1.5	F	F	F	Y	F	G	G	DM	F	F	L	СВ	Ν	F	L	Р	2,500	Р
Flax	30	.2575	F	F	F	N	Р	Р	Р	SM	F	Р	F	СВ	N	Р	н	н	80,000	Р
Lentils	30	1 - 1.5	Р	Р	Р	Y	Р	Р	Р	SL	Р	Р	F	СВ	Ν	Р	L	м	20,000	Р
Millet, hay	15	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	Ν	Р	м	н	180,000	Р
Millet, proso	25	.5 - 1.0	G	G	G	N	G	G	G	SL	G	F	F	WG	Ν	Р	м	н	80,000	Р
Mustard	6	.2575	F	F	F	N	G	F	Р	MH	G	F	F	СВ	N	Р	L	Ν	140,000	Р
Oats	70	.5 - 1.5	G	G	G	N	G	G	G	ММ	G	F	L	CG	Ν	F	м	н	16,000	F
Peas	70	1.5 - 3.0	F	Р	Р	Y	F	G	G	SL	F	F	L	СВ	N	Р	L	м	3,500	F
Phacelia	4	.255	F	F	F	N	Р	Р	Р	DH	F	Р	F	СВ	N	Р	L	м	225,000	F
Radishes	8	.2575	F	F	G	N	G	Р	G	DH	G	G	F	СВ	N	Р	L	N	25,000	Р
Rapeseed	5	.2575	F	F	G	N	G	F	G	MM	G	G	F	СВ	s	G	L	N	140,000	F
Rye, Cereal planted after Oct.1	60	.75 - 2.0	G	G	G	N	G	G	G	мн	G	G	L	CG	Y	G	н	М	18,000	G
Rye, Cereal planted			_	_	_		_	_	_		-	_								
on or before Oct.1	40	.75 - 2.1	G	G	G	N	G	G	G	MH	G	G	L	CG	Y	G	H	M	18,000	G
Ryegrass, Annual	15	.5 - 1.5	G	G	G	N	F	G	G	MM	G	F	F	CG	S	F	M	M	190,000	G
Safflowers	30	.5 - 1.0	F	F	G	N	F	Р	Р	DM	F	G	L	WB	N	F	М	M	15,000	Р
Sorghum, Forage and Sudan Hybrids	15	.5 - 1.5	G	G	G	N	G	G	G	ММ	G	G	L	WG	N	F	м	н	17,000	Р
Sorghum, Grain	5	.5 - 1.5	G	G	G	N	G	G	G	MM	G	G	L	WG	N	F	M	н	17,000	
Soybeans	35	1 - 1.5	F	P	F	Y	F	F	F	SM	F	F	L	WB	N	Р	L	м	3,000	
Sudangrass	20	.5 - 1.5	G	G	G	N	г G	G	G	MM	G	G	L	WG	N	F	M	H	25,000	
Sugar beets	4	.255	F	P	G	N	F	P	G	DH	G	G	F	CB	N	G	L	N	23,000	
Sugar beets Sunflowers	7	.255	F	F	G	N	F	P P	G	DH	F	G	F	WB	N	F	M	M	8,000	
	15		F	F	F	Y	F	P	F	DM	F	F	L	WB	N	Р Р	L	M		
Sunn hemp Toff grass		1.5 - 2.0			F		F						F			P P			15,000	
Teff grass	5	.1325	G	G		N		G	G	SM	G	F		WG	N		M	H	1M	
Triticale	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	M	M	15,000	
Turnips	4	.255	F	P	G	N	G	P	G	DH	G	G	F	CB	S	Р	L	N	175,000	
Vetch, Chickling	50	.5 - 1.5	F	F	F	Y	F	F	Р	SL	F	F	L	CB	N	Р	L	M	2,500	
Vetch, Common	25	.5 - 1.5	F	F	F	Y	F	F	G	SM	F	F	L	CB	N	Р	L	M	8,000	
Vetch, Hairy	15	.5 - 1.5	G	F	F	Y	F	F	F	SM	G	F	L	СВ	Y	Р	L	м	14,000	
Wheat, Spring	60	.5 - 1.5	G	G	G	N	G	G	G	MH	G	F	L	CG	N	G	М	M	15,000	
Wheat, Winter	60	.75 - 2.0	G	G	G	N	G	G	G	MH	G	F	L	CG	Y	G	М	М	15,000	F



Seeding Dates:

May 1 through August 5 - warm-season winter kill species

Early spring through August 20 – cool-season winter kill species

August 1 through Winter – species that do not winter kill

Seeding dates fluctuate annually. Seeding dates may be adjusted up to 15 days by the District Conservationist based on local weather and site conditions. Be cognizant of potential green bridge that could happen if cool-season grass cover crop species such as wheat, oats, barley, or rye are planted near commercially grown winter wheat fields.

Key to Rating Descriptions:

Rooting Depth/Water Use:	Ratings:	Crop Types:
SL = Shallow rooted/Low water use SM = Shallow rooted/Medium water use SH = Shallow rooted/High water use ML = Medium rooted/Low water use MH = Medium rooted/Medium water use DL = Deep rooted/Low water use DM = Deep rooted/Medium water use DH = Deep rooted/Medium water use	G = Good F = Fair P = Poor L = Low M = Medium H = High Y = Yes N = No S = Sporadic N/A = Not Applicable	CG = Cool season grass CB = Cool season broadleaf WG = Warm season grass WB = Warm season broadleaf
Shallow = 6-18 inches Medium = 18-24 inches Deep = 24+ inches		

For more information on cover crops, visit the South Dakota NRCS cover crop Web page: https://www.nrcs.usda.gov/conservation-basics/ conservation-by-state/south-dakota/cover-crops-in-south-dakota

South Dakota Natural Resources Conservation Service nrcs.usda.gov/