



CONSERVATION ENHANCEMENT ACTIVITY

E646136Z2

CONSERVATION STEWARDSHIP PROGRAM

Extend retention of captured rainfall to provide enhanced food sources for late winter habitat for migratory waterfowl and shorebirds

Conservation Practice 646: Shallow Water Development and Management

APPLICABLE LAND USE: Crop (Annual & Mixed)

RESOURCE CONCERN ADDRESSED: Fish & Wildlife – Inadequate Habitat

PRACTICE LIFE SPAN: 5 years

Enhancement Description:

When flooded to shallow depths during fall and winter, agricultural fields provide ideal foraging habitat for myriad species of waterfowl and wading birds. Harvested and idled agricultural lands, notably those occurring within rice rotations, contain high densities of residual (i.e., waste) grain and natural seeds following harvest. In addition, flooded conditions promote establishment of aquatic invertebrate populations, thus providing protein-rich food sources for shorebirds as well as waterfowl and wading birds. Benefits may become greatest during late winter and early spring as birds are assimilating nutrient and fat reserves in preparation for northward migration. However, agricultural fields flooded during fall-winter are typically drained during late January or February in advance of spring planting. This often results in a rapid reduction in available habitat, and may constrain ability of migratory birds to adequately prepare for migration, with greatest impacts likely occurring during years of low winter precipitation. Retention of water on agricultural lands into early spring will produce maximum benefits to migratory waterfowl and shorebirds by providing high quality habitat during a time when habitat may otherwise be in low abundance.

Criteria:



CONSERVATION STEWARDSHIP PROGRAM

This enhancement applies to crop land use acres with leveed fields capable of holding water at an average depth of 6 to 18 inches for the duration of the activity.

- Develop a wildlife habitat management plan for the targeted species suite.
- Water control structures affecting the subject land use are to be closed by mid-fall and remain closed until late winter to early spring.
 - Water depths of 6 to 10 inches provide maximum benefit to targeted species.
 - Water depths shall not exceed 18 inches for any extended period.
- A Wildlife Habitat Evaluation Guide (WHEG) specific to shallow water habitat on cropland must be used to show that implementation of the Enhancement will improve wildlife habitat value from fair (planning criteria = 0.5) to good (planning criteria greater than 0.5 and less than or equal to 0.6) or from good to very good (planning criteria greater than 0.6).

Note: This Enhancement may be grouped with E647136Z1-Manipulate vegetation on fields where rainfall is to be captured and retained. If not grouped with E647136Z1-Manipulate vegetation on fields where rainfall is to be captured and retained, this Enhancement may also be grouped with E646136Z3-Shorebird habitat, late season shallow water with manipulation or E646136Z4-Shorebird habitat, extended late season shallow water with manipulation.

Documentation Requirements:

- Copy of the wildlife habitat management plan for the targeted species suite (including the WHEG scores for present and planned conditions).
- Field log to include:
 - Crops grown and the harvest date for the crops grown on the applicable acres;
 - Date/time the water control structure was closed;
 - Date/time of each field visit and observed water levels;



- Date/time when the water control structures were opened.
- Digital photographs of the condition of the structures and the habitat provided.

CONSERVATION STEWARDSHIP PROGRAM

The enhancement is considered adopted when the water control structure affecting the land use acre remains closed beyond mid to late winter.

