

Rangeland Soil Quality Compaction

What is compaction?

Soil compaction occurs when moist or wet soil aggregates are pressed together and the pore space between them is reduced. Compaction changes soil structure, reduces the size and continuity of pores, and increases soil density (bulk density). Wheel traffic or pressure (weight per unit area) exerted on the soil surface by large animals, vehicles, and people can cause soil compaction. In areas of rangeland, compacted soil layers are generally at, or within 6 inches of the surface. They can be as deep as 2 feet under heavily used tracks and roads. Increases in density can be small to large.



Platy soil surface structure indicating management induced compaction

When is compaction a problem?

Compaction changes several structural characteristics and functions of the soil. It is a problem when the decreased pore space limits water infiltration, percolation, and storage; plant growth; or nutrient cycling.

Water movement and storage.—Compaction reduces the capacity of the soil to hold water and the rate of water movement through soil. It also limits water infiltration into the soil, causing increased runoff and, in some areas, increased erosion. Compacted wheel tracks or trails can concentrate runoff that can create rills or gullies, especially on steep slopes. When the amount of water that enters the soil is reduced, less water is available for plant growth and percolation to deep root zones.

Water entering the soil can perch on a subsurface compacted layer, saturating the soil to or near the surface or ponding on

the surface. An increase in the amount of water stored near the soil surface and a decrease in the amount of water deeper in the soil may favor the shallower rooted annuals over the deeper rooted plant species, such as shrubs.

Plant growth.—Where compaction increases significantly, it limits plant growth by physically restricting root growth. Severe compaction can limit roots to the upper soil layers, effectively cutting off access to the water and nutrients stored deeper in the soil. Anaerobic conditions (lack of oxygen) can develop in or above the compacted layer during wet periods, further limiting root growth. Even in arid climates, anaerobic conditions can occur where water accumulates.

Nutrient cycling.—Compaction alters soil moisture and temperature which control microbial activity in the soil and the release of nutrients to plants. The depth and pattern of root growth is also changed which affects the contributions of roots to soil organic matter and nutrients. Compaction compresses the soil, reducing the number of large pores. This reduction can restrict the habitat for the larger soil organisms that play a role in nutrient cycling thus reducing the number of these organisms. Where anaerobic conditions exist, loss of soil nitrogen through microbial activity increases.

How can compacted soil layers be identified?

The following features may indicate a compacted soil layer: platy, blocky, dense, or massive appearance;

- significant resistance to penetration with a metal rod;
- high soil bulk density; and
- restricted, flattened, turned, horizontal, or stubby plant roots.



Because some soils that are not compacted exhibit these features, refer to a soil survey report for information about the inherent characteristics of the soil. Each soil texture has a minimum bulk density (weight of soil divided by its volume) at which root-restricting conditions may occur, although the restriction also depends on the plant species.

Soil Texture	Root-restricting bulk density (g/cm ³)*
Coarse, medium, and fine sand and loamysand other than loamy very fine sand	1.80
Very fine sand, loamy very fine sand	1.77
Sandy loam	1.75
Loam, sandy clay loam	1.70
Clay loam	1.65
Sandy clay	1.60
Slit, slit loam	1.55
Slity clay loam	1.50
Slity clay	1.45
Clay	1.40

* Grams per cubic centimeter

What affects the ability of soil to resist compaction?

Moisture.—Dry soils are much more resistant to compaction than moist or wet soils. Soils that are wet for long periods, such as those on north-facing slopes and those on the lower parts of the landscape, where they receive runoff, are susceptible to compaction for longer periods than other soils. Saturated soils lose the strength to resist the deformation caused by trampling and wheeled traffic. They become fluid and turn into “mud” when compressed.

Texture.—Sandy loams, loams, and sandy clay loams are more easily compacted than other soils. Gravelly soils are less susceptible to compaction than nongravelly soils.

Soil structure.—Soils with well developed structure and high aggregate stability have greater strength to resist compression than other soils.

Plants and soil organic matter.—Near-surface roots, plant litter, and above-ground plant parts reduce the susceptibility to compaction by helping to cushion impacts. Vegetation also adds soil organic matter, which helps strengthen the soil, making it more resistant to compaction.

What breaks up a compacted layer?

Natural recovery is often slow, taking years to decades or more. Cycles of wetting, drying, shrinking and swelling can break down compacted layers, especially in clays and clay loams. Deep compaction occurs in smaller areas than shallow compaction, but it persists longer because it is less affected by the soil expansion caused by freezing. Shallow compaction may be very persistent in areas that are not subject to freezing and thawing.

Roots help to break up compacted layers by forcing their way between soil particles. Plants with large taproots are more effective at penetrating and loosening deep compacted layers, while shallow, fibrous root systems can break up compacted layers near the surface. Large soil organisms move soil particles as they burrow through the soil. Small mammals that tunnel through and mix the soil may also help loosen compacted layers.

Management strategies that minimize compaction

- Minimize concentrated grazing, recreational use, and vehicular traffic when the soils are wet.
- Use only designated trails or roads; reduce the number of trips.
- Maintain or increase the content of organic matter in the soil by improving the plant cover and plant production.

For more information, check the following: <http://soils.usda.gov/sqi> and <http://www.nrcs.usda.gov/wps/portal/nrcs/main/national/soils/health/> (Adapted from Rangeland Sheet #4, May 2001, developed by the Soil Quality Institute, Grazing Lands Technology Institute, and National Soil Survey Center, Natural Resources Conservation Service, USDA; the Jornada Experimental Range, Agricultural Research Service, USDA; and Bureau of Land Management, USDI)