

RUSLE2 Related Attributes

Worcester County, Massachusetts, Northwestern Part

[This report shows only the major soils in each map unit]

| Map symbol and soil name | Pct. of map unit | Hydrologic group | Kf | T factor | Representative value | | |
|--------------------------|------------------|------------------|-----|----------|----------------------|--------|--------|
| | | | | | % Sand | % Silt | % Clay |
| 1: Water | 100 | --- | --- | --- | --- | --- | --- |
| 19A: Lim | 80 | B/D | .32 | 3 | 64.5 | 25.5 | 10.0 |
| 28A: Searsport | 80 | A/D | --- | 1 | 5.0 | 0.0 | 0.0 |
| 29B: Naumburg | 80 | A/D | .17 | 5 | 71.1 | 21.4 | 7.5 |
| 31A: Walpole | 80 | B/D | --- | 4 | --- | --- | --- |
| 36A: Saco | 75 | B/D | .32 | 4 | 64.1 | 26.4 | 9.5 |
| 43A: Scarboro | 80 | A/D | --- | 2 | --- | --- | --- |
| 56A: Greenwood | 45 | A/D | --- | 2 | 8.0 | 0.0 | 0.0 |
| Chocorua | 35 | A/D | --- | 1 | 8.0 | 0.0 | 0.0 |
| 59A: Bucksport | 48 | B/D | --- | 2 | --- | --- | --- |
| Wonsqueak | 41 | B/D | --- | 1 | --- | --- | --- |
| 253A: Hinckley | 85 | A | --- | 3 | --- | --- | --- |
| 253B: Hinckley | 85 | A | --- | 3 | --- | --- | --- |
| 253C: Hinckley | 85 | A | --- | 3 | --- | --- | --- |
| 253D: Hinckley | 85 | A | --- | 3 | --- | --- | --- |
| 253E: Hinckley | 85 | A | --- | 3 | --- | --- | --- |

RUSLE2 Related Attributes

Worcester County, Massachusetts, Northwestern Part

| Map symbol and soil name | Pct. of map unit | Hydrologic group | Kf | T factor | Representative value | | |
|------------------------------|------------------|------------------|-----|----------|----------------------|--------|--------|
| | | | | | % Sand | % Silt | % Clay |
| 254A: Merrimac | 85 | A | .28 | 4 | 60.1 | 34.5 | 5.4 |
| 254B: Merrimac | 85 | A | .28 | 4 | 60.1 | 34.5 | 5.4 |
| 254C: Merrimac | 85 | A | .28 | 4 | 60.1 | 34.5 | 5.4 |
| 255B: Windsor, loamy sand | 85 | A | --- | 5 | --- | --- | --- |
| 255D: Windsor | 90 | A | --- | 5 | --- | --- | --- |
| 260A: Sudbury | 75 | B | .20 | 3 | 65.0 | 27.5 | 7.5 |
| 260B: Sudbury | 75 | B | .20 | 3 | 65.0 | 27.5 | 7.5 |
| 280A: Adams | 85 | A | .10 | 5 | 80.5 | 17.0 | 2.5 |
| 280B: Adams | 85 | A | .10 | 5 | 80.5 | 17.0 | 2.5 |
| 280C: Adams | 80 | A | .10 | 5 | 80.5 | 17.0 | 2.5 |
| 281B: Allagash | 85 | B | .28 | 3 | 70.1 | 21.9 | 8.0 |
| 281C: Allagash | 80 | B | .28 | 3 | 70.1 | 21.9 | 8.0 |
| 282A: Colton | 80 | A | .10 | 3 | 80.2 | 16.8 | 3.0 |
| 282B: Colton | 85 | A | .10 | 3 | 80.2 | 16.8 | 3.0 |
| 282C: Colton | 80 | A | .10 | 3 | 80.2 | 16.8 | 3.0 |
| 282D: Colton | 80 | A | .10 | 3 | 80.2 | 16.8 | 3.0 |

RUSLE2 Related Attributes

Worcester County, Massachusetts, Northwestern Part

| Map symbol and soil name | Pct. of map unit | Hydrologic group | Kf | T factor | Representative value | | |
|-----------------------------|------------------|------------------|-----|----------|----------------------|--------|--------|
| | | | | | % Sand | % Silt | % Clay |
| 282E: | | | | | | | |
| Colton | 75 | A | .10 | 3 | 80.2 | 16.8 | 3.0 |
| 284A: | | | | | | | |
| Croghan | 85 | A/D | .17 | 5 | 80.7 | 16.8 | 2.5 |
| 284B: | | | | | | | |
| Croghan | 85 | A/D | .17 | 5 | 80.7 | 16.8 | 2.5 |
| 305B: | | | | | | | |
| Paxton | 80 | C | .28 | 3 | 61.0 | 32.0 | 7.0 |
| 305C: | | | | | | | |
| Paxton | 85 | C | .28 | 3 | 61.0 | 32.0 | 7.0 |
| 307D: | | | | | | | |
| Paxton, extremely stony | 85 | C | --- | 3 | --- | --- | --- |
| 310B: | | | | | | | |
| Woodbridge, fine sandy loam | 82 | C/D | .28 | 3 | 61.0 | 32.0 | 7.0 |
| 315B: | | | | | | | |
| Scituate | 75 | C | .20 | 3 | 70.8 | 21.7 | 7.5 |
| 351B: | | | | | | | |
| Becket | 80 | C | .28 | 3 | 65.0 | 27.5 | 7.5 |
| 351C: | | | | | | | |
| Becket | 80 | C | .28 | 3 | 65.0 | 27.5 | 7.5 |
| 351D: | | | | | | | |
| Becket | 85 | C | .28 | 3 | 65.0 | 27.5 | 7.5 |
| 355B: | | | | | | | |
| Marlow | 85 | C | .28 | 3 | 57.0 | 38.0 | 5.0 |
| 355C: | | | | | | | |
| Marlow | 84 | C | .28 | 3 | 57.0 | 38.0 | 5.0 |
| 360B: | | | | | | | |
| Peru | 84 | C/D | .37 | 3 | 47.0 | 47.0 | 6.0 |
| 365B: | | | | | | | |
| Skerry | 85 | C | .24 | 3 | 65.0 | 27.5 | 7.5 |
| 420B: | | | | | | | |
| Canton | 80 | B | .24 | 3 | 61.2 | 31.5 | 7.3 |

RUSLE2 Related Attributes

Worcester County, Massachusetts, Northwestern Part

| Map symbol and soil name | Pct. of map unit | Hydrologic group | Kf | T factor | Representative value | | |
|-----------------------------|------------------|------------------|-----|----------|----------------------|--------|--------|
| | | | | | % Sand | % Silt | % Clay |
| 420C: | | | | | | | |
| Canton | 80 | B | .24 | 3 | 61.2 | 31.5 | 7.3 |
| 600: | | | | | | | |
| Pits, gravel | 96 | --- | .10 | --- | 91.5 | 7.0 | 1.5 |
| 651: | | | | | | | |
| Udorthents | 99 | --- | --- | --- | --- | --- | --- |
| 652: | | | | | | | |
| Udorthents | 95 | --- | --- | --- | --- | --- | --- |
| 900E: | | | | | | | |
| Becket | 40 | C | .28 | 3 | 65.0 | 27.5 | 7.5 |
| Monadnock | 30 | B | .24 | 4 | 64.6 | 27.9 | 7.5 |
| 901E: | | | | | | | |
| Berkshire, extremely stony | 55 | B | --- | 5 | --- | --- | --- |
| Marlow, extremely stony | 30 | C | --- | 3 | --- | --- | --- |
| 902E: | | | | | | | |
| Charlton, extremely stony | 45 | B | --- | 5 | --- | --- | --- |
| Paxton, extremely stony | 40 | C | --- | 3 | --- | --- | --- |
| 905C: | | | | | | | |
| Peru, extremely stony | 61 | C/D | --- | 3 | --- | --- | --- |
| Marlow, extremely stony | 20 | C | --- | 3 | --- | --- | --- |
| 908C: | | | | | | | |
| Becket | 40 | C | .28 | 3 | 65.0 | 27.5 | 7.5 |
| Skerry | 30 | C | .24 | 3 | 65.0 | 27.5 | 7.5 |
| 910C: | | | | | | | |
| Woodbridge, extremely stony | 55 | C/D | --- | 3 | --- | --- | --- |
| Paxton, extremely stony | 35 | C | --- | 3 | --- | --- | --- |
| 913C: | | | | | | | |
| Lyman | 35 | D | --- | 1 | 5.0 | 0.0 | 0.0 |
| Tunbridge | 25 | B | --- | 2 | 5.0 | 0.0 | 0.0 |
| Berkshire | 20 | A | --- | 5 | 5.0 | 0.0 | 0.0 |

RUSLE2 Related Attributes

Worcester County, Massachusetts, Northwestern Part

| Map symbol and soil name | Pct. of map unit | Hydrologic group | Kf | T factor | Representative value | | |
|---------------------------|------------------|------------------|-----|----------|----------------------|--------|--------|
| | | | | | % Sand | % Silt | % Clay |
| 913E: | | | | | | | |
| Lyman | 35 | D | --- | 1 | 5.0 | 0.0 | 0.0 |
| Tunbridge | 25 | B | --- | 2 | 5.0 | 0.0 | 0.0 |
| Berkshire | 15 | A | --- | 5 | 5.0 | 0.0 | 0.0 |
| 915E: | | | | | | | |
| Montauk, extremely stony | 45 | C | --- | 3 | --- | --- | --- |
| Canton, extremely stony | 40 | B | --- | 3 | --- | --- | --- |
| 917B: | | | | | | | |
| Pillsbury | 45 | C/D | .24 | 3 | 63.5 | 29.0 | 7.5 |
| Peacham | 35 | D | --- | 1 | 5.0 | 0.0 | 0.0 |
| 918B: | | | | | | | |
| Ridgebury | 50 | D | .24 | 2 | 71.1 | 21.4 | 7.5 |
| Whitman | 30 | D | .28 | 3 | 46.9 | 45.6 | 7.5 |
| 924C: | | | | | | | |
| Tunbridge | 25 | B | --- | 2 | 5.0 | 0.0 | 0.0 |
| Lyman | 20 | D | --- | 1 | 5.0 | 0.0 | 0.0 |
| Berkshire | 15 | A | --- | 5 | 5.0 | 0.0 | 0.0 |
| 925E: | | | | | | | |
| Charlton | 40 | A | .20 | 5 | 63.9 | 28.6 | 7.5 |
| Chatfield | 20 | B | .17 | 2 | 70.9 | 21.6 | 7.5 |
| Hollis | 15 | D | .28 | 1 | 71.1 | 21.4 | 7.5 |
| 926C: | | | | | | | |
| Charlton | 45 | A | .20 | 5 | 63.9 | 28.6 | 7.5 |
| Chatfield | 20 | B | .17 | 2 | 70.9 | 21.6 | 7.5 |
| 927C: | | | | | | | |
| Montauk, extremely stony | 35 | C | --- | 3 | --- | --- | --- |
| Scituate, extremely stony | 35 | C/D | --- | 3 | --- | --- | --- |
| Canton, extremely stony | 25 | B | --- | 3 | --- | --- | --- |