

HARDIN COUNTY, KENTUCKY

Highly Erodible Land Legend

Approved – WHC – 11/4/86

Frozen: 1/1/90

<u>SYMBOL</u>	<u>CLASS*</u>	<u>NAME</u>
AID	HEL	Allegheny-Lenberg-Caneyville complex, 12 to 20 percent slopes
As	NHEL	Ashton silt loam
CnD	HEL	Caneyville-Rock outcrop complex, 6 to 20 percent slopes
CnE	HEL	Caneyville-Rock outcrop complex, 20 to 30 percent slopes
CrB	NHEL	Crider silt loam, 2 to 6 percent slopes
CrC	HEL	Crider silt loam, 6 to 12 percent slopes
CrD	HEL	Crider silt loam, 12 to 20 percent slopes
CsC	HEL	Cumberland silt loam, 6 to 12 percent slopes
CsD	HEL	Cumberland silt loam, 12 to 20 percent slopes
CtC3	HEL	Cumberland silty clay loam, 6 to 12 percent slopes, severely eroded
CtD3	HEL	Cumberland silty clay loam, 12 to 20 percent slopes, severely eroded
Dn	NHEL	Dunning silty clay loam
EIB	NHEL	Elk silt loam, 2 to 6 percent slopes
EIC	PHEL	Elk silt loam, 6 to 12 percent slopes
FdC	HEL	Fredonie-Rock outcrop complex, 6 to 20 percent slopes
FrC	HEL	Frondorf-Lenberg silt loams, 6 to 12 percent slopes
FrD	HEL	Frondorf-Lenberg silt loams, 12 to 20 percent slopes
GmE	HEL	Garmon silt loam, 25 to 60 percent slopes
GnB	HEL	Gatton silt loam, 2 to 6 percent slopes
Gu	HEL	Gullied land
HnB	NHEL	Hagerstown silt loam, 2 to 6 percent slopes
HnC	HEL	Hagerstown silt loam, 6 to 12 percent slopes
HnD	HEL	Hagerstown silt loam, 12 to 20 percent slopes
Hu	NHEL	Huntington silt loam
Lc	NHEL	Lawrence silt loam
LfE	HEL	Lenberg-Frondorf complex, 20 to 30 percent slopes
Ln	NHEL	Lindside silt loam
MdC3	HEL	Markland silty clay, 6 to 12 percent slopes, severely

		eroded
Mr	NHEL	McGary silt loam
Mv	NHEL	Melvin silt loam
Nb	NHEL	Newark silt loam
NcA	NHEL	Nicholson silt loam, 0 to 2 percent slopes
NcB	HEL	Nicholson silt loam, 2 to 6 percent slopes
No	NHEL	Nolin silt loam
Nv	NHEL	Nolin variant fine sandy loam
OtA	NHEL	Otwell silt loam, 0 to 2 percent slopes
OtB	NHEL	Otwell silt loam, 2 to 6 percent slopes
PmB	NHEL	Pembroke silt loam, 2 to 6 percent slopes
PmC	HEL	Pembroke silt loam, 6 to 12 percent slopes
RaE	HEL	Ramsey-Steinsburg-Allegheny complex, 20 to 40 percent slopes
RbC	HEL	Riney loam, 6 to 12 percent slopes
RbD	HEL	Riney loam, 12 to 20 percent slopes
RbE	HEL	Riney loam, 20 to 30 percent slopes
RcD3	HEL	Riney sandy clay loam, 6 to 20 percent slopes, severely eroded
Rd	NHEL	Robertsville silt loam
RoE	HEL	Rock outcrop-Corydon complex, 12 to 30 percent slopes
SdA	NHEL	Sadler silt loam, 0 to 2 percent slopes
SdB	HEL	Sadler silt loam, 2 to 6 percent slopes
SdC	HEL	Sadler silt loam, 6 to 12 percent slopes
Sg	NHEL	Sensabaugh silt loam
SnB	HEL	Sonora silt loam, 2 to 6 percent slopes
SnC	HEL	Sonora silt loam, 6 to 12 percent slopes
SnC3	HEL	Sonora silt loam, 6 to 12 percent slopes, severely eroded
VrC	HEL	Vertrees silt loam, 6 to 12 percent slopes
VrD	HEL	Vertrees silt loam, 12 to 20 percent slopes
VrE	HEL	Vertrees silt loam, 20 to 30 percent slopes
VrD3	HEL	Vertrees silty clay loam, 6 to 20 percent slopes, severely eroded
WbC	HEL	Waynesboro loam, 6 to 12 percent slopes
WbD	HEL	Waynesboro silt loam, 12 to 20 percent slopes
WbE	HEL	Waynesboro loam, 20 to 30 percent slopes
WcC3	HEL	Waynesboro clay loam, 6 to 12 percent slopes,

WcD3	HEL	severely eroded Waynesboro clay loam, 12 to 20 percent slopes, severely eroded
WIB	HEL	Wellston silt loam, 2 to 6 percent slopes
WIC	HEL	Wellston silt loam, 6 to 12 percent slopes
WIC3	HEL	Wellston silt loam, 6 to 12 percent slopes, severely eroded

***CLASS**

- HEL = Highly Erodible Land
- NHEL = Not Highly Erodible Land
- PHEL = Potentially Highly Erodible Land