

MARION COUNTY, KENTUCKY

Highly Erodible Land Legend

Approved – WHC – 12/3/86

Frozen: 1/1/90

<u>SYMBOL</u>	<u>CLASS*</u>	<u>NAME</u>
BaB	HEL	Beasley silt loam, 2 to 6 percent slopes
BcC2	HEL	Beasley silty clay loam, 6 to 12 percent slopes, eroded
BeC3	HEL	Beasley silty clay, 6 to 12 percent slopes, severely eroded
BeD3	HEL	Beasley silty clay, 12 to 25 percent slopes, severely eroded
CaB	HEL	Carpenter gravelly silt loam, 2 to 6 percent slopes
CaC	HEL	Carpenter gravelly silt loam, 6 to 12 percent slopes
CaD	HEL	Carpenter gravelly silt loam, 12 to 20 percent slopes
CbF2	HEL	Carpenter-Lenberg complex, 20 to 45 percent slopes, eroded
CeB	HEL	Chenault gravelly silt loam, 2 to 8 percent slopes
ChC2	HEL	Christian silt loam, 6 to 12 percent slopes, eroded
ChD2	HEL	Christian silt loam, 12 to 20 percent slopes, eroded
CnD3	HEL	Christian clay loam, 12 to 20 percent slopes, severely eroded
CrB	HEL	Crider silt loam, 2 to 6 percent slopes
CrC2	HEL	Crider silt loam, 6 to 12 percent slopes, eroded
CyF2	HEL	Cynthiana-Faywood-Rock outcrop complex, 20 to 60 percent slopes, eroded
EdD2	HEL	Eden silty clay loam, 6 to 20 percent slopes, eroded
EeE2	HEL	Eden-Cynthiana complex, 20 to 35 percent slopes, eroded, rocky
EkB	HEL	Elk silt loam, 2 to 6 percent slopes
EkC2	HEL	Elk silt loam, 6 to 12 percent slopes, eroded
FaC2	HEL	Faywood silty clay loam, 6 to 12 percent slopes, eroded
FaD2	HEL	Faywood silty clay loam, 12 to 20 percent slopes, eroded
FcE2	HEL	Faywood-Cynthiana complex, 20 to 30 percent slopes, eroded, very rocky
FrC	HEL	Frankstown gravelly silt loam, 6 to 12 percent slopes
FrD	HEL	Frankstown gravelly silt loam, 12 to 20 percent slopes
GaF	HEL	Garmon channery silt loam, 30 to 80 percent slopes, very rocky
GrB	HEL	Greenbriar silt loam, 2 to 6 percent slopes
JeB	HEL	Jessietown-Trappist complex, 2 to 6 percent slopes
La	NHEL	Lawrence silt loam
LbC2	HEL	Lenberg silt loam, 6 to 12 percent slopes, eroded
LbD2	HEL	Lenberg silt loam, 12 to 20 percent slopes, eroded
LeE2	HEL	Lenberg silty clay loam, 20 to 30 percent slopes, eroded
LnB	HEL	Lowell silt loam, 2 to 6 percent slopes
LoC2	HEL	Lowell silty clay loam, 6 to 12 percent slopes, eroded
LoD2	HEL	Lowell silty clay loam, 12 to 20 percent slopes, eroded
LpC3	HEL	Lowell silty clay, 6 to 12 percent slopes, severely eroded

LpD3	HEL	Lowell silty clay, 12 to 20 percent slopes, severely eroded
LsC2	HEL	Lowell silty clay loam, shale substratum, 2 to 12 percent slopes, eroded
Me	NHEL	Melvin silt loam, occasionally flooded
Mn	NHEL	Melvin silt loam, frequently flooded
MoB	HEL	Mountview silt loam, 2 to 6 percent slopes
MoC2	HEL	Mountview silt loam, 6 to 12 percent slopes, eroded
Na	NHEL	Newark silt loam, occasionally flooded
Ne	NHEL	Newark silt loam, frequently flooded
NhB	HEL	Nicholson silt loam, 2 to 6 percent slopes
NhC2	HEL	Nicholson silt loam, 6 to 12 percent slopes, eroded
No	NHEL	Nolin silt loam, frequently flooded
OtB	HEL	Otwell silt loam, 2 to 8 percent slopes
RaC	HEL	Riney loam, 6 to 12 percent slopes
RcE2	HEL	Riney-Christian complex, 20 to 35 percent slopes, eroded
Ro	NHEL	Robertsville silt loam
RtD2	HEL	Rohan-Trappist complex, 6 to 20 percent slopes, eroded, very rocky
RtF2	HEL	Rohan-Trappist complex, 20 to 50 percent slopes, eroded, very rocky
SaB	HEL	Sandview silt loam, 2 to 6 percent slopes
SaC2	HEL	Sandview silt loam, 6 to 12 percent slopes, eroded
Se	NHEL	Sensabaugh gravelly silt loam, frequently flooded
SrE2	HEL	Shrouds-Brassfield complex, 12 to 30 percent slopes, eroded, very rocky
TbA	NHEL	Tilsit-Berea silt loams, 0 to 2 percent slopes
TbB	HEL	Tilsit-Berea silt loams, 2 to 6 percent slopes
TbC2	HEL	Tilsit-Berea silt loams, 6 to 12 percent slopes, eroded
TeC2	HEL	Trappist-Jessietown complex, 6 to 12 percent slopes, eroded
TrD2	HEL	Trappist-Rohan-Greenbriar complex, 12 to 20 percent slopes, eroded, rocky
TrD3	HEL	Trappist-Rohan-Greenbriar complex, 12 to 25 percent slopes, severely eroded, rocky

\*CLASS

- HEL = Highly Erodible Land
- NHEL = Not Highly Erodible Land
- PHEL = Potentially Highly Erodible Land