



United States Department of Agriculture

CRP Practice CP4B & CP4D

Natural Resources Conservation Service - Indiana – March 2016 (ver. 1.3)

Permanent Wildlife Habitat Program Job Sheet



PURPOSE

Permanent wildlife habitat plantings are used to reduce soil erosion, improve water quality, and create or enhance wildlife habitat by providing food and cover.

WHERE PRACTICE APPLIES

Apply this practice on fields that meet eligibility requirements for the Conservation Reserve Program (CRP) as determined by the Farm Service Agency (FSA).

PLANNING

A wildlife conservation plan will be developed in consultation with the Indiana Department of Natural Resources (IDNR) District Wildlife Biologist. When selecting plant species, consider the requirements for optimum wildlife habitat, and adaptability to the soils, and moisture conditions in the field being established.

CRP POLICY

General Signup: To award 50 points for National Ranking Factor N1a, existing vegetation or seeding mixes must have at least five (5) **native** grasses, forbs, legumes, plus 20% trees and/or shrubs planted in blocks or strips. To award 40 points, existing vegetation or seeding mixes must have at least four (4) native or introduced species of grasses, forbs, legumes, plus 20% trees and/or shrubs planted in blocks or strips. The

selected species will be planted in mixes, blocks or strips best suited for various wildlife species in the area.

Conservation Reserve Enhancement Program (CP4D only): Seeding mixes must have at least five (5) **native** grasses, forbs, legumes, trees and shrubs, 20% of which must be planted to trees and/or shrubs.

CREP Permanent Wildlife Habitat are only eligible as buffers on cropland that is adjacent and parallel to streams, sinkholes and karst areas, wetlands, and permanent bodies of water such as lakes/ponds.

The width of CREP Permanent Wildlife Habitat will be a minimum of 35 feet, and a maximum of 180 feet, from the edge of the eligible body of water.

NOTE: An average maximum width of 300 feet is allowed when area to be enrolled involves a predominance of alluvial soils. If the site already contains existing vegetation, these acres will be included in the calculation of maximum width and included in the CREP Plan, but will not be eligible for payments.

Highly Erodible Land Initiative (CP4D only) and Source Water Protection Program: See 40-point criteria.

For CP4B (corridors), the minimum width is 66 feet, with a maximum width of 200 feet. CP4D does not have any size requirements.

SEEDING RATES AND SPECIES

The [IN Natural Resources Conservation Service \(NRCS\) Seeding Tool](#) and/or tables found in the Field Office Technical Guide (FOTG) [Standards Conservation Cover \(327\)](#) - (primary purpose wildlife) and Tree/Shrub Establishment (612) will be used when developing seeding mixes and to determine tree and shrub species for this practice. Commonly used, suitable species can also be found in the [IN Biology Technical Note - Upland Wildlife Habitat](#).

Any prepackaged mixes must be approved before seeding. Site-specific requirements are listed on the attached Specifications Sheet.

GRASS PLANTINGS

Companion/Nurse Crops

A companion/nurse crop will be used when erosion control and weed suppression are needed.

Companion/nurse crops include Winter Wheat (after the Hessian Fly-free dates in Table 2), Oats, Barley, Cereal Rye or Annual Ryegrass; native Wildryes (i.e. – *Elymus* sp. such as Canada, Riverbank, and Virginia Wildrye) and other species are also effective.

Companion crops will be clipped after jointing, but before seed head pollination unless otherwise directed (control of Wildrye species is not necessary so that they persist as part of native seedings). A second and subsequent clipping is necessary if re-growth provides competition. Clipping height should be above developing seedlings. Where excessive growth has accumulated, the vegetation will be chopped rather than swathed.

Lime and fertilizer

Lime and fertilizer should be based on a current soil test (less than four years old). In areas with existing vegetation that shows signs of nutrient deficiencies, or if the soil test shows phosphorus (P) and potassium (K) are in the low to very low range, apply enough fertilizer (organic or inorganic) to raise N, P and K to a level needed for a 1 ton/ac yield goal. Do not apply any nitrogen (N) for warm season grasses. Use Purdue University recommendations from the *Crop Fertilizer Recommendation Calculator*, or the Indiana NRCS Seeding Tool – *Indiana Fertilizer Calculator*.

If the pH is 6.0 or less, apply enough lime per acre to bring pH to meet the tolerance range of the planned plant species. Soil amendments will be incorporated during seedbed preparation, or applied before planting if a no-till drill is used. Apply lime according to *Tri-State Fertilizer Recommendations - PU AY-9-32*, Extension Bulletin E-2567, or Indiana NRCS Seeding Tool.

Site Preparation

It is very important to plant the vegetation into a weed-free seedbed. Use herbicides and/or tillage to eliminate competing vegetation. Weed control efforts should begin as early as 12 months prior to planting, and may require multiple applications or operations in both the fall and spring prior to planting.

Pay particular attention to sites where noxious and potentially invasive species are likely. Many of these species are perennials that spread through seed and

roots, and many have rhizomatous root systems that will persist and negatively impact the planting.

Cool season weeds (i.e. Canada thistle, quack grass) are best controlled in the fall (mid-September to early November) with a translocation herbicide. Plants should be actively growing at the time of application. Avoid herbicide application after 3:00 pm if overnight temperatures are expected to drop below 50 degrees(F).

Warm season weeds (i.e. Johnsongrass) are best controlled prior to flower with a follow-up application prior to first frost. Plants should be actively growing at the time of application. Contact your local Purdue University Cooperative Extension Service for specific herbicides to use. **Apply all herbicides according to the label.**

Use a nurse/companion crop to control potential weed issues and/or a temporary cover for erosion control.

If prescribed burning is used for site preparation, it must be conducted according to IN NRCS FOTG Standard [*Prescribed Burning \(338\)*](#).

Seeding Dates

Selected species will be planted within the dates in the specification sheet that will be provided for the site.

Seed preparation

Inoculate legume seed before seeding with the proper rhizobia bacteria specific for the species. Re-inoculate seed if it was pre-inoculated more than 60 days prior to seeding or beyond dates specified on the seed / inoculant tag. Inoculant left in the sun, even for a short period of time can significantly reduce the viability and effectiveness. Pre-inoculated seed will have a coating that changes the pure live seed per pound and thus the bulk seeding rate per acre.

Be aware that blending seed of varying size, shape and weight can make calibration of equipment and seeding uniformity difficult.

Some seeding mixtures contain seed that is extremely small and thus have very low seeding rates. This may make it difficult to set seeding equipment to uniformly seed these low rates. To add enough volume to the mix for proper metering, a **carrier** or coated seed may be desirable. The carrier should be no larger than the largest seed species and have similar shape, density and texture to the majority of seeds in the mix. The carrier can be an inert material (i.e. cracked corn) that does not have abrasive properties that may cause damage to the equipment or the seed. Inexpensive seed (unimproved

varieties) that will have no significant negative impact on the purpose of the seeding may also be used.

Planting Methods

No-Till seeding: Use a no-till drill with seven (7) inch or less row spacing. Ensure the drill is designed to handle the type of seed being planted (especially important for native grasses). Set the no-till drill to provide good seed-to-soil contact and a planting depth preferred for the desired species (see table below). Soils that are too wet or too dry can also cause improper seed placement.

Conventional Seeding: Prepare a fine firm seedbed to a depth of three (3) to four (4) inches. Incorporate lime and fertilizer during seedbed preparation. Use a drill with seven (7) inch or less row spacing or a culti-packer seeder designed for the seed to be planted. Seed should be drilled uniformly at a proper seeding depth for the desired species.

Seeding Depth Guidance

Groups	Seed Size (seeds/lb.)	Optimum (inches)	Max. (in.)
Brassicas, clovers, small seeded legumes & grasses, native forbs	150,000 – 500,000	1/4	1/2
Vetches, sorghums, wildryes, trefoils, native legumes, radishes	50,000 – 150,000	1/2	3/4
Cereal grains	12,000 – 50,000	3/4	1
Beans, peas, corn	1500 – 12,000	1 1/2	2

Broadcast Seeding: Seed may be broadcast if completed in a uniform manner. Pre-mix the seed with 200 pounds per acre of pelletized lime if using an airflow applicator. Seedbeds should be worked to a minimum depth of three (3) inches and firmed before seeding. The seedbed should be culti-packed before and after seeding. It is acceptable to see up to one-third (1/3) of the seed on the soil surface. Wind speed should be 15 miles per hour or less when broadcasting.

Inter-seeding:

- Legumes/Forbs (frost seeding):** No-till drill or broadcast as above into existing vegetation or residues. Broadcasting relies on freeze/thaw cycles, rain and/or snow to incorporate the seed. This method does not include a seedbed preparation. This is most commonly used during the dormant seeding period.
- Cover Crops:** No-till drill or broadcast as above into existing vegetation or residues. Broadcasting relies on freeze/thaw cycles, rain and/or snow to incorporate the seed. Inter-seeding does not include a seedbed preparation. This method can be used to establish cover crop species or combination mixes into relatively light (e.g. soybean) and weed free

crop residues, or to establish vegetation into standing crops.

- Grasses:** No-till drill into existing covers only if prior-treated with herbicides or tillage, or if existing cover is diminishing (i.e. – older alfalfa plantings).

Weed Control during Establishment

Control competing vegetation as needed until a Final Status Review is issued or three (3) years (whichever comes first). Mow, burn, or apply herbicides as needed to control unwanted vegetation for up to three (3) years after planting. Mow when competing weeds are taller than the planted vegetation, and at a height above the planted vegetation. Use selective herbicides and/or spot spraying to protect the desired species. Refer to the Purdue/Ohio State [Weed Control Guide for Ohio and Indiana](#) for herbicide timing and treatment.

TREES AND SHRUBS

Weed Control

It is important to plant tree seedlings into a weed free area to help ensure survival and maximum growth of the trees. Use herbicides or tillage to eliminate competing vegetation as recommended in a site specific plan. Weed control efforts should begin prior to planting and may require multiple applications in both the fall and the spring prior to planting and during tree planting. Cost share for weed control is available as part of the tree establishment expenses. Mowing is usually not recommended for weed control because it encourages grass competition and mowing equipment can damage tree seedlings. However in some specific instances it may be recommended in a site specific plan.

For information on herbicide controls, contact a local consultant or Purdue Extension Specialist. Always apply herbicides according to labeled directions.

If necessary for weed suppression or erosion control, seed a temporary cover of native *Elymus* species (Wildrye) adapted to the site.

Cost share is authorized for an additional weed control application within 24 months after planting.

Operation and Maintenance

After the Final Status Review or three (3) years (whichever comes first), maintain the planting according to your CRP conservation plan. Maintenance activities are allowed only if necessary to maintain stand health, or to control pests, noxious weeds or any plant species whose presence or overpopulation may jeopardize the CRP cover, or have detrimental effects to the surrounding land.

The presence of annual weeds (such as foxtail, common ragweed, and perennial forbs) is not a concern, as these plants are important sources of food for wildlife, especially bobwhite quail. Maintenance may be needed to control excessive density of these annuals, especially during the establishment years, but is not intended to eliminate this group of plants.

Maintenance activities will not occur from **April 1 through August 1** to protect ground-nesting wildlife. If maintenance activities are needed during the April 1 – August 1 time frame, the FSA County Committee **must** approve the maintenance activity **prior to** the activity occurring, and it may **only be on a spot basis**.

Mowing for generic weed control or for cosmetic purposes is prohibited.

Native grasses will not be mowed lower than eight (8) to 12 inches.

Inspect the vegetation annually and after storm events, and repair any gullies that have formed; remove unevenly deposited sediment and/or crop residues that will disrupt the function or kill desired vegetation; and reseed high mortality and disturbed areas.

The contract area cannot be used for field roads or other uses that will damage or destroy the cover.

Apply supplemental nutrients as needed to maintain the desired species composition and stand density. Check survivability of planted species after 3 years to insure that the desired stocking rate for the site is present, usually 70% survival of the planted rate. Additional planting will be completed if it is determined that additional natural regeneration will not be sufficient to colonize the site within an acceptable time frame (usually 5 years) so that 300 acceptable woody plants per acre are established.

MID-CONTRACT MANAGEMENT

Mid-Contract Management (MCM) is required on this practice. If the CRP acres are less than 5 acres, the entire acreage can be managed in a single year; otherwise, the maximum amount that can be disturbed during any one year is ½ of the contract acreage. For maximum habitat value, disturb no more than 1/3 of the contract acreage in any given year.

MCM activities will be avoided on environmentally sensitive areas including:

- a) Concentrated flow areas,
- b) Critical areas,
- c) Within the first 20 feet of a practice that borders a water resource to avoid water quality resource concerns, and

- d) Other areas where gully erosion is likely.

Environmentally sensitive areas will be marked on the plan map to ensure Mid-Contract Management activities are avoided on these areas.

Areas devoted to grass have the following options:

- Prescribed Burning
- Strip Disking
- Strip Spraying
- Inter-seeding forbs/legumes/pollinator habitat

Areas devoted to trees have the following options:

- Inner Seedling Planting (re-enrollments)
- Inter-seeding forbs/legumes/pollinator habitat
- Follow-up Weed Control
- Pruning
- Thinning (re-enrollments)

MCM activities operations will not be performed from April 1 through August 1 for contracts starting in 2008, to protect the primary nesting period for grassland bird species. It is also recommended, but is not required, to delay MCM activities until after August 15 to reduce the chance of harming fledgling birds and other young wildlife.

MCM activities operations will be performed along field contours, or across the slope, when practical.

Strips will parallel brushy or woody escape cover when feasible.

Managed Haying and Grazing

Only CP4D plantings may be used for managed haying or grazing. CP4B (corridors) is not eligible for managed haying and grazing.

The managed haying and grazing period for CRP is August 2 through September 15. All hay and animals must be removed from CRP acreage no later than 10 days after the end of the managed haying and grazing period.

Managed haying and grazing activities must be performed according to NRCS Standards and Specifications as found in the FOTG and CRP policy. The same acreage may not be hayed or grazed more than once every three (3) years. Annual CRP rental payments will be reduced based on FSA policy. All managed haying and grazing activities must be approved by FSA and included in the conservation plan prior to harvesting the forage.