


TAMA -- IOWA STATE SOIL



Tama Soil Profile

Surface layer: very dark brown silty clay loam
Subsurface layer: very dark grayish brown silty clay loam
Subsoil - upper: brown silty clay loam
Subsoil - lower: dark yellowish brown silty clay loam
Substratum: yellowish brown silty clay loam

The Tama series is considered one of the most productive of the soils in Iowa that are used for agricultural purposes. It makes up about 825,000 acres in east-central and eastern Iowa. The series was first identified in Black Hawk County, Iowa, in 1917. It has been identified in 26 counties in Iowa. It also has been identified in Illinois, Minnesota, and Wisconsin.

Tama soils formed in 48 or more inches of silty loess; under tall prairie grasses with a deep, fibrous root system; and under relatively humid climatic conditions. Over hundreds of years, the grasses have added organic matter to the soils, producing a relatively thick, dark surface layer. In some areas, erosion has significantly affected the properties of the soils. Eroded Tama soils have less total nitrogen and organic matter and more clay in the surface layer than uneroded Tama soils.



The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). *USDA is an equal opportunity provider and employer.*

USDA NRCS
Natural Resources Conservation Service