


# 2013-17 Cumulative Sediment Load Reductions


## 1,372,892 Tons



Since 2013, voluntary conservation efforts from Indiana's private landowners, with support from the ICP, have reduced sediment and nutrients from entering Indiana's waterways.



**1,372,892 tons of sediment.**  
**A football field covered to a depth of 596 feet, which is almost as tall as the Space Needle.**



### Sediment (tons)



Based on EPA Region 5 Model analyses conducted on 21,957 conservation practices installed by the Indiana Conservation Partnership January 2013 thru December 2017. This effort does not include the many unassisted practices designed and installed solely by a private landowner without ICP assistance.

The cumulative analysis encompassed a breakdown of 2013 thru 2017 conservation practices by lifespan including 1, 5, 10, 15, 20 and 40 years. The map reflects all of the practices minus the 2013 thru 2016 practices with a lifespan of one year and 2013 practices with a lifespan of 5 years.

To learn more about Indiana's Nutrient Reduction Strategy visit <http://www.in.gov/isda/2991.htm>  
 For questions and comments email [ISDANutrientReduction@isda.in.gov](mailto:ISDANutrientReduction@isda.in.gov)

March 1, 2018  
 Deb Fairhurst, ISDA Program Manager  
 Trevor Laureys, ISDA Program Manager