Approved
State Soil Scientist
5/20/09

TECH' GUIDE
'ION II
CROPLAND INTE. ATION

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
14A	Nuley Variant loam	N	Y	Y	Y	Y	Y	Y	N	N
14C	Nuley Variant loam	N	Y	Y	Y	Y	Y	Y	N	N
14D	Nuley Variant loam	N	Y	Y	Y	Y	Y	Y	N	N
18A	Comer Family Gravelly coarse sandy loam	N	Y	Y	Y	Y	Y	Y	N	N
20B	Anaconda Family-Comer Family complex	N	N	N	N	N	N	Y	N	N
24A	Beaverell cobbly loam	N	Y	Y	Y	Y	Y	Y	N	N
24D	Beaverell cobbly loam	N	Y	Y	Y	Y	Y	Y	N	N
26A	Gledive Variant sandy loam	N	N	Y	Y	Y	Y	Y	N	N
27A	Attewan Variant loam	N	N	Y	Y	Y	Y	Y	N	N
28A	Fubartoo silt loam	Y	Y	Y	Y	Y	Y	Y	N	N
29C	Dinnen coarse sandy loam	N	Y	Y	Y	Y	Y	Y	N	N
29D	Dinnen coarse sandy loam	N	Y	Y	Y	Y	Y	Y	N	N
32E	Brocko family cobbly silt loam	N	N	N	N	N	Y	Y	N	Y
37C	Brownsto gravelly loam	N	N	Y	Y	Y	Y	Y	N	N
47E	Supervisor family-Merino-Rock outcrop complex	N	Y	Y	Y	Y	Y	Y	N	Y
49C	Libeg gravelly loam	N	N	Y	Y	Y	Y	Y	N	N
50A	Bonebasin, occasionally flooded-Meadowcreek, rarely flooded complex, 0 to 4 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
50C	Buffork loam	N	N	Y	Y	Y	Y	Y	N	N
54C	Little Horn family cobbly silt loam	N	N	Y	Y	Y	Y	Y	N	N
54D	Little Horn family cobbly silt loam	N	N	Y	Y	Y	Y	Y	N	N
111E	Sebud-Philipsburg complex, 8 to 25 percent slopes, bouldery	N	N	N	N	N	N	Y	N	N
112C	Varney Varient-Radersburg Variant complex	N	N	Y	Y	Y	Y	Y	N	N
141E	Nuley Variant very cobbly loam	N	Y	Y	Y	Y	Y	Y	N	Y
143C	Nuley Variant stony loam	N	Y	Y	Y	Y	Y	Y	N	N
291E	Dinnen-Branham-Rock outcrop complex	Y	Y	Y	Y	Y	Y	Y	N	Y
301D	Hauz-Tolman complex	Y	Y	Y	Y	Y	Y	Y	N	Y
301E	Philipsburg-Beeftrail, bouldery-Danielvil complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	Y
371E	Brownsto cobbly loam	N	N	Y	Y	Y	Y	Y	N	Y
501D	Buffork-Sawcreek complex	N	N	Y	Y	Y	Y	Y	N	N
521C	Whitecap-Sawcreek complex	Y	Y	Y	Y	Y	Y	Y	N	N
10A	Slickens-Aquic Cumulic Haplustolls, occasionally flooded-Aridic Ustifluvents, occasionally flooded complex, 0 to 2 percent slopes, severely impacted	N	N	Y	Y	Y	Y	Y	N	N
11A	Mannixlee-Bonebasin complex, 0 to 4 percent slopes, frequently flooded	N	N	Y	Y	Y	Y	Y	N	N
12A	Riverrun, occasionally flooded-Mannixlee, frequently flooded complex, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
15B	Anamac-Varney-Rivra, rarely flooded complex, 0 to 6 percent slopes	N	N	N	N	Y	Y	Y	N	N
21C	Kilgore, occasionally flooded-Foolhen, rarely flooded-Philipsburg complex, 2 to 12 percent slopes	N	N	N	Y	Y	Y	Y	N	N
22B	Philipsburg loam, 1 to 4 percent slopes, stony	N	N	N	N	N	Y	Y	N	N
24B	Kilgore, rarely flooded-Mooseflat, occasionally flooded-Philipsburg complex, 0 to 6 percent slopes	N	N	N	Y	Y	Y	Y	N	N
25D	Work, stony-Adel-Kilgore, rarely flooded complex, 2 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
26C	Mooseflat, occasionally flooded-Bridger-Kilgore, rarely flooded complex, 0 to 8 percent slopes	N	N	N	N	N	N	N	N	N
27D	Savenac-Bridger complex, 4 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
28C	Kilgore-Danielvil complex, 2 to 8 percent slopes	N	N	N	Y	Y	Y	Y	N	N
31A	Philipsburg cobbly loam, 0 to 4 percent slopes, very stony	N	N	N	Y	Y	Y	Y	N	N
32B	Kilgore, occasionally flooded-Danielvil complex, 1 to 4 percent slopes	N	N	N	N	N	N	N	N	N
33B	Sebud very cobbly loam, 2 to 6 percent slopes, very stony	N	N	N	N	N	N	N	N	N
34A	Danielvil-Danielvil, rarely flooded complex, 0 to 4 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
36A	Foolhen-Monaberg-Mooseflat complex, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
37B	Mooseflat, occasionally flooded-Monaberg, rarely flooded complex, 1 to 4 percent slopes	N	N	N	N	N	N	N	N	N
38B	Rivra gravelly sandy loam, 0 to 4 percent slopes, moderately impacted	Y	Y	Y	Y	Y	Y	Y	N	N
40B	Cetrack loam, 0 to 4 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
42D	Varney-Varney, stony-Anamac, stony complex, 2 to 12 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
43B	Meadowcreek-Mannixlee, rarely flooded complex, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
44B	Meadowcreek-Anamac complex, 1 to 6 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
46A	Varney-Rivra complex, 0 to 4 percent slopes	N	N	N	Y	Y	Y	Y	N	N
47B	Foolhen-Monaberg-Kilgore, occasionally flooded complex, 0 to 6 percent slopes	N	N	N	N	N	Y	Y	N	N
51D	Foxgulch-Libeg complex, 6 to 25 percent slopes, stony	N	N	N	N	Y	Y	Y	N	N
53C	Sixbeacon-Sieben complex, 2 to 8 percent slopes	N	N	N	N	N	Y	Y	N	N
55A	Bonebasin, occasionally flooded-Monaberg, very rarely flooded-Meadowcreek, rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
56B	Bonebasin, occasionally flooded-Meadowcreek, rarely flooded complex, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
58B	Varney-Anaconda loams, 0 to 4 percent slopes, moderately impacted	N	N	N	N	N	N	Y	N	N
59A	Anamac-Meadowcreek complex, 0 to 2 percent slopes	N	N	N	N	N	N	Y	N	N
60A	Riverrun-Rivra complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
61A	Rivra, stony-Riverrun complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
62A	Vendome-Sieben complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
64A	Work clay loam, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
65B	Sebud-Bearmouth complex, 1 to 4 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	N	N
67A	Foxgulch-Bearmouth, very stony complex, 0 to 4 percent slopes	N	N	N	N	Y	Y	Y	N	N
68B	Bearmouth, rarely flooded-Foxgulch, occasionally flooded complex, 0 to 4 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	N	N
69B	Foxgulch loam, 1 to 4 percent slopes	N	N	N	Y	Y	Y	Y	N	N
71D	Sebud-Ratiopeak complex, 4 to 15 percent slopes, stony	N	N	N	N	N	N	Y	N	N
72B	Tibson gravelly loam, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
74A	Bearmouth very cobbly sandy loam, 0 to 2 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
75A	Danielvil loam, 0 to 2 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
76A	Bearmouth very cobbly sandy loam, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
80A	Water-Riverwash complex	0	0	0	0	0	0	0	0	0
81A	Varney sandy loam, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
82D	Philipsburg-Monaberg-Kilgore, occasionally flooded complex, 0 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
82E	Elve gravelly loam, 15 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
85E	Loberg gravelly loam, 15 to 35 percent slopes	N	N	N	N	N	N	Y	N	N
101B	Philipsburg sandy loam, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
102D	Varney, noncalcic-Philipsburg-Varney complex, 2 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
106E	Gnojek-Libeg-Philipsburg complex, 8 to 30 percent slopes, stony	Y	Y	Y	Y	Y	Y	Y	Y	Y
107E	Anaconda-Varney-Work, stony complex, 6 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
108D	Varney-Work complex, 4 to 15 percent slopes, stony	N	N	N	N	N	N	Y	N	N
109A	Bearmouth coarse sandy loam, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
110E	Danielvil-Bearmouth complex, 2 to 25 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
112E	Monaberg-Bridger-Libeg, stony complex, 8 to 25 percent slopes	N	N	N	N	N	N	Y	N	N
114B	Varney loam, 0 to 4 percent slopes, moderately impacted	N	N	N	N	N	N	Y	N	N
115D	Philipsburg-Ratiopeak complex, 8 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
116C	Monaberg-Philipsburg complex, 2 to 12 percent slopes	N	N	N	Y	Y	Y	Y	N	N
118E	Reedpoint-Rock outcrop-Sixbeacon complex, 6 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
119C	Varney-Udecide complex, 2 to 12 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
120C	Varney-Anamac complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
122C	Varney-Work complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
123E	Sebud, stony-Danielvil-Monaberg, very stony complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
124E	Monaberg-Bridger complex, 8 to 30 percent slopes, very stony	N	N	N	N	N	Y	Y	N	N
125E	Libeg-Monaberg complex, 15 to 35 percent slopes, very bouldery	N	N	N	N	N	N	Y	N	N
126E	Danielvil, very bouldery-Monaberg, bouldery-Adel complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
128D	Monaberg loam, 4 to 15 percent slopes, bouldery	N	N	N	N	N	Y	Y	N	N
129D	Philipsburg-Danielvil complex, 4 to 15 percent slopes	N	N	N	Y	Y	Y	Y	N	N
131E	Dumps, placer mining-Philipsburg, stony complex, 4 to 15 percent slopes	N	N	N	Y	Y	Y	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
136D	Varney-Varney, stony-Anamac complex, 2 to 12 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
139D	Beeftrail, occasionally flooded-Adel-Monaberg complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
140D	Monaberg-Adel complex, 4 to 20 percent slopes	N	N	N	N	N	N	Y	N	N
141F	Danielvil-Danielvil, cool-Philipsburg complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
142E	Ratiopeak-Philipsburg complex, 15 to 35 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
145D	Danielvil-Philipsburg complex, 4 to 12 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
147B	Philipsburg-Ratiopeak complex, 1 to 6 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
148C	Philipsburg-Monaberg complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
149D	Bridger-Libeg complex, 8 to 25 percent slopes, very stony	N	N	N	N	N	N	N	N	N
150D	Sebud, very stony-Ratiopeak, stony-Bridger, stony complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
152E	Sieben-Varney complex, 15 to 45 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
153D	Sebud, very stony-Philipsburg-Ratiopeak, very stony complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
158C	Varney-Anamac-Anaconda complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
159E	Monaberg-Varney-Philipsburg complex, 8 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
160E	Bridger-Patouza, stony-Varney complex, 8 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
161C	Philipsburg-Philipsburg, stony-Danielvil complex, 2 to 8 percent slopes	N	N	N	Y	Y	Y	Y	N	N
162B	Monaberg loam, 1 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
164B	Danielvil loam, moist, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
165A	Mooseflat, rarely flooded-Foxgulch, very rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	N	N	N
166B	Kilgore-Foolhen-Monaberg complex, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
167C	Monaberg loam, 2 to 8 percent slopes, stony	N	N	N	N	N	N	Y	N	N
168E	Sebud-Sieben complex, 8 to 30 percent slopes, stony	N	N	N	N	N	N	Y	N	N
169E	Sebud-Sieben complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
170E	Ratiopeak, very stony-Bridger-Adel complex, 8 to 30 percent slopes	N	N	N	N	Y	Y	Y	N	N
171D	Varney-Sieben, stony complex, 4 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
172F	Ratiopeak-Sixbeacon-Tiban complex, 15 to 45 percent slopes, extremely stony	N	N	N	N	Y	Y	Y	N	N
173B	Monaberg-Foolhen complex, 1 to 6 percent slopes	N	N	N	N	N	N	Y	N	N
174C	Philipsburg loam, 2 to 8 percent slopes	N	N	N	N	N	Y	Y	N	N
175B	Foolhen-Kilgore complex, 0 to 4 percent slopes	N	N	N	Y	Y	Y	Y	N	N
177C	Udecide-Gnojek, stony complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
178E	Rootel-Varney complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
179F	Vendome, extremely stony-Sixbeacon complex, 25 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
182C	Varney, stony-Anamac, very stony-Rivra, extremely stony complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
183B	Anamac-Varney complex, 1 to 6 percent slopes	N	N	N	N	N	N	Y	N	N
235B	Anaconda sandy loam, 0 to 4 percent slopes, severely impacted	N	N	Y	Y	Y	Y	Y	N	N
272F	Rencot, stony-Spudbar-Rock outcrop complex, 25 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
303D	Moosejaw, occasionally flooded-Highrye-Silas, occasionally flooded complex, 2 to 12 percent slopes	N	N	N	N	N	N	Y	N	N
305D	Beeftrail-Branham-Minestope complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
306E	Wissikihon-Branham-Highrye complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
312D	Oro Fino-Highrye-Branham complex, 4 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
313E	Beeftrail-Dinnen-Highrye complex, 8 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
314F	Basincreek-Comad complex, 20 to 50 percent slopes	N	N	N	Y	Y	Y	Y	Y	Y
315F	Stecum-Hiore complex, 20 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
316F	Stecum, very bouldery-Rock outcrop-Zonite, very bouldery complex, 20 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
317E	Stecum-Caseypeak-Rock outcrop complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
319D	Silas, stony-Branham, stony-Tepete complex, 2 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
321E	Beeftrail-Bavdark-Zonite complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
323E	Highrye-Tuggle, very stony-Moosejaw complex, 2 to 25 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
324E	Beeftrail, stony-Branham, stony-Fleecer complex, 8 to 20 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
326C	Fleecer-Branham-Passmore complex, 2 to 8 percent slopes	N	N	N	Y	Y	Y	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
		Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
mu_sym	muname									
327E	Highrye-Stecum-Wissikihon complex, 15 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
328E	Stecum-Zonite-Basincreek complex, 8 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
331C	Mooseflat, occasionally flooded-Foolhen, occasionally flooded-Fleecer complex, 2 to 8 percent slopes	N	N	N	N	N	N	N	N	N
332D	Bobowic-Goldflint complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
333E	Stecum-Hiore-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
335E	Stecum-Goldflint-Branham complex, 12 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	N	Y
340E	Peeler gravelly sandy loam, 8 to 25 percent slopes, stony	N	N	N	Y	Y	Y	Y	N	N
344A	Valleyflat complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
345D	Adit, moderately impacted, stony-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
347D	Nuley-Modess complex, 2 to 12 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
348E	Adit, moderately impacted, stony-Rock outcrop-Modess, moderately impacted complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
349D	Modess-Anaconda complex, 2 to 15 percent slopes, moderately impacted	N	Y	Y	Y	Y	Y	Y	N	N
353C	Highrye-Wissikihon-Beeftrail complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
356E	Peeler-Bobowic-Zonite complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
357D	Beeftrail-Dinnen-Tuggle complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
360B	Tepete mucky peat, 1 to 4 percent slopes, rarely flooded	N	N	N	N	N	N	N	N	N
361G	Rock outcrop-Goldflint-Rubble land complex, 45 to 80 percent slopes	N	N	N	N	N	N	N	Y	Y
362E	Comad-Stecum complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
363C	Tepete mucky peat, sandy substratum, 2 to 6 percent slopes	N	N	N	N	N	N	N	N	N
364F	Comad-Goldflint-Rock outcrop complex, 20 to 50 percent slopes	N	N	N	Y	Y	Y	Y	N	N
365F	Stecum-Rock outcrop-Goldflint complex, 30 to 60 percent slopes	N	N	N	N	N	N	N	Y	Y
366F	Stecum-Rock outcrop-Basincreek complex, 25 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
367G	Rubble land-Stecum-Zonite complex, 35 to 75 percent slopes	N	N	N	N	N	N	N	Y	Y
369E	Rubick, bouldery-Comad, very bouldery complex, 8 to 30 percent slopes	N	N	N	Y	Y	Y	Y	N	N
370C	Comad-Bobowic complex, 2 to 8 percent slopes, bouldery	Y	Y	Y	Y	Y	Y	Y	N	N
371G	Stecum-Rock outcrop-Comad complex, 35 to 70 percent slopes	N	N	N	N	N	N	N	Y	Y
372E	Basincreek-Peeler-Stecum complex, 15 to 45 percent slopes	N	N	N	Y	Y	Y	Y	N	Y
373E	Hiore-Caseypeak-Rock outcrop complex, 15 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
374B	Valleyflat sandy loam, 1 to 4 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
375E	Valleyflat-Pitchstone complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
376A	Cometcrik, rarely flooded-Histic Endoaquolls, rarely flooded-Passmore, very rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	N	N	N
377B	Valleyflat-Pitchstone complex, 1 to 4 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
378C	Valleyflat-Pitchstone complex, 4 to 8 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
379F	Pitchstone-Rockerjohn complex, 15 to 50 percent slopes, extremely bouldery	N	N	N	N	N	N	N	Y	Y
380D	Beeftrail-Dinnen-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
382B	Varney-Anaconda-Varney, sandy substratum complex, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
383C	Highrye-Beeftrail-Oro Fino complex, 1 to 8 percent slopes	N	N	N	N	Y	Y	Y	N	N
384E	Minestope, extremely bouldery-Branham, extremely bouldery-Rock outcrop complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
385D	Highrye-Beeftrail complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
386D	Varney-Con complex, 8 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
389C	Anaconda-Rockerjohn complex, 1 to 8 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
390D	Rockerjohn-Valleyflat complex, 4 to 20 percent slopes, very stony, moderately impacted	Y	Y	Y	Y	Y	Y	Y	N	N
392F	Bobowic, very bouldery-Comad, very bouldery-Rock outcrop complex, 20 to 50 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
394E	Minestope, very stony-Beeftrail, very stony-Rock outcrop complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	Y
395E	Beeftrail-Stecum-Wissikihon complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
397E	Anaconda, sandy substratum-Anaconda-Valleyflat complex, 8 to 45 percent slopes, moderately impacted	N	N	N	Y	Y	Y	Y	N	Y
400C	Anaconda-Varney complex, 2 to 8 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
401D	Modess-Adit-Rockerjohn complex, 4 to 12 percent slopes, moderately impacted	N	Y	Y	Y	Y	Y	Y	N	N
402E	Typic Cryorthents, reclaimed area-Beeftrail, severely impacted complex, 4 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
404E	Beeftrail-Minestope-Dinnen complex, 4 to 25 percent slopes, moderately impacted	Y	Y	Y	Y	Y	Y	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
405E	Minestope-Beeftrail, very bouldery-Nuley complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
406E	Stecum, very bouldery-Comad-Rock outcrop complex, 8 to 30 slopes	Y	Y	Y	Y	Y	Y	Y	N	N
407D	Adit-Adit, moist-Rock outcrop complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
408E	Stecum-Mooseflat-Basincreek complex, 4 to 30 percent slopes, very bouldery	N	Y	Y	Y	Y	Y	Y	N	N
409E	Stecum, moderately impacted, very stony-Rock outcrop-Zonite, moderately impacted, very bouldery complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
410E	Bobowic-Peeler-Caseypeak complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
411D	Modess-Nuley complex, 4 to 12 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
412E	Stecum, stony-Zonite, stony-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
413E	Modess-Adit-Rock outcrop complex, 4 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
414G	Rock outcrop-Stecum-Comad complex, 30 to 90 percent slopes	N	N	N	N	N	N	N	N	N
415F	Stecum-Goldflint-Basincreek complex, 20 to 50 percent slopes, extremely stony	Y	Y	Y	Y	Y	Y	Y	Y	Y
416E	Beeftrail-Fleecer-Stecum complex, 8 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
417E	Fleecer-Tuggle complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
419E	Peeler-Comad complex, 8 to 30 percent slopes, very stony	N	Y	Y	Y	Y	Y	Y	N	N
420B	Dinnen-Wissikihon-Shewag complex, 1 to 6 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
421B	Shewag-Shewag, moderately well drained complex, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
423D	Fleecer-Dinnen complex, 4 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
501C	Patouza-Chinasprings-Dutton complex, 2 to 8 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
502E	Hungryhill-Savenac complex, 8 to 30 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
503F	Bridger-Eastridge-Hungryhill complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	Y	Y	N	N
504E	Bullrey-Hungryhill-Larkspur complex, 8 to 30 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	N	Y
505E	Chinasprings-Euell-Nissler complex, 8 to 30 percent slopes, stony	N	N	N	Y	Y	Y	Y	N	N
507C	Chinasprings ashy sandy loam, 2 to 8 percent slopes, stony, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
508F	Eastridge-Judco complex, 20 to 60 percent slopes	N	N	N	N	N	N	Y	N	N
509D	Bridger-Nissler complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
510F	Euell, very stony-Hungryhill, very stony-Rock outcrop complex, 30 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
511E	Hungryhill-Euell-Larkspur complex, 8 to 30 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
512E	Euell-Larkspur complex, 8 to 30 percent slopes, stony, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
513E	Euell, moderately impacted, very stony-Illiano, moderately impacted, very stony-Rock outcrop complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
514E	Hungryhill-Euell-Poin complex, 8 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
515F	Eastridge-Hungryhill-Poin complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	N
516F	Eastridge-Germangulch complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	Y	Y	Y	Y
517D	Poin-Larkspur complex, 4 to 20 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
518F	Germangulch-Eastridge-Euell complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	Y	Y	Y	Y
519E	Eastridge-Euell complex, 15 to 30 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
522D	Foolhen, rarely flooded-Silas-Vitroff complex, 2 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
523E	Nissler-Euell complex, 12 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
525G	Eastridge gravelly ashy loam, 45 to 75 percent slopes	N	N	N	N	N	N	Y	Y	Y
527E	Patouza-Nivean-Chinasprings complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
528E	Poin-Rock outcrop-Euell complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
530E	Bigbutte, stony-Poin, stony-Brownsgulch complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
532E	Hungryhill-Poin-Larkspur complex, 15 to 45 percent slopes, stony	N	Y	Y	Y	Y	Y	Y	N	N
534D	Chinasprings-Travona complex, 4 to 15 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
536F	Hungryhill, very stony-Euell, very stony-Poin extremely stony complex, 25 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
537E	Bigbutte-Hungryhill-Poin complex, 8 to 30 percent slopes, very stony	Y	Y	Y	Y	Y	Y	Y	N	Y
540D	Evaro-Germangulch complex, 4 to 25 percent slopes, extremely stony	N	N	N	N	N	Y	Y	N	N
541F	Hungryhill-Euell complex, 20 to 50 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	Y	Y
542D	Euell-Hungryhill-Bullrey complex, 4 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
543F	Evaro-Vitroff-Germangulch, very stony complex, 20 to 50 percent slopes	N	N	N	N	N	Y	Y	N	Y
545G	Illiano, very stony-Rock outcrop-Rubble land complex, 45 to 75 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
546E	Hungryhill-Euell-Bullrey complex, 8 to 45 percent slopes, stony	N	N	Y	Y	Y	Y	Y	Y	Y
548F	Evaro-Eastridge-Vitroff complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	N
550E	Evaro-Vitroff complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
551D	Brownsgulch gravelly sandy loam, 4 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
552E	Nissler-Brownsgulch complex, 8 to 25 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
554E	Bridger-Poin, very stony complex, 8 to 30 percent slopes	N	N	N	N	N	Y	Y	N	N
557D	Savenac-Pappascreek-Mooseflat complex, 4 to 25 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
559E	Eastridge-Euell, cool complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
561F	Euell-Bigbutte complex, 20 to 50 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
562G	Poin, very stony-Rubble land-Eastridge, very stony complex, 25 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
564D	Chinasprings-Patouza-Nivean complex, 4 to 15 percent slopes, moderately impacted, stony	N	N	Y	Y	Y	Y	Y	N	N
567F	Evaro-Eastridge complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	N
568F	Evaro-Coslaw-Hungryhill complex, 15 to 50 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
569E	Evaro-Savenac complex, 15 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
570E	Eastridge-Euell complex, 15 to 40 percent slopes	N	N	N	N	N	N	Y	N	N
572D	Euell-Nissler-Poin complex, 4 to 20 percent slopes, stony	N	Y	Y	Y	Y	Y	Y	N	N
573E	Whitlash, very stony-Rock outcrop-Hungryhill, very stony complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
574F	Bullrey-Brownsgulch complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	Y	N	Y
578D	Whitlash, very stony-Rock outcrop-Reedpoint, very stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
579D	Chinasprings-Dutton complex, 4 to 15 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
580E	Patouza-Chinasprings-Nivean complex, 8 to 30 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
584F	Nivean, very stony-Rock outcrop-Whitlash, very stony complex, 20 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
586D	Chinasprings-Patouza-Chinasprings, deep complex, 4 to 15 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
587E	Nivean very gravely sandy loam, 15 to 30 percent slopes, moderately impacted	Y	Y	Y	Y	Y	Y	Y	Y	Y
589E	Nivean-Patouza complex, 8 to 35 percent slopes, very stony, moderately impacted	Y	Y	Y	Y	Y	Y	Y	Y	Y
593F	Nivean-Patouza-Whitlash complex, 20 to 50 percent slopes, very stony, moderately impacted	Y	Y	Y	Y	Y	Y	Y	Y	Y
594D	Evaro-Savenac complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
595D	Mooseflat, very stony-Pappascreek, very stony-Euell complex, 2 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
596D	Bigbutte, stony-Euell, stony-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
597C	Kilgore-Foolhen-Philipsburg complex, 2 to 12 percent slopes	N	N	N	Y	Y	Y	Y	N	N
598E	Chinasprings-Whitlash-Dutton complex, 8 to 30 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
601D	Nissler-Kilgore-Pappascreek, very stony complex, 2 to 15 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
608E	Illiano, very stony-Euell, very stony-Rock outcrop complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
612A	Kilgore, frequently flooded-Foxgulch, rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	N	N	N
614B	Brownsgulch-Foxgulch complex, 2 to 6 percent slopes	N	N	N	N	N	Y	Y	N	N
616D	Silas-Vitroff complex, 2 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
701E	Rubick-Stecum complex, 15 to 45 percent slopes	N	N	N	N	N	N	Y	N	N
702E	Maurice, very stony-Maurice-Sigbird, very stony complex, 12 to 35 percent slopes	N	N	N	N	N	Y	Y	N	N
703G	Surdal, very stony-Rubble land complex, 30 to 70 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
706E	Rubick-Libeg complex, 8 to 35 percent slopes	N	N	N	N	N	N	N	N	N
708D	Rubick-Moosejaw complex, 4 to 20 percent slopes	N	N	N	N	N	N	N	N	N
709E	Libeg-Monad-Copenhaver complex, 15 to 35 percent slopes	N	N	N	N	N	N	Y	N	Y
710E	Worock gravelly loam, dry, 15 to 35 percent slopes	N	N	N	N	N	N	N	N	N
711E	Worock gravelly loam, 15 to 35 percent slopes	N	N	N	N	N	N	Y	N	N
712F	Rubick-Maurice complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	N
713F	Elve gravelly loam, 35 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
715G	Worock, dry-Rubick complex, 30 to 70 percent slopes	N	N	N	N	N	N	Y	N	N
716G	Tiban-Sigbird-Maurice complex, 35 to 75 percent slopes	N	N	N	N	N	N	N	Y	Y
717E	Libeg-Loberg complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
718E	Maurice-Libeg complex, 8 to 30 percent slopes, very stony	N	N	N	N	N	N	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
719G	Tiban, very stony-Sigbird, very stony-Rubble land complex, 35 to 75 percent slopes	N	N	N	N	N	N	N	N	Y
721E	Ratiopeak-Sigbird complex, 15 to 40 percent slopes, very stony	N	N	N	N	N	N	N	N	N
724E	Winspect gravelly loam, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
725E	Winspect-Wilspring complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
726C	Anaconda sandy loam, 0 to 4 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
727D	Pappascreek-Monaberg-Mooseflat complex, 2 to 15 percent slopes, stony	N	N	N	N	N	Y	Y	N	N
728F	Sebud, very bouldery-Sebud, extremely bouldery complex, 20 to 50 percent slopes	N	N	N	N	N	N	N	N	N
729E	Maurice, very stony-Sigbird, rubbly-Rock outcrop complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
730E	Maurice, bouldery-Sigbird, very bouldery complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
731F	Rubick, stony-Worock complex, 20 to 50 percent slopes	N	N	N	N	N	N	N	N	N
732D	Maurice-Mawspring-Sigbird complex, 6 to 20 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
734D	Bullrey-Maurice, very stony-Libeg complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
737D	Libeg-Bridger complex, 6 to 20 percent slopes	N	N	N	N	N	N	Y	N	N
738E	Rubick-Surdal complex, 15 to 35 percent slopes, very stony	N	N	N	N	N	N	N	N	N
739E	Maurice-Surdal-Mawspring complex, 12 to 35 percent slopes, stony	N	N	N	N	N	N	N	N	N
740F	Tiban-Sigbird complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	N
741F	Maurice-Sigbird-Surdal complex, 20 to 50 percent slopes, stony	N	N	Y	Y	Y	Y	Y	Y	Y
742F	Trimad, very stony-Frenchcreek, very stony-Rubble land complex, 25 to 60 percent slopes	N	N	N	N	N	N	N	N	Y
744E	Sigbird, very shallow-Sigbird-Surdal complex, 8 to 25 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
747F	Rubick-Surdal complex, 30 to 70 percent slopes, very stony	N	N	N	N	N	N	N	N	Y
749E	Surdal-Mawspring, bouldery complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
750F	Poin, extremely stony-Hungryhill, very stony complex, 20 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
751F	Sigbird-Tiban-Maurice complex, 20 to 50 percent slopes, very stony	N	Y	Y	Y	Y	Y	Y	N	Y
756D	Passcreek-Poin, stony complex, 6 to 20 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
757F	Surdal-Sigbird-Mawspring complex, 15 to 60 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	Y	Y
761E	Quincreek, very stony-Whitlash, very stony-Rock outcrop complex, 12 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
801F	Skaggs, extremely stony-Skaggs, very stony-Rock outcrop complex, 20 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
802E	Starley-Tropal complex, 12 to 40 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
803E	Sieben very cobbly loam, 12 to 30 percent slopes, stony	N	N	N	N	N	N	N	N	N
805C	Sixbeacon, stony-Bronec complex, 2 to 8 percent slopes	N	N	N	N	N	N	Y	N	N
806D	Sieben very cobbly loam, 4 to 12 percent slopes	N	N	N	N	N	N	N	N	N
807E	Pensore, stony-Roto complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
808F	Pensore, stony-Rock outcrop complex, 20 to 45 percent slopes	N	N	N	N	N	N	N	Y	Y
809G	Pensore, stony-Rock outcrop complex, 45 to 70 percent slopes	N	N	N	N	N	N	N	Y	Y
810F	Whitore, very stony-Skaggs-very stony-Rock outcrop complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	Y
811G	Whitore-Skaggs complex, 40 to 70 percent slopes, very stony	N	N	N	N	N	N	Y	Y	Y
812F	Whitore, moist-Skaggs complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	Y	N	Y
813F	Whitore, stony-Whitore complex, 20 to 60 percent slopes	N	N	N	N	N	N	Y	Y	Y
814E	Whitore complex, 12 to 45 percent slopes, stony	N	N	N	N	N	N	Y	N	Y
816G	Whitore, very stony-Tropal, very stony-Rock outcrop complex, 45 to 80 percent slopes	N	N	N	N	N	N	Y	Y	Y
817E	Whitore, very stony-Raynesford complex, 15 to 40 percent slopes	N	N	N	N	N	N	Y	N	N
818D	Raynesford-Whitore, very stony complex, 4 to 20 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
819D	Raynesford-Whitore complex, 4 to 15 percent slopes, stony	N	Y	Y	Y	Y	Y	Y	N	N
820E	Whitore, stony-Tropal, very stony-Raynesford, stony complex, 12 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
823E	Skaggs-Raynesford-Tropal, very stony complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
825E	Skaggs-Whitore complex, 12 to 35 percent slopes, stony	N	N	Y	Y	Y	Y	Y	Y	Y
826E	Raynesford-Whitore-Skaggs complex, 12 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
830D	Kalsted gravelly sandy loam, 8 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
901E	Sebud-Poin-Tiban complex, 15 to 45 percent slopes, extremely stony	Y	Y	Y	Y	Y	Y	Y	N	Y
904D	Sebud, stony-Redchief complex, 8 to 25 percent slopes	N	N	N	N	N	N	N	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
905E	Tigeron, stony-Rubick, very stony complex, 15 to 45 percent slopes	N	N	N	N	Y	Y	Y	N	N
906E	Rubick, very stony-Tigeron, stony complex, 15 to 45 percent slopes	N	N	N	N	Y	Y	Y	N	N
907F	Sebud, extremely stony-Rubick, very stony complex, 25 to 60 percent slopes	N	N	N	N	N	N	N	N	Y
908E	Sebud-Tiban-Ratiopeak complex, 12 to 30 percent slopes, stony	N	N	N	N	N	N	N	N	N
909G	Rubick, rubbly-Rubble land complex, 40 to 75 percent slopes	N	N	N	N	Y	Y	Y	N	Y
910F	Sebud-Ratiopeak complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	Y
911E	Sebud, stony-Adel complex, 12 to 30 percent slopes	N	N	N	N	N	N	N	N	N
912D	Ratiopeak-Redchief complex, 4 to 15 percent slopes, stony	N	N	N	N	N	N	N	N	N
913E	Rubick gravelly sandy loam, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
920G	Poin, rubbly-Rubble land-Rock outcrop complex, 40 to 80 percent slopes	N	N	N	N	N	N	N	Y	Y
921F	Whitlash-Frenchcreek complex, 20 to 50 percent slopes, rubbly	Y	Y	Y	Y	Y	Y	Y	Y	Y
923F	Whitore, rubbly-Poin, rubbly-Rock outcrop complex, 25 to 60 percent slopes	N	N	N	N	N	N	Y	N	Y
924F	Whitlash, extremely stony-Gnojek, extremely stony-Rock outcrop complex, 20 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
926F	Rubick-Tigeron complex, 30 to 60 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	N	Y
927E	Tigeron, very stony-Rubick, very stony-Tigeron, stony complex, 15 to 45 percent slopes	N	N	N	N	N	Y	Y	N	N
928E	Mawspring-Maurice complex, 15 to 45 percent slopes, very stony	N	N	N	N	N	N	N	N	N
929F	Rubick, very stony-Poin, extremely bouldery complex, 25 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
930F	Ratiopeak-Tiban complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	N	N	N	N
931E	Ratiopeak-Monaberg complex, 8 to 30 percent slopes, very stony	N	N	N	N	N	N	N	N	N
932D	Tigeron-Rubick complex, 2 to 15 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
933B	Foxgulch, occasionally flooded-Bearmouth, rarely flooded complex, 0 to 4 percent slopes, stony	N	N	N	N	Y	Y	Y	N	N
935D	Anaconda sandy loam, 8 to 15 percent slopes, moderately impacted	N	Y	Y	Y	Y	Y	Y	N	N
936B	Foxgulch, rarely flooded-Kilgore, occasionally flooded complex, 0 to 4 percent slopes, very stony	N	N	N	N	Y	Y	Y	N	N
939F	Evaro, stony-Tigeron complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	N
940E	Ratiopeak, stony-Tiban, very stony complex, 8 to 25 percent slopes	N	N	N	N	N	N	Y	N	N
941E	Bridger gravelly loam, 12 to 30 percent slopes, stony	N	N	N	N	N	N	N	N	N
942F	Whitlash-Gnojek complex, 20 to 60 percent slopes, very stony	Y	Y	Y	Y	Y	Y	Y	Y	Y
943D	Bridger-Adel complex, 4 to 15 percent slopes	N	N	N	N	N	N	N	N	N
992	Pits, borrow	0	0	0	0	0	0	0	0	0
994	Pits, mine	0	0	0	0	0	0	0	0	0
995	Dumps, garbage	0	0	0	0	0	0	0	0	0
997	Dumps, mine	0	0	0	0	0	0	0	0	0
998	Rock outcrop and Rubble land	0	0	0	0	0	0	0	0	0
998	Rock outcrop and Rubble land	0	0	0	0	0	0	0	0	0
GP	Gravel pits	0	0	0	0	0	0	0	0	0
M-W	Miscellaneous water	0	0	0	0	0	0	0	0	0
W	Water	0	0	0	0	0	0	0	0	0