Approved
State Soil Scientist
5/20/09

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
14A	Nuley Variant loam	N	Y	Y	Y	Y	Y	Y	N	N
14C	Nuley Variant loam	N	Y	Y	Y	Y	Y	Y	N	N
14D	Nuley Variant loam	N	Y	Y	Y	Y	Y	Y	N	N
18A	Corner Family Gravelly coarse sandy loam	N	Y	Y	Y	Y	Y	Y	N	N
20B	Anaconda Family-Corner Family complex	N	N	N	N	N	N	Y	N	N
24A	Beaverell cobbly loam	N	Y	Y	Y	Y	Y	Y	N	N
24D	Beaverell cobbly loam	N	Y	Y	Y	Y	Y	Y	N	N
26A	Gledive Variant sandy loam	N	N	Y	Y	Y	Y	Y	N	N
27A	Attewan Variant loam	N	N	Y	Y	Y	Y	Y	N	N
28A	Fubartoo silt loam	Y	Y	Y	Y	Y	Y	Y	N	N
29C	Dinnen coarse sandy loam	N	Y	Y	Y	Y	Y	Y	N	N
29D	Dinnen coarse sandy loam	N	Y	Y	Y	Y	Y	Y	N	N
32E	Brocko family cobbly silt loam	N	N	N	N	N	Y	Y	N	Y
37C	Brownsto gravelly loam	N	N	Y	Y	Y	Y	Y	N	N
47E	Supervisor family-Merino-Rock outcrop complex	N	Y	Y	Y	Y	Y	Y	N	Y
49C	Libeg gravelly loam	N	N	Y	Y	Y	Y	Y	N	N
50A	Bonebasin, occasionally flooded-Meadowcreek, rarely flooded complex, 0 to 4 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
50C	Buffork loam	N	N	Y	Y	Y	Y	Y	N	N
54C	Little Horn family cobbly silt loam	N	N	Y	Y	Y	Y	Y	N	N
54D	Little Horn family cobbly silt loam	N	N	Y	Y	Y	Y	Y	N	N
111E	Sebud-Philipsburg complex, 8 to 25 percent slopes, bouldery	N	N	N	N	N	N	Y	N	N
112C	Varney Varient-Radersburg Variant complex	N	N	Y	Y	Y	Y	Y	N	N
141E	Nuley Variant very cobbly loam	N	Y	Y	Y	Y	Y	Y	N	Y
143C	Nuley Variant stony loam	N	Y	Y	Y	Y	Y	Y	N	N
291E	Dinnen-Branham-Rock outcrop complex	Y	Y	Y	Y	Y	Y	Y	N	Y
301D	Hauz-Tolman complex	Y	Y	Y	Y	Y	Y	Y	N	Y
301E	Philipsburg-Beeftrail, bouldery-Danielvil complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	Y
371E	Brownsto cobbly loam	N	N	Y	Y	Y	Y	Y	N	Y
501D	Buffork-Sawcreek complex	N	N	Y	Y	Y	Y	Y	N	N
521C	Whitecap-Sawcreek complex	Y	Y	Y	Y	Y	Y	Y	N	N
10A	Slickens-Aquic Cumulic Haplustolls, occasionally flooded-Aridic Ustifluvents, occasionally flooded complex, 0 to 2 percent slopes, severely impacted	N	N	Y	Y	Y	Y	Y	N	N
11A	Mannixlee-Bonebasin complex, 0 to 4 percent slopes, frequently flooded	N	N	Y	Y	Y	Y	Y	N	N
12A	Riverrun, occasionally flooded-Mannixlee, frequently flooded complex, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
15B	Anamac-Varney-Rivra, rarely flooded complex, 0 to 6 percent slopes	N	N	N	N	Y	Y	Y	N	N
21C	Kilgore, occasionally flooded-Foolhen, rarely flooded-Philipsburg complex, 2 to 12 percent slopes	N	N	N	Y	Y	Y	Y	N	N
22B	Philipsburg loam, 1 to 4 percent slopes, stony	N	N	N	N	N	Y	Y	N	N
24B	Kilgore, rarely flooded-Mooseflat, occasionally flooded-Philipsburg complex, 0 to 6 percent slopes	N	N	N	Y	Y	Y	Y	N	N
25D	Work, stony-Adel-Kilgore, rarely flooded complex, 2 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
26C	Mooseflat, occasionally flooded-Bridger-Kilgore, rarely flooded complex, 0 to 8 percent slopes	N	N	N	N	N	N	N	N	N
27D	Savenac-Bridger complex, 4 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
28C	Kilgore-Danielvil complex, 2 to 8 percent slopes	N	N	N	Y	Y	Y	Y	N	N
31A	Philipsburg cobbly loam, 0 to 4 percent slopes, very stony	N	N	N	Y	Y	Y	Y	N	N
32B	Kilgore, occasionally flooded-Danielvil complex, 1 to 4 percent slopes	N	N	N	N	N	N	N	N	N
33B	Sebud very cobbly loam, 2 to 6 percent slopes, very stony	N	N	N	N	N	N	N	N	N
34A	Danielvil-Danielvil, rarely flooded complex, 0 to 4 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
36A	Foolhen-Monaberg-Mooseflat complex, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
37B	Mooseflat, occasionally flooded-Monaberg, rarely flooded complex, 1 to 4 percent slopes	N	N	N	N	N	N	N	N	N
38B	Rivra gravelly sandy loam, 0 to 4 percent slopes, moderately impacted	Y	Y	Y	Y	Y	Y	Y	N	N
40B	Cetrack loam, 0 to 4 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
42D	Varney-Varney, stony-Anamac, stony complex, 2 to 12 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
43B	Meadowcreek-Mannixlee, rarely flooded complex, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
44B	Meadowcreek-Anamac complex, 1 to 6 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
46A	Varney-Rivra complex, 0 to 4 percent slopes	N	N	N	Y	Y	Y	Y	N	N
47B	Foolhen-Monaberg-Kilgore, occasionally flooded complex, 0 to 6 percent slopes	N	N	N	N	N	Y	Y	N	N
51D	Foxgulch-Libeg complex, 6 to 25 percent slopes, stony	N	N	N	N	Y	Y	Y	N	N
53C	Sixbeacon-Sieben complex, 2 to 8 percent slopes	N	N	N	N	N	N	Y	N	N
55A	Bonebasin, occasionally flooded-Monaberg, very rarely flooded-Meadowcreek, rarely flooded complex, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
56B	Bonebasin, occasionally flooded-Meadowcreek, rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
58B	Varney-Anaconda loams, 0 to 4 percent slopes, moderately impacted	N	N	N	N	N	N	Y	N	N
59A	Anamac-Meadowcreek complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
60A	Riverrun-Rivra complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
61A	Rivra, stony-Riverrun complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
62A	Vendome-Sieben complex, 0 to 2 percent slopes	N	N	N	N	N	N	Y	N	N
64A	Work clay loam, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
65B	Sebud-Bearmouth complex, 1 to 4 percent slopes, very stony	N	N	N	N	Y	Y	Y	N	N
67A	Foxgulch-Bearmouth, very stony complex, 0 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
68B	Bearmouth, rarely flooded-Foxgulch, occasionally flooded complex, 0 to 4 percent slopes, very stony	N	N	N	Y	Y	Y	Y	N	N
69B	Foxgulch loam, 1 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
71D	Sebud-Ratiopeak complex, 4 to 15 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	N
72B	Tibson gravelly loam, 1 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
74A	Bearmouth very cobbly sandy loam, 0 to 2 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	N	N
75A	Danielvil loam, 0 to 2 percent slopes	N	N	N	N	N	N	Y	N	N
76A	Bearmouth very cobbly sandy loam, 0 to 4 percent slopes	0	0	0	0	0	0	0	0	0
80A	Water-Riverwash complex	N	N	Y	Y	Y	Y	Y	N	N
81A	Varney sandy loam, 0 to 4 percent slopes	N	N	N	N	N	Y	Y	N	N
82D	Philipsburg-Monaberg-Kilgore, occasionally flooded complex, 0 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
82E	Elve gravelly loam, 15 to 35 percent slopes	N	N	N	N	N	N	Y	N	N
85E	Loberg gravelly loam, 15 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
101B	Philipsburg sandy loam, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
102D	Varney, noncalcic-Philipsburg-Varney complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
106E	Gnojek-Libeg-Philipsburg complex, 8 to 30 percent slopes, stony	Y	Y	Y	Y	Y	Y	Y	N	Y
107E	Anaconda-Varney-Work, stony complex, 6 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
108D	Varney-Work complex, 4 to 15 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	N
109A	Bearmouth coarse sandy loam, 0 to 4 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
110E	Danielvil-Bearmouth complex, 2 to 25 percent slopes	N	N	N	N	N	N	Y	N	N
112E	Monaberg-Bridger-Libeg, stony complex, 8 to 25 percent slopes	N	N	N	N	N	N	Y	N	N
114B	Varney loam, 0 to 4 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
115D	Philipsburg-Ratiopeak complex, 8 to 15 percent slopes	N	N	N	Y	Y	Y	Y	N	N
116C	Monaberg-Philipsburg complex, 2 to 12 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
118E	Reedpoint-Rock outcrop-Sixbeacon complex, 6 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
119C	Varney-Udecide complex, 2 to 12 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
120C	Varney-Anamac complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
122C	Varney-Work complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
123E	Sebud, stony-Danielvil-Monaberg, very stony complex, 8 to 30 percent slopes	N	N	N	N	N	Y	Y	N	N
124E	Monaberg-Bridger complex, 8 to 30 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
125E	Libeg-Monaberg complex, 15 to 35 percent slopes, very bouldery	N	Y	Y	Y	Y	Y	Y	Y	Y
126E	Danielvil, very bouldery-Monaberg, bouldery-Adel complex, 8 to 35 percent slopes	N	N	N	N	N	Y	Y	N	N
128D	Monaberg loam, 4 to 15 percent slopes, bouldery	N	N	N	Y	Y	Y	Y	N	N
129D	Philipsburg-Danielvil complex, 4 to 15 percent slopes	N	N	N	Y	Y	Y	Y	N	N
131E	Dumps, placer mining-Philipsburg, stony complex, 4 to 15 percent slopes									

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
136D	Varney-Varney, stony-Anamac complex, 2 to 12 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
139D	Beeftrail, occasionally flooded-Adel-Monaberg complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
140D	Monaberg-Adel complex, 4 to 20 percent slopes	N	N	N	N	N	N	Y	N	N
141F	Danielvil-Danielvil, cool-Philipsburg complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
142E	Ratiopeak-Philipsburg complex, 15 to 35 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
145D	Danielvil-Philipsburg complex, 4 to 12 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
147B	Philipsburg-Ratiopeak complex, 1 to 6 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
148C	Philipsburg-Monaberg complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
149D	Bridger-Libeg complex, 8 to 25 percent slopes, very stony	N	N	N	N	N	N	N	N	N
150D	Sebud, very stony-Ratiopeak, stony-Bridger, stony complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
152E	Sieben-Varney complex, 15 to 45 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
153D	Sebud, very stony-Philipsburg-Ratiopeak, very stony complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
158C	Varney-Anamac-Anaconda complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
159E	Monaberg-Varney-Philipsburg complex, 8 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
160E	Bridger-Patouza, stony-Varney complex, 8 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
161C	Philipsburg-Philipsburg, stony-Danielvil complex, 2 to 8 percent slopes	N	N	N	Y	Y	Y	Y	N	N
162B	Monaberg loam, 1 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
164B	Danielvil loam, moist, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
165A	Mooseflat, rarely flooded-Foxgulch, very rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	N	N	N
166B	Kilgore-Foolhen-Monaberg complex, 0 to 4 percent slopes	N	N	N	N	N	N	Y	N	N
167C	Monaberg loam, 2 to 8 percent slopes, stony	N	N	N	N	N	N	Y	N	N
168E	Sebud-Sieben complex, 8 to 30 percent slopes, stony	N	N	N	N	N	N	Y	N	N
169E	Sebud-Sieben complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
170E	Ratiopeak, very stony-Bridger-Adel complex, 8 to 30 percent slopes	N	N	N	N	Y	Y	Y	N	N
171D	Varney-Sieben, stony complex, 4 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
172F	Ratiopeak-Sixbeacon-Tiban complex, 15 to 45 percent slopes, extremely stony	N	N	N	N	Y	Y	Y	N	N
173B	Monaberg-Foolhen complex, 1 to 6 percent slopes	N	N	N	N	N	N	Y	N	N
174C	Philipsburg loam, 2 to 8 percent slopes	N	N	N	N	N	Y	Y	N	N
175B	Foolhen-Kilgore complex, 0 to 4 percent slopes	N	N	N	Y	Y	Y	Y	N	N
177C	Udecide-Gnojek, stony complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
178E	Rootel-Varney complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
179F	Vendome, extremely stony-Sixbeacon complex, 25 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
182C	Varney, stony-Anamac, very stony-Rivra, extremely stony complex, 2 to 8 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
183B	Anamac-Varney complex, 1 to 6 percent slopes	N	N	N	N	N	N	Y	N	N
235B	Anaconda sandy loam, 0 to 4 percent slopes, severely impacted	N	N	Y	Y	Y	Y	Y	N	N
272F	Rencot, stony-Spudbar-Rock outcrop complex, 25 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
303D	Moosejaw, occasionally flooded-Highrye-Silas, occasionally flooded complex, 2 to 12 percent slopes	N	N	N	N	N	N	Y	N	N
305D	Beeftrail-Branham-Minestope complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
306E	Wissikihon-Branham-Highrye complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
312D	Oro Fino-Highrye-Branham complex, 4 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
313E	Beeftrail-Dinnen-Highrye complex, 8 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
314F	Basincreek-Comad complex, 20 to 50 percent slopes	N	N	N	Y	Y	Y	Y	Y	Y
315F	Stecum-Hiore complex, 20 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
316F	Stecum, very bouldery-Rock outcrop-Zonite, very bouldery complex, 20 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
317E	Stecum-Caseypeak-Rock outcrop complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
319D	Silas, stony-Branham, stony-Tepete complex, 2 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
321E	Beeftrail-Bavdark-Zonite complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
323E	Highrye-Tuggle, very stony-Moosejaw complex, 2 to 25 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
324E	Beeftrail, stony-Branham, stony-Fleecer complex, 8 to 20 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
326C	Fleecer-Branham-Passmore complex, 2 to 8 percent slopes	N	N	N	Y	Y	Y	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
327E	Highrye-Stecum-Wissikihon complex, 15 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
328E	Stecum-Zonite-Basincreek complex, 8 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
331C	Mooseflat, occasionally flooded-Foolhen, occasionally flooded-Fleecer complex, 2 to 8 percent slopes	N	N	N	N	N	N	N	N	N
332D	Bobowic-Goldflint complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
333E	Stecum-Hiore-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
335E	Stecum-Goldflint-Branham complex, 12 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	N	Y
340E	Peeler gravelly sandy loam, 8 to 25 percent slopes, stony	N	N	N	Y	Y	Y	Y	N	N
344A	Valleyflat complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
345D	Adit, moderately impacted, stony-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
347D	Nuley-Modess complex, 2 to 12 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
348E	Adit, moderately impacted, stony-Rock outcrop-Modess, moderately impacted complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
349D	Modess-Anaconda complex, 2 to 15 percent slopes, moderately impacted	N	Y	Y	Y	Y	Y	Y	N	N
353C	Highrye-Wissikihon-Beeftrail complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
356E	Peeler-Bobowic-Zonite complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
357D	Beeftrail-Dinnen-Tuggle complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
360B	Tepete mucky peat, 1 to 4 percent slopes, rarely flooded	N	N	N	N	N	N	N	N	N
361G	Rock outcrop-Goldflint-Rubble land complex, 45 to 80 percent slopes	N	N	N	N	N	N	N	Y	Y
362E	Comad-Stecum complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
363C	Tepete mucky peat, sandy substratum, 2 to 6 percent slopes	N	N	N	N	N	N	N	N	N
364F	Comad-Goldflint-Rock outcrop complex, 20 to 50 percent slopes	N	N	N	Y	Y	Y	Y	N	N
365F	Stecum-Rock outcrop-Goldflint complex, 30 to 60 percent slopes	N	N	N	N	N	N	N	Y	Y
366F	Stecum-Rock outcrop-Basincreek complex, 25 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
367G	Rubble land-Stecum-Zonite complex, 35 to 75 percent slopes	N	N	N	N	N	N	N	Y	Y
369E	Rubick, bouldery-Comad, very bouldery complex, 8 to 30 percent slopes	N	N	N	Y	Y	Y	Y	N	N
370C	Comad-Bobowic complex, 2 to 8 percent slopes, bouldery	Y	Y	Y	Y	Y	Y	Y	N	N
371G	Stecum-Rock outcrop-Comad complex, 35 to 70 percent slopes	N	N	N	N	N	N	N	Y	Y
372E	Basincreek-Peeler-Stecum complex, 15 to 45 percent slopes	N	N	N	Y	Y	Y	Y	N	Y
373E	Hiore-Caseypeak-Rock outcrop complex, 15 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
374B	Valleyflat sandy loam, 1 to 4 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
375E	Valleyflat-Pitchstone complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
376A	Cometcrik, rarely flooded-Histic Endoaquolls, rarely flooded-Passmore, very rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	N	N	N
377B	Valleyflat-Pitchstone complex, 1 to 4 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
378C	Valleyflat-Pitchstone complex, 4 to 8 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
379F	Pitchstone-Rockerjohn complex, 15 to 50 percent slopes, extremely bouldery	N	N	N	N	N	N	N	Y	Y
380D	Beeftrail-Dinnen-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
382B	Varney-Anaconda-Varney, sandy substratum complex, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
383C	Highrye-Beeftrail-Oro Fino complex, 1 to 8 percent slopes	N	N	N	N	Y	Y	Y	N	N
384E	Minestope, extremely bouldery-Branham, extremely bouldery-Rock outcrop complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
385D	Highrye-Beeftrail complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
386D	Varney-Con complex, 8 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
389C	Anaconda-Rockerjohn complex, 1 to 8 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
390D	Rockerjohn-Valleyflat complex, 4 to 20 percent slopes, very stony, moderately impacted	Y	Y	Y	Y	Y	Y	Y	N	N
392F	Bobowic, very bouldery-Comad, very bouldery-Rock outcrop complex, 20 to 50 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
394E	Minestope, very stony-Beeftrail, very stony-Rock outcrop complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	Y
395E	Beeftrail-Stecum-Wissikihon complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
397E	Anaconda, sandy substratum-Anaconda-Valleyflat complex, 8 to 45 percent slopes, moderately impacted	N	N	N	Y	Y	Y	Y	N	Y
400C	Anaconda-Varney complex, 2 to 8 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
401D	Modess-Adit-Rockerjohn complex, 4 to 12 percent slopes, moderately impacted	N	Y	Y	Y	Y	Y	Y	N	N
402E	Typic Cryorthents, reclaimed area-Beeftrail, severely impacted complex, 4 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
404E	Beeftrail-Minestope-Dinnen complex, 4 to 25 percent slopes, moderately impacted	Y	Y	Y	Y	Y	Y	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
405E	Minestope-Beeftrail, very bouldery-Nuley complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
406E	Stecum, very bouldery-Comad-Rock outcrop complex, 8 to 30 slopes	Y	Y	Y	Y	Y	Y	Y	N	N
407D	Adit-Adit, moist-Rock outcrop complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
408E	Stecum-Mooseflat-Basincreek complex, 4 to 30 percent slopes, very bouldery	N	Y	Y	Y	Y	Y	Y	N	N
409E	Stecum, moderately impacted, very stony-Rock outcrop-Zonite, moderately impacted, very bouldery complex, 8 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
410E	Bobowic-Peeler-Caseypeak complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
411D	Modess-Nuley complex, 4 to 12 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
412E	Stecum, stony-Zonite, stony-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
413E	Modess-Adit-Rock outcrop complex, 4 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
414G	Rock outcrop-Stecum-Comad complex, 30 to 90 percent slopes	N	N	N	N	N	N	N	N	N
415F	Stecum-Goldflint-Basincreek complex, 20 to 50 percent slopes, extremely stony	Y	Y	Y	Y	Y	Y	Y	Y	Y
416E	Beeftrail-Fleecer-Stecum complex, 8 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
417E	Fleecer-Tuggle complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
419E	Peeler-Comad complex, 8 to 30 percent slopes, very stony	N	Y	Y	Y	Y	Y	Y	N	N
420B	Dinnen-Wissikihon-Shewag complex, 1 to 6 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
421B	Shewag-Shewag, moderately well drained complex, 1 to 4 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
423D	Fleecer-Dinnen complex, 4 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
501C	Patouza-Chinasprings-Dutton complex, 2 to 8 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
502E	Hungryhill-Savenac complex, 8 to 30 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
503F	Bridger-Eastridge-Hungryhill complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	Y	Y	N	N
504E	Bullrey-Hungryhill-Larkspur complex, 8 to 30 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	N	Y
505E	Chinasprings-Euell-Nissler complex, 8 to 30 percent slopes, stony	N	N	N	Y	Y	Y	Y	N	N
507C	Chinasprings ashy sandy loam, 2 to 8 percent slopes, stony, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
508F	Eastridge-Judco complex, 20 to 60 percent slopes	N	N	N	N	N	N	Y	N	N
509D	Bridger-Nissler complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
510F	Euell, very stony-Hungryhill, very stony-Rock outcrop complex, 30 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
511E	Hungryhill-Euell-Larkspur complex, 8 to 30 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
512E	Euell-Larkspur complex, 8 to 30 percent slopes, stony, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
513E	Euell, moderately impacted, very stony-Illiano, moderately impacted, very stony-Rock outcrop complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
514E	Hungryhill-Euell-Poin complex, 8 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
515F	Eastridge-Hungryhill-Poin complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	N
516F	Eastridge-Germangulch complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	Y	Y	Y	Y
517D	Poin-Larkspur complex, 4 to 20 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
518F	Germangulch-Eastridge-Euell complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	Y	Y	Y	Y
519E	Eastridge-Euell complex, 15 to 30 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
522D	Foolhen, rarely flooded-Silas-Vitroff complex, 2 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
523E	Nissler-Euell complex, 12 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
525G	Eastridge gravelly ashy loam, 45 to 75 percent slopes	N	N	N	N	N	N	Y	Y	Y
527E	Patouza-Nivean-Chinasprings complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
528E	Poin-Rock outcrop-Euell complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
530E	Bigbutte, stony-Poin, stony-Brownsgulch complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
532E	Hungryhill-Poin-Larkspur complex, 15 to 45 percent slopes, stony	N	Y	Y	Y	Y	Y	Y	N	N
534D	Chinasprings-Travona complex, 4 to 15 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
536F	Hungryhill, very stony-Euell, very stony-Poin extremely stony complex, 25 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
537E	Bigbutte-Hungryhill-Poin complex, 8 to 30 percent slopes, very stony	Y	Y	Y	Y	Y	Y	Y	N	Y
540D	Evaro-Germangulch complex, 4 to 25 percent slopes, extremely stony	N	N	N	N	N	Y	Y	N	N
541F	Hungryhill-Euell complex, 20 to 50 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	Y	Y
542D	Euell-Hungryhill-Bullrey complex, 4 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
543F	Evaro-Vitroff-Germangulch, very stony complex, 20 to 50 percent slopes	N	N	N	N	N	Y	Y	N	Y
545G	Illiano, very stony-Rock outcrop-Rubble land complex, 45 to 75 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
546E	Hungryhill-Euell-Bullrey complex, 8 to 45 percent slopes, stony	N	N	Y	Y	Y	Y	Y	Y	Y
548F	Evaro-Eastridge-Vitroff complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	N
550E	Evaro-Vitroff complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
551D	Brownsgulch gravelly sandy loam, 4 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
552E	Nissler-Brownsgulch complex, 8 to 25 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
554E	Bridger-Poin, very stony complex, 8 to 30 percent slopes	N	N	N	N	N	Y	Y	N	N
557D	Savenac-Pappascreek-Mooseflat complex, 4 to 25 percent slopes, very stony	N	N	N	N	N	Y	Y	N	N
559E	Eastridge-Euell, cool complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
561F	Euell-Bigbutte complex, 20 to 50 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
562G	Poin, very stony-Rubble land-Eastridge, very stony complex, 25 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
564D	Chinasprings-Patouza-Nivean complex, 4 to 15 percent slopes, moderately impacted, stony	N	N	Y	Y	Y	Y	Y	N	N
567F	Evaro-Eastridge complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	N
568F	Evaro-Coslaw-Hungryhill complex, 15 to 50 percent slopes	N	N	Y	Y	Y	Y	Y	N	Y
569E	Evaro-Savenac complex, 15 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
570E	Eastridge-Euell complex, 15 to 40 percent slopes	N	N	N	N	N	N	Y	N	N
572D	Euell-Nissler-Poin complex, 4 to 20 percent slopes, stony	N	Y	Y	Y	Y	Y	Y	N	N
573E	Whitlash, very stony-Rock outcrop-Hungryhill, very stony complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
574F	Bullrey-Brownsgulch complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	Y	N	Y
578D	Whitlash, very stony-Rock outcrop-Reedpoint, very stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
579D	Chinasprings-Dutton complex, 4 to 15 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
580E	Patouza-Chinasprings-Nivean complex, 8 to 30 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
584F	Nivean, very stony-Rock outcrop-Whitlash, very stony complex, 20 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
586D	Chinasprings-Patouza-Chinasprings, deep complex, 4 to 15 percent slopes, moderately impacted	N	N	Y	Y	Y	Y	Y	N	N
587E	Nivean very gravely sandy loam, 15 to 30 percent slopes, moderately impacted	Y	Y	Y	Y	Y	Y	Y	Y	Y
589E	Nivean-Patouza complex, 8 to 35 percent slopes, very stony, moderately impacted	Y	Y	Y	Y	Y	Y	Y	Y	Y
593F	Nivean-Patouza-Whitlash complex, 20 to 50 percent slopes, very stony, moderately impacted	Y	Y	Y	Y	Y	Y	Y	Y	Y
594D	Evaro-Savenac complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
595D	Mooseflat, very stony-Pappascreek, very stony-Euell complex, 2 to 15 percent slopes	N	N	N	N	N	Y	Y	N	N
596D	Bigbutte, stony-Euell, stony-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
597C	Kilgore-Foolhen-Philipsburg complex, 2 to 12 percent slopes	N	N	N	Y	Y	Y	Y	N	N
598E	Chinasprings-Whitlash-Dutton complex, 8 to 30 percent slopes, stony	N	N	Y	Y	Y	Y	Y	N	Y
601D	Nissler-Kilgore-Pappascreek, very stony complex, 2 to 15 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
608E	Illiano, very stony-Euell, very stony-Rock outcrop complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
612A	Kilgore, frequently flooded-Foxgulch, rarely flooded complex, 0 to 4 percent slopes	N	N	N	N	N	N	N	N	N
614B	Brownsgulch-Foxgulch complex, 2 to 6 percent slopes	N	N	N	N	N	Y	Y	N	N
616D	Silas-Vitroff complex, 2 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
701E	Rubick-Stecum complex, 15 to 45 percent slopes	N	N	N	N	N	N	Y	N	N
702E	Maurice, very stony-Maurice-Sigbird, very stony complex, 12 to 35 percent slopes	N	N	N	N	N	Y	Y	N	N
703G	Surdal, very stony-Rubble land complex, 30 to 70 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
706E	Rubick-Libeg complex, 8 to 35 percent slopes	N	N	N	N	N	N	N	N	N
708D	Rubick-Moosejaw complex, 4 to 20 percent slopes	N	N	N	N	N	N	N	N	N
709E	Libeg-Monad-Copenhaver complex, 15 to 35 percent slopes	N	N	N	N	N	N	Y	N	Y
710E	Worock gravelly loam, dry, 15 to 35 percent slopes	N	N	N	N	N	N	N	N	N
711E	Worock gravelly loam, 15 to 35 percent slopes	N	N	N	N	N	N	Y	N	N
712F	Rubick-Maurice complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	N
713F	Elve gravelly loam, 35 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
715G	Worock, dry-Rubick complex, 30 to 70 percent slopes	N	N	N	N	N	N	Y	N	N
716G	Tiban-Sigbird-Maurice complex, 35 to 75 percent slopes	N	N	N	N	N	N	N	Y	Y
717E	Libeg-Loberg complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
718E	Maurice-Libeg complex, 8 to 30 percent slopes, very stony	N	N	N	N	N	N	Y	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
719G	Tiban, very stony-Sigbird, very stony-Rubble land complex, 35 to 75 percent slopes	N	N	N	N	N	N	N	N	Y
721E	Ratiopeak-Sigbird complex, 15 to 40 percent slopes, very stony	N	N	N	N	N	N	N	N	N
724E	Winspect gravelly loam, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	Y
725E	Winspect-Wilspring complex, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
726C	Anaconda sandy loam, 0 to 4 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
727D	Pappascreek-Monaberg-Mooseflat complex, 2 to 15 percent slopes, stony	N	N	N	N	N	Y	Y	N	N
728F	Sebud, very bouldery-Sebud, extremely bouldery complex, 20 to 50 percent slopes	N	N	N	N	N	N	N	N	N
729E	Maurice, very stony-Sigbird, rubbly-Rock outcrop complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
730E	Maurice, bouldery-Sigbird, very bouldery complex, 8 to 30 percent slopes	N	N	N	N	N	N	Y	N	N
731F	Rubick, stony-Worock complex, 20 to 50 percent slopes	N	N	N	N	N	N	N	N	N
732D	Maurice-Mawspring-Sigbird complex, 6 to 20 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
734D	Bullrey-Maurice, very stony-Libeg complex, 4 to 15 percent slopes	N	N	N	N	N	N	Y	N	N
737D	Libeg-Bridger complex, 6 to 20 percent slopes	N	N	N	N	N	N	Y	N	N
738E	Rubick-Surdal complex, 15 to 35 percent slopes, very stony	N	N	N	N	N	N	N	N	N
739E	Maurice-Surdal-Mawspring complex, 12 to 35 percent slopes, stony	N	N	N	N	N	N	N	N	N
740F	Tiban-Sigbird complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	N
741F	Maurice-Sigbird-Surdal complex, 20 to 50 percent slopes, stony	N	N	Y	Y	Y	Y	Y	Y	Y
742F	Trimad, very stony-Frenchcreek, very stony-Rubble land complex, 25 to 60 percent slopes	N	N	N	N	N	N	N	N	Y
744E	Sigbird, very shallow-Sigbird-Surdal complex, 8 to 25 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
747F	Rubick-Surdal complex, 30 to 70 percent slopes, very stony	N	N	N	N	N	N	N	N	Y
749E	Surdal-Mawspring, bouldery complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y	N	N
750F	Poin, extremely stony-Hungryhill, very stony complex, 20 to 50 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
751F	Sigbird-Tiban-Maurice complex, 20 to 50 percent slopes, very stony	N	Y	Y	Y	Y	Y	Y	N	Y
756D	Passcreek-Poin, stony complex, 6 to 20 percent slopes	N	Y	Y	Y	Y	Y	Y	N	N
757F	Surdal-Sigbird-Mawspring complex, 15 to 60 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	Y	Y
761E	Quincreek, very stony-Whitlash, very stony-Rock outcrop complex, 12 to 35 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
801F	Skaggs, extremely stony-Skaggs, very stony-Rock outcrop complex, 20 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
802E	Starley-Tropal complex, 12 to 40 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
803E	Sieben very cobbly loam, 12 to 30 percent slopes, stony	N	N	N	N	N	N	N	N	N
805C	Sixbeacon, stony-Bronec complex, 2 to 8 percent slopes	N	N	N	N	N	N	Y	N	N
806D	Sieben very cobbly loam, 4 to 12 percent slopes	N	N	N	N	N	N	N	N	N
807E	Pensore, stony-Roto complex, 8 to 30 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
808F	Pensore, stony-Rock outcrop complex, 20 to 45 percent slopes	N	N	N	N	N	N	N	Y	Y
809G	Pensore, stony-Rock outcrop complex, 45 to 70 percent slopes	N	N	N	N	N	N	N	Y	Y
810F	Whitore, very stony-Skaggs-very stony-Rock outcrop complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	Y
811G	Whitore-Skaggs complex, 40 to 70 percent slopes, very stony	N	N	N	N	N	N	Y	Y	Y
812F	Whitore, moist-Skaggs complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	Y	N	Y
813F	Whitore, stony-Whitore complex, 20 to 60 percent slopes	N	N	N	N	N	N	Y	Y	Y
814E	Whitore complex, 12 to 45 percent slopes, stony	N	N	N	N	N	N	Y	N	Y
816G	Whitore, very stony-Tropal, very stony-Rock outcrop complex, 45 to 80 percent slopes	N	N	N	N	N	N	Y	Y	Y
817E	Whitore, very stony-Raynesford complex, 15 to 40 percent slopes	N	N	N	N	N	N	Y	N	N
818D	Raynesford-Whitore, very stony complex, 4 to 20 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
819D	Raynesford-Whitore complex, 4 to 15 percent slopes, stony	N	Y	Y	Y	Y	Y	Y	N	N
820E	Whitore, stony-Tropal, very stony-Raynesford, stony complex, 12 to 45 percent slopes	N	N	Y	Y	Y	Y	Y	Y	Y
823E	Skaggs-Raynesford-Tropal, very stony complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
825E	Skaggs-Whitore complex, 12 to 35 percent slopes, stony	N	N	Y	Y	Y	Y	Y	Y	Y
826E	Raynesford-Whitore-Skaggs complex, 12 to 30 percent slopes	N	Y	Y	Y	Y	Y	Y	Y	Y
830D	Kalsted gravelly sandy loam, 8 to 15 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
901E	Sebud-Poin-Tiban complex, 15 to 45 percent slopes, extremely stony	Y	Y	Y	Y	Y	Y	Y	N	Y
904D	Sebud, stony-Redchief complex, 8 to 25 percent slopes	N	N	N	N	N	N	N	N	N

Erodible Land Index
Survey: MT670-Silverbow Co. Area and, Part of Beaverhead and Jefferson Co.'s
0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind							Water	
		C-20	C-30	C-50	C-60	C-70	C-80	C-90	R-15	R-20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel	Hel
905E	Tigeron, stony-Rubick, very stony complex, 15 to 45 percent slopes	N	N	N	N	Y	Y	Y	N	N
906E	Rubick, very stony-Tigeron, stony complex, 15 to 45 percent slopes	N	N	N	N	Y	Y	Y	N	N
907F	Sebud, extremely stony-Rubick, very stony complex, 25 to 60 percent slopes	N	N	N	N	N	N	N	N	Y
908E	Sebud-Tiban-Ratiopeak complex, 12 to 30 percent slopes, stony	N	N	N	N	N	N	N	N	N
909G	Rubick, rubbly-Rubble land complex, 40 to 75 percent slopes	N	N	N	N	Y	Y	Y	N	Y
910F	Sebud-Ratiopeak complex, 20 to 50 percent slopes, very stony	N	N	N	N	N	N	N	N	Y
911E	Sebud, stony-Adel complex, 12 to 30 percent slopes	N	N	N	N	N	N	N	N	N
912D	Ratiopeak-Redchief complex, 4 to 15 percent slopes, stony	N	N	N	N	N	N	N	N	N
913E	Rubick gravelly sandy loam, 8 to 30 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
920G	Poin, rubbly-Rubble land-Rock outcrop complex, 40 to 80 percent slopes	N	N	N	N	N	N	N	Y	Y
921F	Whitlash-Frenchcreek complex, 20 to 50 percent slopes, rubbly	Y	Y	Y	Y	Y	Y	Y	Y	Y
923F	Whitore, rubbly-Poin, rubbly-Rock outcrop complex, 25 to 60 percent slopes	N	N	N	N	N	N	Y	N	Y
924F	Whitlash, extremely stony-Gnojek, extremely stony-Rock outcrop complex, 20 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y	Y	Y
926F	Rubick-Tigeron complex, 30 to 60 percent slopes, very stony	N	N	Y	Y	Y	Y	Y	N	Y
927E	Tigeron, very stony-Rubick, very stony-Tigeron, stony complex, 15 to 45 percent slopes	N	N	N	N	N	Y	Y	N	N
928E	Mawspring-Maurice complex, 15 to 45 percent slopes, very stony	N	N	N	N	N	N	N	N	N
929F	Rubick, very stony-Poin, extremely bouldery complex, 25 to 60 percent slopes	N	N	Y	Y	Y	Y	Y	N	N
930F	Ratiopeak-Tiban complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	N	N	N	N
931E	Ratiopeak-Monaberg complex, 8 to 30 percent slopes, very stony	N	N	N	N	N	N	N	N	N
932D	Tigeron-Rubick complex, 2 to 15 percent slopes, very stony	N	N	N	N	N	N	Y	N	N
933B	Foxgulch, occasionally flooded-Bearmouth, rarely flooded complex, 0 to 4 percent slopes, stony	N	N	N	N	Y	Y	Y	N	N
935D	Anaconda sandy loam, 8 to 15 percent slopes, moderately impacted	N	Y	Y	Y	Y	Y	Y	N	N
936B	Foxgulch, rarely flooded-Kilgore, occasionally flooded complex, 0 to 4 percent slopes, very stony	N	N	N	N	Y	Y	Y	N	N
939F	Evaro, stony-Tigeron complex, 20 to 50 percent slopes	N	N	N	N	N	N	Y	N	N
940E	Ratiopeak, stony-Tiban, very stony complex, 8 to 25 percent slopes	N	N	N	N	N	N	Y	N	N
941E	Bridger gravelly loam, 12 to 30 percent slopes, stony	N	N	N	N	N	N	N	N	N
942F	Whitlash-Gnojek complex, 20 to 60 percent slopes, very stony	Y	Y	Y	Y	Y	Y	Y	Y	Y
943D	Bridger-Adel complex, 4 to 15 percent slopes	N	N	N	N	N	N	N	N	N
992	Pits, borrow	0	0	0	0	0	0	0	0	0
994	Pits, mine	0	0	0	0	0	0	0	0	0
995	Dumps, garbage	0	0	0	0	0	0	0	0	0
997	Dumps, mine	0	0	0	0	0	0	0	0	0
998	Rock outcrop and Rubble land	0	0	0	0	0	0	0	0	0
998	Rock outcrop and Rubble land	0	0	0	0	0	0	0	0	0
GP	Gravel pits	0	0	0	0	0	0	0	0	0
M-W	Miscellaneous water	0	0	0	0	0	0	0	0	0
W	Water	0	0	0	0	0	0	0	0	0

Approved
State Soil Scientist
5/20/09

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1101E	Monaberg, stony-Libeg, bouldery, complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y
1120E	Quaint channery loam, 8 to 35 percent slopes, very stony	Y	Y	Y	Y	Y	Y	Y
1121E	Quaint-Rock outcrop-Redfist complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y
1122D	Quaint-Redfist channery loams, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1131C	Redfist-Quaint channery loams, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
1131D	Redfist-Quaint channery loams, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
1132D	Redfist, bouldery-Perma, bouldery-Rock outcrop complex, 2 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
113C	Amesha-Amesha, stony, complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
113D	Amesha-Amesha, stony, complex, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1141D	Devilfence very channery loam, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1142E	Devilfence-Rock outcrop-Wilspring complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
114D	Amesha-Crago-Shoddy complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
114E	Amesha-Crago-Shoddy complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	Y	Y
1151C	Willowsprings channery loam	Y	Y	Y	Y	Y	N	N
1152D	Wilspring-Devilfence complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1152E	Wilspring-Devilfence-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1153C	Wilspring-Quincreek-Devilfence complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
115A	Amesha gravelly loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
115C	Amesha gravelly loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
115D	Amesha gravelly loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
115E	Amesha gravelly loam, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
1161E	Yreka, bouldery-Hoyt, bouldery-Shaboom, very bouldery, complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	N	Y
116A	Amesha loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
116C	Amesha loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
116D	Amesha loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
117E	Amesha-Cabbart-Bronec cobbly loams, 4 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
1180E	Farnuf loam, 15 to 35 percent slopes, stony	N	N	N	Y	Y	N	Y
1181D	Shawa-Yegan-Castner complex	N	Y	Y	Y	Y	N	Y
118C	Amesha cobbly loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
1191E	Silverchief very cobbly clay loam, 8 to 35 percent slopes, bouldery	N	Y	Y	Y	Y	N	Y
11B	Amesha silt loam, 1 to 4 percent slopes	N	Y	Y	Y	Y	N	N
11C	Amesha silt loam, 4 to 8 percent slopes	N	Y	Y	Y	Y	N	N
11D	Amesha silt loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1210C	Ferball clay loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
121E	Maiden-Lap-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1220C	Quigley sandy loam	N	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1221C	Quigley-Duffy complex	N	Y	Y	Y	Y	N	N
122D	Maiden-Lap-Windham complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
122E	Maiden-Lap-Windham complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
1231D	Vebar Variant-Castner-Tally complex	Y	Y	Y	Y	Y	N	Y
123E	Maiden, very stony-Rock outcrop-Lap, very stony, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
1241D	Baxendale Variant-Castner complex	Y	Y	Y	Y	Y	N	Y
1250E	Work very cobbly clay loam, 8 to 25 percent slopes	N	N	N	N	Y	N	N
1251C	Work-Gregroy complex	N	Y	Y	Y	Y	N	N
1260D	Lolo gravelly sandy loam	Y	Y	Y	Y	Y	N	N
1271D	Placerton-Farnuf-Connieo complex, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1281D	Crackerville-Catgulch, bouldery-Rock outcrop complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
1282D	Crackerville-Bielenberg-Catgulch, bouldery, complex, 8 to 20 percent slopes	Y	Y	Y	Y	Y	N	N
1283E	Crackerville-Bielenberg-Catgulch, bouldery, complex, 20 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1284D	Boperan-Perma-Castner complex	Y	Y	Y	Y	Y	N	Y
1291D	Broadmoor-Sharrott-Rock outcrop complex	Y	Y	Y	Y	Y	N	Y
1292E	Boradmoor-Broadmoor-Sharrott complex	N	Y	Y	Y	Y	N	Y
1321B	Beaverell, very stony-Beaverell-Sieberell, stony, complex, 1 to 4 percent slopes	Y	Y	Y	Y	Y	N	N
1321D	Beaverell, very stony-Sieberell, stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1341D	Windham very gravelly loam, 2 to 15 percent slopes, very stony	Y	Y	Y	Y	Y	N	N
1351D	Perma stony loam, 2 to 15 percent slopes, very bouldery	Y	Y	Y	Y	Y	N	N
1361E	Lumpgulch, bouldery-Rock outcrop-Elmark, bouldery, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1371D	Harrol-Tolman Variant complex	Y	Y	Y	Y	Y	N	Y
1381D	Jeffcity, stony-Connieo, stony-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
1391B	Bronec fine sandy loam, 1 to 4 percent slopes	Y	Y	Y	Y	Y	N	N
151D	Delpoint-Abor complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
182D	Brocko-Amesha complex, 4 to 15 percent slopes	N	Y	Y	Y	Y	N	N
183E	Brocko-Rock outcrop-Rencot complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
18A	Brocko silt loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
18C	Brocko silt loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
18D	Brocko silt loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
18E	Brocko silt loam, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
191C	Cabbart-Shoddy-Amesha complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
191D	Cabbart-Shoddy-Amesha complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
191E	Cabbart-Shoddy-Amesha complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
192C	Cabbart clay loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
193E	Cabbart-Haxby loams, 8 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
201E	Windham-Rock outcrop-Lap, very stony, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
202D	Windham-Judell complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
202E	Windham-Judell complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
203B	Windham very gravelly loam	Y	Y	Y	Y	Y	N	N
203D	Windham gravelly loam, 4 to 15 percent slopes, stony	Y	Y	Y	Y	Y	N	N
203E	Windham gravelly loam, 15 to 35 percent slopes, stony	Y	Y	Y	Y	Y	Y	Y
204D	Windham-Maiden-Lap complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
204E	Windham, stony-Maiden, very stony-Lap, very stony, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
205E	Windham very cobbly loam, 4 to 35 percent slopes, very stony	Y	Y	Y	Y	Y	N	Y
206E	Widnam complex	Y	Y	Y	Y	Y	N	Y
207E	Windham, stony-Lap, very stony-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
208D	Windham-Judell gravelly loams, 8 to 25 percent slopes, stony	Y	Y	Y	Y	Y	N	Y
208D	Windham-Judell gravelly loams, 8 to 25 percent slopes, stony	Y	Y	Y	Y	Y	N	N
209C	Windham-Judell complex	Y	Y	Y	Y	Y	N	N
209D	Windham-Judell complex	Y	Y	Y	Y	Y	N	Y
20C	Windham gravelly loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
20D	Windham gravelly loam, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
20E	Windham very gravelly loam, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
221A	Zatony clay loam, 0 to 2 percent slopes, wet	N	Y	Y	Y	Y	N	N
222A	Zatoville silt loam	N	Y	Y	Y	Y	N	N
22A	Zatoville clay loam	Y	Y	Y	Y	Y	N	N
22C	Zatony clay loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
241E	Whitlash, very stony-Rock outcrop-Perma, stony, complex, 2 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
242E	Castner-Blaincreek complex	Y	Y	Y	Y	Y	N	Y
243E	Castner-Boperan-Rock outcrop complex	Y	Y	Y	Y	Y	N	Y
244D	Castner-Baxendale-Rock outcrop complex	Y	Y	Y	Y	Y	N	N
245E	Tolbert, very stony-Rock outcrop-Absarook, stony, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
251B	Cozberg sandy loam, 1 to 4 percent slopes, stony	Y	Y	Y	Y	Y	N	N
251D	Cozberg sandy loam, 4 to 15 percent slopes, stony	Y	Y	Y	Y	Y	N	N
25A	Cozberg sandy loam	Y	Y	Y	Y	Y	N	N
25C	Cozberg sandy loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
261D	Crago-Brocko complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
261E	Crago-Brocko complex, 15 to 60 percent slopes	Y	Y	Y	Y	Y	N	Y
263D	Crago-Rock outcrop-Pensore complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	N	N
264C	Crago-Amesha complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
264D	Crago-Amesha complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
264E	Crago-Amesha complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
265B	Crago gravelly loam, 1 to 4 percent slopes, stony	Y	Y	Y	Y	Y	N	N
265D	Crago gravelly loam, 4 to 15 percent slopes, stony	Y	Y	Y	Y	Y	N	N
265E	Crago very cobbly loam, 15 to 45 percent slopes, very stony	Y	Y	Y	Y	Y	N	Y
266D	Crago, stony-Crago complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
266E	Crago, stony-Crago complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
268C	Crago-Amesha cobbly loams, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
269D	Crago, Very stony, and Crago, rubbly, soils, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
26C	Crago gravelly loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
26D	Crago gravelly loam, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
26E	Crago gravelly loam, 15 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
271C	Bronec-Amesha complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
271D	Bronec-Amesha complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
271E	Bronec-Amesha-Bronec complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
272E	Bronec-Geohrock-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
273D	Bronec-Shoddy-Amesha complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
273E	Bronec-Shoddy-Amesha complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
274A	Bronec complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
274C	Bronec complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
274D	Bronec complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
274E	Bronec-Bronec, very stony, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
275A	Bronec gravelly loam, 1 to 4 percent slopes, stony	Y	Y	Y	Y	Y	N	N
275D	Bronec very gravelly loam, 4 to 15 percent slopes, very stony	Y	Y	Y	Y	Y	N	N
275E	Bronec very gravelly loam, 15 to 35 percent slopes, very stony	Y	Y	Y	Y	Y	N	Y
276C	Bronec gravelly loam, 2 to 8 percent slopes, saline	Y	Y	Y	Y	Y	N	N
277C	Bronec-Amesha cobbly loams, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
278E	Bronec-Rencot-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
27A	Bronec gravelly loam, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
27C	Bronec gravelly loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
27D	Bronec gravelly loam, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
27E	Bronec gravelly loam, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
291C	Sieben complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
291D	Sieben complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
292C	Sieben-Varney cobbly loams, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
293D	Sieben cobbly loam, 4 to 15 percent slopes, stony	N	Y	Y	Y	Y	N	N
294C	Sieben, stony-Sieberell, very stony, complex, 2 to 15 percent slopes	N	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
295D	Sieben cobbly loam, 4 to 15 percent slopes, bouldery	N	Y	Y	Y	Y	N	N
296D	Sieberell-Sieben-Beaverell complex, 4 to 15 percent slopes, stony	N	Y	Y	Y	Y	N	N
311D	Beenom, stony-Wimper-Whitlash, very stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
31C	Beenom loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	Y
321A	Fairway-Meadowcreek complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
325A	Fairway-Nestley clay loams, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
326A	Fairway-Moltoner complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
331C	Geohrock-Bronec gravelly loams, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
331D	Geohrock-Bronec gravelly loams, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
332D	Geohrock-Sappington complex, 4 to 15 percent slopes, stony	N	Y	Y	Y	Y	N	N
333B	Geohrock cobbly loam	N	Y	Y	Y	Y	N	N
334D	Geohrock, stony-Bronec, very stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
33B	Geohrock gravelly loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
33E	Geohrock cobbly clay loam, 15 to 35 percent slopes, stony	N	Y	Y	Y	Y	N	N
341A	Pieriver-Cardwell-Riverrun loams, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
342A	Handke fine sandy loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
34A	Glendive Variant sandy loam	N	Y	Y	Y	Y	N	N
35A	Hanly sandy loam	Y	Y	Y	Y	Y	N	N
361D	Udecide-Varney-Walbert complex, 4 to 25 percent slopes	N	Y	Y	Y	Y	N	N
362C	Udecide-Varney sandy clay loams, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
371A	Havre-Ryell-Handke complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
372A	Havre loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
37A	Pieriver silt loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
381C	Kalsted gravelly sandy loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
381D	Kalsted gravelly sandy loam, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
382D	Kalsted gravelly sandy loam, 4 to 15 percent slopes, stony	Y	Y	Y	Y	Y	N	N
38A	Kalsted sandy loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
38C	Kalsted sandy loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
38D	Kalsted sandy loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
391C	Musselshell-Crago gravelly loams, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
391D	Musselshell-Crago complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
392D	Musselshell-Crago complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
393B	Musselshell-Amesha complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
394B	Musselshell-Crago cobbly loams, 1 to 4 percent slopes	Y	Y	Y	Y	Y	N	N
401A	Moltoner silty clay loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
40A	Moltoner loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
41	Bronec, Amesha, and Riverrun, channeled, soils, 0 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
411A	Nestley loam, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
412A	Nesda Variant-Rivra complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
413A	Nestley-Riverrun-Pieriver complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
41A	Nesda Variant, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
421E	Perma, stony-Whitlash, very stony, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
422F	Perma, very stony-Whitlash, very stony-Rock outcrop complex, 15 to 45 percent slopes, moist	Y	Y	Y	Y	Y	Y	Y
423C	Wimper loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
423D	Wimper loam, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
423E	Wimper loam, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
424D	Wimper-Wimper, stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
424E	Wimper-Wimper, stony, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
425E	Wimper gravelly loam, 8 to 35 percent slopes, stony	Y	Y	Y	Y	Y	N	Y
427E	Perma-Whitlash complex, 15 to 35 percent slopes, bouldery	Y	Y	Y	Y	Y	N	Y
428D	Perma-Shawa-Castner complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
429C	Perma-Quigley Varient complex	Y	Y	Y	Y	Y	N	N
42D	Perma cobbly loam, 4 to 15 percent slopes, stony	Y	Y	Y	Y	Y	N	N
42E	Perma cobbly loam, 15 to 25 percent slopes, stony	Y	Y	Y	Y	Y	N	Y
441F	Warneke-Warneke, very stony-Rock outcrop association, 8 to 60 percent slopes	Y	Y	Y	Y	Y	N	Y
451A	Geohrock cobbly clay loam, 1 to 4 percent slopes, stony	N	Y	Y	Y	Y	N	N
461D	Absarook-Beenom complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
471A	Cardwell-Pieriver complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
481A	Riverrun gravelly sandy loam, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
482A	Riverrun-Cardwell complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
483A	Riverrun, Handke, and Ryell soils, 0 to 2 percent slopes, channeled	Y	Y	Y	Y	Y	N	N
48A	Riverrun sandy loam, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
492D	Roto-Pensore-Crago complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
492E	Roto-Pensore-Crago complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
493D	Pensore-Rock outcrop-Roto complex, 2 to 25 percent slopes	Y	Y	Y	Y	Y	N	N
511C	Haxby-Amesha-Rencot complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
511E	Rootel-Amesha-Rencod complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
521A	Cardwell-Riverrun complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
522A	Ryell-Riverrun complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
523A	Cardwell-Riverrun-Pieriver complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
524A	Cardwell loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
525A	Cardwell-Riverrun complex, 0 to 2 percent slopes, saline	Y	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
52A	Ryell loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
532C	Sappington-Amesha complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
534C	Sappington-Geohrock complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
534D	Sappington-Geohrock complex, 8 to 15 percent slopes	N	N	N	N	Y	N	N
536A	Sappington-Amesha complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
536C	Sappington-Amesha complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
539B	Sappington-Amesha complex, 2 to 8 percent slopes, cobbly	N	Y	Y	Y	Y	N	N
539C	Sappington-Amesha complex, 2 to 8 percent slopes, stony	N	Y	Y	Y	Y	N	N
541D	Whitlash, very stony-Brickner, stony-Rock outcrop complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	N	N
562C	Trudau-Bronec, saline, complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
563A	Trudau Variant, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
56A	Trudau loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
56B	Trudau loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
582E	Whitecow, bouldery-Shawmut, very bouldery-Rock outcrop complex, 15 to 45 percent slopes, warm	Y	Y	Y	Y	Y	N	Y
583E	Whitecow-Warneke complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
58E	Whitecow Channery loam, 15 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y
631D	Rencot-Rock outcrop-Rencot, stony, complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y
632C	Rencot-Lahood-Rock outcrop complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
632D	Rencot-Lahood-Rock outcrop complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
645B	Varney-Sieben complex, 1 to 4 percent slopes, stony	N	Y	Y	Y	Y	N	N
645D	Varney, stony-Sieben, very stony, complex, 4 to 15 percent slopes	N	Y	Y	Y	Y	N	N
645E	Varney, stony-Sieben, very stony, complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
651C	Judell gravelly loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
652C	Judell cobbly loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
661C	Judith loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
662B	Judell gravelly loam, 1 to 4 percent slopes, very stony	N	Y	Y	Y	Y	N	N
66C	Judith gravelly loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
66D	Judith gravelly loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
67A	Manhattan Variant silt loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
691A	Meadowcreek, Clunton, and Cardwell soils, 0 to 2 percent slopes, channeled	N	Y	Y	Y	Y	N	N
692A	Meadowcreek-Nestley-Riverrun complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
69A	Meadowcreek silty clay loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
70C	Benz loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
711D	Assiniboine sand loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
712E	Rock River Variant sandy loam, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
71C	Raghorn sandy loam, 4 to 8 percent slopes	N	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
71D	Raghorn sandy loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
721E	Zbart-Bondoe-Brocko complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
732D	Martinsdale-Shawmut, stony-Martinsdale, bouldery, complex, 4 to 25 percent slopes	N	N	N	N	Y	N	N
735C	Martinsdale-Absarook-Whitlash complex, 2 to 8 percent slopes, stony	Y	Y	Y	Y	Y	N	N
741D	Hilger complex, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
742E	Shawmut, stony-Martinsdale, very stony, complex, 4 to 25 percent slopes	N	N	N	N	Y	N	N
743E	Hilger Variant-Farnuf complex, 15 to 35 percent slopes, bouldery	N	Y	Y	Y	Y	N	Y
744E	Shawmut, bouldery-Shawmut, stony-Tolbert, bouldery, complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
74D	Shawmut gravelly loam, 4 to 15 percent slopes, bouldery	N	Y	Y	Y	Y	N	N
751C	Sixbeacon gravelly sandy loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
752B	Sixbeacon-Vendome complex, 1 to 4 percent slopes	Y	Y	Y	Y	Y	N	N
752D	Sizbeacon-Vendome complex, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
753C	Sixbeacon-Cozberg complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
754D	Sixbeacon-Cozberg, stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
755A	Vendome very cobbly loam, 0 to 4 percent slopes, very stony	N	Y	Y	Y	Y	N	N
756C	Sixbeacon Variant complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
75C	Sixbeacon loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
75D	Sixbeacon loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
76A	Absay silty clay loam, 0 to 4 percent slopes	N	Y	Y	Y	Y	N	N
781A	Vendome sandy loam, 0 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
781C	Vendome sandy loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
782A	Vendome sandy loam, 0 to 8 percent slopes, stony	Y	Y	Y	Y	Y	N	N
782C	Vendome sandy loam, 2 to 8 percent slopes, stony	Y	Y	Y	Y	Y	N	N
78C	Vendome very cobbly loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
78E	Vendome very cobbly loam, 15 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
79A	Chinook sandy loam, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
79C	Chinook sandy loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
79D	Chinook sandy loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
79E	Chinook sandy loam, 15 to 25 percent slopes	N	Y	Y	Y	Y	N	N
811C	Ethridge, saline-Zatony clay loams, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
821C	Rothiemay loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
82C	Rothiemay very gravelly loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
83C	Shoddy silty clay loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
83D	Shoddy silty clay loam, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
851D	Walbert-Shoddy-Cabbart complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
851F	Walbert-Shoddy-Cabbart complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
852C	Walbert sandy clay loam, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
859E	Yetull loamy fine sand, 15 to 30 percent slopes	Y	Y	Y	Y	Y	N	Y
85D	Walbert coarse sandy loam, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
86C	Marias silty clay loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
871C	Kobar cobbly clay loam	N	Y	Y	Y	Y	N	N
88C	Lahood loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
88D	Lahood loam, 8 to 15 percent slopes	Y	Y	Y	Y	Y	N	Y
9	Bronec, Riverrun, Channeled, and Amesha soils, 0 to 8 percent slopes	N	Y	Y	Y	Y	N	N
901A	Attewan-Nippt complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
911C	Absarook-Martinsdale, stony, complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
915C	Quincreek channery loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
941E	Tigeron, bouldery-Tigeron, very bouldery, complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	N	N
942E	Tigeron extremely gravelly loam, 15 to 35 percent slopes, bouldery	N	Y	Y	Y	Y	N	Y
99E	Libeg very gravelly loam, 15 to 35 percent slopes, bouldery	N	Y	Y	Y	Y	N	Y
1	Riverwash	0	0	0	0	0	0	0
2	Rubble land-Rock outcrop association	0	0	0	0	0	0	0
3	Dumps, mine	0	0	0	0	0	0	0
4	Bronec, Clunton, Channeled, and Amesha soils, 0 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
5	Borrow areas and Gravel pits	0	0	0	0	0	0	0
6	Wetsand, Cardwell, and Clunton soils, 0 to 8 percent slopes, channeled	Y	Y	Y	Y	Y	N	N
13A	Anamac loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
13C	Anamac loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
21A	McKenton silty clay loam, 0 to 2 percent slopes	N	N	N	N	N	N	N
23A	McKenton silt loam, 0 to 2 percent slopes	N	N	N	N	N	N	N
30A	Dougcliff mucky peat, 0 to 1 percent slopes	N	Y	Y	Y	Y	N	N
32A	Fairway silt loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
47A	Cardwell silty clay loam, 0 to 2 percent slopes	N	N	Y	Y	Y	N	N
53C	Sappington gravelly clay loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
60C	Watne loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
64A	Varney sandy clay loam, 0 to 2 percent slopes	N	N	N	Y	Y	N	N
64C	Varney sandy clay loam, 2 to 8 percent slopes	N	N	N	Y	Y	N	N
65C	Judell loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
68D	Bondoe channery loam, 4 to 15 percent slopes	N	N	N	N	Y	N	N
72F	Zbart-Rock outcrop association, 25 to 70 percent slopes	N	N	N	N	N	Y	Y
73C	Martinsdale loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
77F	Rock outcrop-Pensore, stony-Crago, stony, association, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
80A	Floweree silt loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
80C	Floweree silt loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
81A	Ethridge clay loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
81C	Ethridge clay loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
84A	Clunton-Faith-Dougcliff complex, 1 to 4 percent slopes	N	N	N	N	Y	N	N
87C	Kobarter clay loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
87D	Kobarter clay loam, 8 to 15 percent slopes	N	N	N	N	Y	N	N
92D	Clunton, Cometcrik, and Perma, stony, soils, 0 to 15 percent slopes	N	N	N	Y	Y	N	N
99D	Libeg gravelly loam, 4 to 15 percent slopes, bouldery	N	N	N	N	Y	N	N
121F	Maiden-Lap-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
122F	Maiden-Lap-Windham complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
123F	Maiden, very stony-Rock outcrop-Lap, very stony, complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
125D	Maiden-Lap-Windham complex, 4 to 15 percent slopes, warm	Y	Y	Y	Y	Y	N	N
125E	Maiden-Lap-Windham complex, 15 to 35 percent slopes, warm	Y	Y	Y	Y	Y	Y	Y
126F	Maiden, very stony-Rock outcrop-Lap, very stony, complex, 35 to 60 percent slopes, warm	Y	Y	Y	Y	Y	Y	Y
132A	Anamac silt loam, 0 to 2 percent slopes, saline	N	N	N	N	Y	N	N
171D	Branham-Opitz-Tuggle complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
171E	Branham-Opitz-Tuggle complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
172D	Branham-Clugulch-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
172E	Branham-Clugulch-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
173F	Branham, stony-Tuggle, very stony-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
184E	Brocko-Rock outcrop-Bronec, very stony, complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	N	Y
194E	Cabbart-Bronec, stony-Rencot, very stony complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
195E	Cabbart, very stony-Bronec, stony-Rock outcrop complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
195F	Cabbart, very stony-Rock outcrop-Bronec, very stony, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
201F	Windham, very stony-Rock outcrop-Lap, very stony, complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
204F	Windham, very stony-Maiden, very stony-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
205F	Windham very cobbly loam, 35 to 60 percent slopes, very stony	Y	Y	Y	Y	Y	Y	Y
206F	Windham-Windham, stony, complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
207F	Windham, stony-Lap, very stony-Rock outcrop complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
211A	Clunton silty clay loam, 0 to 2 percent slopes	N	N	N	N	N	N	N
231A	Ledger-Moltoner-McKenton complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
232A	Clunton-Wetsand-Bonebasin complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
233A	Ledger-Wetsand, saline, complex, 0 to 2 percent slopes	Y	Y	Y	Y	Y	N	N
241F	Whitlash, very stony-Rock outcrop-Perma, very stony, complex, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
263F	Crago, stony-Rock outcrop-Pensore, stony, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
264F	Crago-Amesha complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
267F	Crago, very stony-Pensore, stony-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
271F	Bronec-Amesha-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
297D	Sieben, very stony-Sieben, rubbly, complex, 2 to 25 percent slopes	N	N	N	N	Y	N	N
297F	Sieben, rubbly-Sieben, very stony, complex, 15 to 60 percent slopes	N	N	N	N	Y	N	N
311F	Beenom, stony-Wimper-Whitlash, very stony, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
322A	Fairway loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
323A	Fairway-McKenton silt loams, 0 to 2 percent slopes	N	N	N	N	Y	N	N
324A	Fairway clay loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
327A	Faith loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
328A	Faith loam, 0 to 2 percent slopes, cool	N	N	N	N	Y	N	N
329C	Faith-Slickens complex, 0 to 8 percent slopes, impacted	Y	Y	Y	Y	Y	N	N
421F	Perma-Whitlash complex, 35 to 60 percent slopes, very stony	N	N	N	Y	Y	Y	Y
426F	Wimper-Whitlash association, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
429E	Perma, very stony-Perma, rubbly-Rock outcrop complex, 8 to 35 percent slopes	N	N	N	N	Y	N	N
492F	Roto-Pensore-Crago complex, 35 to 60 percent slopes, stony	Y	Y	Y	Y	Y	Y	Y
531C	Sappington very cobbly clay loam, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
533A	Sappington clay loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
533C	Sappington clay loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
533D	Sappington clay loam, 8 to 15 percent slopes	N	N	N	N	Y	N	N
537B	Sappington loam, 1 to 4 percent slopes, stony	N	N	N	N	Y	N	N
537D	Sappington loam, 4 to 15 percent slopes, stony	N	N	N	N	Y	N	N
538C	Sappington gravelly loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
541E	Whitlash, very stony-Brickner, stony-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
551E	Brickner, stony-Whitlash, very stony-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
552F	Brickner, very bouldery-Rock outcrop-Tolbert, very bouldery, association, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
553F	Brickner, very stony-Wickes, very bouldery-Rock outcrop complex, 15 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
554F	Brickner, very stony-Rock outcrop-Mocmont, stony, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
564C	Trudau-Benz clay loams, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
581E	Whitecow, stony-Warneke, very stony-Rock outcrop complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
581F	Whitecow, very stony-Warneke, very stony-Rock outcrop complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
584F	Whitecow-Whitecow, stony-Warneke complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
585E	Whitecow, bouldery-Shawmut, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
591F	Windham-Rock outcrop-Warneke complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
631E	Rencot-Rencot, very stony-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
631F	Rencot-Bronec-Rock outcrop complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
632E	Rencot, very stony-Lahood, stony-Rock outcrop complex, 25 to 45 percent slopes	N	N	N	N	N	Y	Y
633D	Rencot, very stony-Bronec, very stony-Rock outcrop complex, 4 to 25 percent slopes	N	N	N	N	N	N	N
634E	Rencot, very stony-Rock outcrop-Bronec, very stony, complex, 25 to 45 percent slopes	N	N	N	N	N	Y	Y
641C	Varney gravelly loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
642C	Varney clay loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
642D	Varney clay loam, 8 to 15 percent slopes	N	N	N	N	Y	N	N
643A	Varney cobbly loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
643C	Varney cobbly loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
644C	Varney complex, 2 to 15 percent slopes, gullied	N	N	N	N	Y	N	N
646C	Varney loam, 2 to 8 percent slopes, stony	N	N	N	N	Y	N	N
655C	Judell gravelly loam, 2 to 8 percent slopes, warm	Y	Y	Y	Y	Y	N	N
713E	Raghorn-Ethridge-Kalsted complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	N
713F	Raghorn-Ethridge-Kalsted complex, 35 to 70 percent slopes	N	Y	Y	Y	Y	Y	Y
731C	Martinsdale, stony-Martinsdale-Hilger complex, 2 to 8 percent slopes	N	N	N	N	Y	N	N
734D	Martinsdale loam, 4 to 15 percent slopes, very stony	N	N	N	N	Y	N	N
736C	Martinsdale-Work complex, 2 to 8 percent slopes	N	N	N	N	Y	N	N
745D	Shawmut-Wickes-Gnojek complex, 2 to 15 percent slopes, bouldery	Y	Y	Y	Y	Y	N	N
745E	Shawmut, bouldery-Shawmut, very bouldery-Tolbert, bouldery, complex, 15 to 45 percent slopes, dry	N	N	N	Y	Y	N	N
746E	Shawmut-Tolbert complex, 8 to 35 percent slopes	N	N	N	N	Y	N	N
747E	Shawmut, stony-Tolbert, very stony, complex, 15 to 35 percent slopes	N	N	N	N	Y	N	N
748E	Shawmut, stony-Wickes, very stony, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
773F	Rock outcrop-Pensore association, 15 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
774F	Rock outcrop-Whitlash, bouldery, association, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
775F	Rock outcrop-Lap-Lap, very stony, association, 15 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
776D	Rock outcrop-Devilfence association, 2 to 25 percent slopes	Y	Y	Y	Y	Y	N	N
777E	Rock outcrop-Clugulch-Bobowic complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
777F	Rock outcrop-Clugulch-Bobowic complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
778E	Rock outcrop-Kounter, very bouldery-Jeffcity, bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
791C	Chinook sandy clay loam, 2 to 8 percent slopes, saline	N	N	N	Y	Y	N	N
831E	Shoddy-Cabbart-Kobarter complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y
832E	Shoddy-Rock outcrop-Delpoint complex, 2 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y
841A	Clunton loam, 0 to 2 percent slopes	N	N	N	N	Y	N	N
858E	Yetull-Yetull, stony, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
859C	Yetull loamy fine sand, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
872E	Kobarter-Abor, stony, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
943F	Tigeron, stony-Tigeron very stony, complex, 25 to 60 percent slopes	N	N	N	N	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
944E	Tigeron, very bouldery-Redfern, bouldery-Rock outcrop complex, 15 to 45 percent slopes, warm	N	N	N	N	N	N	N
945E	Tigeron, very bouldery-Redfern, bouldery-Rock outcrop complex, 15 to 45 percent slopes, dry	N	N	N	N	N	N	N
946F	Tigeron, very stony-Redfern, rubbly-Rock outcrop complex, 25 to 60 percent slopes	N	N	N	N	Y	N	N
947F	Tigeron, very stony-Redfern, rubbly-Rock outcrop complex, 25 to 60 percent slopes, dry	N	N	N	N	Y	N	N
952F	Redfern, bouldery-Rock outcrop-Tigeron, very bouldery, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
953F	Redfern, rubbly-Rock outcrop-Rubble land association, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
954F	Redfern, rubbly-Rock outcrop-Tigeron, very bouldery, complex, 35 to 60 percent slopes	N	N	N	N	N	Y	Y
963E	Elve-Warwood complex, 15 to 45 percent slopes, stony	N	Y	Y	Y	Y	N	N
964E	Elve, very stony-Elve, rubbly-Cowood, rubbly, complex, 15 to 35 percent slopes	N	N	N	N	N	N	N
964F	Elve, very stony-Cowood, rubbly-Rock outcrop complex, 35 to 60 percent slopes, cool	N	N	N	N	N	N	Y
965E	Elve, very stony-Cowood, rubbly, complex, 15 to 35 percent slopes	N	N	N	N	N	N	N
965F	Elve, very stony-Cowood, rubbly-Rock outcrop complex, 35 to 60 percent slopes, dry	N	N	N	N	N	N	Y
966E	Elve, very stony-Rock outcrop-Rubble land complex, 8 to 35 percent slopes	N	N	N	N	N	N	N
967F	Elve, very stony-Cowood, rubbly-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	N	N	Y
968E	Elve, stony-Worock complex, 15 to 35 percent slopes	N	N	N	Y	Y	N	Y
968F	Elve, stony-Worock complex, 35 to 60 percent slopes	N	N	N	Y	Y	Y	Y
969F	Elve, bouldery-Worock-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	N	N	Y
971F	Cowood, rubbly-Rock outcrop association, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
972F	Cowood, very bouldery-Kimpton, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	N	N	N	N	N	Y	Y
973D	Cowood, very stony-Elve, very stony-Rock outcrop complex, 4 to 25 percent slopes	N	N	N	N	N	N	N
982F	Kimpton, very bouldery-Rock outcrop-Tiban, very bouldery, complex, 25 to 50 percent slopes	N	N	N	N	N	Y	Y
991E	Libeg loam, 15 to 35 percent slopes, bouldery	N	Y	Y	Y	Y	Y	Y
992E	Libeg, very bouldery-Libeg, bouldery-Nieman, bouldery, complex, 15 to 45 percent slopes	N	N	N	N	Y	N	N
993D	Ratiopeak-Tiban gravelly loams, 4 to 15 percent slopes, bouldery	N	N	N	N	Y	N	N
994E	Libeg, stony-Nieman, bouldery, complex, 15 to 45 percent slopes	N	N	N	N	Y	N	N
996D	Libeg-Monaberg gravelly loams, 2 to 15 percent slopes, bouldery	N	N	N	N	Y	N	N
997E	Libeg, stony-Monaberg-Adel complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y
998E	Libeg-Nieman, stony, complex, 8 to 25 percent slopes	N	Y	Y	Y	Y	N	N
999E	Libeg, very stony-Libeg, very bouldery, complex, 4 to 25 percent slopes	N	N	N	N	N	N	N
999F	Libeg, very stony-Libeg, rubbly, association, 25 to 60 percent slopes	N	N	N	N	N	N	N
1003E	Tiban, bouldery-Cheadle, very bouldery, complex, 15 to 35 percent slopes	N	N	N	N	Y	N	N
1004E	Tiban, rubbly-Tiban, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	N	N	N	N	N	N	N
1102B	Adel-Libeg, stony, complex, 1 to 4 percent slopes	N	N	N	N	Y	N	N
1103D	Adel-Libeg complex, 4 to 15 percent slopes, stony	N	N	N	N	Y	N	N
1104C	Adel loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
1121F	Quaint-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1142F	Devilfence-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1143F	Deville-Wilde-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1146E	Deville-Rock outcrop-Wilde complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1154E	Wilde-Deville-Vigilante complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
1154F	Wilde, stony-Vigilante-Deville, very stony, complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
1162E	Yreka very cobbly loam, 15 to 35 percent slopes, bouldery	N	N	N	N	Y	N	N
1163F	Ymark, very bouldery-Elmark, very bouldery-Rock outcrop complex, 25 to 60 percent slopes	N	N	N	N	N	N	Y
1164F	Yreka-Brickner, stony, complex, 35 to 70 percent slopes	N	Y	Y	Y	Y	N	Y
1170E	Whitlash-Whitlash, stony-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1171F	Castner, bouldery-Rock outcrop complex, 25 to 50 percent slopes	N	N	N	N	N	Y	Y
1172F	Castner, very bouldery-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	N	Y	Y
1182C	Breeton gravelly loam, 2 to 8 percent slopes	N	N	N	Y	Y	N	N
1222C	Martinsdale-Martinsdale, stony-Shawmut complex, 2 to 8 percent slopes	N	N	N	Y	Y	N	N
1222E	Martinsdale-Martinsdale, stony-Shawmut complex, 15 to 35 percent slopes	N	N	N	Y	Y	N	N
1223D	Martinsdale-Shawmut complex, 2 to 15 percent slopes, bouldery	N	N	N	Y	Y	N	N
1242D	Baxton-Connieo, very bouldery-Rock outcrop complex, 4 to 15 percent slopes, moist	Y	Y	Y	Y	Y	N	N
1242E	Baxton-Connieo, very bouldery-Rock outcrop complex, 15 to 35 percent slopes, moist	Y	Y	Y	Y	Y	N	N
1243D	Baxton-Connieo coarse sandy loams, 4 to 15 percent slopes, bouldery	Y	Y	Y	Y	Y	N	N
1244D	Baxton-Connieo, very bouldery-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1244E	Baxton-Connieo-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
1245E	Baxton-Connieo complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
1246E	Baxton, stony-Breeton, bouldery-Catgulch, very stony, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
1247E	Baxton-Kellygulch-Connieo complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1250C	Work cobbly clay loam, 2 to 8 percent slopes, stony	N	N	N	N	Y	N	N
1272D	Placerton-Connieo-Jeffcity complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1273E	Placerton-Farnuf-Breeton complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
1275D	Placerton-Farnuf-Connieo gravelly sandy clay loams, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1275E	Placerton-Farnuf-Connieo complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y
1276D	Placerton-Connieo-Jeffcity complex, 4 to 15 percent slopes, warm	Y	Y	Y	Y	Y	N	N
1277C	Placerton-Jeffcity complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
1280D	Crackerville-Catgulch complex, 2 to 15 percent slopes, bouldery	Y	Y	Y	Y	Y	N	N
1286E	Crackerville-Bielenberg-Catgulch, bouldery, complex, 15 to 35 percent slopes, warm	Y	Y	Y	Y	Y	N	Y
1287E	Clancy, very stony-Crampton, bouldery-Bielenberg, very stony, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1322D	Beaverell, stony-Beaverell, rubbly-Sieberell, stony, complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1331A	Bonebasin-Wetsand complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
1342E	Windham-Lap very cobbly loams, 15 to 45 percent slopes, bouldery	Y	Y	Y	Y	Y	N	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1343D	Windham-Judell very cobbly loams, 4 to 15 percent slopes, bouldery	Y	Y	Y	Y	Y	N	N
1352E	Perma-Whitlash complex, 15 to 35 percent slopes, bouldery, warm	N	N	N	Y	Y	N	N
1353F	Perma, very stony-Whitlash, very stony-Rock outcrop complex, 15 to 45 percent slopes	N	N	N	Y	Y	N	N
1355D	Wimper-Wimper, stony, complex, 4 to 15 percent slopes, warm	N	N	N	Y	Y	N	N
1355E	Wimper-Wimper, stony, complex, 15 to 35 percent slopes, warm	N	N	N	Y	Y	N	Y
1356E	Wimper gravelly loam, 8 to 35 percent slopes, stony, moist	N	N	N	Y	Y	N	N
1357F	Perma, very bouldery-Shaboom, extremely bouldery-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	N	N	N
1362F	Lumpgulch, bouldery-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1372D	Burtoner-Connieo, bouldery-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	Y	Y
1373E	Burtoner-Elmark-Connieo complex, 8 to 25 percent slopes, very bouldery	Y	Y	Y	Y	Y	N	N
1374D	Burtoner-Clancy-Connieo complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1375D	Burtoner-Connieo, bouldery-Rock outcrop complex, 4 to 15 percent slopes, warm	Y	Y	Y	Y	Y	N	N
1375E	Burtoner, very stony-Connieo, bouldery-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1376F	Burtoner, very stony-Connieo, very stony-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1377E	Burtoner, very stony-Crampton, bouldery-Catgulch, bouldery complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1378E	Burtoner-Elmark-Shaboom, very bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1451F	Blaincreek, very stony-Sawicki, very stony-Tolbert, very bouldery, complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
1460C	Clasoil loam, 2 to 8 percent slopes	N	N	N	Y	Y	N	N
1461D	Bielenberg-Burtoner, very stony-Catgulch, bouldery, complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
1540F	Shaboom, extremely bouldery-Rock outcrop-Elmark, very bouldery, association, 35 to 60 percent slopes	N	N	N	N	N	Y	Y
1541E	Shaboom, bouldery-Lumpgulch, very bouldery-Rock outcrop complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1542E	Shaboom, very bouldery-Rock outcrop-Kellygulch, very bouldery, complex, 8 to 35 percent slopes	N	N	N	N	N	Y	Y
1543E	Shaboom, very bouldery-Kellygulch, very bouldery-Rock outcrop complex, 15 to 35 percent slopes	N	N	N	N	N	Y	Y
1543F	Shaboom, extremely bouldery-Kellygulch, extremely bouldery-Rock outcrop complex, 35 to 60 percent slope	N	N	N	N	N	Y	Y
1544E	Shaboom, bouldery-Kellygulch, bouldery-Rock outcrop complex, 8 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1563D	Hilger, rubbly-Hilger complex, 8 to 25 percent slopes	N	N	N	N	Y	N	N
1564E	Hilger, very stony-Hilger, rubbly-Rock outcrop complex, 8 to 35 percent slopes	N	N	N	N	Y	N	N
1591E	Catgulch, bouldery-Crackerville-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1595E	Connieo, bouldery-Crackerville-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1596C	Connieo-Rock outcrop-Placerton complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
1602C	Farnuf-Placerton sandy clay loams, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
1603C	Farnuf sandy loam, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
1604D	Farnuf-Farnuf, stony-Burtoner complex, 4 to 15 percent slopes	N	N	N	Y	Y	N	N
1605C	Farnuf-Placerton sandy clay loams, 2 to 8 percent slopes, warm	N	Y	Y	Y	Y	N	N
1606D	Farnuf loam, 2 to 15 percent slopes	N	N	N	Y	Y	N	N
1607D	Farnuf-Placerton-Martinsdale complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1621D	Connieo, stony-Baxton, stony-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1621E	Connieo, very stony-Baxton, stony-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1622D	Connieo, moist-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1623D	Connieo-Burtoner complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1623E	Connieo-Baxton-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1624F	Connieo, very stony-Baxton, bouldery-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1625F	Connieo, extremely bouldery-Rock outcrop-Burtoner, extremely stony, complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1626D	Connieo, bouldery-Burtoner, bouldery-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1627E	Connieo, very bouldery-Burtoner-Rock outcrop complex, 8 to 35 percent slopes, moist	Y	Y	Y	Y	Y	Y	Y
1628D	Connieo, bouldery-Ashbray, very bouldery-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1629C	Connieo-Catgulch-Rock outcrop complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
1640D	Nieman, very stony-Rock outcrop-Libeg, stony, complex, 2 to 15 percent slopes	N	N	N	N	N	N	N
1641E	Nieman, very stony-Rock outcrop-Libeg, bouldery, complex, 15 to 45 percent slopes	N	N	N	N	N	Y	Y
1641F	Nieman, very stony-Rock outcrop-Libeg, very stony, complex, 45 to 70 percent slopes	N	N	N	N	N	Y	Y
1642F	Nieman, bouldery-Rock outcrop-Libeg, very bouldery, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1643E	Nieman, stony-Libeg complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1643F	Nieman, stony-Libeg-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1651C	Sawbuck-Sawbuck, very stony-Clasoil complex, 2 to 8 percent slopes	N	N	N	Y	Y	N	N
1652E	Sawicki-Clasoil complex, 8 to 35 percent slopes, bouldery	N	N	N	Y	Y	N	N
1654E	Sawicki, stony-Blaincreek-Tolbert, very stony, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1655E	Sawicki-Clasoil complex, 8 to 35 percent slopes, bouldery, warm	N	N	N	Y	Y	N	N
1656E	Sawicki-Bielenberg, very stony-Tolbert, very stony, complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	Y	Y
1657E	Sawicki, very bouldery-Crampton, bouldery-Catgulch, bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	N	N
1658D	Sawicki, stony-Blaincreek, very stony, complex, 4 to 15 percent slopes	N	N	N	Y	Y	N	N
1658E	Sawicki, very stony-Blaincreek, very stony-Tolbert, bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1659E	Sawbuck, stony-Sawbuck, bouldery, complex, 15 to 35 percent slopes	N	N	N	N	Y	N	N
1661D	Catgulch-Baxton complex, 2 to 15 percent slopes, stony	Y	Y	Y	Y	Y	N	N
1661E	Catgulch-Baxton complex, 15 to 35 percent slopes, stony	Y	Y	Y	Y	Y	Y	Y
1662D	Catgulch, very stony-Rock outcrop-Burtoner complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1663D	Catgulch, bouldery-Burtoner, bouldery-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1664E	Catgulch, bouldery-Rock outcrop-Ashbray, bouldery, complex, 4 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1665F	Catgulch, very bouldery-Rock outcrop-Connieo, very stony, complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1667E	Catgulch, extremely bouldery-Baxton, extremely bouldery-Burtoner, bouldery, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1671E	Tolbert-Blaincreek complex, 8 to 35 percent slopes, warm	Y	Y	Y	Y	Y	Y	Y
1672E	Tolbert-Blaincreek complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1675E	Tolbert, very stony-Blaincreek, stony-Rock outcrop complex, 8 to 35 percent slopes	N	N	N	N	N	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1675F	Tolbert, very stony-Rock outcrop-Blaincreek, very stony, complex, 35 to 60 percent slopes	N	N	N	N	N	Y	Y
1680D	Raynesford silt loam, 4 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1690F	Cheadle, very stony-Rock outcrop-Tiban, bouldery, complex, 15 to 45 percent slopes	N	N	N	N	N	Y	Y
1721C	Martinsdale loam, 2 to 8 percent slopes, warm	N	N	N	N	Y	N	N
1722C	Martinsdale-Martinsdale, stony-Shawmut complex, 2 to 8 percent slopes, warm	N	N	N	Y	Y	N	N
1722E	Martinsdale-Martinsdale, stony-Shawmut complex, 15 to 35 percent slopes, warm	N	N	N	Y	Y	N	N
1723D	Martinsdale-Shawmut complex, 2 to 15 percent slopes, bouldery, warm	N	N	N	Y	Y	N	N
1724D	Martinsdale-Shawmut, stony-Martinsdale, bouldery, complex, 4 to 25 percent slopes, warm	N	N	N	N	Y	N	N
1731E	Tepecreek, bouldery-Caseypeak, very bouldery-Rock outcrop complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1731F	Tepecreek, very bouldery-Caseypeak, rubbly-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	N	Y	Y
1732F	Tepecreek, very bouldery-Caseypeak, very bouldery-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	N	N	Y
1734F	Hiore, stony-Kurrie, stony-Caseypeak, very stony, complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1735E	Tepecreek, stony-Caseypeak, very bouldery-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
1735F	Tepecreek, stony-Caseypeak, very stony-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	N	Y
1740E	Tropal, bouldery-Hanson, stony-Rock outcrop complex, 8 to 25 percent slopes	Y	Y	Y	Y	Y	N	Y
1741F	Tropal, bouldery-Rock outcrop-Whitore, bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1742F	Tropal, very bouldery-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1750F	Whitore, bouldery-Tropal, very bouldery-Rock outcrop complex, 25 to 45 percent slopes	Y	Y	Y	Y	Y	N	Y
1751F	Whitore, very stony-Tropal, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1752E	Whitore, stony-Helmville, bouldery-Firada, very stony, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1753E	Whitore, stony-Tropal, very stony-Firada, very stony, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1760E	Hanson, stony-Whitore, bouldery, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
1770E	Helmville, rubbly-Tiban, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	N	N	N	N	N	Y	Y
1781E	Firada, stony-Tropal, very stony-Rock outcrop complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	N	N
1790F	Sigbird, very bouldery-Sigbird, stony-Rock outcrop complex, 25 to 70 percent slopes	N	N	N	N	N	Y	Y
1800D	Breeton coarse sandy loam, 4 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1802D	Breeton-Baxton-Connieo complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1803C	Breeton-Cometcrik complex, 2 to 8 percent slopes	N	Y	Y	Y	Y	N	N
1810F	Hoyt, very stony-Ymark, bouldery-Shaboom, very bouldery, complex, 25 to 60 percent slopes	N	N	N	Y	Y	Y	Y
1821F	Kellygulch, bouldery-Rock outcrop-Bielenberg complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
1822F	Kellygulch, stony-Shaboom, very bouldery-Rock outcrop association, 45 to 75 percent slopes	Y	Y	Y	Y	Y	Y	Y
1823E	Kellygulch, stony-Shaboom, very bouldery-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1823F	Kellygulch, stony-Shaboom, very bouldery-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1830E	Clancy-Bielenberg-Breeton complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	Y
1831D	Clancy-Burtoner, bouldery, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1832D	Clancy-Burtoner, bouldery-Rock outcrop complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1833D	Clancy-Connieo complex, 2 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1835D	Clancy-Bielenberg-Connieo complex, 4 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1836E	Clancy, bouldery-Bielenberg, stony-Catgulch, bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1837E	Clancy-Bielenberg-Connieo complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y
1838D	Clancy-Clancy, very stony-Bielenberg complex, 4 to 15 percent slopes	N	Y	Y	Y	Y	N	N
1841D	Tuggle-Branham-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1842D	Caseypeak-Branham-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1842E	Caseypeak-Branham-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1842F	Caseypeak, bouldery-Branham, bouldery-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
1851D	Branham-Lowder loams, 0 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
1851E	Branham-Lowder loams, 8 to 25 percent slopes	Y	Y	Y	Y	Y	Y	Y
1853D	Branham-Tuggle complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1853E	Branham-Tuggle complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1861F	Clugulch-Bobowic-Rock outcrop complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
1871E	Hiore, stony-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
1871F	Hiore, stony-Rock outcrop complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
1872E	Hiore-Clugulch-Rock outcrop complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
1872F	Hiore-Clugulch-Rock outcrop complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
1901F	Warwood-Tigeron, very stony-Cowood, very stony, complex, 25 to 60 percent slopes	N	N	N	Y	Y	N	N
1902D	Warwood, very bouldery-Warwood, very stony-Tigeron, very bouldery, complex, 2 to 15 percent slopes	N	N	N	N	Y	N	N
1910F	Elmark, very bouldery-Rock outcrop-Shaboom, extremely bouldery, complex, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
1921D	Judell-Lap, very stony, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1921E	Judell-Lap, very stony, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1930E	Elmark-Kellygulch, very bouldery-Rock outcrop complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y
1933E	Elmark, bouldery-Breeton-Shaboom, bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1940E	Elmark, bouldery-Lumpgulch, very bouldery-Rock outcrop complex, 8 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y
1945E	Elmark, bouldery-Lumpgulch, very bouldery-Rock outcrop complex, 8 to 35 percent slopes, dry	N	Y	Y	Y	Y	Y	Y
1946E	Elmark, bouldery-Hoyt-Shaboom, very bouldery, complex, 8 to 35 percent slopes, dry	N	Y	Y	Y	Y	Y	Y
1947E	Elmark, bouldery-Burtoner-Rock outcrop complex, 8 to 45 percent slopes	Y	Y	Y	Y	Y	N	N
1948E	Elmark, very bouldery-Skyview, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	N	N	N	N	N	N	N
1950F	Franconi, very bouldery-Warwood-Caseypeak, very bouldery, complex, 25 to 60 percent slopes	N	N	N	Y	Y	Y	Y
1960D	Lumpgulch, bouldery-Hoyt-Shaboom, very bouldery, complex, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
1961E	Lumpgulch, bouldery-Hoyt-Shaboom, very bouldery, complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1962E	Lumpgulch, bouldery-Yreka, very bouldery-Shaboom, very bouldery, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
1963F	Lumpgulch, very bouldery-Rock outcrop-Kellygulch, very bouldery, complex, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
1964E	Lumpgulch, very bouldery-Shaboom, very bouldery-Rock outcrop complex, 8 to 25 percent slopes	N	N	N	N	N	N	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
1965E	Lumpgulch, bouldery-Ymark, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
1970E	Bignell, stony-Yreka, very stony, complex, 15 to 35 percent slopes	N	N	N	N	Y	N	N
1980F	Stemple cobbly loam, 35 to 60 percent slopes, very stony	N	N	N	N	Y	Y	Y
1990F	Bobowic, very bouldery-Rock outcrop-Tepecreek, very bouldery, complex, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
1991D	Bobowic-Clugulch, bouldery-Rock outcrop complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	N	N
2000E	Skyview, very bouldery-Rock outcrop-Roegulch, very bouldery, complex, 8 to 35 percent slopes	N	N	N	N	N	N	Y
2001E	Skyview, very bouldery-Elmark, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	N	N	N	N	N	N	N
2011D	Shawmut gravelly loam, 4 to 15 percent slopes, bouldery, warm	N	N	N	N	Y	N	N
2012D	Shawmut, stony-Martinsdale, very stony, complex, 4 to 15 percent slopes, warm	N	N	N	N	Y	N	N
2012E	Shawmut, stony-Martinsdale, very stony, complex, 15 to 25 percent slopes, warm	N	N	N	N	Y	N	N
2013E	Shawmut, bouldery-Wickes, stony-Tolbert, bouldery, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
2014E	Shawmut-Tolbert complex, 8 to 35 percent slopes, warm	N	N	N	N	Y	N	N
2020D	Shawmut, stony-Shawmut, bouldery, complex, 4 to 15 percent slopes	N	N	N	N	Y	N	N
2030E	Kokoruda-Elmark, very bouldery-Rock outcrop complex, 8 to 35 percent slopes	N	N	N	N	Y	N	N
2031D	Eagleton, stony-Kokoruda-Cometcrik complex, 2 to 25 percent slopes	N	N	Y	Y	Y	N	N
2040F	Shaboom, extremely bouldery-Rock outcrop-Rubble land association, 35 to 70 percent slopes	N	N	N	N	N	Y	Y
2041F	Rock outcrop-Catgulch, bouldery, complex, 15 to 70 percent slopes	0	0	0	0	0	0	0
2042F	Rock outcrop-Cheadle, very bouldery-Tiban, very bouldery, complex, 15 to 50 percent slopes	N	N	N	N	N	N	N
2043F	Rencot, very stony-Rencot, bouldery-Rock outcrop association, 15 to 60 percent slopes	N	N	N	N	N	Y	Y
2045F	Caseypeak, very stony-Rock outcrop-Rubble land association, 15 to 60 percent slopes, dry	Y	Y	Y	Y	Y	Y	Y
2046F	Caseypeak, very bouldery-Rock outcrop-Rubble land association, 15 to 60 percent slopes, cool	Y	Y	Y	Y	Y	Y	Y
2051E	Opitz, bouldery-Branham, very bouldery-Tuggle, very bouldery, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
2081F	Windham, very stony-Rock outcrop-Lap, very stony, complex, 35 to 70 percent slopes, warm	Y	Y	Y	Y	Y	Y	Y
2082D	Windham-Judell complex, 8 to 15 percent slopes, warm	Y	Y	Y	Y	Y	N	Y
2084E	Windham, stony-Maiden, very stony-Lap, very stony, complex, 15 to 35 percent slopes, warm	Y	Y	Y	Y	Y	Y	Y
2086E	Windham-Windham, stony, complex, 15 to 35 percent slopes, warm	Y	Y	Y	Y	Y	Y	Y
2088D	Windham-Judell gravelly loams, 8 to 25 percent slopes, stony, warm	Y	Y	Y	Y	Y	N	Y
2090F	Caseypeak, very bouldery-Franconi, very bouldery-Rock outcrop complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
2091E	Caseypeak, bouldery-Franconi, bouldery-Rock outcrop complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
2110D	Sebud very cobbly loam, 4 to 15 percent slopes, very stony	N	N	N	N	Y	N	N
2111E	Sebud, very stony-Hapgood complex, 8 to 45 percent slopes	N	Y	Y	Y	Y	Y	Y
2112D	Sebud-Marcel complex, 4 to 25 percent slopes, bouldery	N	N	N	N	Y	N	N
2121F	Hapgood-Hanson-Tiban complex, 25 to 60 percent slopes, very stony	N	N	N	N	N	N	N
2122F	Hapgood-Tiban complex, 35 to 70 percent slopes, very stony	N	N	N	N	Y	Y	Y
2123F	Hapgood-Sebud-Arrowpeak complex, 35 to 60 percent slopes, very stony	N	N	N	N	Y	N	N
2125F	Coweed, rubbly-Elve, very stony-Rock outcrop complex, 25 to 60 percent slopes	N	N	N	N	N	Y	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
2151E	Releep, very bouldery-Kurrie, very bouldery-Rock outcrop complex, 15 to 35 percent slopes	N	N	N	N	N	N	N
2161E	Ellena, bouldery-Worock, very bouldery-Rock outcrop complex, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
2161F	Kurrie, very bouldery-Ellena, very bouldery-Rock outcrop complex, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
2171F	Hiore-Kurrie, stony, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
2172F	Rubick, very stony-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	N	N	N
2173F	Rubick gravelly sandy loam, 35 to 60 percent slopes, stony	Y	Y	Y	Y	Y	Y	Y
2181F	Repkie, very stony-Yreka, stony-Skyview, very bouldery, complex, 25 to 60 percent slopes	N	Y	Y	Y	Y	N	Y
2211E	Sebud-Arrowpeak, stony, complex, 8 to 45 percent slopes	N	N	N	N	Y	N	N
2211F	Sebud, very stony-Arrowpeak, very stony-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	N	Y	N	N
2212D	Sebud, very stony-Libeg-Arrowpeak, stony, complex, 4 to 15 percent slopes	N	N	N	N	Y	N	N
2212E	Sebud, very stony-Libeg, stony-Arrowpeak, stony, complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	Y	Y
2213E	Sebud, stony-Surdal, stony-Arrowpeak, very stony, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
2213F	Sebud, stony-Surdal, stony-Arrowpeak, very stony, complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
2214E	Sebud, bouldery-Surdal, very bouldery-Arrowpeak, very bouldery, complex, 15 to 35 percent slopes	N	N	N	N	Y	N	N
2215D	Sebud-Tibkey cobbly loams, 2 to 15 percent slopes, bouldery	N	N	N	N	Y	N	N
2216D	Sebud-Surdal complex, 4 to 25 percent slopes, stony	Y	Y	Y	Y	Y	N	N
2221E	Arrowpeak, very stony-Surdal, stony-Rock outcrop complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
2222F	Arrowpeak, very stony-Sebud, stony-Surdal, very stony, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
2230B	Tineman cobbly loam, 2 to 8 percent slopes, very stony	N	Y	Y	Y	Y	N	N
2251F	Nivean, very stony-Rock outcrop-Rubble land complex, 25 to 60 percent slopes	N	N	N	N	N	Y	Y
2252E	Nivean, very stony-Macabre, stony-Rock outcrop complex, 15 to 35 percent slopes	N	N	N	N	N	Y	Y
2261D	Lowland loam, 4 to 15 percent slopes, stony	N	Y	Y	Y	Y	N	N
2261E	Lowland loam, 15 to 35 percent slopes, stony	N	Y	Y	Y	Y	Y	Y
2261F	Lowland, stony-Rock outcrop-Rubble land complex, 35 to 60 percent slopes	N	Y	Y	Y	Y	Y	Y
2270F	Macabre, very stony-Rock outcrop-Rubble land complex, 35 to 60 percent slopes	N	Y	Y	Y	Y	N	Y
2271D	Macabre gravelly loam, 8 to 15 percent slopes	N	Y	Y	Y	Y	N	N
2281F	Judco, stony-Torpy, stony-Rock outcrop complex, 35 to 60 percent slopes	N	N	Y	Y	Y	N	N
2291F	Mocmont-Kadygulch cobbly loams, 35 to 60 percent slopes, very stony	N	N	N	N	Y	Y	Y
2301F	Mocmont, bouldery-Roegulch, rubbly-Rock outcrop complex, 25 to 60 percent slopes	N	N	N	N	Y	Y	Y
2311F	Worock, stony-Cowood, very stony-Rock outcrop complex, 35 to 60 percent slopes	N	Y	Y	Y	Y	N	N
2312F	Worock, stony-Elve, stony-Rock outcrop complex, 35 to 60 percent slopes	N	Y	Y	Y	Y	N	N
2321D	Torpy loam, 4 to 15 percent slopes	N	N	N	Y	Y	N	N
2321E	Torpy gravelly loam, 15 to 35 percent slopes	N	N	N	Y	Y	N	N
2321F	Torpy gravelly loam, 35 to 60 percent slopes	N	N	N	Y	Y	Y	Y
2322E	Lowland-Torpy complex, 15 to 35 percent slopes	N	N	N	Y	Y	N	Y
2322F	Lowland-Torpy complex, 35 to 60 percent slopes	N	N	N	Y	Y	N	Y

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
2331B	Mooseflat loam, 1 to 4 percent slopes	N	Y	Y	Y	Y	N	N
2332B	Mooseflat-Elvick loams, 1 to 4 percent slopes	N	Y	Y	Y	Y	N	N
2350D	Clasoil, very stony-Sawicki, bouldery, complex, 4 to 15 percent slopes	N	Y	Y	Y	Y	N	N
2360E	Gnojek, stony-Wickes, stony-Shawmut complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
2360F	Gnojek, stony-Wickes, stony-Rock outcrop complex, 35 to 70 percent slopes	Y	Y	Y	Y	Y	Y	Y
2361E	Gnojek, stony-Wickes, stony-Rock outcrop complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
2361F	Gnojek, stony-Rock outcrop-Wickes, stony, complex, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
2391C	Marcel, very bouldery-Tibkey, bouldery, complex, 2 to 8 percent slopes	N	N	N	N	Y	N	N
2411E	Ashbray, bouldery-Rock outcrop-Rubble land complex, 8 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
2412F	Ashbray, rubbly-Rock outcrop-Kellygulch, very stony, complex, 35 to 70 percent slopes	N	N	N	N	N	Y	Y
2421E	Surdal, stony-Arrowpeak, very stony, complex, 4 to 25 percent slopes	Y	Y	Y	Y	Y	N	N
2431C	Foolhen, stony-Tibkey, bouldery, complex, 0 to 8 percent slopes	N	N	N	N	Y	N	N
2441E	Tineman, very stony-Franconi, bouldery-Rock outcrop complex, 4 to 25 percent slopes	N	Y	Y	Y	Y	N	N
2450E	Kounter, bouldery-Rock outcrop-Cedric, bouldery, complex, 8 to 35 percent slopes	Y	Y	Y	Y	Y	Y	Y
2451D	Kounter, bouldery-Rock outcrop-Cedric, bouldery, complex, 4 to 25 percent slopes, dry	Y	Y	Y	Y	Y	N	N
2452E	Kounter, very bouldery-Rock outcrop-Jeffcity, bouldery, complex, 15 to 35 percent slopes	Y	Y	Y	Y	Y	N	N
2460D	Cedric, bouldery-Jeffcity, bouldery-Rock outcrop complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
2461D	Cedric, bouldery-Rock outcrop-Jeffcity, bouldery, complex, 2 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
2471F	Elve, stony-Worock, stony-Rock outcrop complex, 35 to 60 percent slopes	N	N	N	Y	Y	Y	Y
2472E	Elvick-Lowder complex, 8 to 25 percent slopes, very bouldery	N	N	N	N	N	N	N
2473E	Elve-Cowood very cobbly loams, 8 to 35 percent slopes, very stony	N	N	N	N	N	N	N
2481F	Brickner, very bouldery-Rock outcrop-Rubble land complex, 15 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
2483F	Elve, rubbly-Rock outcrop-Rubble land association, 25 to 60 percent slopes, cool	N	N	N	N	N	N	N
2484F	Rock outcrop-Rubble land-Redfern, rubbly, association, 35 to 70 percent slopes	N	N	N	N	N	Y	Y
2485F	Redfern, rubbly-Rock outcrop-Tigeron, very bouldery, association, 25 to 70 percent slopes	N	N	N	N	N	Y	Y
2486F	Elve, rubbly-Rock outcrop-Rubble land complex, 35 to 60 percent slopes	N	N	N	N	N	N	Y
2487F	Torpy, rubbly-Rock outcrop-Rubble land complex, 35 to 60 percent slopes	N	N	N	N	N	N	Y
2488D	Elve, very stony-Rock outcrop-Rubble land complex, 4 to 35 percent slopes	N	N	N	N	N	N	N
2501D	Lowder-Elvick very cobbly loams, 2 to 15 percent slopes, very bouldery	N	N	N	N	N	N	N
2511C	Monaberg loam, 2 to 8 percent slopes, bouldery	N	N	N	N	Y	N	N
2581E	Worock, very bouldery-Elve, very stony, complex, 15 to 35 percent slopes	N	N	N	N	N	N	N
2582D	Worock-Elve very cobbly loams, 2 to 15 percent slopes, very bouldery, cool	N	N	N	N	N	N	N
2582E	Worock, very bouldery-Worock, rubbly, complex, 8 to 35 percent slopes	N	N	N	N	N	N	N
2582F	Worock, rubbly-Rock outcrop-Rubble land complex, 35 to 60 percent slopes	N	N	N	N	N	N	N
2583D	Worock, stony-Worock, very bouldery, complex, 2 to 15 percent slopes	N	N	N	N	Y	N	N
2583F	Worock, very bouldery-Worock, rubbly, complex, 35 to 60 percent slopes	N	N	N	N	N	N	N

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

Erodible Land Index
Survey Area: MT627-Jefferson Co. and part of Silverbow Co., Montana

0=DATA NOT AVAILABLE Y=HIGHLY ERODIBLE P=POTENTIALLY HIGHLY ERODIBLE N=NOT HIGHLY ERODIBLE

		Wind					Water	
		C - 40	C - 60	C - 70	C - 80	C - 100	R - 15	R - 20
mu_sym	muname	Hel	Hel	Hel	Hel	Hel	Hel	Hel
2584E	Worock, very bouldery-Worock, rubbly, complex, 15 to 45 percent slopes, dry	N	N	N	N	N	N	N
2591F	Kadygulch-Roegulch, stony, complex, 35 to 60 percent slopes	N	N	N	N	Y	Y	Y
2661F	Elve-Cowood complex, 45 to 70 percent slopes	N	Y	Y	Y	Y	N	N
2662E	Elve-Cowood complex, 15 to 45 percent slopes	N	Y	Y	Y	Y	N	N
2681E	Sawbuck-Catgulch, stony, complex, 8 to 45 percent slopes	N	Y	Y	Y	Y	N	N
2682E	Sawbuck, stony-Yreka, stony-Catgulch, very stony, complex, 15 to 45 percent slopes	N	N	N	Y	Y	N	N
2691F	Connieo, very stony-Crackerville, stony-Rock outcrop complex, 35 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
2695E	Macabre-Nivean complex, 15 to 35 percent slopes	N	Y	Y	Y	Y	N	N
2695F	Macabre-Judco-Rock outcrop complex, 35 to 60 percent slopes	N	Y	Y	Y	Y	N	Y
2701F	Crampton-Catgulch complex, 35 to 60 percent slopes, very stony	Y	Y	Y	Y	Y	Y	Y
2705F	Vitroff-Torpy loams, 35 to 60 percent slopes	N	N	Y	Y	Y	Y	Y
2711E	Libeg very gravelly loam, 15 to 45 percent slopes	N	N	N	N	Y	N	N
2712D	Libeg-Mooseflat loams, 4 to 25 percent slopes	N	Y	Y	Y	Y	N	N
3001B	Aridic Ustifluvents, 0 to 4 percent slopes	N	Y	Y	Y	Y	N	N
3033B	Sappington-Amesha loams, 1 to 4 percent slopes	Y	Y	Y	Y	Y	N	N
3033C	Sappington-Amesha loams, 4 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
3061E	Holter-Castner channery loams, 8 to 45 percent slopes	N	N	N	N	Y	N	N
3064D	Windham channery loam, 4 to 15 percent slopes	Y	Y	Y	Y	Y	N	N
3137B	Musselshell-Crago complex, 2 to 8 percent slopes	Y	Y	Y	Y	Y	N	N
3141E	Crago-Pensore channery loams, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
3218A	Meadowcreek-Fairway complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
3233C	Geohrock-Crago very cobbly loams, 2 to 8 percent slopes	N	N	N	N	Y	N	N
3433E	Crago-Musselshell gravelly loams, 4 to 35 percent slopes	Y	Y	Y	Y	Y	N	Y
3486F	Peeler-Rock outcrop complex, 15 to 60 percent slopes	N	N	N	N	N	N	N
3501B	Fluvaquents-Fluvaquentic Haplustolls complex, 0 to 4 percent slopes	N	Y	Y	Y	Y	N	N
3513A	Attewan-Nippt complex, 0 to 2 percent slopes	N	Y	Y	Y	Y	N	N
3532B	Geohrock gravelly loam, 2 to 8 percent slopes	N	N	N	N	Y	N	N
3664E	Windham-Whitecow-Lap channery loams, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
3685F	Whitecow channery loam, 25 to 60 percent slopes	Y	Y	Y	Y	Y	Y	Y
3885F	Whitecow-Warneke channery loams, 15 to 45 percent slopes	Y	Y	Y	Y	Y	Y	Y
DAM	Dam	0	0	0	0	0	0	0
M-W	Miscellaneous water	0	0	0	0	0	0	0
W	Water	0	0	0	0	0	0	0

*Data for these map units did not exist at the origination of this Highly Erodible Map Unit Listing. Data was developed as the survey progressed and was included upon correlation.

ERODIBLE LAND INDEX
SURVEY NUMBER 670 SURVEY NAME SILVER BOW COUNTY
0=DATA NOT AVAILABLE 1=HIGHLY ERODIBLE 2=POTENTIALLY HIGHLY ERODIBLE 3=NOT HIGHLY ERODIBLE

Superseded by HEL Map Updated Unit List dated 5/6/93 per TO Notice 193 dated 5/21/89 JC

MUID	MAP SYMBOL	SOIL NAME		%	WIND C 20 HEL CLASS	C 30 HEL CLASS	C 50 HEL CLASS	C 60 HEL CLASS	C 70 HEL CLASS	C 80 HEL CLASS	C 90 HEL CLASS	WATER R 15 HEL CLASS	R 20 HEL CLASS	I VALUE	K VALUE	T VALUE	SLOPE-PERCEN MIN.	SLOPE-PERCEN MAX.	SLOPE-LENGTH MIN.	SLOPE-LENGTH MAX.
670111E	111E	VARNEY VAR CBL	1	100	3	3	3	3	3	3	1	2	2	48	.28	5	15	35	25	400
670112C	112C	VARNEY VAR L	1	55	3	3	3	3	3	3	1	3	3	48	.37	5	2	8	25	400
670112C	112C	RADERSBURG VARGRCL	2	45	3	3	1	1	1	1	1	3	3	48	.24	3	2	8	25	400
670141E	141E	NULEY VAR CBVL	1	100	3	1	1	1	1	1	1	2	2	56	.24	2	15	35	25	400
670143C	143C	NULEY VAR STL	1	100	3	1	1	1	1	1	1	3	3	56	.24	2	2	8	25	400
67014A	14A	NULEY VAR LOAM	1	100	3	1	1	1	1	1	1	3	3	56	.28	2	0	2	25	400
67014C	14C	NULEY VAR LOAM	1	100	3	1	1	1	1	1	1	3	3	56	.28	2	2	8	25	400
67014D	14D	NULEY VAR LOAM	1	100	3	1	1	1	1	1	1	2	2	56	.28	2	8	15	25	400
67018A	18A	COMER FAM GRCOSL	1	100	3	1	1	1	1	1	1	3	3	86	.17	3	0	2	25	400
67020B	20B	ANACONDA FAM L	1	50	3	3	3	3	3	3	1	3	3	48	.37	5	1	4	25	400
67020B	20B	COMER FAM GRCOSL	2	50	3	1	1	1	1	1	1	3	3	86	.17	3	1	4	25	400
67024A	24A	BEAVERELL CB L	1	100	3	1	1	1	1	1	1	3	3	56	.20	2	0	2	25	400
67024D	24D	BEAVERELL CB L	1	100	3	1	1	1	1	1	1	3	2	56	.20	2	8	15	25	400
67026A	26A	GLENDIVE VAR SL	1	100	3	3	1	1	1	1	1	3	3	86	.20	5	0	2	25	400
67027A	27A	ATTEWAN VAR L	1	100	3	3	1	1	1	1	1	3	3	56	.37	3	0	2	25	400
67028A	28A	FUBARTOO SIL	1	100	1	1	1	1	1	1	1	3	3	86	.32	2	0	1	25	400
670291E	291E	DINNEN COSL	1	50	3	1	1	1	1	1	1	2	2	86	.24	3	4	35	25	400
670291E	291E	BRANHAM GRCOSL	2	35	1	1	1	1	1	1	1	2	2	86	.15	2	4	35	25	400
670291E	291E	ROCK OUTCROP	3	15	0	0	0	0	0	0	0	0	0	38	.00	1	4	35	25	400
67029C	29C	DINNEN COSL	1	100	3	1	1	1	1	1	1	3	3	86	.24	3	2	8	25	400
67029D	29D	DINNEN COSL	1	100	3	1	1	1	1	1	1	3	2	86	.24	3	8	15	25	400
670301D	301D	HAUZ GRCOSCL	1	50	3	1	1	1	1	1	1	3	2	56	.20	2	4	15	25	400
670301D	301D	TOLMAN GRCOSCL	2	35	1	1	1	1	1	1	1	2	2	56	.20	1	4	15	25	400
670301D	301D	ROCK OUTCROP	3	15	0	0	0	0	0	0	0	0	0	38	.00	1	4	15	25	400
670301E	301E	HAUZ GRCOSCL	1	45	3	1	1	1	1	1	1	2	2	56	.20	2	15	35	25	400
670301E	301E	TOLMAN GRCOSCL	2	30	1	1	1	1	1	1	1	2	2	56	.20	1	15	35	25	400
670301E	301E	ROCK OUTCROP	3	25	0	0	0	0	0	0	0	0	0	38	.00	1	4	15	25	400
670301E	301E	HAUZ GRCOSCL	1	45	3	1	1	1	1	1	1	2	2	56	.20	2	15	35	25	400
670301E	301E	TOLMAN GRCOSCL	2	30	1	1	1	1	1	1	1	2	2	56	.20	1	15	35	25	400
670301E	301E	ROCK OUTCROP	3	25	0	0	0	0	0	0	0	0	0	38	.00	1	4	15	25	400
67032E	32E	BROCKO FAM CBSIL	1	100	3	3	3	3	3	1	1	2	2	56	.24	5	15	25	25	400
670371E	371E	BROWNSTO CBL	1	100	3	3	1	1	1	1	1	2	2	86	.20	4	15	35	25	400
67037C	37C	BROWNSTO GRL	1	100	3	3	1	1	1	1	1	3	3	86	.20	4	2	8	25	400
67047E	47E	SUPERVISOR FAM GRL	1	50	3	1	1	1	1	1	1	2	2	56	.24	2	8	25	25	400
67047E	47E	MERINO VCBL	2	30	1	1	1	1	1	1	1	2	2	56	.15	1	8	25	25	400
67047E	47E	ROCK OUTCROP	3	20	0	0	0	0	0	0	0	0	0	38	.00	1	8	25	25	400
67049C	49C	LIBEG GRL	1	100	3	3	1	1	1	1	1	3	3	48	.15	3	2	8	25	400
670501D	501D	BUFFORK L	1	60	3	3	1	1	1	1	1	2	2	48	.32	3	4	15	25	400
670501D	501D	SAWCREEK FAM L	2	40	3	3	1	1	1	1	1	2	2	56	.32	3	4	15	25	400
67050A	50A	BUFFORK L	1	100	3	3	1	1	1	1	1	3	3	48	.32	3	0	2	25	400
67050C	50C	BUFFORK L	1	100	3	3	1	1	1	1	1	3	3	48	.32	3	2	8	25	400

SURVEY NAME SILVER BOW COUNTY

MUID	MAP SYMBOL	SOIL NAME	%	WIND C 20 HEL CLASS	C 30 HEL CLASS	C 50 HEL CLASS	C 60 HEL CLASS	C 70 HEL CLASS	C 80 HEL CLASS	C 90 HEL CLASS	WATER R 15 HEL CLASS	R 20 HEL CLASS	I VALUE	K VALUE	T VALUE	SLOPE-PERCEN MIN.	MAX.	SLOPE-LENGTH MIN.	MAX.
670521C	521C	WHITECAP FAMGRCOSL 1	50*	1	1	1	1	1	1	1	3	3	86	.10	1	2	8	25	400
670521C	521C	SAWCREEK FAM L 2	50*	3	3	1	1	1	1	1	3	3	56	.32	3	2	8	25	400
670521C	521C	WHITECAP FAMGRCOSL 1	50	1	1	1	1	1	1	1	3	3	86	.10	1	2	8	25	400
670521C	521C	SAWCREEK FAM L 2	50	3	3	1	1	1	1	1	3	3	56	.32	3	2	8	25	400
67054C	54C	LTL.HORN FAM CBSIL 1	100	3	3	1	1	1	1	1	3	3	48	.24	2	2	8	25	400
67054D	54D	LTL.HORN FAM CBSIL 1	100	3	3	1	1	1	1	1	2	2	48	.24	2	8	15	25	400

*These are undifferentiated or highly variable map units. Percent composition of the soil component in this unit will vary from field to field. The determination of the actual composition requries on-site investigation.